

UNCLASSIFIED

AD 297 213

*Reproduced
by the*

ARMED SERVICES TECHNICAL INFORMATION AGENCY
ARLINGTON HALL STATION
ARLINGTON 12, VIRGINIA



UNCLASSIFIED

NOTICE: When government or other drawings, specifications or other data are used for any purpose other than in connection with a definitely related government procurement operation, the U. S. Government thereby incurs no responsibility, nor any obligation whatsoever; and the fact that the Government may have formulated, furnished, or in any way supplied the said drawings, specifications, or other data is not to be regarded by implication or otherwise as in any manner licensing the holder or any other person or corporation, or conveying any rights or permission to manufacture, use or sell any patented invention that may in any way be related thereto.

CATALOGUED BY ASTIA
AS AD NO. 297213

297 213



Research Translation

T-R-392+

A New Radiation Chart

HELGI NILISK

RESEARCH LIBRARY

Requests for additional copies by Agencies of the Department of Defense, their contractors, and other government agencies should be directed to the:

Armed Services Technical Information Agency
Arlington Hall Station
Arlington 12, Virginia

Department of Defense contractors must be established for ASTIA services, or have their 'need-to-know' certified by the cognizant military agency of their project or contract.

All other persons and organizations should apply to the:

U. S. DEPARTMENT OF COMMERCE
OFFICE OF TECHNICAL SERVICES,
WASHINGTON 25, D. C.

AMERICAN METEOROLOGICAL SOCIETY
45 BEACON STREET
BOSTON 8, MASSACHUSETTS

TRANSLATION OF

A NEW RADIATION CHART

(Novaia radiatsionnaia nomogramma)

by

Helgi Niilisk

Akademiia Nauk Estonskoi SSR. Izvestiia, Seriiia
Fiziko-Matematicheskikh i Tekhnicheskikh
Nauk, 1961

This translation has been made by the
American Meteorological Society under
Contract AF 19(604)-6113, through the
support and sponsorship of the

AIR FORCE CAMBRIDGE RESEARCH LABORATORIES

L. G. HANSCOM FIELD

BEDFORD, MASSACHUSETTS

1. "Novaia radiatsionnaia nomogramma"
2. "A new radiation chart"
3. Niilisk, Helgi. Akademiia Nauk Estonskoi SSR. Izvestiia, Seriia Fiziko-Matematicheskikh i Tekhnicheskikh Nauk, No. 4: 329-339, 1961.
4. 19 typewritten pages
5. Date of translation: August 1962
6. Translator: Myron Ricci
7. Translated for Air Force Cambridge Research Laboratories, Office of Aerospace Research, United States Air Force, L. G. Hanscom Field, Bedford, Massachusetts, by the American Meteorological Society, Contract number AF 19(604)-6113.
8. Unclassified
9. Complete

A NEW RADIATION CHART

by

Helgi Niilisk

At present, graphic methods of computing the fluxes of the thermal radiation of the atmosphere with the aid of radiation charts are well known. The charts of F. Brooks [1], A. A. Dmitriev [2], R. Mugge and F. Moller [3, 4], G. Robinson [5, 6], F. N. Shekhter [7, 8], W. Elsasser [9], and G. Yamamoto [10] have been the most widely used. Unfortunately, all these charts contain certain deficiencies [11-14], viz., the use of insufficiently reliable characteristics of the absorption of long-wave radiation by atmospheric gases and the analysis of the integral function dependent only on the effective content of water vapor. The effect of carbon dioxide on absorption, moreover, is considered to be very approximate.¹⁾ The charts mentioned do not at all consider the absorption of thermal radiation by atmospheric ozone.

It has been established [13, 14] that one of the most important factors which determines the value of radiation fluxes, calculated from any chart, is the transmission function. Therefore, to make radiation charts more accurate, it is first of all necessary to study the available quantitative characteristics of long-wave radiation in the atmosphere, and on that basis to obtain the most reliable transmission function.

Proceeding from what has been said above, the purpose of our work is to construct the integral transmission function of the atmosphere by considering the mutual effect of water vapor, carbon dioxide, and ozone²⁾ on the absorption of thermal radiation in the atmosphere. On the basis of this new transmission function, it is possible to solve the problem of constructing a radiation chart which is sufficiently accurate and convenient from a practical point of view. This work completes a number of studies made by the author [14-18] which were devoted to the problem of calculating the fluxes of thermal radiation in the atmosphere.

Quantitative data on the absorption of long-wave radiation, used in the present work to determine the integral transmission function, are, for the most part, given for 300°K, and also for the usual room temperatures of the order of 285-295°K. Since at surface temperatures the integral transmission function is slightly dependent on temperature [10, 13], we can consider the transmission function obtained on the basis of the mentioned data to be

-
- 1) Consideration of the mutual influence of H₂O and CO₂ on the absorption of thermal radiation in the atmosphere is correct in principle only in G. Yamamoto's chart.
 - 2) As is known [13], the remaining atmospheric gases play an insignificant role in the absorption of long-wave radiation by the atmosphere.

sufficiently reliable for the temperature interval 270-310°K. It should be noted that the integral transmission function was not calculated for the entire spectrum (0-∞ μ), and for the 2.27-250 μ region (in parts of the spectrum, 0-2.27 and 250-∞ μ, the amount of energy of thermal radiation is negligible and, consequently, practically has no effect on the value of the transmission function at atmospheric temperatures).

The determination of the quantitative characteristics of radiation absorption by water vapor is the subject of numerous works [1, 12, 13, 20-42], the results of which were obtained either by theoretical calculations or on the basis of an experiment (under natural or laboratory conditions). Let us note that there are important discrepancies in the data of various authors for almost all the regions of the spectrum. The results of the most recent and complete works are used when possible to determine the integral transmission function in the present work.

The values of the absorption of radiation by water vapor in the near infrared region (spectral intervals: 2.27-2.99, 2.99-3.57 and 4.88-8.7 μ) are calculated on the basis of the corresponding formulas of D. Howard, D. Burch, and D. Williams [20]. There are no experimental data for the 3.57-4.88 μ region. It is only known that in this region of the spectrum, there are no bands of absorption by water vapor and the absorption is caused by the overlapping of this interval by the wings of the absorption bands situated in the neighboring regions of the spectrum. We can assume that the logarithmic coefficient of absorption k_w here does not exceed 0.5 cm²/g. Calculations have shown that a change in the absorption coefficient in the limits 0-0.5 cm²/g has almost no effect on the value of the transmission function 2.27-8.7 μ, and on the value of the integral transmission function. Using this as a basis, in the present work the transmission function P for the interval 3.57-4.88 μ was calculated from the formula

$$P = e^{-k_w w^*}, \quad (1)$$

using a certain average value $k_w = 0.2 \text{ cm}^2/\text{g}$. (here w^* - the effective content of water vapor).

To determine the transmission function in the 8.7-12 μ region of the spectrum, there are no satisfactory reasons to prefer the results of one or other author (works [12, 13, 21, 22, 26, 28, 35, 37-40]). Therefore, in the present work the transmission function for the mentioned interval of the spectrum was calculated from formula (1), using certain average values for the coefficients of absorption by water vapor (Table 1).

TABLE 1

The Absorption Coefficients of Water Vapor for the
8.7-12 μ Region of the Spectrum

Spectrum Region	$k_w, \text{ cm}^2/\text{g}$
8.7 - 9.0	0.15
9.0 - 11.5	0.10
11.5 - 12.0	0.20

Let us note that according to the most recent data [37-39], k_w about 9.6 and 11.1 μ , is approximately 0.1.

In the 9.0-10.3 μ interval, it is necessary to consider the effect of a strong absorption band of ozone. We previously found the transmission function for the 9.0-10.3 μ region [16].

In the 12-18 μ region, the gases which absorb the thermal radiation of the atmosphere are water vapor and carbon dioxide, since a strong absorption band of carbon dioxide¹⁾ is found in this interval. The transmission function for 12-18 μ region obtained in works [15, 18] on the basis of data [19] and [22] is used in the present work.

1) According to [13] we can disregard the effect of the remaining bands of absorption by carbon dioxide on the transfer of long-wave radiation in the atmosphere.

Data are given in works [13, 21, 22, 27-30, 36, 41, 42] on the absorption of long-wave radiation by water vapor in the far infrared region of the spectrum. A comparison of the results of these works shows that the results of the theoretical calculations of G. Yamamoto [22] coincide most closely with the most recent data of the laboratory measurements of C. Palmer [36]. Proceeding from this, the transmission function for the 18-250 μ interval in the present work is determined on the basis of the usage of generalized coefficients of absorption by water vapor according to the data of work [22].

Let us note that for all the examined intervals of the spectrum, the transmission function for diffuse radiation was determined from the following known relationship [13]:

$$P_T(\omega^*) = 2 \int_0^{\pi/2} P(\omega^* \sec \theta) \sin \theta \cos \theta d\theta, \quad (2)$$

where P_T is the transmission function for diffuse radiation, P is the transmission function for the directed radiation, θ is the zenith angle.

By using the results of the determination of the transmission functions for various parts of the spectrum, the integral transmission function for diffuse radiation was calculated from the following formula:

$$P_T = \frac{1}{f} \sum_{\Delta} f_{\Delta} P_{T\Delta}. \quad (3)$$

Here the index Δ designates the examined intervals of the spectrum, f is a fraction of the integral radiation of an absolutely black body arriving on the 2.27-250 μ part of the spectrum, f_{Δ} is a fraction of the radiation of an absolutely black body arriving at the interval of the spectrum Δ .

Tables 2 and 3 give the calculation results. The tables also give the values of the functions $\Delta P_1(\omega^*, u^*)$ and $\Delta P_2(\omega^*, m^*)$ with which the sought integral transmission function can be calculated from the relationship:

$$P_T(\omega^*, u^*, m^*) = 0,001 (\Delta P_1 + \Delta P_2), \quad (4)$$

where u^* is the effective content of carbon dioxide, m^* is the effective content of ozone.¹

Since during the compilation of Tables 2 and 3 approximate methods were also used (numerical integration, interpolation), we can consider the transmission function obtained reliable to an accuracy of 1/100.

Having the data of an aerological sounding, it is easy to determine the values of the fluxes of thermal radiation in the atmosphere by using Tables 2 and 3 and the usual graphic method based on the use of relationships (see [13]):

$$G = \int P_T dB, \quad (5)$$

where G is the flux of thermal radiation of the atmosphere, $B = \sigma T^4$ is the integral flux of radiation of an absolutely black body.

In other words, G is numerically equal to the area in the coordinate system (P_F , B).

Figure 1 gives the chart for the determination of G .

It should be noted that formula (5) was derived on the assumption of the independence of the integral transmission function on temperature [13]. But, as is known [2, 4, 9, 10, 13], the transmission function actually depends not only on the content of absorbing matter in the atmosphere, but also on pressure and temperature. The dependence P_F on pressure is usually calculated with the aid of an effective absorbing mass, (see [1-10, 13, 14]). However, the problem of calculating the temperature dependence has not yet been finally solved, and, to a considerable degree, this problem may be considered debatable [13].

1. The values w^* , u^* , and m^* are expressed in "cm" (the thickness of the layer of the precipitated matter in centimeters under normal pressure and temperature).




Fig. 1. Chart for the determination of fluxes of long-wave radiation in the atmosphere.

With a radiation chart, the present work attempts to evaluate the effect of the dependence $P_F = P_F(T)$ on the determination of the fluxes of the thermal radiation of the atmosphere.

By solving the general equations for the transfer of long-wave radiation in the atmosphere, we have the following expression for the intensity of the descending monochromatic radiation which propagates in the direction θ [13]:

$$I_{\lambda \downarrow}(z, \theta) = \int_z^{\infty} \frac{k_{\lambda}(T, p)}{\cos \theta} Q(\eta) E_{\lambda}(T) e^{-\frac{1}{\cos \theta} \int_z^{\eta} k_{\lambda}(T, p)(z) dz} d\eta. \quad (6)$$

Here $I_{\lambda \downarrow}(z, \theta)$ - is the intensity of the descending long-wave monochromatic radiation at the level z , $k_{\lambda}(T, p)$ is the absorption coefficient for the wave length λ , $T = T(\eta)$ is the temperature of the absorbing (radiating)

medium, $p = p(\eta)$ is the total pressure in the atmosphere, ρ is the density of the matter absorbing the radiation, $E_\lambda(T)$ is the intensity of the radiation of an absolutely black body for wave-length λ , θ is the zenith angle.

Let us express the absorption coefficient in the following form:

$$k_\lambda(T, p) = k_{0\lambda} F(T, p), \quad (7)$$

where $F(T, p)$ is a certain function of temperature and pressure ($k_{0\lambda}$ is considered independent of T and p).

With the calculation of (7), expression (6) will have the following form:

$$I_{\lambda \downarrow} = \int_z^\infty \frac{k_{0\lambda}}{\cos^2 \theta} F(T, p) \varrho(\eta) E_\lambda(T) e^{-\frac{k_{0\lambda}}{\cos \theta} \int_z^\eta F(T, p) \varrho(\xi) d\xi} d\eta. \quad (8)$$

Let us introduce the effective absorbing mass w^* which is determined in the following fashion:

$$w^* = \int_z^\infty F(T, p) \varrho(\xi) d\xi. \quad (9)$$

In such a case

$$I_{\lambda \downarrow} = \int_0^{w^*} \frac{k_{0\lambda}}{\cos^2 \theta} E_\lambda(T) e^{-\frac{k_{0\lambda}}{\cos \theta} w^*} d(w^*). \quad (10)$$

Integrating (10) according to λ with the aid of the Ambartsumian-Lebedinskii method [13], we have

$$I_{\downarrow} = \int_0^{w^*} E(T) d(w^*) \int_0^\infty \frac{k_0}{\cos^2 \theta} f(k_0, T) e^{-\frac{k_0}{\cos \theta} w^*} dk_0. \quad (11)$$

It was assumed here that in regions of the spectrum, for which the inequality $k_0 < k_{0\lambda} < k_0 + dk_0$, the value of the absorption coefficient may in practice be considered as constant (k_0). The function $f(k_0, T)$ determines a

TABLE 2
Values of the function ΔP_1 (w*, u*)

lg u*	lg w*	2.00	1.90	1.80	1.70	1.60	1.50	1.40	1.30	1.20	1.10	1.00	0.90	0.80	0.70	0.60	0.50	0.40	0.30	0.20	0.10	0.00	1.80	1.60	1.40	1.20	1.00	0.80	0.60	0.40	0.20	0.00	2.30	2.40	2.50																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																		
4.00	868	851	838	811	822	823	812	805	790	787	781	777	771	765	758	752	746	739	733	727	722	717	712	707	701	695	689	683	677	671	665	659	653	647	641	635	629	623	617	611	605	599	593	587	581	575	569	563	557	551	545	539	533	527	521	515	509	503	497	491	485	479	473	467	461	455	449	443	437	431	425	419	413	407	401	395	389	383	377	371	365	359	353	347	341	335	329	323	317	311	305	299	293	287	281	275	269	263	257	251	245	239	233	227	221	215	209	203	197	191	185	179	173	167	161	155	149	143	137	131	125	119	113	107	101	95	89	83	77	71	65	59	53	47	41	35	29	23	17	11	5	-1	-7	-13	-19	-25	-31	-37	-43	-49	-55	-61	-67	-73	-79	-85	-91	-97	-103	-109	-115	-121	-127	-133	-139	-145	-151	-157	-163	-169	-175	-181	-187	-193	-199	-205	-211	-217	-223	-229	-235	-241	-247	-253	-259	-265	-271	-277	-283	-289	-295	-301	-307	-313	-319	-325	-331	-337	-343	-349	-355	-361	-367	-373	-379	-385	-391	-397	-403	-409	-415	-421	-427	-433	-439	-445	-451	-457	-463	-469	-475	-481	-487	-493	-499	-505	-511	-517	-523	-529	-535	-541	-547	-553	-559	-565	-571	-577	-583	-589	-595	-601	-607	-613	-619	-625	-631	-637	-643	-649	-655	-661	-667	-673	-679	-685	-691	-697	-703	-709	-715	-721	-727	-733	-739	-745	-751	-757	-763	-769	-775	-781	-787	-793	-799	-805	-811	-817	-823	-829	-835	-841	-847	-853	-859	-865	-871	-877	-883	-889	-895	-901	-907	-913	-919	-925	-931	-937	-943	-949	-955	-961	-967	-973	-979	-985	-991	-997	-1003	-1009	-1015	-1021	-1027	-1033	-1039	-1045	-1051	-1057	-1063	-1069	-1075	-1081	-1087	-1093	-1099	-1105	-1111	-1117	-1123	-1129	-1135	-1141	-1147	-1153	-1159	-1165	-1171	-1177	-1183	-1189	-1195	-1201	-1207	-1213	-1219	-1225	-1231	-1237	-1243	-1249	-1255	-1261	-1267	-1273	-1279	-1285	-1291	-1297	-1303	-1309	-1315	-1321	-1327	-1333	-1339	-1345	-1351	-1357	-1363	-1369	-1375	-1381	-1387	-1393	-1399	-1405	-1411	-1417	-1423	-1429	-1435	-1441	-1447	-1453	-1459	-1465	-1471	-1477	-1483	-1489	-1495	-1501	-1507	-1513	-1519	-1525	-1531	-1537	-1543	-1549	-1555	-1561	-1567	-1573	-1579	-1585	-1591	-1597	-1603	-1609	-1615	-1621	-1627	-1633	-1639	-1645	-1651	-1657	-1663	-1669	-1675	-1681	-1687	-1693	-1699	-1705	-1711	-1717	-1723	-1729	-1735	-1741	-1747	-1753	-1759	-1765	-1771	-1777	-1783	-1789	-1795	-1801	-1807	-1813	-1819	-1825	-1831	-1837	-1843	-1849	-1855	-1861	-1867	-1873	-1879	-1885	-1891	-1897	-1903	-1909	-1915	-1921	-1927	-1933	-1939	-1945	-1951	-1957	-1963	-1969	-1975	-1981	-1987	-1993	-1999	-2005	-2011	-2017	-2023	-2029	-2035	-2041	-2047	-2053	-2059	-2065	-2071	-2077	-2083	-2089	-2095	-2101	-2107	-2113	-2119	-2125	-2131	-2137	-2143	-2149	-2155	-2161	-2167	-2173	-2179	-2185	-2191	-2197	-2203	-2209	-2215	-2221	-2227	-2233	-2239	-2245	-2251	-2257	-2263	-2269	-2275	-2281	-2287	-2293	-2299	-2305	-2311	-2317	-2323	-2329	-2335	-2341	-2347	-2353	-2359	-2365	-2371	-2377	-2383	-2389	-2395	-2401	-2407	-2413	-2419	-2425	-2431	-2437	-2443	-2449	-2455	-2461	-2467	-2473	-2479	-2485	-2491	-2497	-2503	-2509	-2515	-2521	-2527	-2533	-2539	-2545	-2551	-2557	-2563	-2569	-2575	-2581	-2587	-2593	-2599	-2605	-2611	-2617	-2623	-2629	-2635	-2641	-2647	-2653	-2659	-2665	-2671	-2677	-2683	-2689	-2695	-2701	-2707	-2713	-2719	-2725	-2731	-2737	-2743	-2749	-2755	-2761	-2767	-2773	-2779	-2785	-2791	-2797	-2803	-2809	-2815	-2821	-2827	-2833	-2839	-2845	-2851	-2857	-2863	-2869	-2875	-2881	-2887	-2893	-2899	-2905	-2911	-2917	-2923	-2929	-2935	-2941	-2947	-2953	-2959	-2965	-2971	-2977	-2983	-2989	-2995	-3001	-3007	-3013	-3019	-3025	-3031	-3037	-3043	-3049	-3055	-3061	-3067	-3073	-3079	-3085	-3091	-3097	-3103	-3109	-3115	-3121	-3127	-3133	-3139	-3145	-3151	-3157	-3163	-3169	-3175	-3181	-3187	-3193	-3199	-3205	-3211	-3217	-3223	-3229	-3235	-3241	-3247	-3253	-3259	-3265	-3271	-3277	-3283	-3289	-3295	-3301	-3307	-3313	-3319	-3325	-3331	-3337	-3343	-3349	-3355	-3361	-3367	-3373	-3379	-3385	-3391	-3397	-3403	-3409	-3415	-3421	-3427	-3433	-3439	-3445	-3451	-3457	-3463	-3469	-3475	-3481	-3487	-3493	-3499	-3505	-3511	-3517	-3523	-3529	-3535	-3541	-3547	-3553	-3559	-3565	-3571	-3577	-3583	-3589	-3595	-3601	-3607	-3613	-3619	-3625	-3631	-3637	-3643	-3649	-3655	-3661	-3667	-3673	-3679	-3685	-3691	-3697	-3703	-3709	-3715	-3721	-3727	-3733	-3739	-3745	-3751	-3757	-3763	-3769	-3775	-3781	-3787	-3793	-3799	-3805	-3811	-3817	-3823	-3829	-3835	-3841	-3847	-3853	-3859	-3865	-3871	-3877	-3883	-3889	-3895	-3901	-3907	-3913	-3919	-3925	-3931	-3937	-3943	-3949	-3955	-3961	-3967	-3973	-3979	-3985	-3991	-3997	-4003	-4009	-4015	-4021	-4027	-4033	-4039	-4045	-4051	-4057	-4063	-4069	-4075	-4081	-4087	-4093	-4099	-4105	-4111	-4117	-4123	-4129	-4135	-4141	-4147	-4153	-4159	-4165	-4171	-4177	-4183	-4189	-4195	-4201	-4207	-4213	-4219	-4225	-4231	-4237	-4243	-4249	-4255	-4261	-4267	-4273	-4279	-4285	-4291	-4297	-4303	-4309	-4315	-4321	-4327	-4333	-4339	-4345	-4351	-4357	-4363	-4369	-4375	-4381	-4387	-4393	-4399	-4405	-4411	-4417	-4423	-4429	-4435	-4441	-4447	-4453	-4459	-4465	-4471	-4477	-4483	-4489	-4495	-4501	-4507	-4513	-4519	-4525	-4531	-4537	-4543	-4549	-4555	-4561	-4567	-4573	-4579	-4585	-4591	-4597	-4603	-4609	-4615	-4621	-4627	-4633	-4639	-4645	-4651	-4657	-4663	-4669	-4675	-4681	-4687	-4693	-4699	-4705	-4711	-4717	-4723	-4729	-4735	-4741	-4747	-4753	-4759	-4765	-4771	-4777	-4783	-4789	-4795	-4801	-4807	-4813	-4819	-4825	-4831	-4837	-4843	-4849	-4855	-4861	-4867	-4873	-4879	-4885	-4891	-4897	-4903	-4909	-4915	-4921	-4927	-4933	-4939	-4945	-4951	-4957	-4963	-4969	-4975	-4981	-4987	-4993	-4999	-5005	-5011	-5017	-5023	-5029	-5035	-5041	-5047	-5053	-5059	-5065	-5071	-5077	-5083	-5089	-5095	-5101	-5107	-5113	-5119	-5125	-5131	-5137	-5143	-5149	-5155	-5161	-5167	-5173	-5179	-5185	-5191	-5197	-5203	-5209	-5215	-5221	-5227	-5233	-5239	-5245	-5251	-5257	-5263	-5269	-5275	-5281	-5287	-5293	-5299	-5305	-5311	-5317	-5323	-5329	-5335	-5341	-5347	-5353	-5359	-5365	-5371	-5377	-5383	-5389	-5395	-5401	-5407	-5413	-5419	-5425	-5431	-5437	-5443	-5449	-5455	-5461	-5467	-5473	-5479	-5485	-5491	-5497	-5503	-5509	-5515	-5521	-5527	-5533	-5539	-5545	-5551	-5557	-5563	-5569	-5575	-5581	-5587	-5593	-5599	-5605	-5611	-5617	-5623	-5629	-5635	-5641	-5647	-5653	-5659	-5665	-5671	-5677	-5683	-5689	-5695	-5701	-5707	-5713	-5719	-5725	-5731	-5737	-5743	-5749	-5755	-5761	-5767	-5773	-5779	-5785	-5791	-5797	-5803	-5809	-5815	-5821	-5827	-5833	-5839	-5845	-5851	-5857	-5863	-5869	-5875	-5881	-5887	-5893	-5899	-5905	-5911	-5917	-5923	-5929	-5935	-5941	-5947	-5953	-5959	-5965	-5971	-5977	-5983	-5989	-5995	-6001	-6007	-6013	-6019	-6025	-6031	-6037	-6043	-6049	-6055	-6061	-6067	-6073	-6079	-6085	-6091	-6097	-6103	-6109	-6115	-6121	-6127	-6133	-6139	-6145	-6151	-6157	-6163	-6169	-6175	-6181	-6187	-6193	-6199	-6205	-6211	-6217	-6223	-6229	-6235	-6241	-6247	-6253	-6259	-6265	-6271	-6277	-6283	-6289	-6295	-6301	-6307	-6313	-6319	-6325	-6331	-6337	-6343	-6349	-6355	-6361	-6367	-6373	-6379	-6385	-6391	-6397	-6403	-6409	-6415	-6421	-6427	-6433	-6439	-6445	-6451	-6457	-6463	-6469	-647

TABLE 3

Values of the function $\Delta P_1 (w^*, m^*)$

w^*	1.00	3.00	3.50	2.00	2.10	2.20	2.30	2.40	2.50	2.60	2.70	2.80	2.90	1.00	1.10	1.20	1.30	1.40	1.50	1.60	1.70	1.80	1.90	0.00
1.00	87	87	86	82	81	81	80	78	76	73	71	68	65	62	58	55	51	48	45	42	40	38	36	34
1.05	86	85	84	81	80	79	78	76	73	71	68	65	62	59	55	51	47	44	41	38	36	34	32	30
1.10	81	81	81	80	79	78	77	75	73	71	68	65	62	59	55	51	47	44	41	38	36	34	32	30
1.15	82	82	82	81	80	79	78	76	73	71	68	65	62	59	55	51	47	44	41	38	36	34	32	30
1.20	80	80	80	79	78	77	75	73	71	68	65	62	59	55	51	47	44	41	38	36	34	32	30	28
1.25	77	77	77	76	75	74	72	70	68	65	62	59	55	52	49	46	43	40	37	35	33	31	29	27
1.30	76	76	75	74	73	72	70	68	65	62	59	55	52	49	46	43	40	37	35	33	31	29	27	25
1.35	75	75	74	73	72	71	69	67	64	61	58	54	51	48	45	42	39	36	34	32	30	28	26	24
1.40	73	73	72	71	70	69	67	64	61	58	54	51	48	45	42	39	36	34	32	30	28	26	24	22
1.45	72	72	71	70	69	68	66	63	60	57	53	50	47	44	41	38	35	33	31	29	27	25	23	21
1.50	71	71	70	69	68	67	65	62	59	55	52	49	46	43	40	37	35	33	31	29	27	25	23	21
1.55	70	70	69	68	67	66	64	61	58	54	51	48	45	42	39	36	34	32	30	28	26	24	22	20
1.60	69	69	68	67	66	65	63	60	57	53	50	47	44	41	38	35	33	31	29	27	25	23	21	19
1.65	68	68	67	66	65	64	62	59	55	52	49	46	43	40	37	35	33	31	29	27	25	23	21	19
1.70	67	67	66	65	64	63	61	58	54	51	48	45	42	39	36	34	32	30	28	26	24	22	20	18
1.75	66	66	65	64	63	62	60	57	53	50	47	44	41	38	35	33	31	29	27	25	23	21	19	17
1.80	65	65	64	63	62	61	59	56	52	49	46	43	40	37	35	33	31	29	27	25	23	21	19	17
1.85	64	64	63	62	61	60	58	55	51	48	45	42	39	36	34	32	30	28	26	24	22	20	18	16
1.90	63	63	62	61	60	59	57	54	50	47	44	41	38	35	33	31	29	27	25	23	21	19	17	15
1.95	62	62	61	60	59	58	56	53	49	46	43	40	37	35	33	31	29	27	25	23	21	19	17	15
2.00	61	61	60	59	58	57	55	52	48	45	42	39	36	34	32	30	28	26	24	22	20	18	16	14
2.05	60	60	59	58	57	56	54	51	47	44	41	38	35	33	31	29	27	25	23	21	19	17	15	13
2.10	59	59	58	57	56	55	53	50	46	43	40	37	35	33	31	29	27	25	23	21	19	17	15	13
2.15	58	58	57	56	55	54	52	49	45	42	39	36	34	32	30	28	26	24	22	20	18	16	14	12
2.20	57	57	56	55	54	53	51	48	44	41	38	35	33	31	29	27	25	23	21	19	17	15	13	11
2.25	56	56	55	54	53	52	50	47	43	40	37	35	33	31	29	27	25	23	21	19	17	15	13	11
2.30	55	55	54	53	52	51	49	46	42	39	36	34	32	30	28	26	24	22	20	18	16	14	12	10
2.35	54	54	53	52	51	50	48	45	41	38	35	33	31	29	27	25	23	21	19	17	15	13	11	9
2.40	53	53	52	51	50	49	47	44	40	37	35	33	31	29	27	25	23	21	19	17	15	13	11	9
2.45	52	52	51	50	49	48	46	43	39	36	34	32	30	28	26	24	22	20	18	16	14	12	10	8
2.50	51	51	50	49	48	47	45	42	38	35	33	31	29	27	25	23	21	19	17	15	13	11	9	7
2.55	50	50	49	48	47	46	44	41	37	34	32	30	28	26	24	22	20	18	16	14	12	10	8	6
2.60	49	49	48	47	46	45	43	40	36	33	31	29	27	25	23	21	19	17	15	13	11	9	7	5
2.65	48	48	47	46	45	44	42	39	35	32	30	28	26	24	22	20	18	16	14	12	10	8	6	4
2.70	47	47	46	45	44	43	41	38	34	31	29	27	25	23	21	19	17	15	13	11	9	7	5	3
2.75	46	46	45	44	43	42	40	37	33	30	28	26	24	22	20	18	16	14	12	10	8	6	4	2
2.80	45	45	44	43	42	41	39	36	32	29	27	25	23	21	19	17	15	13	11	9	7	5	3	1
2.85	44	44	43	42	41	40	38	35	31	28	26	24	22	20	18	16	14	12	10	8	6	4	2	0
2.90	43	43	42	41	40	39	37	34	30	27	25	23	21	19	17	15	13	11	9	7	5	3	1	0
2.95	42	42	41	40	39	38	36	33	29	26	24	22	20	18	16	14	12	10	8	6	4	2	0	0
3.00	41	41	40	39	38	37	35	32	28	25	23	21	19	17	15	13	11	9	7	5	3	1	0	0

fraction of the intensity of an absolutely black body arriving on the total number of regions of the spectrum to which the absorption coefficient k_0 corresponds. $E(T)$ in expression (11) designates the integral intensity of the radiation of an absolutely black body.

On the other hand, it is easy to show that by using an effective absorbing mass which is determined from formula (9), the integral transmission function of the atmosphere is expressed as follows:

$$P(\omega^*, T_1) = \int_0^{\infty} j(k_0, T_1) e^{-\frac{k_0}{\cos \theta} \omega^*} dk_0. \quad (12)$$

Here T_1 designates the temperature of the source of the radiation; and it is assumed that this source is an absolutely black body.

In addition, let us introduce a new function P^* which is determined by the relationship

$$P^*[\omega^*, T(\omega^*)] = P[\omega^*, T_1 = T(\omega^*)]. \quad (13)$$

In other words,

$$P^*[\omega^*, T(\omega^*)] = \int_0^{\infty} j[k_0, T(\omega^*)] e^{-\frac{k_0}{\cos \theta} \omega^*} dk_0. \quad (14)$$

Now let us find the derivatives P and P^* with respect to w^* :

$$\frac{dP}{d\omega^*} = - \int_0^{\infty} \frac{k_0}{\cos \theta} j(k_0, T_1) e^{-\frac{k_0}{\cos \theta} \omega^*} dk_0. \quad (15)$$

Here it is assumed that temperature T_1 does not change.

$$\begin{aligned} \frac{dP^*}{d\omega^*} = & - \int_0^{\infty} \frac{k_0}{\cos \theta} j(k_0, T) e^{-\frac{k_0}{\cos \theta} \omega^*} dk_0 + \\ & + \frac{dT}{d\omega^*} \int_0^{\infty} \frac{\partial j(k_0, T)}{\partial T} e^{-\frac{k_0}{\cos \theta} \omega^*} dk_0. \end{aligned} \quad (16)$$

We see that the integral $S_1 = \int_0^{\infty} \frac{k_0}{\cos \theta} j(k_0, T) e^{-\frac{k_0}{\cos \theta} \omega^*} dk_0$ in expression (11) is equal to the first term of expression (16). Therefore, using $dP^*/d\omega^*$ and $dP/d\omega^*$ instead of S_1 , creates errors. In the first case we do not consider the second term of expression (16), and in the second case the "displacement effect" is not considered.

Let us try to make an approximate evaluation of the magnitude of these errors. Under average conditions in the troposphere¹, the relationship between T and w^* is well described by the formula

$$T = -8(w^*)^8 - 10w^* + 20, \quad (17)$$

where w^* is expressed in "cm" and T in C° .

To simplify the calculations, let us use instead of expressions (12) and (14) the approximate formulas

$$P(w^*, T_1) = \sum_{j=1}^n f_j(T_1) e^{-k_j w^*}, \quad (18)$$

where T_1 is considered constant, and

$$P^*(w^*, T) = \sum_{j=1}^n f_j(T) e^{-k_j w^*}, \quad (19)$$

where $T = T(w^*)$.

Formulas (18) and (19) are taken from the monograph of K. Ia. Kondrat'ev [13].

This work also gives the absorption coefficients k_j and the connection between f_j and the temperature:

$$f_j = a_j + b_j T. \quad (20)$$

Here a_j and b_j are certain constants for each region j .

1. It is assumed that the vertical temperature gradient is equal to 6 deg/km, and the vapor density ρ_w decreases with height according to exponential law. At the earth's surface $T = 20^\circ C$ and $\rho_w = 7 \times 10^{-6} g/cm^3$.

On the basis of the data given above, the following magnitudes were calculated in the present work: S_1 , $S_2 = dP^*/dw^*$, $S_3 = dP(T_1 = 260^\circ\text{K})/dw^*$, $S_4 = dP(T_1 = 300^\circ\text{K})/dw^*$. Figure 2 shows the results of these calculations. As this figure shows, the use of dP^*/dw^* instead of S_1 causes many great errors, especially in the upper layers of the troposphere. The differences between S_1 and S_3 , and also between S_1 and S_4 are rather small. Therefore, it is necessary to consider as completely substantiated the use of $dP(w^*, T_1)/dw^*$ instead of S_1 in formula (11) (assuming that T_1 is a certain average temperature in the atmosphere). In other words, we can disregard the influence of the displacement effect in the given case.




Fig. 2. A comparison of the functions S_n : 1- S_1 , 2- S_2 , 3- S_3 , 4- S_4 .

As is known [13], the dependence of the absorption coefficient on temperature and the "displacement effect" act in opposite directions. Therefore, it should be considered as proven that the relationship $k_\lambda = k_\lambda(T)$ is not considered. In such a case formula (9) has the following form:

$$\omega^* = \int F(p) Q(z) dz, \quad (21)$$

where $F(p)$ is a certain function of pressure.

On the basis of what has been said above, instead of (11) we will get

$$I \downarrow = \int_{\omega_{\infty}^*}^0 E(T) d\omega^* \frac{dP}{d\omega^*} = \int_{P(\omega_{\infty}^*)}^1 E(T) dP. \quad (22)$$

Partially integrating (22), we get an expression analogous to formula (5):

$$I \leftarrow = \oint P dE. \quad (23)$$

We can obtain an expression even for $I \uparrow$ in this way.

We should note that the results given above are justified even for fluxes of long-wave radiation in the atmosphere. As is known [13], the radiation fluxes are calculated by integrating the corresponding intensities according to the solid angles corresponding to a semisphere. In such a case, we will have formula (5) instead of formula (23).

Summarizing the results given above, we can say that on the basis of the approximate formula $G = \int P_F(\omega^*) dB$, we can determine with sufficient accuracy the long-wave fluxes of the radiation of the atmosphere, but the use of the formula $G = \int P_F[\omega^*, T(\omega^*)] dB$ cannot be justified.

As was shown in works [16] and [43], atmospheric ozone plays only an insignificant role in the determination of the values of the integral fluxes of thermal radiation in the lower troposphere. Therefore, the effect of ozone on the absorption of long-wave radiation in the atmosphere can be disregarded when calculating descending fluxes ($G \downarrow$) in the lower troposphere and ascending fluxes ($G \uparrow$) to 15-20 km. In such a case, we can use only one table instead of Tables 2 and 3, in which the values of the integral transmission function $P_F(\omega^*, u^*)$ are given.

As an example, we calculated the descending flux, the ascending flux, and the effective radiation ($F = G \uparrow - G \downarrow$) at the levels 0.3 and 8 km for certain latitudinal zones of the earth. In addition, stratifications of temperature, humidity, and pressure, taken from works [10, 44, 45] were used. Data on the vertical distribution of ozone were taken from work [46]. The concentration of carbon dioxide in the atmosphere was taken equal to 0.03% (by volume) for all the zones, since there were no data on the latitudinal variation of CO_2 . When calculating the effective absorbing masses of water vapor, we applied the correction which is usually used $(p/p_0)^{0.5}$, and for carbon dioxide and ozone this correction was $(p/p_0)^{0.8}$ and $(p/p_0)^{0.2}$ respectively (see [15, 16, 18]). Table 4 gives the results of these calculations.

TABLE 4
Values of the Fluxes of Thermal Radiation in the
Atmosphere ($\text{cal/cm}^2 \text{ min}$)

latitudinal zone ($^{\circ}\text{N}$)	$z=0$		$z=3$		$z=8$	
	$G \downarrow$	F	$G \downarrow$	F	$G \downarrow$	F
0-10	0,574	0,093	0,392	0,194	0,164	0,296
10-20	0,545	0,101	0,348	0,209	0,132	0,317
20-30	0,507	0,114	0,312	0,220	0,117	0,313
30-40	0,429	0,117	0,275	0,208	0,097	0,293
40-50	0,359	0,122	0,240	0,198	0,082	0,270
50-60	0,296	0,125	0,201	0,187	0,076	0,246
60-70	0,236	0,129	0,169	0,172	0,077	0,218

Analogous calculations are carried out in works [14, 17] with the aid of the charts of F. Brooks, A. A. Dmitriev, R. Mugge and F. Moller, G. Robinson, F. N. Shekhter, V. Elsasser and G. Yamamoto. Comparing the results of the determination of $G \downarrow$, $G \uparrow$ and F which were obtained in the present work and in works [14, 17], we see that near the earth's surface and in the lower part of the troposphere, our results agree rather closely with the data obtained from the charts of F. Shekhter and F. Brooks. However,

when $z = 8$ km, there is a closer agreement with the results of the calculations from the charts of F. Møller and G. Robinson.

Unfortunately, at present there is no satisfactorily complete complex of experimental data, which makes it impossible to make a direct comparison of the results with the experiment.

Since the radiation nomogram chart in this work was constructed on the basis of a most careful analysis of contemporary data on the absorption of infrared radiation in the atmosphere, it may be considered most reliable.

LITERATURE

1. Brooks, F. A. "Atmospheric radiation and its reflection from the ground," Journal of Meteorology, 9(1): 41-52, 1952.
2. Dmitriev, A. A. "K voprosu o radiatsionnom balanse atmosfery" (The problem of the radiation balance of the atmosphere), Meteorologiya i Gidrologiya, No. 11, 1940.
3. Moller, F. "Grundlagen eines Diagramms zur Berechnung langwelliger Strahlungsströme" (Basis of the diagram for calculating long-wave radiation), Meteorologische Zeitschrift, 61(2), 1944.
4. Moller, F. "Long-wave radiation," Compendium of Meteorology, Boston, AMS, pp. 34-49, 1951.
5. Robinson, G. D. "Notes on the measurement and estimation of atmospheric radiation - 1," Royal Meteorological Society, Quarterly Journal, 73: (315-316), 1947.
6. Robinson, G. D. "Notes on the measurement and estimation of atmospheric radiation-2," Royal Meteorological Society, Quarterly Journal, 76(327): 37-51, 1950.
7. Shekhter, F. N. "K vychisleniiu luchistykh potokov tepla v atmosfere" (Calculation of the radiational heat flux in the atmosphere), Glavnaia Geofizicheskaya Observatoriya, Trudy, 22(84); 38-51, 1950.
8. Shekhter, F. N. "Raschet luchistykh potokov dlinnovolnovo radiatsii iz ogranichennogo telesnogo ugla i poluprostranstva" (Calculation of the radiation fluxes of long-wave radiation from a limited solid angle and semispace), Glavnaia Geofizicheskaya Observatoriya, Trudy, 39(101), 1953.
9. Elsasser, W. M. "Heat transfer by infrared radiation in the atmosphere," Harvard Meteorological Station, No. 6, 1942.
10. Yamamoto, G. "On a radiation chart", Tohoku University, Science Reports, 5th Series, Geophysics, 4(1): 9-23, 1952.
11. Kondrat'ev, K. Ia. "Nekotorye voprosy luchistogo teploobmena v atmosfere" (Several problems of radiant heat exchange in the atmosphere), Uchrezhdenie zapiski, LGU, Seriya Fizicheskaya, No. 7, 1947.

12. Kondrat'ev, K. Ia. "Perenos dlinnovolnovogo izlucheniia v atmosfere" (The transfer of long-wave radiation in the atmosphere). Moscow, Gostekhizdat, 1950. 287 pp.
13. Kondrat'ev, K. Ia. Luchistyi teploobmen v atmosfere (Radiant heat exchange in the atmosphere) Moscow, Gidrometeoizdat, 1956. 419 pp.
14. Kh. Iu. Niilisk. "K voprosu o raschetakh teplovogo izlucheniia atmosfery, issledovaniia po fizike atmosfery" (The problem of calculating the thermal radiation of the atmosphere, investigations of the physics of the atmosphere), Institut Fiziki i Astronomii, Akademii Nauk ESSR, No. 2, 1960.
15. Kondrat'ev, K. Ia., and Kh. Iu. Niilisk. "K voprosu o teplovom izluchenii uglekislogo gaza v atmosfere; o teplovom izluchenii 9.6 μ polosy pogloshcheniia ozona v atmosfere," Nauchnoe soobshchenie Instituta Geologii i Geografii Akademii Nauk Lit. SSR" (The problem of the thermal radiation of carbon dioxide in the atmosphere; the thermal radiation of the 9.6 μ absorption band of ozone in the atmosphere." A scientific report of the Institute of Geology and Geography of the Academy of Sciences of the Lithuanian Soviet Socialist Republic), No. 2, 1960 in print.
16. Same as 15.
17. Kondrat'ev, K. Ia., and H. J. Niilisk. "Comparison of radiation charts," Geofisica pura e applicata, 46, 1960/II.
18. Kondrat'ev, K. Ia., and H. J. Niilisk. "The problem of carbon dioxide heat radiation in the atmosphere," Geofisica pura e applicata, 46, 1960/II.
19. Yamamoto, G. and T. Sasamori. "Calculation of the absorption of the 15 μ carbon dioxide band." Tohoku University, Science Reports, 5th Series, Geophysics, 10(2): 37-57, 1958.
20. Howard, J. N., D. E. Burch, and D. Williams. "Near infrared transmission through synthetic atmospheres," U. S. Air Force, Cambridge Research Center, Geophysical Research Paper, No. 40, 1955. 244 pp.
21. Elsasser, W. "New values for the infrared absorption coefficient of atmospheric water vapor," Monthly Weather Review, 66(6), 1938.

22. Yamamoto, G. and G. Onishi. "Absorption coefficient of water vapor in the far infrared region," Tohoku University, Science Reports, 5th Series, 1(1), 1949.
23. Yamamoto, G. and G. Onishi. "Absorption coefficient of water vapor in the near infrared region," Tohoku University, Science Reports, 5th Series, 1(2), 1949.
24. Adel, A. "Atmospheric absorption of infrared solar radiation at the Lowell Observatory I", Astrophysical Journal, 89(1), 1939.
25. Adel, A. and C. D. Lampland. "Atmospheric absorption of infrared solar radiation at the Lowell Observatory II", Astrophysical Journal, 91(1), 1940.
26. Adel, A. and C. D. Lampland. "Atmospheric absorption of infrared solar radiation at the Lowell Observatory III, IV, Astrophysical Journal, 91(5), 1940.
27. Adel, A. "An estimate of transparency of the atmospheric window 16 to 24 μ ". Optical Society of America, Journal, 37(10), 1947.
28. Anthony, R. "Atmospheric absorption of solar infrared radiation", Physical Review, 85(4): 674, 1952.
29. Cowling, T. C. "The absorption of water vapor in the far infrared," Report on Progress of Physics, V(IX), 1943.
30. Cowling, T. C. "Atmospheric absorption of heat radiation by water vapor," Philosophical Magazine, 7th Series, 41(313), 1950.
31. Elder, T. and J. Strong. "The infrared transmission of atmospheric windows," Franklin Institute, Journal, 255(3), 1953.
32. Fowle, F. E. "The spectroscopic determination of aqueous vapor," Astrophysical Journal, 35: 149-162, 1912.
33. Fowle, F. E. "The transparency of aqueous vapor," Astrophysical Journal, 42: 394-411, 1915.
34. Gebbie, H. A., W. K. Harding, C. Hilsum, A. W. Price, and V. Roberts. "Atmospheric transmission in the 1-14 μ region," Royal Society of London, Proceedings A, 206: 87-107, 1951.
35. Goody, R. M. and W. T. Roach. "Absorption and emission in the atmospheric window from 770-1250 cm^{-1} ," Royal Meteorological Society, Quarterly Journal, 84(362): 319-333, 1958.

36. Palmer, C. H. "Long path water vapor spectra with pressure broadening I, II", Optical Society of America, Journal, 47(11), 1958.
37. Saiedy, F. "Background (continuum) extinction in the atmosphere," Oxford Radiation Symposium, Abstract, No. 30, 1959.
38. Saiedy, F. and R. M. Goody. "The solar emission intensity at 11μ ," Royal Astronomical Society, Monthly Notes, 119(3), 1959.
39. Vigroux, E. "Emission continue de l'atmosphère terrestre à 9.6μ " (Continuous emission of the terrestrial atmosphere at 9.6μ), Annales de Geophysique, 15(4): 453-460, 1959.
40. Kastrov, V. G. "Izmereniia effektivnogo izlucheniia v oblasti maksimal'noi prozrachnosti vodianogo para" (Measurements of the effective radiation in a region of the maximum transparency of water vapor), Meteorologiya i Gidrologiya, No. 1, 1946.
41. Kastrov, V. G. "Izmereniia infrakrasnogo izlucheniia atmosfery v trekh uchastakh spektra" (Measurements of the infrared radiation of the atmosphere in three parts of the spectrum), Tsentral'naia Aerologicheskaya Observatoriya, Trudy, No. 3, 1948.
42. Iaroslavskii, N. G. and A. E. Stanevich. "Vrashchatel'nyi spektr H_2O v dlinnovolnovoi infrakrasnoi oblasti 50-1500 μ " (The rotary spectrum of H_2O in the long-wave infrared region 50-1500 μ), Optika i Spektroskopiya, 5(4), 1958.
43. Kondrat'ev, K. Ia. and T. D. Matreshina. "O vliianii dlinnovolnovogo izlucheniia ozona na radiatsionnyi balans zemnoi poverkhnosti i atmosfery" (The influence of the long-wave radiation of ozone on the radiation balance of the earth's surface and atmosphere), Glavnaia Geofizicheskaya Observatoriya, Trudy, 41(103), 1953.
44. London, J. "The distribution of radiational temperature change in the Northern Hemisphere during March", Journal of Meteorology, 9(2): 145-151, 1952.
45. Ohring, G. "The radiation budget of the stratosphere," N. Y. Univ. College of Engineering, Research Div. Dept. of Meteorology and Oceanography, Contract AF 19(604)-1738, Scientific Report, No. 1, 1957.
46. Ramanathan, K. R. and R. N. Kulkarni. "Mean meridional distributions of ozone in different seasons calculated from UMKEHR observations and probable vertical transport mechanisms", Royal Meteorological Society, Quarterly Journal, 86(368), 1960.