

UNCLASSIFIED

AD 4 2 1 0 6 4

DEFENSE DOCUMENTATION CENTER

FOR

SCIENTIFIC AND TECHNICAL INFORMATION

CAMERON STATION, ALEXANDRIA, VIRGINIA



UNCLASSIFIED

NOTICE: When government or other drawings, specifications or other data are used for any purpose other than in connection with a definitely related government procurement operation, the U. S. Government thereby incurs no responsibility, nor any obligation whatsoever; and the fact that the Government may have formulated, furnished, or in any way supplied the said drawings, specifications, or other data is not to be regarded by implication or otherwise as in any manner licensing the holder or any other person or corporation, or conveying any rights or permission to manufacture, use or sell any patented invention that may in any way be related thereto.

CATALOGED BY DDC
421064
AS AD No. _____

Physical Studies, Inc.

48 Maple Avenue

Centerville 59, Ohio

PROTON AND ELECTRON SCATTERING BY DIATOMIC MOLECULES

by

R. G. BREENE, Jr.

1 April 1963

Final Report

Contract Number AF19(628)-476

Project 8658

Task 86580



Prepared For

AIR FORCE CAMBRIDGE RESEARCH LABORATORIES

OFFICE OF AEROSPACE RESEARCH

UNITED STATES AIR FORCE

BEDFORD, MASSACHUSETTS

Notices:

Request for additional copies by Agencies of the Department of Defense, their contractors, and other Government agencies should be directed to the:

DEFENSE DOCUMENTATION CENTER (DDC)
CAMERON STATION
ALEXANDRIA, VIRGINIA

Department of Defense contractors must be established for DDC services or have their 'need-to-know' certified by the cognizant military agency of their project or contract.

U. S. DEPARTMENT OF COMMERCE
OFFICE OF TECHNICAL SERVICES
Washington 25, D. C.

PHYSICAL STUDIES, INC.
48 Maple Avenue
Centerville 59, Ohio

PROTON AND ELECTRON SCATTERING BY DIATOMIC MOLECULES

by

R. G. Breene, Jr.

1 April 1963

Final Report

Contract Number AF19(628)-476

Project 8658

Task 86580

Prepared
for


~~GEOPHYSICS RESEARCH DIRECTORATE~~
AIR FORCE CAMBRIDGE RESEARCH LABORATORIES
• OFFICE OF AEROSPACE RESEARCH
UNITED STATES AIR FORCE
BEDFORD, MASSACHUSETTS

INTRODUCTION

The first, short section gives a brief introduction to the scattering matrix. In the next two sections we consider the heart of our present problem. Section II deals with the scattering of electrons by diatomic molecules, specifically oxygen. The treatment is restricted in such a way that electronic computing machinery is not requisite to the final solution. In the electron case the Born approximation is, of course, applicable and we carry the problem to the point where a numerical solution, which could readily be carried out on a desk calculator, is required. In Section III the proton-molecule scattering problem is discussed, specifically proton scattering by oxygen. The Born approximation is not applicable in this case. We begin by carrying a distorted wave treatment to the point where a very small set of one dimensional differential equations are obtained. The equations are not solved although their hand solution would pose no particular problem. We then consider the impact parameter treatment of this same proton-oxygen problem. For the familiar two-state case we carry through the problem to obtain the total cross-sections. In addition, we introduce a general multi-state treatment of this impact parameter problem which is not specifically applied. In both these latter sections target distortion by the projectile particle is included in the scattering potential. The last section is concerned with additions to Professor Takayanagi's bibliography.

SECTION I
THE SCATTERING MATRIX*

We consider first the scattering of electrons by diatomic molecules in the energy range from one to fifty thousand electron volts. One here encounters elastic, inelastic, and charge exchange scattering where the inelastic scattering may involve electronic, vibrational, and rotational excitation, ionization, and dissociation either alone or in combination. As an example of what may occur we remark the following reactions :



which may of course also occur in any combination.

In considering the problem we may begin by indicating the scattering matrix form although the individual elements therein will not be evaluated until a later stage in the development. The wave function for the electron in the absence of the neutral molecule will have the asymptotic form

$$\phi_x(r) \sim \frac{1}{r} A_x \left[\exp\left\{-i(k_x r - l\frac{\pi}{2})\right\} - \exp\left\{+i(k_x r - l\frac{\pi}{2})\right\} \right] \quad (1)$$

*The material in this section is of a review nature. References are provided in Section IV and in the Takayanagi review.

in the case where the electron is in channel λ . The normalization is to unit amplitude at infinity. The first term in the bracket represents the incoming electron while the second term represents the outgoing. We shall define the term "channel" after our introduction of the scatterer. When we introduce the scatterer we have the following expression for the free electron in the presence of the molecule:

$$\phi_{\lambda}(r) \sim v_{\lambda}^{-1/2} r^{-1} [A_{\lambda} \exp\{-i(k_{\lambda}r - l\frac{\pi}{2})\} - B_{\lambda} \exp\{i(k_{\lambda}r - l\frac{\pi}{2})\}] \quad (2)$$

We now define "channel" as a specific set of parameters defining the translational state of the electron and the electronic-vibration-rotation state of the molecule. In an elastic collision the electron-molecule system will re-enter the entry channel after collision. We now consider the exit channel parameter, B_{λ} , for the case of an elastic collision.

In order to do so we consider the familiar asymptotic solution for the l -th partial wave for a potential falling off more rapidly than r^{-2} .

$$\frac{1}{kr} \sin(kr - l\frac{\pi}{2} + \delta_l) \quad (3)$$

This may be re-written as

$$\begin{aligned} & \frac{1}{2kr} [e^{i\delta_l} e^{i(kr - l\frac{\pi}{2})} - e^{-i\delta_l} e^{-i(kr - l\frac{\pi}{2})}] \\ & = -\frac{e^{-i\delta_l}}{2kr} [e^{-i(kr - l\frac{\pi}{2})} - S_l e^{i(kr - l\frac{\pi}{2})}] \end{aligned} \quad (4)$$

where we now define the factor S_l as the scattering matrix element. It is apparent that it gives us the ratio of the outgoing to the incoming waves thus yielding at least one description of the scattering. In terms of the channel dispersion parameters, A_l and B_l , this may be written:

$$S_l = \frac{B_l}{A_l} \quad (5)$$

Now we remark that in our elastic case above the exit channel parameter was of amplitude equal to that of the entrance channel parameter. This, of course, would not be the case in inelastic scattering. In that situation the system sometimes departs partially through the entrance channel. However it departs partially through at least one additional channel so that entry and exit amplitudes must differ. Here again, however, the matrix element is still defined by the channel parameter ratios. The diagonal elements correspond to the ratios between the elastic parameters (A_l and B_l). The inelastically scattered electron has only an outgoing wave viz:

$$\varphi_{l'}(r) \sim r^{-1/2} B_{l'} \exp[i(k_{l'} r - Et)] \quad (6)$$

The off-diagonal elements in the scattering matrix then are given by the ratio $B_{l'}/A_l$. We discuss the scattering matrix no further but refer the reader to the extensive work which has been done in this field in recent years.

In introducing the scattering matrix we have in Eq. (1) assumed the partial wave form for the free electron. The reader will, of course,

recall that the partial wave is specified by the momentum quantum number " l " in the above equations. Although the partial wave analysis is used quite extensively in actual scattering problems, we revert to a more general form in what follows:

SECTION II

ELECTRON SCATTERING BY DIATOMICS*

A. General Formulation of the Problem

First, we are going to assume that the Franck-Condon principle is operative, that is, we suppose that the molecular wave function can be written as a product of electronic, vibrational, and rotational wave functions. Therefore we are going to somewhat simplify the following by considering only the electronic function, the vibrational and rotational being possible of later consideration.

The wave function for the molecule - electron system is taken as:

$$\Psi = \sum_{ie} P_{ie} \Phi_{An}(r_i) \chi(r_e) \psi_{n^+}(r) + \sum_{ie} P_{ie} \Phi_{An}(r_i) \chi(r_e) \psi_{n^-}(r) \quad (7)$$

In Eq. (6) P_{ie} is the permutation operator; $\Phi_{An}(r_i)$ is the molecular wave function and consists of a simple product of the molecular orbitals. We have, of course, assumed a single determinant LCAO-MO form for the molecular wave function.

We take H_{mol} to be the Hamiltonian for the molecule in the absence of any external interaction. The interaction Hamiltonian will be

$$H_{int} = -\frac{Z_a}{r_{ae}} - \frac{Z_b}{r_{be}} + \sum_{i=1}^{Z_a+Z_b} \frac{1}{r_{ie}} \quad (8a)$$

*The general development general case and Born approximation are discussed in Theory of Atomic Collisions, Quantum Theory I. Elements etc.

which leads to:

$$\left(H_{mol} - \frac{1}{2} \nabla_e^2 - \frac{k_n^2}{2} - \frac{Z_a}{r_{ae}} - \frac{Z_b}{r_{be}} + \sum \frac{1}{r_{ie}} - E_n \right) \Psi = 0 \quad (8b)$$

Under the constraint

$$(H_{mol} - E_n) \Phi_{An}(r_i) = 0$$

and after substitution for Ψ Eq (8b) becomes:

$$\begin{aligned} & \left[\nabla_e^2 + k_n^2 + 2 \left(\frac{Z_a}{r_{ae}} + \frac{Z_b}{r_{be}} - \sum \frac{1}{r_{ie}} \right) \right] \left[\sum_{ie} P \Phi_{An}(r_i) \chi(r_e) \psi_n(r) \right. \\ & \left. + \sum_{ie} P \Phi_{An}(r_i) \chi^-(r_e) \psi_n(r) \right] = 0 \end{aligned} \quad (9)$$

The χ_n^\pm are spin functions for the free electron, χ_n^+ referring to a free electron spin parallel to the molecular spin, χ_n^- referring to the anti-parallel situation. Now multiply through on the left by the plus spin function and integrate appropriately to get

$$\sum_{ie} P \Phi_{An}(r_i) \left[\nabla_e^2 + k_n^2 + 2 \left(\frac{Z_a}{r_{ae}} + \frac{Z_b}{r_{be}} - \sum \frac{1}{r_{ie}} \right) \right] \psi_n(r) = 0 \quad (10)$$

Further we now suppose that the molecule is stationary so that \underline{r} becomes \underline{r}_e , the coordinate of the free electron referred to the center-of-mass of the molecule as a coordinate center. Next the specific product nature of Φ_{An} is introduced.

$$\bar{\Phi}_{An}(r_i) = \frac{1}{\sqrt{Z_a + Z_b}} T_{\alpha} \Phi_{n\alpha}$$

and we multiply Eq. (10) through on the left by $\bar{\Phi}_{An}$ and integrate over all space. The result is:

$$\begin{aligned} \left\{ \nabla_j^2 + k_n^2 - U_{nn} \right\} \psi_{ne}(j) - \sum_{\alpha} \int K_{nn}(r_i, r_j) \psi_{n\alpha}(i) d\tau_i \\ = \sum_{m \neq n} \left\{ U_{nm} \psi_{me}(j) + \sum_{\alpha} \int K_{nm}(r_i, r_j) \psi_{m\alpha}(i) d\tau_i \right\} \end{aligned} \quad (11a)$$

$$U_{nm} = -\frac{1}{Z_a + Z_b} \sum_{\alpha} \int \phi_{n\alpha}^*(i) \phi_{m\alpha}(i) \sum H_{ij} d\tau_i \quad (11b)$$

$$\sum_{\alpha} \int K_{nm}(r_i, r_j) \psi_{m\alpha}(i) d\tau_i = \sum_{\alpha} \frac{1}{Z_a + Z_b} \int \phi_{m\alpha}(j) \phi_{n\alpha}^*(i) \left\{ H_i - \sum H_{ij} \right\} \psi_{m\alpha}(i) d\tau_i \quad (11c)$$

where

$$H_j = \nabla_j^2 + k_m^2 \quad (12a)$$

$$H_{ij} = 2 \left(\frac{Z_a}{r_{aj}} + \frac{Z_b}{r_{bj}} - \sum \frac{1}{r_{ij}} \right) \quad (12b)$$

Now note that in Eq. (11c) the individual molecular orbitals must have their spins parallel to that of the free electron. Finally, one is able to apply Green's Theorem to Eq. (11c) with the result:

$$\sum_{\alpha} \int K_{nm}(r_i, r_j) \psi_{m\alpha}(i) d\tau_i = \frac{1}{Z_a + Z_b} \sum_{\alpha} \int \phi_{m\alpha}(j) \phi_{n\alpha}^*(i) \left\{ k_m^2 - 2 \left(E_{nd} + \sum \frac{1}{r_{ij}} \right) \right\} \psi_{m\alpha}(i) d\tau_i \quad (11c')$$

Eqs. (11a), (11b), and (11c') are the general forms for the free elec-

tron wave function where spin-orbit type interactions have been left out of the original Hamiltonian. These rather fine interactions appear to have no particular place in this treatment. Some simplification of these equations has to be made in order to carry their solution any further. A number of such approximations have been made most of them having to do with the relative magnitude of the U_{nm} and K_{nm} .

The most drastic assumptions lead to the Born approximation where, in effect, one supposes that the scattering molecule is hardly there at all. We shall show that this appears adequate for the molecule-electron scattering and inadequate for the molecule-proton scattering case.

B. The Born Approximation for Electron Excitation

As we remarked above, the Born approximation is basically predicated on a rather small interaction between electron and molecule. Under these conditions, we suppose the diagonal elements in Fqs. (11) to be zero, and we replace ψ on the right hand side of the equation by the plane wave:

$$\exp(ik_m \cdot r_j)$$

If we now eliminate exchange from the equation; the result is:

$$\{\nabla_j^2 + k_n^2\} \psi_{ne}(j) = U_{nm} \exp(ik_m \cdot r_j) \quad (13a)$$

On the other hand if the exchange term is maintained we obtain:

$$\begin{aligned} \{\nabla_j^2 + k_n^2\} \psi_{ne}(j) = & U_{nm} \exp(ik_m \cdot r_j) + \frac{1}{E_a + E_b} \sum_a \int \psi_{na}(j) \psi_{na}^*(i) \{k_n^2 \\ & - 2(E_{na} + \sum \frac{1}{r_{ij}})\} \exp(ik_m \cdot r_i) d\tau_i \end{aligned} \quad (13b)$$

We consider Eq. (13a). It is well known that the solution to the equation,

$$\nabla^2 \psi + k^2 \psi = F(r)$$

is:

$$\psi = G(r) - \frac{1}{4\pi} \int \frac{e^{ik|r-r'|} F(r') dr'}{|r-r'|}$$

where:

$$\nabla^2 G + k^2 G = 0$$

In quite precise analogy we may write the solution to Eq. (13a) as

$$\psi_{ne} = e^{ik_n \underline{n} \cdot \underline{r}_j} - \frac{1}{4\pi} \int \frac{e^{ik_n |\underline{r}_j - \underline{r}'|} U_{nm}(\underline{r}') e^{ik_m \underline{n}_m \cdot \underline{r}'} d\tau'}{|\underline{r}_j - \underline{r}'|} \quad (14)$$

We may now use this expression to obtain the scattering cross section for the general case. It should be remarked here that the source of difficulty in Eq. (14) is, of course going to be U_{nm} to which we return shortly. For now, however, we recall that the cross section makes use of the scattering amplitude which in turn is available from the asymptotic form of the wave function. The asymptotic form of Eq. (14) is:

$$\psi_{ne} \sim \frac{1}{4\pi r} e^{ik_n r} \int e^{i(k_m \underline{n}_m - k_n \underline{n}_n) \cdot \underline{r}'} U_{nm}(\underline{r}') d\tau' \quad (15)$$

from which we obtain the cross section for scattering of the system into the n-th channel, the electron undergoing a deflection through an angle ϑ :

$$\sigma_n(\vartheta) = \frac{k_n}{k_m} \frac{1}{4\pi^2} \left| \int e^{i(k_m \underline{n}_m - k_n \underline{n}_n) \cdot \underline{r}'} U_{nm}(\underline{r}') d\tau' d\tau \right|^2 \quad (16)$$

Eq. (16) is written in atomic units, and k_n refers to the entry channel.

In essence the collision has induced a transition from molecular state "m" to molecular state "n" and a change of free electron linear momentum from k_m to k_n . Eq. (16) is, of course the required result. Before considering it in more detail we inquire of its validity.

The requirement which must be fulfilled in this electron - molecule case is: the velocity of the free electron must be much larger than the velocity of the orbital electrons. If we suppose something like one-half of the orbital energy is associated with kinetic energy of the orbital electrons then we may compare this energy to the translational energy of the incident electron.

The $2\sigma_g$, $2\sigma_u$ and $3\sigma_g$ electrons in oxygen, for example, have kinetic energies of around ten or twenty electron volts. Thus, one thousand volt electrons should be sufficiently energetic. We therefore accept the Born approximation for this problem and return our attention to what we may designate "the potential", namely U_{nm} .

We are using a simple determinant LCAO-MO form for the molecular wave function. Let us suppose that:

- (1) The state "n" differs from the state "m" in only one orbital and
- (2) The orbitals that remain the same are orthonormal and the orbital which changes is orthogonal to the one which does not.

Actually we can check with little difficulty on the two electron-jump case at some later point. We shall consider Eq. (11b) later; we now write

the diagonal element as:

$$\frac{2(Z_a + Z_b - 1)}{(Z_a + Z_b)} \left[\frac{Z_a}{r_{ij}} + \frac{Z_b}{r_{ij}} \right] - 2 \sum_{\alpha} \sum_{\beta} \int \frac{|\varphi_{n\alpha}^*(r_i)|^2}{r_{ij}} d^3r_i = U_{nn} \quad (17)$$

We consider Eq. (17). If for the $\varphi_{n\alpha}$ we take the un-perturbed molecular orbital, Eq. (17) will yield, in essence, the Coulomb potential produced by the isolated molecule. It has been shown recently¹, in the case of two oxygen molecules that this potential is particularly unrealistic. In order to get the proper repulsive form for the potential it was necessary to introduce the reciprocal effect of the other molecule on this potential. The Coulomb potential alone yielded a nice attractive potential. Therefore, there does not appear to be any point in taking the $\varphi_{n\alpha}$ as unperturbed orbitals. On the other hand we do not wish to go back beyond the Born beginning to the coupled set through which polarization can be introduced automatically. Therefore we may sensibly begin with molecular orbitals already distorted by the presence of the free electron. This we have done recently in the case of molecule-molecule collisions.¹

On the other hand we might take the type potential used by Fisk² in his treatment of the scattering of slow electrons. If "d" is the internuclear separation in the molecule and "r₁", and "r₂" the free electron separation from the two nuclei then the following coordinates may be adopted:

$$\xi = \frac{r_1 + r_2}{d} \quad \eta = \frac{r_1 - r_2}{d}$$

Then the potential due to the nuclei may be written:

$$\Phi_c = 2Z \left(\frac{1}{r_1} + \frac{1}{r_2} \right) = \frac{8Z}{d} \frac{r^2}{r^2 - \gamma^2} \quad (18)$$

In order for this to approximately represent the field of the molecule, it must go to zero at some boundary, say, $r = r_0 = \text{constant}$. Fisk selects functions $f(r)$ to multiply Φ_c and produce a potential function which meets the following criteria

$$\begin{aligned} f(1) &= 1, & f'(r_0) &\leq 0 \\ f(r_0) &= 0, & f'(1) &\leq 0 \end{aligned} \quad (19)$$

Of several forms for $f(r)$ considered Fisk settled on:

$$f(r) = 1 - \left[\frac{r_0}{(r_0 - 1)^2} \right] \left[\frac{(r_0 - 1)^2}{r} \right] \quad (20)$$

Stier³ had previously appealed to the following form for this function

$$f(r) = \left[\frac{(r_0 - r)}{(r_0 - 1)} \right]^2$$

At any rate a value for r_0 is chosen, and, after the evaluation of the integral over the charge density, a factor is determined to correct the atomic number in Eq. (18). This essentially completes Fisk's evaluation of his potential. He then proceeds to evaluate the elastic scattering cross section for a number of molecules including oxygen and nitrogen. This evaluation itself is, of course, available and could be extended to the energies of our interest. On the one hand it has the advantage over the Born

approximation of yielding precise results for the scattering while on the other hand it has the disadvantage which arises from the approximate nature of its potential.

We could develop a Coulomb potential for the isolated molecule which would be quite general. However, it becomes somewhat more complicated when the polarization of the molecular charge cloud by the free electron is included. In this case one must first obtain expressions for the molecular orbitals distorted by the free electron. This can, of course, be accomplished for the general case on a digital computer. Since we do not intend to appeal to computing machinery at this stage we restrict the position of the free electron to either the internuclear axis of the molecule or the plane perpendicular to this axis and containing the center-of-mass of the molecule. For the present we shall restrict ourselves to the latter path.

In arriving at our potential we first determine the form for the perturbed MO's. We begin with the on-axis case. The form for the distorted molecular orbital is as follows:^{1, 4}

$$\chi_i = [1 + a_i r_i \cos \psi_i] \chi_i^{(0)} \quad (21a)$$

where $\chi_i^{(0)}$ is the undistorted molecular orbital and "a_i" is a parameter. The angle ψ_i is referred to the center-of-mass of the molecule. The parameter a_i is given by:

$$a_i = \frac{(\chi_j^{(0)} | H' | \chi_i^{(0)})}{(\chi_j^{(0)} | \cos \vartheta_i | \chi_i^{(0)}) (E_i^0 - E_j^0)} \quad (21b)$$

where:

$$H' = -\frac{1}{R} + \frac{1}{r} \quad (21c)$$

The matrix elements of $\cos \vartheta_i$ have already been evaluated.⁴ The evaluation of the matrix elements of H' remain to be carried out. Having obtained this distortion we shall now apply it to each of the two atomic orbitals in the molecule. This will yield the potential in which our free electron exists. This potential in turn is to be used in Eq. (16) to obtain the scattering cross section.

In evaluating these matrix elements we first specifically restrict ourselves to oxygen. Next we note the MO's whose distortion we consider;⁵

$$2\sigma_g = .5641\sigma_g 2s + .1677\sigma_g 2p \quad (22a)$$

$$3\sigma_g = -.3117\sigma_g 2s + .6148\sigma_g 2p \quad (22b)$$

$$2\sigma_u = .7500\sigma_u 2s - .1617\sigma_u 2p \quad (22c)$$

$$3\sigma_u = .5345\sigma_u 2s + .9275\sigma_u 2p \quad (22d)$$

The matrix elements between pairs of these orbitals will then consist of obvious linear combinations of matrix elements over the orbitals $\sigma_g 2s$ and $\sigma_g 2p$. We remark that there can be no difference between such grade

and ungerade matrix elements. The reason for this may be explained as follows.

In the on-axis case one can readily demonstrate that the contribution to the matrix element from the nether nucleus is negligible. This means in essence:

$$\sigma_{\delta} 2s = A(e^{-\delta r_a} + e^{-\delta r_b}) = Ae^{-\delta r_b}, \text{ etc.}$$

so that our integrals simplify considerably. We shall follow through the evaluation of the first of these matrix elements. The general expression will be:

$$(\sigma_{\delta} 2s | H' | \sigma_{\delta} 2s) = \frac{\delta^5}{3\pi} \iiint \frac{r_a^2 e^{-2\delta r}}{r_{ij}} r^2 \sin \nu \, dr \, d\nu \, d\varphi = \frac{1}{R} \quad (23)$$

wherein the r_{ij} represents the separation of the electron from the element of the molecular charge cloud under consideration. The "r" represents the MO coordinate referred to the nucleus closest to the perturbing proton. The effective nuclear charge has a value of 2.275. The familiar Legendre expansion reduces as follows for this case:

$$\frac{1}{r_{ij}} = \sum \frac{(k-1m)!}{(k+1m)!} \frac{r_<^k}{r_>^{k+1}} P_k^{1m}(\cos \nu) \exp [im(\varphi - \Phi)] = \sum \frac{r_<^k}{r_>^{k+1}} P_k^0(\cos \nu) \quad (24)$$

For the $\sigma_{\delta} 2s$ case all terms in Eq. (24) save the first ($k = 0$) will disappear during angular integration so that there results:

$$(\sigma_{\delta} 2s | H' | \sigma_{\delta} 2s) = \frac{4\delta^5}{3} \int_0^{\infty} e^{-2\delta R} \left[\frac{R^2}{(2\delta)^2} + \frac{6R}{(2\delta)^3} + \frac{18}{(2\delta)^4} + \frac{24}{(2\delta)^5 R} \right] \delta R \, dR \quad (25)$$

where R is the separation of the electron from the near nucleus .

In like manner we obtain:

$$\begin{aligned}
 (0_2 2s | H' | 0_2 2s) &= \frac{4}{3\sqrt{3}} \delta^5 \left\{ -e^{-2\delta R} \left[\frac{3R^2}{(2\delta)^2} + \frac{18R}{(2\delta)^3} + \frac{60}{(2\delta)^4} \right. \right. \\
 &\quad \left. \left. + \frac{120}{(2\delta)^5 R} + \frac{120}{(2\delta)^6 R^2} \right] + \frac{120}{(2\delta)^6 R^2} \right\} \quad (26)
 \end{aligned}$$

$$\begin{aligned}
 (0_2 2p | H' | 0_2 2p) &= \frac{8}{15} \delta^5 \left\{ -e^{-2\delta R} \left[\frac{15R^2}{2(2\delta)^2} + \frac{90R}{2(2\delta)^3} + \frac{165}{(2\delta)^4} \right. \right. \\
 &\quad \left. \left. + \frac{420}{(2\delta)^5 R} + \frac{720}{(2\delta)^6 R^2} + \frac{720}{(2\delta)^7 R^3} \right] + \frac{720}{(2\delta)^7 R^3} \right\} \quad (27)
 \end{aligned}$$

The matrix elements of $\cos \nu$ have been obtained by us previously for oxygen.⁴ Hence all information is now available for the evaluation of the a_i in Eq. (21b) and hence the distorted orbitals. In Figure 1 we indicate the a_i of interest as a function of electron separation from the near nucleus. The perturbation scheme appealed to in distortion coefficient calculation may be expected to break down at some minimum proton separation.

We are now in a position to evaluate the U_{nn} of Eq. (15) for the ground electronic state of the oxygen molecule. We now make use of the appropriate distorted orbitals in Eq. (17) to obtain:

$$\begin{aligned}
 U_{nn} &= V^{(0)} + 2a \left\{ [.86842 a_{2\sigma_g} + .13084 a_{3\sigma_g} + .88962 a_{2\sigma_u}] V_{2s}^{(0)} \right. \\
 &\quad \left. + [.13076 a_{2\sigma_g} + .86923 a_{3\sigma_g} + .11090 a_{2\sigma_u}] V_{2p}^{(0)} \right\} \quad (28a)
 \end{aligned}$$

$$V_{2s}^{(0)} = -(19.165R^2 + 9.184R + 4.465 + \frac{1}{R})e^{-5.718R} + \frac{1}{R} \quad (28b)$$

$$V_{2p}^{(0)} = -(3.523R^2 + 4.816R + 3.292 + \frac{1}{R})e^{-4.389R} + \frac{1}{R} \quad (28c)$$

$$V^{(0)} = -(15.322 + \frac{3}{R})e^{-15.322R} - (38.330R^2 + 18.368R + 8.930 + \frac{2}{R})e^{-5.718R} - (14.092R^2 + 19.269R + 13.168 + \frac{4}{R})e^{-4.389R} \quad (28d)$$

In Eq. (28a) "a" is one-half the internuclear separation in the molecule.

We next evaluate Eq. (28a) numerically, the result being displayed as Figure II. We are able to make a reasonable fit to this figure with the much simplified function:

$$U_{nn} = e^{-1.209R} \left(\frac{3.14012}{R^2} - \frac{.46219}{R} + .030832 \right) \quad (28e)$$

We now recall that our objective is the evaluation of Eq. (16) for some pair of electronic states and that U_{nm} not U_{nn} is needed for this evaluation, although the latter will be utilized in Eqs. (45). We now discuss our reason for the U_{nn} evaluation at this point.

The development leading to Eq. (29) illustrates our method of including polarization, a phenomena which is generally of some importance. The effects of polarization cannot be included in our derivation of U_{nm} where n and m correspond to the Schumann-Runge system. However, these effects likewise cannot be included in the corresponding contribution to U_{nn} above

or to the U_{nn} of another problem we discuss below. Even so these latter results are in reasonable agreement with experiment. The inference then is that our U_{nm} is nevertheless reasonably good. We now discuss this in more detail.

We have computed the interaction potential between two aligned oxygen molecules whose charge clouds are undistorted. The result is

$$V = -108.3 r^2 e^{-5r}$$

Vanderslice, Mason and Maisch¹³ on the other hand have obtained the following from experiment:

$$V = 812 e^{-3.565r}$$

When we polarize certain molecular orbitals we compute the following potential:

$$V = 321 e^{-3.26r}$$

Since this is effectively doubled for the off-axis case it is apparent that the improvement through the introduction of polarization is a considerable one. Even so, only the sigma orbitals were polarized; it was not possible to polarize the pi orbitals.

The evaluation of the a_i of Eq. (21b) is carried out between two states which must, of course, be of the same symmetry. What we shall call our basis set consists of eight orbitals, three of one sigma symmetry ($1\sigma_g, 2\sigma_g, 3\sigma_g$), three of the other ($1\sigma_u, 2\sigma_u, 3\sigma_u$) one of one pi symmetry ($1\pi_u$) and one of the other ($1\pi_g$). It is obvious then that we can compute no a_i for the pi while we can compute them for the sigma. Even so our results for $O_2 - O_2$ are as given above,

that is, they are in reasonable agreement with experiment. We shall take this as support for the unpolarized orbitals to be used in computing U_{nm} .

We suppose that the inelastic collision of the electron with the oxygen molecule leads to the excitation of the upper state of the Schumann-Runge system. Hence, we will deal with an upper ${}^3\Sigma_u^-$ state and a lower ${}^3\Sigma_g^-$ state. The basic difference is that what was a π_u^- orbital in the ground state is a π_g^- orbital in the upper state. Our potential is then:

$$U_{nm} = \int \frac{\Psi({}^3\Sigma_u^-) \Psi({}^3\Sigma_g^-)}{r_{ie}} dt_i = \int \frac{1\pi_u / 1\pi_g}{r_{ie}} dt_i$$

The rather extreme reduction from Eq. (17) is due to the one particle nature of the interaction operator. This, coupled with the orbital orthogonality leads to the simple (and necessarily unpolarized) result. The entire molecular orbital (i.e., we do not reflect the contribution from the nether molecular atom) is here considered and only r_{ie}^{-1} is taken in the r_{ie}^{-1} expansion:

$$\begin{aligned} U_{nm} &= (.7631)(.6619) \frac{\delta^5}{\pi} \left\{ \int \frac{r_a^2 e^{-2\delta r_a}}{r_i} r_a^2 dr_a \int d\varphi \int \sin^3 \nu_a d\nu_a - \int \frac{r_b^2 e^{-2\delta r_b}}{r_i} r_b^2 dr_b \right. \\ &\quad \left. \int d\varphi \int \sin^3 \nu_b d\nu_b - \int \frac{r_a e^{-\delta r_a} r_b e^{-\delta r_b}}{r_i(r_i)} \sin \nu_a \sin \nu_b dt + \int \frac{r_b e^{-\delta r_b} r_a e^{-\delta r_a}}{r_i(r_i)} \sin \nu_b \sin \nu_a dt \right\} \\ &= -C \left\{ \left[\frac{R^2}{(2\delta)^2} + \frac{6R}{(2\delta)^3} + \frac{18}{(2\delta)^4} + \frac{24}{(2\delta)^5} R \right] e^{-2\delta R} + \left[\frac{(R+a)^2}{(2\delta)^2} + \frac{6(R+a)}{(2\delta)^3} \right. \right. \\ &\quad \left. \left. + \frac{18}{(2\delta)^4} + \frac{24}{(2\delta)^5} (R+a) \right] e^{-2\delta R} e^{-2\delta a} + \frac{24C}{(2\delta)^5 R} - \frac{24C}{(2\delta)^5 (R+a)} \right\} \end{aligned}$$

$$C = 82.0816$$

this may be plotted and fitted to:

$$U_{nm} = \frac{1.00861}{R} - \frac{1.00816}{R+a} \quad \delta = 2.275 \quad a = 2.282 \quad (29)$$

Eq. (29) is now to be used in the evaluation of Eq. (16) with the result:

$$\sigma_{nm}(\nu) = \frac{k_n}{k_m} \frac{1}{4\pi^2} \left| \int [\exp \{ i(k_n - k_m) \cdot r \}] \left\{ \frac{1.00861}{r_j} - \frac{1.00816}{r_j+a} \right\} dt \right|^2 \quad (30)$$

In the close-collision approximation we suppose that the electron follows a straight-line approach path until it collides head on with the molecule at which time it turns abruptly and proceeds out along its departure path. This approximation is not really very drastic for a close collision.

If we now take angles as indicated in Figure III, the exponent in Eq. (30) becomes

$$k_n r_j \cos \delta - k_m r_j \cos \zeta = -k_n r_j \cos \epsilon - k_m r_j (\cos \epsilon \cos \epsilon - \sin \epsilon \sin \epsilon) \quad (31)$$

Thus the integral in Eq. (30) becomes

$$\iiint e^{i [k_n \sin \epsilon \sin \epsilon - (k_n + k_m \cos \epsilon) \epsilon] r_j} \left(\frac{1.00861}{r_j} - \frac{1.00816}{r_j+a} \right) r_j^2 \sin \epsilon \, dy \, dz \, d\phi \quad (32)$$

Certain obvious analytic integrations can be carried out in Eq. (32), but the lion's share would require numerical integration. It is not our purpose to accomplish this evaluation at this time.

Eqs. (30) and (32) yield the cross section for the excitation of the Schumann-Runge upper state through an electron collision along the molecular

symmetry axis. We now consider the ionization of the oxygen molecule by electron collision.

C. The Born Approximation for Electron Ionization

The only real difference in the Born cross section for ionization is, of course, in U_{nm} . The initial state of the molecule system is still the ${}^3\Sigma_g^-$ ground state while the final state is the free electron and the O_2^+ ion in the state:

$${}^2\Pi_g : \pi_g^+ (\pi_u^-)^2 (\pi_u^+)^2 (\sigma_{3g})^2 (\sigma_{2u})^2 (\sigma_{2g})^2 (\sigma_{1u})^2 (\sigma_{1g})^2 \quad (33)$$

The U_{nm} is here the integral over U of the product of the $1/\pi_g^-$ orbital and the ejected electron wave function. The problem then is first to determine the wave function for the ejected electron.

Bates et al⁶ and others,⁷ have shown that in the case of the oxygen atom, the hydrogenic cross section for photo-ionization serves quite well to describe the oxygen phenomenon. The important point is not the method of ionization but the fact that the hydrogenic free electron wave function serves reasonably well. We shall use it in our evaluation. The orbitals are thus

$$\chi(1/\pi_g^-) = .7631 \sqrt{\frac{5}{\pi}} r e^{-\frac{1}{2}r} \sin \theta e^{i\varphi} \quad (34a)$$

$$\chi(kl) = (-)^{l+1} \frac{2}{\sqrt{1-e^{-2iml}}} \prod_{s=1}^l \sqrt{s^2+n^2} \frac{1}{2\pi (2kr)^{l+1}} \int e^{-\rho} \rho^{\frac{l}{2}} \left(\frac{\rho}{2}\right)^{-in'-l-1} \left(\frac{\rho}{2}\right)^{in'-l-1} d\rho \cdot P_l^{lm}(\cos\theta), \rho = -2ikr, n' = \frac{1}{k} \quad (34b)$$

As usual our operator may be represented by the Legendre expansion:

$$\frac{1}{r_{ij}} = \sum \frac{(k-|m|)!}{(k+|m|)!} \frac{r_k^k}{r_{k+1}^{k+1}} P_k^{lm}(\cos\theta_i) P_k^{lm}(\cos\theta_j) \exp[im(\varphi_i - \varphi_j)] \quad (35)$$

of which we shall only take the first term. This means that, in order for the angular integrations to be non-zero, we must take a pl angular state for the free electron. The contribution to U_{nm} from this may be written down as follows:

$$\int \chi(l\pi_2) \left(\frac{1}{r_{ij}} - \frac{\delta}{R}\right) \chi(kp) dr = .763 \frac{1}{216} \frac{2\sqrt{1+n'^2}}{\sqrt{1-e^{-2iml}}} \frac{1}{2\pi} \frac{1}{4k^2} \frac{\sqrt{2\delta^5}}{\delta\pi^2} \cdot \int \sin^2\theta e^{i\varphi} e^{-i\varphi} d\Omega \int \left(\frac{\rho}{2}\right)^{-in'-2} \left(\frac{\rho}{2}\right)^{in'-2} d\rho \int e^{2ikr} \frac{\rho}{r^2} e^{-\delta r} \left(\frac{1}{r_2} - \frac{\delta}{R}\right) r^2 dr \quad (36)$$

We now integrate over "r":

$$\frac{1}{R} \int_0^R e^{-(\delta-2ik\rho)r} r dr + \int_0^\infty e^{-(\delta-2ik\rho)r} dr - \frac{\delta}{R} \int_0^\infty e^{-(\delta-2ik\rho)r} r dr = \frac{e^{-(\delta-2ik\rho)R}}{R(\delta-2ik\rho)^2} - \frac{1}{R(\delta-2ik\rho)^2}$$

We next integrate over ρ . When we substitute the above the

-integral becomes

$$- \frac{e^{-\delta k}}{R} \int e^{2ik\rho} \left(\frac{\rho}{2}\right)^{-in'-2} \left(\frac{\rho}{2}\right)^{in'-2} (\delta-2ik\rho) d\rho - \frac{1}{R} \int \left(\frac{\rho}{2}\right)^{-in'-2} \left(\frac{\rho}{2}\right)^{in'-2} (\delta-2ik\rho) d\rho$$

We shall evaluate this by integrating around the closed path in the complex plane continuing its pole and extending the path to infinity. The pole is at $\xi = -\frac{1}{2}i\frac{\delta}{k} = -\frac{1}{2}i\delta n'$. The value of such an integral is equivalent to $2\pi i \mathcal{R}$ when \mathcal{R} is the residue at the pole, here of order two. We recall

$$\mathcal{R} = \frac{df}{d\xi} \quad f = e^{2ikR\xi} \left(\xi + \frac{1}{2}\right)^{-in'-2} \left(\xi - \frac{1}{2}\right)^{in'-2}$$

When we make use of the relation:

$$\left(\frac{i\delta n' - 1}{i\delta n' + 1}\right)^{in'} = e^{-2n' \operatorname{ctn}^{-1} \delta n'}$$

The derivative may be evaluated as:

$$\left. \frac{df}{d\xi} \right|_{\xi = -\frac{1}{2}i\delta n'} = -\frac{2^6 \pi k}{(\delta n'^2 + 1)^2} e^{-2n' \operatorname{ctn}^{-1} \delta n'} - \frac{2^7 \pi i (in' - 2)}{R(\delta n'^2 + 1)} e^{-2n' \operatorname{ctn}^{-1} \delta n'}$$

and when this is substituted in Eq. (36) the result is:

$$U_{nm} = \frac{.7631 \sqrt{\delta^5} 2^4}{3k} \frac{e^{-2n' \operatorname{ctn}^{-1} \delta n'}}{(\delta n'^2 + 1)^2} - \frac{.7631 \sqrt{\delta^5} 2^5 (in' - 2)}{3k^2} \frac{e^{-2n' \operatorname{ctn}^{-1} \delta n'}}{(\delta n'^2 + 1)^3} \quad (37)$$

We carry this no further but simply remark that one gets an equation similar to Eq. (32) and requiring numerical integration.

This ends our consideration of electron scattering for the present. We remark that we have failed to consider (1) elastic scattering, (2) dissociation and (3) the existence of rotational and vibrational motion. Under the Born approximation in use here, these pose no problems which are essentially different from those treated above and can be dealt with in this

general fashion at a later stage. However, it does appear that the excitation of electronic upper states and the ionization of the molecule are phenomena to be treated first in our scattering consideration.

There is a portion of the electronic excitation scattering which we have deliberately ignored and which certainly can be of definite importance. In the above sections we have considered what is generally referred to as the direct scattering. Now let us consider Eq. (16). We remark that in the exponential " r_j " is associated both with the incoming momentum vector and the outgoing momentum vector. This in essence describes the direct scattering. To this must be added a consideration of the exchange scattering, and, if we again refer to Eq. (16), our inbound electron will be associated only with the incoming momentum vector while some one of the molecular electrons will be associated with the outgoing momentum vector. This develops naturally from the fact that an exchange is made in the course of the collision. The treatment of this is certainly of some importance and must be presumed to be carried out in conjunction with and subsequent to the treatment of direct scattering.

SECTION III

PROTON SCATTERING BY DIATOMICS

A. Distorted Wave Treatment of Proton - Oxygen Scattering

This scattering problem cannot be treated by the Born approximation, for the velocity of a 50 KeV proton is a great deal less than that of the outer orbitals in the scattering molecules. Hence, we shall first appeal to what might be called a distorted wave approximation. * For the wave function for our molecule-proton system, we refer to Eq. (7) with the exchange effects dropped therefrom:

$$\underline{\Psi} = \sum_n \underline{\Phi}_{An}(\underline{r}_i) \psi_n(\underline{r}) \quad (38)$$

where now \underline{r} is the inter-particle coordinate. Eq. (9) becomes

$$\left[\nabla_p^2 + k_n^2 - 2 \left(\frac{Z_a}{r_{ap}} + \frac{Z_b}{r_{bp}} - \sum \frac{1}{r_{ip}} \right) \right] \sum \Phi_{An}(\underline{r}_i) \psi_n(\underline{r}) = 0 \quad (39)$$

We now multiply through on the left by $\underline{\Phi}_{An}(\underline{r}_i)$ and integrate over the space of the molecular electrons. The result is:

$$\left\{ \nabla_p^2 + k_n^2 - U_{nn} \right\} \psi_{npj} = \sum_{m \neq n} U_{nm} \psi_{mpj} \quad (40)$$

* In general, the following derivation is the standard one, and the reader is referred to the textual references.

This is the familiar infinite set of coupled equations. We propose to simplify the equation by eliminating those U_{nm} which are of little consequence in the oxygen case.

We begin by eliminating spin flip U_{nm} . Since the Hamiltonian is linear in spin and Coulomb parts, two collisions are necessary for changes in both. Further the proton spin-electron spin interaction may well be small due to the mass relations in S. S. Therefore we need consider only triplet states. Further the symmetry of the integral must be the same as that involved in the dipole selection rules since the operator symmetry will be the same. These considerations eliminate a number of levels and, if we consider the first ten or so known levels, we find combination only with the upper state of the Schumann-Runge system. This is, of course, the level we considered for the electron scattering case.

We shall also include the continuum. Eq. (40) now becomes:

$$\{\nabla^2 + k_1^2 - U_1\} \psi_{1p} = U_{12} \psi_{2p} + \int U_{1c} dk \cdot \psi_{cp} \quad (41a)$$

$$\{\nabla^2 + k_2^2 - U_2\} \psi_{2p} = U_{21} \psi_{1p} + \int U_{2c} dk \cdot \psi_{cp} \quad (41b)$$

$$\{\nabla^2 + k_c^2 - U_c\} \psi_{cp} = \int U_{c1} dk \cdot \psi_{1p} + \int U_{c2} dk \cdot \psi_{2p} \quad (41c)$$

We shall treat these equations in cylindrical coordinates for which

$$\nabla^2 = \frac{1}{\rho} \frac{\partial}{\partial \rho} \rho \frac{\partial}{\partial \rho} + \frac{1}{\rho^2} \frac{\partial^2}{\partial \phi^2} + \frac{\partial^2}{\partial z^2} \quad (42)$$

We now consider the reduction of the wave functions to partial waves. As, for example, Geltman⁸ has shown, one may choose a form for ψ on the basis that, in:

$$[\nabla^2 + k^2 - V]\psi = 0$$

as $V \rightarrow 0, \psi \rightarrow e^{i\mathbf{k} \cdot \mathbf{r}}$, a plane wave. We therefore choose the following form for:

$$\psi_1 = e^{i\mathbf{k} \cdot \mathbf{r}} + \chi_1 \tag{43a}$$

$$\psi_2 = \chi_2 \tag{43b}$$

$$\psi_3 = \chi_3 \tag{43c}$$

$$\chi_i = \sum_{\lambda=-\infty}^{+\infty} i^\lambda e^{-i\lambda\beta} F_{\lambda i}(\rho, z) e^{i\lambda\varphi} \tag{43d}$$

$$e^{i\mathbf{k} \cdot \mathbf{r}} = e^{ikz \cos \alpha} \sum_{\lambda} i^\lambda e^{-i\lambda\beta} J_\lambda(k\rho \sin \alpha) e^{i\lambda\varphi} \tag{43e}$$

wherein α is the angle between \mathbf{k} and the molecular symmetry axis and β is the azimuthal angle of \mathbf{k} .

The number of protons in the scattered wave crossing an element of area dS per second is:

$$v |\chi_i|^2 dS = v |\chi_i|^2 r^2 d\Omega$$

from which we can get our cross section as

$$\sigma = r^2 |\chi_i|^2 \tag{44}$$

We now introduce an approximation which leaves a great deal to be desired but with which we do so little that it is of little moment anyway. We suppose the wave function

independent of the z -coordinate. Eqs. (43d) and (42) are substituted into Eqs. (41) with the result:

$$\left\{ \frac{1}{\rho} \frac{\partial}{\partial \rho} \rho \frac{\partial}{\partial \rho} + k_1^2 - \left(\frac{\lambda^2}{\rho^2} + U_{11} \right) \right\} F_{\lambda 1} = U_{12} F_{\lambda 2} + \int U_{1c} dk \cdot F_{\lambda c} \quad (45a)$$

$$\left\{ \frac{1}{\rho} \frac{\partial}{\partial \rho} \rho \frac{\partial}{\partial \rho} + k_2^2 - \left(\frac{\lambda^2}{\rho^2} + U_{22} \right) \right\} F_{\lambda 2} = U_{21} F_{\lambda 1} + \int U_{2c} dk \cdot F_{\lambda c} \quad (45b)$$

$$\left\{ \frac{1}{\rho} \frac{\partial}{\partial \rho} \rho \frac{\partial}{\partial \rho} + k_c^2 - \left(\frac{\lambda^2}{\rho^2} + U_{cc} \right) \right\} F_{\lambda c} = \int U_{1c} dk \cdot F_{\lambda 1} + \int U_{2c} dk \cdot F_{\lambda 2} \quad (45c)$$

These wave functions will in each case have an asymptotic form corresponding to a cylindrical wave. An iterative method of solution is contemplated. The U_{11} and U_{22} will be given by Eq. (28e), the U_{12} by Eq. (29) and the U_{1c} by Eq. (37).

B. The Two State Impact Parameter Treatment

The central assumption in the impact parameter treatment is that of the classical path. The establishment of the validity of the classical path essentially assures us that we may deal with the projectile-target separation as a classical parameter. This validity is assured if the diffusion of the quantal wave packet representing the proton is such that during proton passage through the interaction region a separation is definable. In the present case, the applicability of the classical path is indeed assured.

Let us begin by writing out the Hamiltonian for our proton-oxygen system

$$H = \frac{\hbar^2}{r_{ab}^2} - \sum_{i=1}^{2Z} \left(\frac{1}{r_{ai}} + \frac{1}{r_{bi}} \right) + \sum_{i>j=1}^{2Z-1} \frac{1}{r_{ij}} + \frac{Z_a}{r_{ap}} + \frac{Z_b}{r_{bp}} - \sum_{i=1}^{2Z} \frac{1}{r_{pi}} \quad (46)$$

where all symbols are the familiar standard ones and we shall work in atomic units.

We shall take ψ_i as the wave function for the initial proton-molecule system. By initial, we mean the system which existed at an infinite time in the past. In the same way ψ_f is the final wave function for our system. In this case the system which will exist at an infinite time in the future. The solution to the Schroedinger equation will assuredly be

$$\psi_i = \varphi_i(r_m) \frac{1}{r} \exp[-i\mathbf{k}\cdot\mathbf{r} - iE_r t] \quad (47a)$$

$$\psi_f = \varphi_f(r_m) \frac{1}{r} \exp[i\mathbf{k}'\cdot\mathbf{r} - iE_r' t] \quad (47b)$$

The φ_i refer to the molecular wave functions for oxygen. We shall continue to use the single determinant LCAO-MO form for the oxygen wave functions as computed by Kotani et al.⁵ The k's in the equations correspond to the proton momentum before and after the collision. The 'r' is the radial coordinate of the proton with respect to the center of mass of the oxygen molecule; the energy indicated is the translational energy of the proton.

Eqs. (47) may be used to obtain the following approximate time-dependent system equation

$$\Psi(r, t) = a_i(t) \psi_i \exp(-iE_i t) + b_f(t) \psi_f \exp(-iE_f t) \quad (48)$$

In this equation the a_i and b_i are the state growth coefficients corresponding to the initial and final state of the system. The complete determination of these coefficients constitutes the two state impact parameter treatment. The treatment of this problem to which we shall appeal was, for all practical purposes, introduced by Gurnee and Magee.⁹ It has been discussed in connection with certain inelastic and charge transfer collisions by Bates¹⁰ and somewhat improved upon by McCarroll.¹¹

It is surely true that after an infinite time, the system will be in the final state. Therefore by the definition of the state growth coefficient the probability that the inelastic collision in question will have taken place is given by the square of this coefficient after such a time. Our objective in the calculation then is to determine this coefficient. When we substitute Eq. (48) into the Schroedinger equation, those terms present at infinite separation will cause a number of terms in the equation to drop out. The result is

$$\begin{aligned} & a_i \exp(-iE_i t) H' \psi_i + b_f \exp(-iE_f t) H' \psi_f \\ & = i \dot{a}_i \psi_i \exp(-iE_i t) + i \dot{b}_f \psi_f \exp(-iE_f t) \end{aligned} \quad (49)$$

wherein the H' has been taken as representing those terms in Eq. (46) relating to proton-molecule interaction.

We are now going to make the rather trivial transformation:

$$\begin{aligned} A_i &= a_i \exp[-i\mathbf{k} \cdot \mathbf{r}] \\ B_i &= b_i \exp[i\mathbf{k}' \cdot \mathbf{r}] \end{aligned}$$

with the obvious effect on Eq. (49).

We multiply Eq. (49) through on the left first by

$$\varphi_i^*(r_m) \exp[iE_i t] \exp[iE_i t] \text{ and then by } \varphi_f^*(r_m) \exp[iE_f t] \exp[iE_f t]$$

to obtain

$$a_i H'_{ii} \exp[-i\mathbf{k} \cdot \mathbf{r}] + b_f H'_{if} \exp[i\mathbf{k}' \cdot \mathbf{r}] = i a_i \exp[-i\mathbf{k} \cdot \mathbf{r}] \quad (50a)$$

$$a_i H'_{if} \exp[-i\mathbf{k} \cdot \mathbf{r}] + b_f H'_{ff} \exp[i\mathbf{k}' \cdot \mathbf{r}] = i b_f \exp[i\mathbf{k}' \cdot \mathbf{r}] \quad (50b)$$

where we define the matrix elements in Eqs. (50) as follows:

$$H'_{ii} = \int \varphi_i^* H' \varphi_i d\tau \quad H'_{ff} = \int \varphi_f^* H' \varphi_f d\tau$$

$$H'_{ij} = H'_{ji} = \int \varphi_i^* H' \varphi_j d\tau \quad (51)$$

Eqs. (50) may surely be added with the obvious result

$$i a_i + i b_f = (a_i + b_f) (H'_{ii} + H'_{if}) \quad (52) *$$

We now introduce the specifics of the impact parameter treatment.

Let us consider Figure IV. The proton is colliding with the molecule,

*Here $H'_{ii} = H'_{ff}$ leads to this result for , say, oxygen.

follows a straight-line path whose minimum separation is the impact parameter f . This straight-line path, for our purposes of consideration here lies entirely in the plane which is perpendicular to the molecular symmetry axis and which contains the molecular center of symmetry. Under these classical path conditions the time may be simply and linearly related to the distance along the path through the velocity.

When we make this approximation Eq. (52) leads to

$$v \frac{d(A_i + B_f)}{dx} = -i(H'_{ii} + H'_{if}) \quad (53)$$

which the reader will immediately see to be equivalent to:

$$\int_1^{(A_i + B_f)} d[\ln(A_i + B_f)] = -i \int_{-\infty}^x \frac{1}{v} [H'_{ii} + H'_{if}] dx \quad (54)$$

of solution:

$$A_i + B_f = e^{-iQ} \quad Q = \int_{-\infty}^x \frac{1}{v} (H'_{ii} + H'_{if}) dx \quad (55)$$

The normality requirement on the wave function of Eq. (48) will lead to the following relationship between the state growth coefficients:

$$(\underline{\Psi}, \underline{\Psi}^*) = a_i^* a_i + b_f^* b_f = 1 \quad (56)$$

The two state growth coefficients will at most be complex quantities so that under any condition we can replace them as follows:

$$A_i = \rho e^{i\omega} e^{-ik \cdot r} \quad B_f = \rho' e^{i\omega'} e^{ik' \cdot r} \quad (57)$$

From Eqs. (56) and (57) we obtain the following relationship between the amplitudes of the two state growth coefficients

$$\rho^2 + |\rho'|^2 = 1 \quad (58)$$

which leaves the phase relationship to be determined. Eq. (57) as modified by Eq. (58) may now be substituted in Eqs. (50) and we obtain the following:

$$\rho e^{i\omega} H'_{ii} + \sqrt{1-\rho^2} e^{i\omega'} H'_{if} \exp[i(k+k') \cdot r] = -\dot{\omega} \rho e^{i\omega} + i\rho' e^{i\omega} \quad (59a)$$

$$\begin{aligned} \rho e^{i\omega} H'_{if} \exp[-i(k+k') \cdot r] + \sqrt{1-\rho^2} e^{i\omega'} H'_{ff} &= \\ = -\dot{\omega}' \sqrt{1-\rho^2} e^{i\omega'} + i\sqrt{1-\rho^2} e^{i\omega'} & \end{aligned} \quad (59b)$$

Eqs. (59) provide us with four equations, two real and two imaginary. The imaginary equation arising from Eq. (59) allows us but two choices

$$i\rho' e^{i\omega} = 0 \quad \text{or} \quad b_+ = i\sqrt{1-\rho^2} e^{i\omega}$$

The solution requiring the time derivative of the initial state amplitude to be zero would mean that no transition could be induced by the collision.

This forces our acceptance of the second equation which yields the final state growth coefficient. Our state growth coefficients are now:

$$a_i = \rho e^{i\omega} \quad b_f = i\sqrt{1-\rho^2} e^{i\omega} \quad (60)$$

We consider the imaginary equation from Eq. (59) as modified by

Eq. (60):

$$\dot{r} = \sqrt{1-p^2} H'_{if} \exp [i(\underline{k} + \underline{k}') \cdot \underline{r}]$$

Again appealing to the classical path relationship between distance and time, we may obtain the following equation from this:

$$\int_{-x}^{+x} \frac{dp}{\sqrt{1-p^2}} = \int_{-\infty}^{\infty} \frac{1}{v} H'_{if} \exp [i(\underline{k} + \underline{k}') \cdot \underline{r}] dx \quad (61)$$

From this equation one may obtain the following expression for the final state growth coefficient:

$$|b_f(\infty)|^2 = |\sqrt{1-p_\infty^2}|^2 = \left| \sin \left[\int_{-\infty}^{+\infty} \frac{1}{v} H'_{if} \exp [i(\underline{k} + \underline{k}') \cdot \underline{r}] dx \right] \right|^2 \quad (62)$$

The probability for transition and hence for the inelastic scattering under consideration here is the square of this state growth coefficient after infinite time. In order to obtain the cross section, we simply average over impact parameter with the following results:

$$\sigma = 2\pi \int_0^{\infty} |b_f(\infty)|^2 p dp \quad (63)$$

The expansion of the sine leads to an expression whose first term Arthurs¹² has shown is equivalent to the Born approximation.

C. The Two State IP Treatment of the Proton Oxygen Case

We write down the interaction Hamiltonian as follows:

$$H' = \frac{\delta}{a} + \frac{\delta}{b} - \sum_{i=1}^{16} \frac{1}{r_i} \quad (64)$$

Our first task is the evaluation of this Hamiltonian over the two molecular states involved in the inelastic collision. This matrix element was defined by our development, and we recall that it involves only the molecular coordinates. A point here is that we should evaluate this matrix element over the states of the molecule as perturbed by the incident proton, but we refer to our discussion in connection with Eq. (28e).

This matrix element is given by Eq. (29) with opposite sign:

$$H'_{if} = - \frac{1.00861}{R} + \frac{1.00816}{R+a} \quad (65)$$

Eq. (65) describes that case where the proton proceeds in a long the molecular figure axis. We shall suppose one-third of the oxygen molecules are so aligned, that is, we shall suppose their internuclear axis are aligned along, say, the z -axis which is taken as the proton path. The other two-thirds of the molecules are aligned one-half along the x -axis and one-half along the y -axis. Let us consider this other two-thirds.

Let us suppose the proton is inbound toward the oxygen molecule in the plane which is perpendicular to the molecular symmetry axis and which contains the molecular center of symmetry. The π orbital is identically zero in this plane with the result that U_{nm} is identically zero. As we move the proton path away from this plane it is apparent that U_{nm} will increase to a maximum at the internuclear axis. But it is also apparent that the majority of collisions will occur more nearly under this

than under the former (perpendicular control plane). Therefore we are simply going to assume the axis collision path to be valid.

Eq. (65) is now substituted into Eq. (63). We suppose that the proton maintains its initial velocity -- which corresponds to the momentum k -- until it reaches a particular point on the trajectory. At this point it instantaneously attains its final velocity -- which corresponds to the momentum k' . The space point selected for the change is $x = 0$. The inner product in the exponential now becomes:

$$\underline{k} \cdot \underline{r} = -kr \cos(\pi - \theta) = kr \left(\frac{x}{r}\right) \quad (66)$$

and the bracketed expression in Eq. (63) becomes

$$\frac{1}{v} \int_{-\infty}^0 H_{if}' \exp[i(k-k')x] dx + \frac{1}{v'} \int_0^{\infty} H_{if}' \exp[-i(k-k')x] dx \quad (67)$$

Eq. (67) may be re-written as:

$$\left(\frac{1}{v} + \frac{1}{v'}\right) \int_0^{\infty} H_{if}' \cos[(k-k')x] dx + i\left(\frac{1}{v} - \frac{1}{v'}\right) \int_0^{\infty} H_{if}' \sin[(k-k')x] dx \quad (68)$$

The right side of Eq. (63) now becomes:

$$\sin^2 \left\{ \left(\frac{1}{v} + \frac{1}{v'}\right) \int_0^{\infty} H_{if}' \cos[(k-k')x] dx \right\} \\ + \sinh^2 \left\{ \left(\frac{1}{v} - \frac{1}{v'}\right) \int_0^{\infty} H_{if}' \sin[(k-k')x] dx \right\} \quad (69)$$

The second term in Eq. (69) becomes important at energies below one thousand electron volts. At one KeV for our case it is about two orders

of magnitude smaller than the first term for all values of the impact parameter. We therefore drop the term.

For one thousand volts and above:

$$\cos [(k - k')x] \doteq 1$$

Thus Eq. (63) becomes:

$$|b_f|^2 = \sin^2 \left[\left(\frac{1}{v} + \frac{1}{v'} \right) \int_0^{\infty} H'_{if} dx \right] \quad (70)$$

which is what would have resulted from an assumption of post and prior momentum equality.

This expression was evaluated on a desk computer for a sufficient number of impact parameters to allow evaluation of Eq. (63) for the Schumann-Runge excitation and for values of the proton translational energy ranging from 1 to 50 KeV. A word on this latter evaluation is in order.

If one traces the behavior of $|b_f|^2$ inward from large ρ one finds that it is a monotonically increasing function until a point a few units from the nucleus is reached. At this point the function passed through a maximum and then oscillates more and more rapidly at lesser separations.

The integration inward is numerical to the first minimum and will be discussed in a moment. From the first minimum, however, we take $|b_f|^2 \doteq \frac{1}{2}$ so that the integral has a value $\frac{1}{4} \rho_{\min}$ for this inner portion.

If one retains the infinite upper limit for the integral in Eq. (63) the form of the interaction will lead to a divergence. This is a formal

difficulty at worst since the upper limit on the interaction will not be infinite; it will be determined by the shielding effect of the other particles. We choose the upper limit in Eq. (63) as the Debye radius. For the present, however, we may not specifically evaluate this since we are working with no particular temperature and density. We choose the limit as fifty units.

In Figure V we illustrate the behavior of the cross section for various energies as a function of impact parameter. The total cross sections which result are given for this energy range in Figure VI.

We show that Eq. (62) reduces to the Born approximation for large v or small H'_{if} . We first make the following approximation:

$$b_f = \sin \left\{ \int \frac{1}{v} H'_{if} \exp [i(\underline{k} + \underline{k}') \cdot \underline{r}] d\underline{r} \right\} \\ \doteq \sin \left\{ \frac{1}{v} \int H'_{if} \exp \left[\frac{i}{v} (E_f - E_i) x \right] dx \right\}$$

which depends for its validity on large v . Next the sum is expanded in a MacLaurin series:

$$b_f = \frac{1}{v} \int H'_{if} \exp \left[\frac{i}{v} (E_f - E_i) x \right] dx \\ - \frac{1}{3! v^3} \left\{ \int H'_{if} \exp \left[\frac{i}{v} (E_f - E_i) x \right] dx \right\}^3 + \dots$$

which reduces to its first term for large v or small H'_{if} . Arthurs¹² has shown that this term is equivalent to the Born approximation.

SECTION IV

THE MULTI STATE IMPACT PARAMETER TREATMENT

We consider the molecule as possessing a large number of bound states. In this development we do not include continuum states for the molecule. We take the following expression as the wave function for the proton-molecule system:

$$\Psi = \sum c_i(t) \varphi_i(r_m) \frac{1}{r} \exp[-ik_i \cdot r - iE_{r_i} t] \quad (71)$$

In this equation each term is an exact solution to the Schroedinger equation for infinite separation of the proton-molecule system. The subscript "i" runs over the states of the molecule. When Eq. (71) is substituted into the Schroedinger equation, all terms save the following drop out:

$$\begin{aligned} & \sum c_i(t) \exp[-iE_i t] H' \varphi_i(r_m) \frac{1}{r} \exp[ik'_m \cdot r - iE'_{r_i} t] \\ & = i \sum \dot{c}_i(t) \exp[-iE_i t] \varphi_i(r_m) \frac{1}{r} \exp[ik'_m \cdot r - iE'_{r_i} t] \end{aligned} \quad (72)$$

We remark here that the k_i is negative for $i = 1$ and positive of various values for all other i . The normality condition for the wave function leads to

$$(\Psi^*, \Psi) = \sum c_i^* c_i = 1 \quad (73)$$

Again we take a general form for the state growth coefficient as follows:

$$c_i = f_i e^{i\omega_i} \quad (74)$$

At minus infinity of time we again suppose the coefficient corresponding to the molecular ground state to be unity. Depending on the transition in which we interest ourselves we then are concerned with the value of the upper state coefficient after an infinite time. This we proceed to determine.

First we make the following rather trivial transformation in Eq. (72):

$$C_i = a_i e^{i k_i \cdot r} \quad (75)$$

We shall now multiply Eq. (72) through on the left, first by $\varphi_1^* \exp[-i(E_1 + E_{r1})t]$ then by $\varphi_2^* \exp[-i(E_2 + E_{r2})t]$ and so on.

The result is:

$$\sum C_i H'_{ii} = i \dot{C}_i \quad (76a)$$

$$\sum C_j H'_{ji} = i \dot{C}_i \quad (76b)$$

$$\sum C_j H'_{ji} = \text{etc.} \quad i \dot{C}_i \quad (76c)$$

We now add Eqs. (76) to obtain:

$$\sum_j \sum_i C_i H'_{ji} = i \sum_i \dot{C}_i \quad (77)$$

Eq. (77) may be rewritten as follows:

$$\sum_i \left[C_i \sum_j H'_{ji} - i \dot{C}_i \right] = 0 \quad (78)$$

There are two ways in which this equation could hold. First the various terms or certain of the various terms could be non-zero while some remain zero under all conditions. This is certainly not true at

minus infinity of time since there the time rate of change of the coefficient is zero while the matrix element of the perturbing Hamiltonian is zero. Hence it follows that Eq. (78) is zero because each term in Eq. (78) is zero. We now introduce the impact parameter treatment and again replace "t" by "x". Under these conditions any term in Eq. (78) may be written as

$$\frac{dC_i}{dx} = -\frac{1}{v} \sum_j H'_{ji} \cdot C_j \quad (79)$$

Eq. (79) has the familiar and immediate solution:

$$C_i = e^{-iQ_i} \quad Q_i = -i \int_{-\infty}^x \frac{1}{v} \sum_j H'_{ji} dx \quad (80)$$

Relationships among the f_i and the ω_i are required now for the completion of the calculation. We begin by forming the column matrix which we term the amplitude matrix and which may be represented as:

$$\underline{a} = (f_i e^{-i\omega_i}) \quad (81)$$

With this definition the Schroedinger equation may be written in matrix form as follows:

$$\underline{H} \underline{a} = i \frac{d}{dt} \underline{a} \quad (82)$$

We multiply this equation through on the right by the associate matrix \underline{a}^+ to obtain:

$$\underline{H} \underline{a} \underline{a}^+ = i \left(\frac{d}{dt} \underline{a} \right) \underline{a}^+ \quad (83)$$

Let us write down the product of the matrix "a" and its associate as follows:

$$\underline{a} \underline{a}^T = \underline{b} \quad (84)$$

This matrix is possessed of no inverse and its adjoint matrix is a null matrix. However, we may form the plinverse matrix which when it multiplies the matrix \underline{b} on the right yields the null matrix. We prove this as follows.

First of all the elements of the plinverse matrix are defined as follows:

$$b_{ii}^{\text{pl}} = \sum_{j \neq i} b_{jj} \quad (85a)$$

$$b_{ij}^{\text{pl}} = -b_{ij} \quad (85b)$$

We now write out a typical element of the $\underline{b} \underline{b}^{\text{pl}}$ product matrix as

$$\begin{aligned} b_{ij} &= a_{i1} a_{j1}^T = \rho_i \rho_j \\ c_{ln} &= (\underline{b} \underline{b}^{\text{pl}})_{ln} = \rho_l \rho_n \sum_{i \neq n} \rho_i^2 - \sum_{i \neq n} \rho_l \rho_i^2 \rho_n \\ &= \rho_l \rho_n [\sum \rho_i^2 - \sum \rho_i^2] \equiv 0 \end{aligned} \quad (86)$$

which demonstrates that multiplication on the right by the plinverse matrix yields the null matrix.

We multiply Eq. (83) through on the right to obtain the following :

$$i \left(\frac{d}{dt} \underline{a} \right) \underline{a}^T \underline{b}^{\text{pl}} = \underline{0} \quad (87)$$

An example we treat the two state case. First of all the coefficient matrix and the product of this matrix with its associate may be written down as:

$$\underline{a} = \begin{pmatrix} \rho_1 e^{-i\omega_1} \\ \rho_2 e^{-i\omega_2} \end{pmatrix} \quad \underline{b} = \underline{a} \underline{a}^\dagger = \begin{pmatrix} \rho_1^2 & \rho_1 \rho_2 \\ \rho_1 \rho_2 & \rho_2^2 \end{pmatrix} \quad (88)$$

We may form $\underline{b}^{\mathcal{Q}}$ through Eqs. (85) as

$$\underline{b}^{\mathcal{Q}} = \begin{pmatrix} \rho_2^2 & -\rho_1 \rho_2 \\ -\rho_1 \rho_2 & \rho_1^2 \end{pmatrix} \quad (89)$$

The matrix manipulation indicated by Eq. (87) next leads to two equations in the amplitudes and phases. First through, the assumption of a constant difference between the two phases leads to the following expression:

$$e^{-i(\omega_1 - \omega_2)} = a + ib = \sqrt{1 - b^2} + ib \quad (90)$$

Eq. (90) now allows us to write down one of the two equations arising from Eq. (87) as:

$$\rho_1 \rho_1 \rho_2^2 [1 - \sqrt{1 - b^2} - ib] + i\omega \rho_1^2 \rho_2^2 [-1 + \sqrt{1 - b^2} + ib] = 0 \quad (91)$$

The solution is:

$$b = 1 \quad (92)$$

The normality condition, of course, leads immediately to the relationship among the amplitudes and we have obtained the same result

which was obtained in the previous section in a somewhat more straightforward, if less generally applicable, manner.

SECTION V

BIBLIOGRAPHICAL REMARKS

We first list those books which are of particular interest to scattering problems in general.

The Theory of Atomic Collision. N. F. Mott and H. S. W. Massey Oxford University Press: 1949.

Theoretical Nuclear Physics. J. M. Blatt and F. B. Weisskopf John Wiley & Sons, New York: 1952

The Theory of Protons and Electrons. M. M. Jauch and F. Rohrlich: Addison-Wesley Publishing Company, Inc., Reading: 1955.

Quantum Theory. I. Elements. Edited by D. R. Bates, Academic Press, Inc. New York: 1961

The Shift and Shape of Spectral Lines. R. G. Breene, Jr., Pergamon Press Inc. London: 1961

Atomic and Molecular Processes. Edited by D. R. Bates, Academic Press. New York: 1962.

Quantum Theory of Scattering. Ta-You Wu and T. Ohmura. Prentiss Hall, Inc. New York: 1962.

For general information we also call the reader's attention to the following review type articles:

"On the Convergence of Born Expansions" W. Kohn Rev. Mod. Phys. 26, 292 (1954).

"Theory of Scattering of Slow Electrons". H. S. W. Massey, Rev. Mod. Phys. 28, 199 (1956).

"Theory of Atomic Collision" H. S. W. Massey, Handbuch der Physik XXXVI, Page 232 (1956).

"Excitation and Ionization of Atoms by Electron Impact". H. S. W. Massey, Handbuch der Physik XXXVI, Page 307 (1956).

Professor Takayanagi has developed a very comprehensive explanatory bibliography of non-nuclear scattering phenomena. It consists of three parts plus a supplement which is dated August 1962. Although No. 1 is no longer available it is understood that the remaining parts may be obtained by writing

Professor Kazuo Takayanagi
Department of Physics
Saitama University
Urawa, Saitama, Japan

The bibliography is broken down as follows:

Collision Theory. Part I

- I. Excitation and Ionization of Atoms by Electron Impact
- II. The collision of Electrons with Molecules
- III. Theory of Inelastic Molecular Collisions

Collision Theory. Part II

- IV. Atom - Ion Atom - Atom Inelastic Collisions
- V. Charge Transfer

Collision Theory. Part III

- VI. Inelastic Collisions of Electrons with Atoms or Molecules (experimental)
- VII. Photoionization and Radiative Recombination
- VIII. Theory of Intermolecular Forces

The supplement lists additional references divided into the same categories.

We now list some additional references which we do not believe appear in the Takayanagi bibliography. No claim to exclusiveness is made for the following listing, and we hope to add to it in future.

- I. EXCITATION AND IONIZATION OF ATOMS BY ELECTRON IMPACT (here to include elastic scattering of electrons)

- 1958F S. E. Frish and V. E. Yakhontova. *Optika i Spektrosk* 4, 402 (1958)
New Data On Excitation Functions of Helium Lines.
- 1959A J. W. Albrecht. *An. Real Soc. Espan. Fis. Quim.* 55, 223 (1959)
Elastic and Inelastic Scattering Cross-Sections.
- 1959B R. G. Breene, Jr. and M. C. Nardone. *Phys. Rev.* 115, 93 (1959)
Wave Functions for the Free Electron. I. The Coulomb Potential.
- 1959H L. G. H. Huxley. *Austral. J. Phys.*, 12, 171 (1959)
The Structure of a Stream of Electrons and Ions Drifting and Diffusing In
A Gas When Ionization By Collision and Molecular Attachment Are Present.
- 1959K B. Kivel. *Phys. Rev.*, 116, 926-7 (1959)
Elastic Scattering of Low-Energy Electrons by Argon.
- 1959K B. Kivel. *Phys. Rev.*, 116, 1484-5 (1959)
Electron Scattering by Noble Gases in the Limit of Zero Energy.
- 1959L B. A. Lippmann, H. Mittleman and K. M. Watson. *Phys. Rev.*, 116, 920-5
(1959)
Scattering of Electrons by Neutral Atoms.
- 1959M M. Morand, M. C. L. Le Gentil and S. Desprez-Rebaud. *C.R Acad. Sci
(Paris)* 249, 2060-1 (1959)
Collisions of an Electron With A Hydrogen Atom.
- 1959P R. Peterkop. *Latv. PSR Zinat. Akad. Vestis.* 10, 91 (1959)
Excitation of the S-Levels of the Hydrogen Atom.
- 1959V V. Veldre, M. Gailitis, R. Damburg and P. Stepin'sh. *Latv. PRS Zinat.
Akad. Vestis.* 5, 73 (1959)
Elastic Scattering of Slow Electrons On Lithium Atoms.
- 1959V V. Veldre and R. Damburg. *Latv. PSR Zinat. Akad. Vestis.* 7, 57 (1959)
On the Choice of Atomic Electron Wave-Functions for Scattering Problems.

- 1960C L. Castillejo, I. C. Percival and M. J. Seaton. Proc. Roy. Soc. A 254, 259 (1960)
On The Theory Of Elastic Collisions Between Electrons and Hydrogen Atoms.
- 1960D B. Dawson. Acta. Cryst. 13, 403 (1960)
Atomic Scattering Factors From Wave Functions Calculated By The Poly-detector Method: Cl, Cl-, S and S-.
- 1960G S. Geltman. Proc. Phys. Soc. 75, 67 (1960)
Electron Detachment From The Negative Hydrogen Ion By Electron Impact.
- 1960H R. P. Hurst. Acta. Cryst. 13, 634 (1960)
Three Term Hylleraas Function Atomic Scattering Factors For The Two Electron Ions.
- 1960K W. Kolos. Bull. Acad. Polon. Sci. Ser. Sci. math. astron. phys 8, 67 (1960)
On Wave Functions For The Problem Of Electron And X-Ray Scattering By Helium Atoms.
- 1960M R. P. McEachran and P. A. Fraser. Canad. J. Phys 38, 317 (1960)
d-Wave Contribution To Electron-Hydrogen Atom Scattering.
- 1960N T. Nagy, L. Nagy and S. Desi. Nuclear Instrumen. and Methods 8, 327 (1960)
Drift Velocities Of Electrons In Argon Nitrogen And Argon Mixtures.
- 1960P P. Peterkop. Latv. PSR Zinat. Akad Vestis, 57 (1960)
Consideration Of Exchange In Ionization.
- 1960R H. van Regemorter. Ann. Astrophys. 23, 817 (1960)
Methods of Calculating Electron Impact Excitation Cross-Sections.
- 1960R L. B. Robinson. Phys. Rev. 117, 1281 (1960)
Elastic Scattering of Low-Energy Electrons By The Thomas-Fermi Atom

- 1960S L. Spruch and L. Rosenberg. Phys. Rev. 117, 143 (1960)
Low-Energy Scattering By A Compound System: Positrons On Atomic Hydrogen.
- 1960S D. P. Stevenson. J. Amer. Chem. Soc. 82, 5961 (1960)
The Franck-Condon Principle And The Ionization And Dissociation Of Hydrogen By Electron Impact
- 1960T T. Tietz. Acta phys. Hungar. 12, 151 (1960)
Mott Polarization In Scattering Of Electrons From Thomas-Fermi and Hartree Atoms.
- 1960T T. Tietz. Ann. Phys. 6, 262 (1960)
Scattering Of Electron Beams From Neutral Atoms For Various Thomas-Fermi Potentials
- 1960T T. Tietz. Zh. eksper. teor. Fiz. 39, 1347 (1960)
Electron Scattering In The Thomas-Fermi Model.
- 1960V L. A. Vainshtein and I. I. Sobel'man. Zh. eksper. teor. Fiz. 39, 767 (1960)
Deduction Of The Radial Equations For Collisions Between Electrons And Atoms
- 1960V L. A. Vainshtein and I. I. Sobel'man. Izv. Akad. Nauk SSSR 24, 943 (1960)
(English trans: Bull. Acad. Sci. USSR 24, 943 (1960))
Methods Of Calculating The Cross-Sections For Excitation Of Atoms By Electrons
- 1960V L. M. Volkova, A. M. Devyatov, A. V. Kuralova. Izv. Akad. Nauk SSR 24, 950 (1960)
(English trans: Bull. Acad. Sci. USSR 24, 952 (1960))
Excitation Cross-Sections For Some Spectrum Lines Of Krypton And Xenon
- 1961A D. C. S. Allison, H. A. J. McIntyre and B. L. Moiseiwitsch. Proc. Phys. Soc. 78, 1169 (1961)
The Elastic Scattering Of Positrons By Atomic Hydrogen And Helium.

- 1961B T. Bhattacharyya. Indian J. Phys. 35, 623 (1961)
Scattering Of Electron By Excited Helium Atom.
- 1961B H. Boersch, J. Geiger and H J Reich. Z. Phys. 161, 296 (1961)
Energy Losses Of 25 keV Electrons In Atomic Hydrogen.
- 1961B R. G. Breene, Jr. Phys. Rev. 123, 1718 (1961)
Wave Functions For The Free Electron. II. The Inclusion of Polarization
And Exchange.
- 1961D H. W. Drawin. Z. Phys. 164, 513 (1961)
Representation By Formula Of The Ionization Cross Section For Electron
Impact
- 1961F R. E. Fox. J. Chem. Phys. 35, 1379 (1961)
Ionization Cross Sections Near Threshold By Electron Impact
- 1961F A. J. Freeman and R. E. Watson. Acta cryst. 14, 231 (1961)
Hartree Fock Atomic Scattering Factors For The Neutral Atom Iron Tran-
sition Series.
- 1961G E. Gerjuoy. Rev. mod. Phys. 33, 544 (1961)
Atomic Scattering Processes.
- 1961H Y. Hara, T. Ohmura and T. Yamanouchi. Progr. theor. Phys. 25, 467.
(1961)
Variational Calculation Of The Scattering Lengths In Electron-Hydrogen
Scattering
- 1961I G. S. Ivanov-Kholodnyi, G. M. Nikol'skii and R. A. Gulyaev. Astron. Zh.
37, 799 (1960)
(English trans: Sov. Astron AJ 4, 754 (1961))
Ionization and Excitation of Hydrogen. I. Elementary Processes For The
Upper Levels
- 1961I G. S. Ivanov-Kholodnyi, G. M. Nikol'skii and R. A. Gulyaev. Astron Zh.
38, 455 (1961). (English trans: Sov. Astron --AJ5, 339 (1961))
Ionization And Excitation Of Hydrogen. II. The Number Of Observed Lines
In A Series.

- 1961J H. S. Johnson and W. Kornegay. *Trasn. Faraday Soc.* 57, 1563 (1961)
Rate and Mechanism of the Thermal Ionization of Xenon.
- 1961K A. E. Kingston and B. G. Skinner. *Proc. Phys. Soc.* 77, 724 (1961)
The Elastic Scattering Of Electrons And Positrons By Hydrogen Atoms
- 1961K W. Kolos and K. Pecul. *Ann. Phys.* 16, 201 (1961)
On The Influence Of Correlation Of The Atomic Electrons On The Elastic Electron Scattering By Helium Atoms.
- 1961L B. A. Lippmann and H. M. Schey. *Phys. Rev.* 121, 1112 (1961)
Optical-Model Analysis Of Low-Energy Electron-Hydrogen Scattering.
- 1961M F. B. Malik and E. Trefftz. *Z. Naturforsch.* 16a, 492 (1961)
The Potential Model Treatment Of The Scattering Of Electrons By Atoms And The Existence Of Negative Ions.
- 1961M F. B. Malik. *Z. Naturforsch.* 16a, 500 (1961)
Elastic Scattering Of Low Energy Positrons By Atoms
- 1961M M. R. C. McDowell and G. Peach. *Phys. Rev.* 121, 1383 (1961)
Ionization Of Lithium By Fast Protons And Electrons
- 1961M S. Meyerson. *J. chem. Phys.* 34, 2046 (1961)
Effect Of Electron Energy On Some Electron-Impact Processes
- 1961M B. L. Moiseiwitsch. *Nature* 192, 545 (1961)
Excitation Of The 2s State Of Atomic Hydrogen By Electron Impact.
- 1961M B. L. Moiseiwitsch. *Proc. Phys. Soc.* 77, 721 (1961)
Elastic Scattering Of Slow Electrons By Helium Atoms
- 1961P R. Peterkop. *Proc. Phys. Soc.* 77, 1220 (1961)
Consideration Of Exchange In Ionization
- 1961P R. K. Peterkop. *Zh. eksper. teor. Fiz.* 41, 1938 (1961)
Ionization Of Hydrogen Atoms By Electrons Taking Account Of Interference

- 1961P A. N. Pilyankevich. Zh. tekhn. Fiz. 31, 224 (1961)
(English trans: Soviet Physics-Technical Physics. 6, 161 (1961)
Calculation Of The Characteristics Of Inelastic Scattering Of Medium-Energy Electrons.
- 1961P L. I. Pivovarov, V. M. Tubaev and M. T. Novikov. Zh. eksper. teor. Fiz. 41, 26 (1961)
Electron Loss And Capture By 200-1500 keV Helium Ions In (Various) Gases
- 1961S A. Salmona. C. R. Acad. Sci. 252, 997 (1961)
Calculation Of The Elastic Scattering Of A Slow Electron By A Neutral Sodium Atom In The Ground State.
- 1961M S. N. Milford. Astrophys. J. 134, 724 (1961)
Approximate Cross-Sections For Inelastic Collisions Of Electrons With Atoms.
- 1961S C. Schwartz. Phys. Rev. 124, 1468 (1961)
Electron Scattering From Hydrogen
- 1961S I. P. Shkarofsky. Proc. Inst. Radio Engrs. 49, 1857 (1961)
Generalized Appleton-Hartree Equation For Any Degree Of Ionization And Application To The Ionosphere
- 1961S K. Smith. Proc. Phys. Soc. 78, 549 (1961)
Partial Wave Theory Of Positron Hydrogen Atom Collisions
- 1961S K. Smith and P. G. Burke. Phys. Rev. 123, 174 (1961)
Effect Of Virtual Excitations On The Elastic Scattering Of Electrons And Positrons By Atomic Hydrogen.
- 1961S J. Stickforth. Z. Phys. 164, 1 (1961)
Theory Of Bremsstrahlung In High-Temperature Plasmas.
- 1961T K. Takayanagi and T. Yonezawa. Rep. Ionosphere Space Res. 15, 51 (1961)
Collision Processes In The Aurorae and Ionization In The Auroral Zone

- 1961T A. Temkin. Phys. Rev. Letters 6, 354 (1961)
Polarization And The Triplet Electron-Hydrogen Scattering Length.
- 1961T A. Temkin. Phys. Rev. 121, 788 (1961)
Application Of The Method Of Polarized Orbitals To The Scattering Of Electrons From Hydrogen
- 1961T A. Temkin. J. math. Phys. 2, 336 (1961)
Absolute Definition Of Phase Shift In The Elastic Scattering Of A Particle From Compound Systems.
- 1961T T. Tietz. Ann. Phys. 7, 258 (1961)
The Effect Of Electron Shells On The Asymmetry Effect Of Scattered Atoms
- 1961T T. Tietz. Ann. Phys. 8, 99 (1961)
Asymptotic Phase Shifts And Differential Cross Sections For The Scattering Of Electrons By Atoms With A Latter Potential.
- 1961W R. T. Van de Walle and C. C. Grosjean. Nuovo Cimento 19, 872 (1961)
On The Theory Of Primary Specific Ionization In Helium
- 1961W A. A. Wyller. Astrophys. J. 134, 718 (1961)
Magneto-Ionic Theory For A Fully Ionized Hydrogen Plasma
- 1962A V. A. Apalin, I. Ye. Kutikov, I. I. Lukashevich, L. A. Mikaelyan, G. V. Smirnov and P. Ye. Spivak. Nuclear Phys. 31, 657 (1962)
Asymmetry In Double Mott Electron Scattering For Energies From 45 To 245 keV
- 1962B O. Bely. C. R. Acad. Sci. 254, 3075 (1962)
Calculation Of Electron-Collision Excitation Cross-Sections Of O VI
- 1962B O. Bely. C. R. Acad. Sci. 254, 3167 (1962)
Calculation Of Some Collision Cross-Sections In Fe XIV

- 1962B R. A. Bonham. J. Chem. Phys. 36, 3260 (1962)
Inelastic Scattering From Atoms At Medium Energies. I Bound States
- 1962B G. O. Brink. Phys. Rev. 127, 1204 (1962)
Ionization Of Alkali Atoms By Electron Bombardment.
- 1962B P. G. Burke and H. M. Schey. Phys. Rev. 126, 147 (1962)
Elastic Scattering Of Low-Energy Electrons By Atomic Hydrogen
- 1962B P. G. Burke and H. M. Schey. Phys. Rev. 126, 163 (1962)
Polarization And Correlation Of Electron Spin In Low-Energy Elastic Electron-Hydrogen Collisions.
- 1962B R. E. Burke and G. H. Smith. Proc. Phys. Soc. 79, 673 (1962)
A New Calculation Of Electron Scattering Cross-Sections And A Theoretical Discussion Of Image Contrast In The Electron Microscope
- 1962B P. G. Burke, V. M. Burke, I. C. Percival and R. McCarroll. Proc. Phys. Soc. 80, 413 (1962)
Electron Scattering By Atomic Hydrogen Of The $1s$ $2s$ or $2p$ State I
- 1962B V. M. Burke and R. McCarroll. Proc. Phys. Soc. 80, 422 (1962)
Electron Scattering By Atomic Hydrogen In The $1s$ $2s$ or $2p$ State. II.
- 1962D M. L. De. Naturwiss 49, 228 (1962)
On The Dependence Of Electron Scattering Cross Sections Of Atoms On Their Atomic Number.
- 1962G E. Gerjuoy and N. A. Krall. Phys. Rev. 127, 2105 (1962)
Dispersion Relations For Electron Scattering From Atomic Helium
- 1962G A. F. Gorsfranova and R. Ya. Damburg. Optika i Spect. 12, 113 (1962)
(English trans: Optics and Spect 12 (1962)
Account Of Strong Coupling In Electron Collisions With Hydrogen Atoms
- 1962H A. E. D. Heylen. Proc. Phys. Soc. 79, 284 (1962)
Calculated Electron Mobility In Nitrogen Oxygen And Air For 0.1
 E/p $100V\text{ cm}^{-1}\text{ mmHg}^{-1}$

- 1962S C. Schwartz. Phys. Rev. 126, 1015 (1962)
Importance Of Angular Correlations Between Atomic Electrons.
- 1962S M. J. Seaton. Proc. Phys. Soc. 79, 1105 (1962)
The Impact Parameter Method For Electron Excitation Of Optically Allowed Atomic Transitions.
- 1962S K. Smith, R. P. McEachran and P. A. Fraser. Phys. Rev. 125, 553 (1962)
Effect Of Virtual Excitation Of The 2s State On The Elastic Scattering Of Electrons By Atomic Hydrogen
- 1962T A. Temkin. Phys. Rev. 126, 130 (1962)
Nonadiabatic Theory Of Electron-Hydrogen Scattering.

II. THE COLLISION OF ELECTRONS WITH MOLECULES

- 1960B D. Beck and O. Osberghaus. Z. Phys. 160, 406 (1960)
The Mass-Spectrum Of Uncharged Fragments From Electron Impact Studies On Molecules.
- 1960L J. C. Lorquet. J. Chim phys. 57, 1078 (1960)
Study Of The Electron-Molecule Interaction: Electronic Transitions Induced By Bombardment
- 1961K K. Kraus. Z. Naturforsch 16a, 1378 (1961)
Determination Of The Critical Electron Energy For The Formation Of Negative Ions By Electron Impact By Means Of A Retarding-Potential-Difference Method
- 1961S D. A. Swick and J. Karle. J. chem. Phys. 35, 2257 (1961)
Pattern Inversion For Coherent Inelastic Electron Scattering By Br₂.
- 1962F S. N. Foner and R. L. Hudson. J. chem. Phys. 36, 2676 (1962)
Ionization And Dissociation Of Hydrogen Peroxide By Electron Impact

1962S H. Schlumbohm. Z. Phys. 166, 192 (1962)

Electron Avalanches In Electronegative Gases: Simultaneous Measurement Of The Shock Ionization And Attachment Coefficients And For Electrons And The Velocities Of Positive And Negative Ions.

III. THEORY OF INELASTIC MOLECULAR COLLISIONS (to include experiment)

1959G H. Gienapp. Z. Naturforsch. 14a, 1084 (1959)

The Total Effective Cross-Section Of The Collision Process Between Potassium Atoms And Chlorine Molecules.

1959N E. E. Nikitin and N. D. Sokolov. J. chem. Phys. 31, 1371 (1959)

Theory Of Thermal Second-Order Decomposition Of Molecules

1959W B. Widom. J. chem. Phys. 31, 1387 (1959)

Mean-First-Passage Times And The Collision Theory Of Bimolecular Reactions

1960C W. H. Cramer and A. B. Marcus. J. chem. Phys. 32, 186 (1960)

Elastic And Inelastic Scattering Of Low-Velocity D^+ and D_2^+ In Deuterium

1961H M. Hoare. Molecular Phys. 4, 465 (1961)

Systems Of Oscillators With Statistical Energy Exchange In Collision

1961I A. P. Irsa and L. Friedman. J. chem. Phys. 34, 330 (1961)

Collision-Induced Dissociation Of HD^+ .

1961K Yu. M. Khirnyi. Zh. tekhn. Fiz. 31, 597 (1961)

(English trans: Soviet Physics-Tech. Phys. 6 427 (1961))

Production Of Negative Hydrogen Ions In The Passage Of Protons Through Gas Targets.

1960O A. I. Osipov and E. V. Stupochenko. Izv. Akad. Nauk SSSR 24, 992 (1960)
(English trans: Bull Acad. Sci. USSR 24, 992 (1960))

Energy Transfer In Molecular Collisions

- 1960P R. L. Platzman. J. Phys. Radium 21, 853 (1960)
Probability Of Ionization By The Transfer Of Energy Of Excited Atoms To Molecules
- 1960R E. M. Reeves, R. W. Nicholls and D. A. Bromley. Proc. Phys. Soc. 76, 217 (1960)
Optical Excitation Of N_2 By 0.5 To 1.5 MeV Hydrogenic Ions
- 1961B R. B. Bernstein. J. chem. Phys. 34, 361 (1961)
Quantum Calculations Of The Velocity Dependence Of The Differential And Total Cross Sections For Elastic Scattering Of Molecular Beams.
- 1961C R. K. Curran. J. chem. Phys. 34, 2007 (1961)
Positive And Negative Ion Formation In $C Cl_3F$.
- 1961H L. Herman, H. I. S. Ferguson and R. W. Nicholls. Canad. J. Phys. 39, 476 (1961)
Excitation Of The First Negative System Of O_2^+ By A Proton Beam In Air And Oxygen.
- 1961H D. R. Herschbach, G. H. Kwei and J. A. Norris. J. chem. Phys. 34, 1842 (1961)
Reactive Scattering In Crossed Molecular Beams. K. Atoms With CH_3I And C_2H_5I
- 1961P L. I. Pivovarov, V. M. Tubaev and M. T. Novikov. Zh. eksper. teor. Fiz. 40, 34 (1961)
Dissociation Of Molecular Hydrogen Ions In Collisions With Gas Molecules
- 1961S B. N. Srivastava and A. K. Barua. Proc. Phys. Soc. 77, 677 (1961)
Rotational-Translational Relaxation Time In H_2 Calculated From Thermal Conductivity.
- 1961U N. G. Utterback and G. H. Miller. Phys. Rev. 124, 1477 (1961)
Ionization Of Nitrogen Molecules By Nitrogen Molecules.

1961W R. E. Walker, L. Monchick, A. A. Westenberg and S. Favin. Planet Space Sci. 3, 221 (1961)

High Temperature Gaseous Diffusion Experiments And Intermolecular Potential Energy Functions

1961W D. J. Wilson, B. Noble and B. Lee. J. chem. Phys. 34, 1392 (1961)

Pressure Dependence Of Fluorescence Spectra

1962F L. Friedman and A. P. Irsa. J. chem. Phys. 36, 1802 (1962)

Collision-Induced Dissociation Of N_2O^+ .

1962H R. Heastie and D. H. Martin. Canad. J. Phys. 40, 122 (1962)

Collision-Induced Absorption of Submillimeter Radiation By Non-Polar Atmospheric Gases

1962L J. C. Light. J. chem. Phys. 36, 1016 (1962)

Dissociation Of Gaseous Diatomic Molecules: Classical Adiabatic Scattering Approach.

1962R A. R. Ruffa and V. Griffing. J. chem. Phys. 36, 1389 (1962)

A Quantum Mechanical Study Of The Hydrogen Bimolecular Exchange Reaction.

1962T G. A. Taylor and G. B. Porter. J. chem. Phys. 34, 1392 (1961)

Kinetics Of Excited Molecules. IV. Unimolecular Dissociation of Ketene.

IV. ATOM-ION, ATOM-ATOM INELASTIC COLLISIONS

1959D C. DeLaet and M. Demeur. Bull. Acad. Roy. Belgique Cl. Sci. 45, 546 (1959)

Note On The Theory Of Atomic Coulomb Excitation

1959F N. V. Fedorenko. Uspekhi fiz. Nauk 68, 481 (1959)
(English trans: Soviet Physics - Uspekhi 2, 526 (1959))

Ionization In Collisions Between Ions And Atoms.

- 1959M N. R. C. McDowell and G. Peach. Proc. Phys. Soc. 47, 463 (1959)
Electron Loss From Fast Negative Ions Of Atomic Hydrogen Passing Through Atomic Hydrogen.
- 1959P H. Pauly. Z. Phys. 157, 54 (1959)
Small-Angle Scattering In Collision Between Neutral Atoms
- 1959S T. J. M. Sluyters and J. Kistemaker. Physica 25, 1389 (1959)
Excitation Mechanism Of Ar⁺ Ions In He, Ne, Ar, Kr And Xe. II.
- 1960B H. Bunke and R. Seiwert. Monat. Deutschen Akad. Wiss. Berlin 2, 723 (1960)
Inelastic Collisions Between Excited And Unexcited Atoms
- 1960F Ya. M. Fogel', V. A. Ankudinov and D. V. Pilipenko. Zh. eksper. teor. Fiz. 38, 26 (1960)
Electron Capture And Loss In Collisions Of Fast Helium, Boron & Fluorine Atoms With Gas Molecules
- 1960F Ya. M. Fogel'. Uspekhi fiz. Nauk. 71, 243 (1960)
(English trans: Soviet Physics-Uspekhi 3 390 (1960)).
The Production Of Negative Ions In Atomic Collisions
- 1960L G. H. Lane and E. Everhart. Phys. Rev. 117, 920 (1960)
Calculations Of Total Cross Sections For Scattering From Coulomb Potentials With Exponential Screening.
- 1960M Yu. P. Mordvinov and O. B. Firsov. Zh. eksper. teor. Fiz. 39, 427 (1960)
Dependence Of The Inelastic Collision Cross-Sections Of Atoms And Ions On Velocity In The Case Of Pseudo-Intersection Of The Levels.
- 1960S G. M. Smith and E. E. Muschlitz, Jr. J. chem. Phys. 33, 1819 (1960)
Scattering Of Metastable Helium Atoms In Helium, Neon And Argon
- 1961B H. W. Berry. Phys. Rev. 121, 1714 (1961)
Energy Distribution Of Electrons From Ionizing Collisions Of Heavy Particles.

- 1961C A. Cassuto. C. R. Acad. Sci. 252, 1311 (1961)
Variations Of Mass Spectra With Temperature Between - 195 and 200°C
- 1961D A. Dalgarno. Planet. Space Sci. 3, 236 (1961)
Transport Properties Of Atomic Hydrogen.
- 1961F F. S. Faizullov and N. N. Sobolev. Optika i Spektrosk 11, 571 (1961)
(English trans: Optics and Spectrosc. 11, 310 (1961))
Determination Of The Effective Cross-Section For Collisions Of Excited Barium Ions With Argon Atoms
- 1961F I. P. Flaks. Zh. tekhn. Fiz. 31, 367 (1961)
(English trans: Soviet Physics-Tech. Physics 6, 263 (1961))
The Ionization Of Inert Gases By Singly-Charged Ions And Atoms Of Neon And Krypton.
- 1961F I. P. Flaks, G. N. Ogurtsov and N. V. Fedorenko. Zh. eksper. teor. Fiz. 41, 1438 (1961)
(English trans: Soviet Physics-JETP)
Ionization In The Collisions Of Ne^{n+} With Xe Atoms And Of Xe^n With Ne Atoms ($n = 0 1 2 3 4$)
- 1961F Ya. M. Fogel', A. G. Koval', Yu. Z. Levchenko and A. F. Khodyachikh. Zh. eksper. teor. Fiz. 39, 548 (1960)
(English trans: Soviet Physics-JETP 12, 384 (1961))
The Composition Of Slow Ions Produced During Ionization Of Gases By Negative Ions
- 1961F Ya. M. Fogel', A. G. Koval' and Yu. G. Levchenko. Zh. eksper. teor. Fiz. 40, 13 (1961)
Formation Of Slow Negative Ions In Single Collisions Between Fast Negative Hydrogen And Oxygen Ions And Gas Molecules.
- 1961H J. B. Hasted. Proc. Phys. Soc. 77, 269 (1961)
The Energy Transferred In Inelastic Ion-Atom Collision
- 1961H K. Hoffmann and R. Seiwert. Ann. Phys. 7, 71 (1961)
Inelastic Collisions Between Excited And Non-Excited Potassium Atoms

- 1961S R. M. St. John and R. G. Fowler. Phys. Rev. 122, 1813 (1961)
New Process Of Excitation Transfer In Helium
- 1961S Z. Sternberg and P. Tomas. Phys. Rev. 124, 810 (1961)
Excitation Processes In Helium Induced By Impact Of Deuterons And Protons.
- 1962B R. J. Bell and B. G. Skinner. Proc. Phys. Soc. 80, 404 (1962)
Excitation Of The 1s-2p Transition In Hydrogen And The 3s=3p Transition
In Na By Nuclei
- 1962B W. R. Bennett, Jr. Ann. Phys. 18, 367 (1962)
Optical Spectra Excited In High Pressure Noble Gases By Alpha Impact.
- 1962B H. W. Berry. Phys. Rev. 127, 1634 (1962)
Energy Distribution Of Electrons From Ionizing Collisions Of Atoms And
Ions.
- 1962B R. A. Buckingham and J. W. Fox. Proc. Roy. Soc. A 267, 102 (1962)
The Coefficient Of Viscosity Of Atomic Hydrogen From 25 to 300^oK
- 1962D I. S. Dmitriev, V. S. Nikolayev, L. N. Fateeva and Ya. A. Teplova. Zh.
eksper. teor. Fiz. 42, 16 (1962)
(English trans: Soviet Physics-JETP)
Experimental Investigation Of Electron Loss By Multicharged Ions In Gases
- 1962K J. G. Kelley, P. E. Steigert and W. W. Watson. Phys. Rev. 126, 2100
(1962)
Scattering Of Oxygen Ions From Argon And Xenon
- 1962L G. J. Lockwood and E. Everhart. Phys. Rev. 125, 567 (1962)
Resonant Electron Capture In Violent Proton-Hydrogen Atom Collisions
- 1962R E. W. Rothe, L. L. Marino, R. H. Neynaber, P. K. Rol and S. M.
Trujillo. Phys. Rev. 126, 598 (1962)
Scattering Of Thermal Rare Gas Beams By Argon Influence Of The Long-
Range Dispersion Forces.

1962S W. P. Sholette and E. E. Muschlitz, Jr. J. chem. Phys. 36, 3368 (1962)

Ionizing Collisions Of Metastable Helium Atoms In Gases

1962S B. M. Smirnov. Zh. eksper. teor. Fiz. 43, 112 (1962)
(English trans: Soviet Physics-JETP)

Inelastic Collisions Of Light Atoms In The Adiabatic Approximation

1962S E. S. Solov'ev, R. N. Il'in, V. A. Oparin and N. V. Fedorenko. Zh. eksper. teor. Fiz. 42, 659 (1962)
(English trans: Soviet Physics - JETP)

Ionization Of Gases By Fast Hydrogen Atoms And Protons

1962T P. Toscheh. F. Naturforsch. 17a, 356 (1962)

Reference Cross Section For The Scattering Of Neutral Molecules

1962Y K. S. Yun and E. A. Mason. Phys. of Fluids 5, 380 (1962)

Collision Integrals For The Transport Properties Of Dissociating Air At High Temperatures.

V. CHARGE TRANSFER.

1958K R. M. Kushnir. Ukrayin. fiz. Zh. 3, 788 (1958)

Charge Exchange Between Caesium Ions And Atoms.

1959S T. J. M. Sluyters, E. de Haas and J. Kistemaker. Physica 25, 1376 (1959)

Charge Exchange Ionization And Electron Loss Cross-Sections In The Energy Range 5 to 24 keV I.

1960B A. M. Bukhteev and Yu. F. Bydin. Izv. Akad. Nauk. SSSR 24, 964 (1960)
(English trans: Bull. Acad. Sci. USSR 24, 964 (1960))

Resonance Charge Exchange Of Alkali Metal Ions And Atoms

1960C D. V. Chkuasdi, U. D. Nikoleishvili, A. I. Guldamashirli. Izv. Akad. Nauk. SSSR 24, 970 (1960)
(English trans: Bull. Acad. Sci. USSR 24, 970 (1960))

Resonance Charge Exchange Of Positive Alkali Metal Ions

- 1960G E. Gustafsson and E. Lindholm. Ark. Fys. 18, 219 (1960)
Ionization And Dissociation Of H₂ N₂ and CO In Charge Exchange Collisions With Positive Ions.
- 1960K W. G. Kauman. Austral. J. Phys. 13, 193 (1960)
On The Flow And Exchange Diffusion Of Electrons And Negative Ions Formed By Molecular Attachment
- 1960K R. M. Kushnir and I. M. Buchma. Izv. Akad. Nauk. SSSR 24, 986 (1960)
(English trans: Bull. Acad. Sci. USSR 24, 986 (1960))
Further Investigation Of Resonance Charge Exchange Of Positive Cesium Atoms
- 1960S F. Schwirzke. Z. Phys. 157, 510 (1960)
Ionization And Exchange Cross-Sections Of Hydrogen Atoms And Ions Of 9-60 keV In Hydrogen
- 1960S S. Sunakawa. Progr. theor. Phys. 24, 963 (1960)
On The Theory Of Rearrangement Collisions
- 1961D T. B. Day, L. S. Rodberg, G. A. Snow and J. Sucher. Phys. Rev. 123, 1051 (1961)
Note On Rearrangement Collisions
- 1961D T. M. Donahue and F. Hushfar. Phys. Rev. 124, 138 (1961)
Formation Of Negative Ions In A Gas By Charge Transfer From A Fast Atomic Hydrogen Beam.
- 1961F Ya. M. Fogel', V. F. Kozlov and G. N. Polyakova. Zh. eksper. teor. Fiz. 39, 1186 (1960)
(English trans: Soviet Physics -- JETP 12, 826 (1961))
Double Charge Exchange Of Alkali Metal Ions.
- 1961M M. H. Mittleman. Phys. Rev. 122, 1930 (1961)
Formal Theory Of Rearrangement Collisions
- 1961M Yu. E. Murakhver. Zh. eksper. teor. Fiz. 13, 1080 (1961)
(English trans: Soviet Physics -- JETP 13, 762 (1961))
Resonance Charge Exchange In Hydrogen And Sodium

- 1961P I. Popescu. Proc. Phys. Soc. 78, 584 (1961)
On Resonant Charge Transfer In Ionized Dense Gases
- 1961R A. Ramakrishnan, G. Ramachandran and V. Devanathan. Nuovo Cimento 21, 145 (1961)
A Time-Dependent Approach To Rearrangement Collisions.
- 1961S A. Schmid. Z. Phys. 161, 550 (1961)
Dissociation And Charge Exchange Of Hydrogen And Deuterium Molecular Ions By Passing Through Hydrogen
- 1962D H. G. Dehmelt and F. G. Major. Phys. Rev. Letters 8, 213 (1962)
Orientation of $(\text{He })^+$ Ions By Exchange Collisions With Cesium Atoms
- 1962F W. L. Fite, A. C. H. Smith, R. F. Stebbings. Proc. Roy. Soc. A268, 527 (1962)
Charge Transfer In Collisions Involving Symmetric And Asymmetric Resonance
- 1962H J. B. Hasted and A. R. Lee. Proc. Phys. Soc. 79, 702 (1962)
The Near-Adiabatic Criterion In Charge Transfer Collisions.
- 1962H A. Henglein and G. A. Muccini. F. Naturforsch 17a, 452 (1962)
Mass Spectrometric Observation Of Electron And Proton Transfer Reactions Between Positive Ions And Neutral Molecules.
- 1962J B. L. Johnson. J appl. Phys. 33, 739 (1962)
Charge Transfer Scattering In High-Velocity Neutral Particle Production
- 1962K M. V. Kazarnovskii and A. V. Stepanov. Zh. eksper. teor. Fiz. 42, 489 (1962) (English trans: Soviet Physics -- JETP)
Theory Of Resonance Scattering On Atomic Systems I. Deduction of General Formulae.
- 1962M Yu. E. Murakhver. Zh. eksper. teor. Fiz. 42, 1241 (1962) (English trans: Soviet Physics - JETP)
Resonance Charge Exchange In Atomic Hydrogen

- 1962O G. N. Ogurtsev and I. P. Flaks. Zh. eksper. teor. Fiz. 42, 721 (1962)
Charge Exchange Of Xe³⁺ and Xe⁴⁺ In Neon

VI. INELASTIC COLLISIONS OF ELECTRONS WITH ATOMS OR MOLECULES
(experimental)

- 1959H M. E. Haine and A. W. Agar. Brit. J. appl. Phys. 10, 341 (1959)

Differential Scattering Cross-Section Of Atoms To Medium Energy
Electrons At Small Angles.

- 1959P A. V. Phelps and J. L. Pack. Phys. Rev. Letters 3, 340 (1959)

Electron Collision Frequencies In Nitrogen And In The Lower Ionosphere

- 1959P A. N. Prasad. Proc. Phys. Soc. 74, 33 (1959)

Measurement Of Ionization And Attachment Coefficients In Dry Air In
Uniform Fields And The Mechanism Of Breakdown

- 1960F R. E. Fox. J. chem. Phys. 32, 385 (1960)

Threshold Ionization Of HCl By Electron Impact

- 1960K I. U. Kubic and Y. Inuishi. Technol. Rep. Osaka Univ. 10, 565 (1960)

Measurement Of Gaseous Discharge Plasma With Microwave Interfer-
ometer.

- 1960S M. S. Sodha. Proc. Phys. Soc. 76, 439 (1960)

Estimation Of Inelastic Collision Cross Section From Electron Mobility
Measurements.

- 1961B M. A. Biondi. Planet. Space Sci. 3, 104 (1961)

Measurements of Electron Attachment Recombination Elastic And In-
elastic Collisions In Atmospheric Gases.

- 1961C R. K. Curran and R. E. Fox. J. chem. Phys. 34, 1590 (1961)

Mass Spectrometer Investigation Of Ionization Of N₂O By Electron Impact

- 1961D K. T. Dolder, M. F. A. Harrison and P. C. Thonemann. Proc Phys. Soc. 264A, 367 (1961)
A Measurement Of The Ionization Cross Section Of Helium Ions By Electron Impact
- 1961N R. H. Neynaber, L. L. Marino, F. W. Rothe and S. M. Trujillo. Phys. Rev. 123, 148 (1961)
Low-Energy Electron Scattering From Atomic Oxygen
- 1961N R. H. Neynaber, L. L. Marino, E. W. Rothe and S. M. Trujillo. Phys. Rev. 124, 135 (1961)
Scattering Of Low Energy Electrons By Atomic Hydrogen
- 1961N V. S. Nikolaev, L. N. Fateeva, I. S. Dmitriev and Ya. A. Teplova. Zh. eksper. teor. Fiz. 41, 89 (1961)
(English trans: Soviet Physics - JETP)
Capture Of Several Electrons By Fast Multicharged Ions.
- 1961S I. P. Shkarofsky, M. P. Bachynski and T. W. Johnston. Planet. Space Sci. 6, 24 (1961)
Collision Frequency Associated With High Temperature Air And Scattering Cross-Sections Of The Constituents
- 1961Z I. P. Zapisochnir, S. M. Kishko, V. S. Shevera, P. F. Fel'tsan, L. L. Shimon. Ukrayin Fiz. Zh. 6, 770 (1961)
A Spectroscopic Study Of The Excitation Functions Of Atoms And Molecules
- 1962A P. Armbruster. Z. Phys. 166, 341 (1962)
Ionization Of The Inner Electron Shells In The Stopping Of Fission Fragments
- 1962H J. W. Hooper, D. S. Harmer, D. W. Martin and E. W. McDaniel. Phys. Rev. 125, 2000 (1962)
Comparison Of Electron And Proton Ionization Data With The Born Approximation Predictions
- 1962K A. F. Klein. Phys. of Fluids 5, 392 (1962)
Measurement Of Hall Electromagnetic Force In Air.

- 1962M C. E. Melton. J. chem. Phys. 37, 562 (1962)
Ionization And Excitation Processes In Argon Krypton And The C₂ Hydrocarbons Produced By Various Means.
- 1962P J. L. Pach, R. E. Voshall and A. V. Phelps. Phys. Rev. 127, 2084 (1962)
Drift Velocities Of Slow Electrons In Krypton Xenon Deuterium CO CO₂ H₂O N₂O and NH₃
- 1962R E. W. Rothe, L. L. Marino, R. H. Neynaber and S. M. Trujillo. Phys. Rev. 125, 582 (1962)
Electron Impact Ionization Of Atomic Hydrogen And Atomic Oxygen

VII. PHOTOIONIZATION AND RADIATIVE RECOMBINATION

- 1960B D. L. Bunker. J. chem. Phys. 32, 1001 (1960)
Mechanics Of Atomic Recombination Reactions.
- 1960D I. S. Dimitriev, V. S. Nikolaev, L. N. Fateeva, Ya. A. Teplova. Izv. Akad. Nauk, SSSR 24, 1169 (1960)
(English trans: Bull. Acad. Sci. USSR 24, 1169 (1960))
Concerning The Mean Charge Of Ions Passing Through Matter
- 1960K M. V. Kenyukov. Izv. Akad. Nauk, SSSR 24, 989 (1960)
(English trans: Bull. Acad. Sci. USSR 24, 989 (1960))
Indirect Evaluation Of Electron Attachment Cross Section
- 1960S D. R. Sweetman. Proc. Phys. Soc. 76, 998 (1960)
The Lifetime Of The H_e Ion
- 1961B D. R. Bates and A. E. Kingston. Nature 189, 652 (1961)
Recombination Through Electron-Electron Collisions
- 1961B M. S. Bhalla and J. D. Craggs. Proc. Phys. Soc. 78, 438 (1961)
Measurement Of Ionization And Attachment Coefficients In Carbon Monoxide In Uniform Fields

1961D N. D'Angelo. Phys. Rev. 121, 505 (1961)

Recombination Of Ions And Electrons

1961M K. Mork and H. Olsen. Phys Norweg 1, 21 (1961)

Forward Emission Of Photoelectrons At High Energies.

1961N V. S. Nikolaev, I. S. Dmitriev, L. N. Fateeva and Ya. A. Teplova. Zh. eksper. teor. Fiz. 40, 989 (1961)
(English trans: Soviet Physics-JETP 13, 695 (1961))

Experimental Investigation Of Electron Capture By Multiply Charged Ions

1962B D. R. Bates, A. E. Kingston and R. W. P. McWhirter. Proc. Roy Soc. 267, 297 (1962)

Recombination Between Electrons And Atomic Ions. I. Optically Thin Plasmas

1962B D. R. Bates. Planet Space Sci. 9, 77 (1962)

Dielectronic Recombination To Normal Nitrogen And Oxygen Ions

1962B L. M. Biberman, G. E. Norman, K. N. Ul'yanov. Astron. Zh. 39, 105 (1962)
(English trans: Soviet Astron - AJ 6, 77 (1962)

The Photoionization Of Excited Complex Atoms And Ions

1962B S. Byron, R. C. Stabler and P. I. Bortz. Phys. Rev. Letters 8, 376 (1962)

Electron-Ion Recombination By Collisional And Radiative Processes

1962G E. P. Gray and D. E. Kerr. Ann. Phys. 17, 276 (1962)

The Diffusion Equation With A Quadratic Loss Term Applied To Electron-Ion Volume Recombination In A Plasma

1962H J. B. Hasted and A. Y. J. Cheng. Proc. Phys. Soc. 80, 441 (1962)

Electron Capture Processes For Multiply Charged Ions

1962K I. R. King. J. chem. Phy. 37, 74 (1962)

Recombination Of Ions In Flames

- 1962P L. P. Pitayevskii. Zh. eksper. teor. Fiz. 42, 1326 (1962)
(English trans: Soviet Physics-JETP)

Recombination Of Electrons In A Monatomic Gas

- 1962P L. I. Pivovarov, M. T. Novikov and V. M. Tulaev. Zh. eksper. teor. Fiz. 42, 1490 (1962)
(English trans: Soviet Physics-JETP)

Electron Capture By Helium Ions In Various Gases In The 300-1500 KeV Energy Range

- 1962R V. E. Romanov. Optika i Spekt. 12, 111 (1962)
(English trans: Optics and Spect. 12, 54 (1962)).

Calculation Of A Cross Section Of The Free Free Transition Of An Electron In The Field Of A Neutral Hydrogen Atom.

VIII. THEORY OF INTERMOLECULAR FORCES

- 1959H H. U. Hostettler and R. B. Bernstein. J. chem. Phys. 31, 1422 (1959)

Comparison Of Exp-6 L -J (12 6) And Sutherland Potential Functions Applied To The Calculation Of Differential Scattering Cross Sections

- 1959K D. D. Konowalow, J. O. Hirschfelder and B. Linder. J. chem. Phys. 31, 1575 (1959)

Low-Temperature Low-Pressure Transport Coefficients For Gaseous Oxygen and Sulfur Atoms

- 1960G R. Gaspar. Acta phys. Hungar. 11, 71 (1960)

A Theoretical Estimation Of The Interaction Energy Of Inert Gas Atoms

- 1960J W. J. Jameson, Jr. and H. Aroeste. J. chem. Phys. 32, 374 (1960)

Short-Range Interaction Between A Hydrogen Molecule And A Hydrogen Atom II

- 1960L G. H. Lane and E. Everhart. Phys. Rev. 120, 2065 (1960)

Ion-Atom Potential Energy Functions Obtained From keV Scattering Data

- 1960S I. S. Srivastava. Indian J. Phys. 34, 539 (1960)

Intermolecular Potential And Properties Of Argon

- 1960Z R. W. Zwanzig. *Molecular Phys.* 3, 305 (1960)
Intermolecular Forces From Optical Spectra Of Impurities In Molecular Crystals
- 1961A I. Amdur, M. S. Longmire and E. A. Mason. *J. Chem. Phys.* 35, 895 (1961)
Scattering Of High Velocity Neutral Particles. XII He--CH₄; He--CF₄; CH₄--CH₄ and CF₄--CF₄ Interactions.
- 1961A I. Amdur. *Planet. Space Sci.* 3, 228 (1961)
An Experimental Approach To The Determination Of Gaseous Transport Properties At Very High Temperatures
- 1961A I. Amdur, J. E. Jordan and S. O. Colgate. *J. chem. Phys.* 34, 1525 (1961)
Scattering Of High-Velocity Neutral Particles. XI. Further Study Of The He-He Potential.
- 1961B A. K. Barua and P. K. Chakraborti. *Physica* 27, 753 (1961)
Krypton Krypton Molecular Interaction
- 1961B R. S. Brokaw. *Phys. of Fluids* 4, 944 (1961)
Estimated Collision Integrals For The Exponential Attractive Potential
- 1961K I. Kay and H. E. Moses. *Nuovo Cimento* 22, 689 (1961)
The Determination Of The Scattering Potential From The Spectral Measure Function V The Gelfand-Levitan Equation For The Three-Dimensional Scattering Problem.
- 1961M M. R. C. McDowell and G. Peach. *Phys. of Fluids* 4, 1332 (1961)
Quantal Study Of Diffusion And Viscosity Cross Sections For A Screened Coulomb Potential.
- 1961S I. B. Srivastava. *Indian J. Phys.* 35, 86 (1961)
Intermolecular Potentials of H₂ and D₂.
- 1962A I. Amdur and R. R. Bertrand. *J. chem Phys.* 36, 1078 (1962)
Redetermination Of The He-He Interaction Between 0.55 and 1.0 Å

- 1962B R. G. Breene, Jr. JQSRT 2, 225 (1962)
Dipole Moment Induction In Homonuclear Diatomic Molecules I. The Coulomb Case.
- 1962C R. D. Cloney and J. T. Vanderslice, J. chem. Phys. 36, 1866 (1962)
Interaction Energies From Scattering Cross-Sections Of Hydrogen Ions In CH₄ CF₄ C₂H₆ and C₂F₆
- 1962C R. D. Cloney, E. A. Mason and J. T. Vanderslice, J. chem. Phys. 36, 1103 (1962)
Binding Energy For Ar₂⁺ From Ion Scattering Data
- 1962C A. Cohen and C. H. Blanchard, J. chem. Phys. 36, 1402 (1962)
New Class Of Potential Functions
- 1962F R. J. Fallon, J. T. Vanderslice and R. D. Cloney, J. chem. Phys. 37, 1097 (1962)
Potential Curves And Rotational Perturbations Of CN
- 1962K I. Kirobkin and Z. I. Slawsky, J. chem. Phys. 37, 226 (1962)
Effects Of Electronic Exchange On The Efficiency Of Vibrational Excitation By Molecular Collisions. I Interaction Potential
- 1962M F. A. Morse, R. B. Bernstein and H. V. Hostettler, J. chem. Phys. 36, 1947 (1962)
Evaluation Of The Intermolecular Potential Well Depth From Observations Of Rainbow Scattering: Co-Hg & K-Hg
- 1962N R. G. Newton, J. math. Phys. 3, 75 (1962)
Construction Of Potentials From The Phase Shifts At Fixed Energy
- 1962S D. Steele, E. R. Lippincott and J. T. Vanderslice, RMP 34, 239 (1962)
Comparative Study Of Empirical Internuclear Potential Function

We add the following categories to those adopted by Professor Takayanagi:

A. BORN APPROXIMATION (General)

1957G M. Gavrilă. Stud. cercetari Fiz. 8, 421 (1957)

Derivation Of Sauter's Formula For The Photoelectric Effect By Means Of Born's Approximation Method.

1960D H. Davies. Nuclear Phys. 14, 464 (1960)

On The Convergence Of The Born Approximation

1961A A. M. Arthurs. Proc. Cambridge Phil. Soc. 57, 904 (1961)

The Mathematical Equivalence Of The Born Approximation And The Method Of Impact Parameters

1961M B. L. Moiseiwitsch. Proc. Phys. Soc. 78, 616 (1961)

Scattering By The Screened Coulomb Potential

1962K M. V. Kazarnovskii and A. V. Stepanov. Acta phys. Hungar. 14, 45 (1962)

Theory Of Resonance Scattering On Atomic Systems.

1962S P. Swan. Nuclear Phys. 29, 281 (1962)

Distorted Wave Approximation For Tensor Force Interactions

B BOUNDS

1959S L. Spruch and L. Rosenberg. Phys. Rev. 116, 1034 (1959)

Upper Bounds On Scattering Lengths For Static Potentials

1960S L. Spruch and L. Rosenberg. Phys. Rev. 117, 1095 (1960)

Upper Bounds On Scattering Lengths For Compound Systems: n-D Quartet Scattering.

1960S L. Spruch and L. Rosenberg. J. appl. Phys. 31, 2104 (1960)

Bounds On Low-Energy Scattering Parameters

1961R L. Rosenberg and L. Spruch. Phys. Rev. 121, 1720 (1961)

Bounds On Scattering Phase Shifts For Compound Systems

1962R L. Rosenberg and L. Spruch. Phys. Rev. 125, 1407 (1962)

Minimum Principle For Multi-Channel Scattering.

C. S-MATRIX

1960B I. Brendush. Rev. de Physique 5, 169 (1960)

Unitarity Relations Between The Parameters Of The Scattering Matrix For Elastic Collisions Of Particles With Arbitrary Spin.

1960E H. Ekstein. Phys. Rev. 117, 1590 (1960)

Equivalent Hamiltonians In Scattering Theory

1960T Y. Takahashi. Nuclear Phys. 20, 621 (1960)

Invariance And The S-Matrix II

1961B A. O. Barut and K. H. Ruei. J. math Phys. 2, 181 (1961)

Analytical Properties Of S Matrix And Uniqueness Of The Scattering Potential

1961K S. Kamefuchi, L. O'Raifeartaigh and A. Salam. Nuclear Phys. 28, 529 (1961)

Change Of Variables And Equivalence Theorems In Quantum Field Theories

1961K V. D. Kukin and A. R. Frenkin. Dokl. Akad. Nauk. SSSR 139, 1089 (1961)
(English trans: Soviet Physics-Koklady 6 722 (1962))

The Construction Of The Scattering Matrix In Nonlocal Theory

1961N R. G. Newton. J. Math. Phys. 2, 188 (1961)

Structure Of The Many-Channel S Matrix.

1961O V. S. Ol'khov s'kii and Yu. V. Tsekhmistrenko. Ukrayin. fiz Zh. 6, 149 (1961)

The Analytical Structure Of The S-Functions Of Elastic Scattering In The Case Of Infinite Potentials.

- 1961S A. D. Sukhanov. Zh. eksper. teor. Fiz. 41, 1915 (1961)
(English trans: Soviet Physics-JETP)
On The Interaction Hamiltonian In Quantum Field Theory
- 1961T S. Tani. Phys. Rev. 121, 346 (1961)
Structure Of The S Matrix In The Presence Of A Bound State
- 1962B R. Bartram and L. Spruch. J. math. Phys. 3, 297 (1962)
Bounds On Elements Of The S Matrix For Elastic Scattering: One-Dimensional Scattering
- 1962B A. Bottino, A. M. Longoni and T. Regge. Nuovo Cimento 23 954 (1962)
Potential Scattering For Complex Energy And Angular Momentum
- 1962B A. Bottino and A. M. Longoni. Nuovo Cimento 24, 353 (1962)
Holomorphy Domain Of The S-Matrix In Potential Scattering.
- 1962C K. Chadan. Nuovo Cimento 24, 379 (1962)
The Left-Hand Cut Discontinuity And Equivalent Potentials
- 1962F O. Favella and M. T. Reineri. Nuovo Cimento 23, 616 (1962)
Properties Of The S-Matrix As A Function Of The Complex Angular Momentum For The Many-Channel Case And For The Dirac Equation
- 1962G O. W. Greenberg. J. math. Phys. 3, 31 (1962)
Two Folk Lemmas On The Expansion Of The S Matrix Or The Out Field In Normal Ordered In Fields
- 1962R T. W. Ruijgrok. Physica 28, 198 (1962)
The S-Matrix For Systems With Bound States
- 1962Y K. Yamamoto. Progr. theor. Phys. 27, 219 (1962)
Physical Meaning Of Redundant Poles Of S-Matrix

D. VARIATIONAL TREATMENTS (General)

- 1960O T. Ohmura. J. math. Phys. 1, 27 (1960)
Minimum Property In The Hulthen-Type Variational Methods
- 1960O T. Ohmura. J. math Phys. 1, 35 (1960)
Stationary Expression For Effective Range
- 1960P I. C. Percival. Proc. Phys. Soc. 76, 206 (1960)
A Variational Principle For Scattering Phases
- 1961O T. Ohmura. Phys. Rev. 124, 130 (1961)
Variational Method For Scattering Length
- 1961S H. E. Saraph. Proc. Phys. Soc. 77, 827 (1961)
Application Of The Schwinger Variational Method To Zero-Energy Electron-Hydrogen Scattering.
- 1961S C. Schwartz Ann. Phys. 16, 36 (1961)
Variational Calculations Of Scattering.
- 1962G G. H. Goedecke and E. Brown. Phys. Rev. 125, 2116 (1962)
Variational Principle For Potential Scattering.

E. MISCELLANEOUS

- 1958T M. L. Ter-Mikaelyan and B. V. Khachatryan Zh. eksper. teor. Fiz. 35, 1287 (1958)
(English trans: Soviet-Physics-JETP 35(8) 898 (1959))
- 1959M A. Martin. Nuovo Cimento 14, 403 (1959)
On The Analytic Properties Of Partial Wave Scattering Amplitudes Obtained From The Schroedinger Equation.
- 1960F L. Favella and E. Predazzi. Nuovo Cimento 18, 570 (1960)
On The Coulomb Scattering By A Spherical Charge Distribution

- 1960K O. V. Konstantinov and V. I. Perel' Zh. eksper. teor. Fiz. 39, 861 (1960)
(English trans: Soviet Physics-JETP, 12 597 (1961)).
Collisions Of Particles In A High-Temperature Plasma.
- 1960P R. Peterkop. Latv. PSR Zinat. Akad Vestis 9, 79 (1960)
Asymptotic Expansions Of Charged-Particle Wave-Functions
- 1960R A. Russek Phys. Rev. 120, 1536 (1960)
Effect Of Target Gas Temperature On The Scattering Cross Section
- 1960S L. Spruch, T. F. O'Malley and L. Rosenberg Phys. Rev. Letters 5,
375 (1960)
Modification Of Effective Range Theory In The Presence Of A Long-Range
Potential
- 1960S P. Swan Nuclear Phys. 21, 233 (1960)
An Improved Approximation For Scattering Problems II
- 1960T S. Tani Phys. Rev. 117, 252 (1960)
Scattering Involving A Bound State.
- 1960T T. Tietz. Ann. Phys. 6, 150 (1960)
Effective Range Theory
- 1961A R. Aaron, R. D. Amado and B. W. Lee. Phys. Rev. 121, 319 (1961)
Divergence Of The Green's Function Series For Rearrangement Collisions
- 1961C T. Carrington J. Chem. Phys. 35, 807 (1961)
Transition Probabilities In Multilevel Systems: Calculation From Impulsive
And Steady State Experiments.
- 1961E L. V. East and P. A. Roys. Amer. J. Phys. 29, 548 (1961)
Classical Approach to Mott Scattering
- 1961E G. Ecker. Phys. of Fluids 4 127 (1961)
Enhanced Interaction In The Positive Column

- 1961F L. Fonda Nuovo Cimento Suppl. 20, 116 (1961)
Inelastic Collisions And Threshold Effects.
- 1961G A. Grossmann and Tai Tsun Wu. J. Math Phys. 2, 710 (1961)
Schroedinger Scattering Amplitude I.
- 1961G A. Grossman and Tai Tsun Wu. J. Math. Phys. 2, 714 (1961)
Schroedinger Scattering Amplitude. II.
- 1961H W. Hunziker. Helv. phys. Acta (Switzerland) 34, 593 (1961)
Regularity Properties In Scattering Amplitude In Case Of Potential Scattering.
- 1961K A. H. Kahn. Amer. J. Phys. 29, 77 (1961)
Phase-Shift Method For One-Dimensional Scattering
- 1961M V. V. Malakhov. Zh. eksper. teor. Fiz. 31, 762 (1961)
The Scattering Equations For Low Energies
- 1961M R. A. Mapleton J. Math. Phys. 2, 478 (1961)
Characteristic Coulomb Green's Function And Its Eigenfunction Expansion
- 1961N N. Nakanishi Progr. theor. Phys. 26, 337 (1961)
Integral Representations For Scattering Amplitudes In Perturbation Theory.
- 1961N L. P. Nizhnik. Dokl. Akad. Nauk SSSR 140, 533 (1961)
The Problem Of Inelastic Scattering
- 1961O T. F. O'Malley L. Spruch and L. Rosenberg. J. math. Phys. 2, 491 (1961)
Modification Of Effective-Range Theory In The Presence Of A Long-Range (r^{-4}) Potential.
- 1961S B. G. Skinner. Proc. Phys. Soc. 77, 551 (1961)
On The Validity Of Two Conjectures Relating To Resonance Collisions
- 1961S P. Swan. Nuclear Phys. 27, 620 (1961)
Distorted Wave Expansion For Scattering Problems

1962O T. F. O'Malley, L. Rosenberg and L. Spruch. Phys. Rev. 125, 1300 (1962)

Low-Energy Scattering Of A Charged Particle By A Neutral Polarizable System

1962W J. Weinstock Phys. Rev. 126, 341 (1962)

Binary Collision Expansion Of The Classical N-Body Green's Function

BIBLIOGRAPHY

1. "Dipole Moment Induction in Homonuclear Diatomic Molecules"
R. G. Breene, Jr., Physical Studies, Inc. Final Report under
AF19(604)-8417. 1962.
2. J. B. Fisk. Phys. Rev. 49, 167 (1936)
3. H. C. Stier. Zeit. f. Physik 76, 439 (1932)
4. R. G. Breene, Jr. JQSRT 2 225 (1962)
5. M. Kotani, Y. Mizuno, K. Kayama and E. Ishiguro. J. Phys. Soc.
Japan 12, 707 (1959)
6. D. R. Bates, R. A. Buckingham, H. S. W. Massey and J. J. Unwin.
Proc. Roy. Soc. (London) A170, 322 (1939)
7. R. G. Breene, Jr. J. Chem. Phys. 35, 625 (1961)
8. S. Geltman. Phys. Rev. 112, 176 (1958)
9. E. F. Gurnee and J. L. Magee. J. Chem. Phys. 26, 1237 (1957)
10. D. R. Bates. Proc. Roy. Soc. A245, 299 (1958)
11. R. McCarroll. Proc. Phys. Soc. A246 547 (1961)
12. A. M. Arthurs. Proc. Cambridge Phil. Soc. 57, 904 (1961)




Figure I. The molecular distortion parameters for the electron-oxygen or proton-oxygen case.




Figure II. The electron or proton oxygen interaction potential to include polarization.




Figure III. The definition of the angles referred to in Eq. (30) and subsequent.




Figure IV. The geometry for the Impact Parameter treatment of the proton-oxygen collision.




Figure V. The cross section integrand for various proton energies.




Figure VI. Total cross section for proton excitation of the Schumann-Runge system in units of $2 \cdot a_0^2$

Air Force Cambridge Research Laboratories, Bedford, Massachusetts. PROTON AND ELECTRON SCATTERING BY DIATOMIC MOLECULES by R. G. Breene, Jr., Physical Studies, Inc., Centerville, Ohio. April 1963. 86 pages. (Project 8658; Task 86580) Contract AF19(628)-476 (AFRL-63-697). The scattering of electrons by diatomic molecules, specifically, oxygen, is first considered. The treatment is restricted in such a way that electronic computing machinery is not required for solution. The Born approximation is applicable, and the appropriate equations are developed but not solved. The proton-molecule problem is next treated for the inelastic scattering of one

(over)

to fifty KeV protons by oxygen resulting in the excitation of the upper state of the Schumann-Runge system. The Born approximation is not applicable. The set of wave equations is reduced to three by arguments which may well not justify neglect of the dissociation equation. Potentials appropriate to the equation are developed and so on, but the equations are not solved. The impact parameter treatment of this proton-oxygen problem is set up for the two state case, and cross sections are computed for the one to fifty KeV region. Next, a general treatment of the multi-state impact parameter treatment is developed. The last portion of the report is concerned with additions to Professor Takayanagi's bibliography.

UNCLASSIFIED

UNCLASSIFIED

UNCLASSIFIED

UNCLASSIFIED

UNCLASSIFIED

UNCLASSIFIED

UNCLASSIFIED

UNCLASSIFIED

Air Force Cambridge Research Laboratories, Bedford, Massachusetts. PROTON AND ELECTRON SCATTERING BY DIATOMIC MOLECULES by R. G. Breene, Jr., Physical Studies, Inc., Centerville, Ohio. April 1963. 86 pages. (Project 8658; Task 86580) Contract AF19(628)-476 (AFRL-63-697). The scattering of electrons by diatomic molecules, specifically, oxygen, is first considered. The treatment is restricted in such a way that electronic computing machinery is not required for solution. The Born approximation is applicable, and the appropriate equations are developed but not solved. The proton-molecule problem is next treated for the inelastic scattering of one

(over)

to fifty KeV protons by oxygen resulting in the excitation of the upper state of the Schumann-Runge system. The Born approximation is not applicable. The set of wave equations is reduced to three by arguments which may well not justify neglect of the dissociation equation. Potentials appropriate to the equation are developed and so on, but the equations are not solved. The impact parameter treatment of this proton-oxygen problem is set up for the two state case, and cross sections are computed for the one to fifty KeV region. Next, a general treatment of the multi-state impact parameter treatment is developed. The last portion of the report is concerned with additions to Professor Takayanagi's bibliography.