

AD 60055 I

DASA-1483



THE EFFECTS OF AMBIENT  
PRESSURE ON THE TOLERANCE OF  
MICE TO AIR BLAST

Edward G. Damon, Donald R. Richmond  
and Clayton S. White

Presented before the  
Symposium on the Biological Effects  
of Blast and Shock  
Headquarters Field Command  
Defense Atomic Support Agency  
Sandia Base, Albuquerque, New Mexico  
April 20, 1963

Technical Progress Report  
on  
Contract No. DA-49-146-XZ-055

1964  
E

This work, an aspect of investigations dealing with  
the Biological Effects of Blast from Bombs, was  
supported by the Defense Atomic Support Agency of  
the Department of Defense.

(Reproduction in whole or in part is permitted for  
any purpose of the United States Government.)

Lovelace Foundation for Medical Education and Research  
Albuquerque, New Mexico  
March 1964

This report has been approved for open publication  
by The Office of the Assistant Secretary of Defense  
for Public Affairs.

FOREWORD

This report describes the development of appropriate apparatus and experiments to explore the relationship between the response of animals to air blast and the ambient pressure existing at the time of exposure. Specifically, the tolerance of mice to overpressure was determined using the expansion chamber of a specially modified shock tube in which the local pre-shot pressures were varied from 7 to 42 psi absolute.

The results and experience from this study will guide the design of appropriate hardware and the initiation of similar investigations, wherein larger animal species may be employed in the most economical manner. The ultimate aim of the work is the accurate prediction of human tolerance to air blast as a function of reduced pressure associated with altitude and increased pressure corresponding to various locations below sea level. Thus the findings have applicability in Aviation, Submarine and Environmental Medicine and are significant in air evacuation of blast-produced casualties, the care and therapy of blast injuries occurring underwater, inside submarines, aloft in aircraft and in other pressurized locations.

## ABSTRACT

↘  
Mice were exposed to overpressures of "long" duration in the expansion chamber of an air-driven shock tube inside which the initial, pre-blast pressures were varied over sixfold. When the animals were held at the initial pressure for one hour following the blast before being returned to the ambient pressure of the laboratory, tolerance values, expressed as LD<sub>50</sub>-1-hour gauge pressures, increased fourfold; they were 20.3, 31.0, 44.5, 55.4, and 91.8 psi for initial pressures of 7, 12, 18, 24, and 42 psia, respectively. When animals were returned to ambient level soon after blast exposure, the LD<sub>50</sub> pressures were lower than the above values for initial pressures greater than ambient and higher for initial pressures lower than ambient. The feasibility of scaling biological blast effects as a function of altitude <sup>was</sup> discussed and one approach suggested by available empirical data <sup>was</sup> regarded as a promising, but tentative procedure. ) ←

## ACKNOWLEDGMENTS

The able technical assistance of Mr. Charles S. Gaylord, Mr. Peter A. Betz, Mr. William Hicks, Mr. Donald E. Pratt, Mr. Dennis D. Branch, Mr. Kabby Mitchell, Jr., Mr. Keith G. Saunders, and Mr. Raymond T. Sanchez is greatly appreciated. Also, the aid of Mr. Robert A. Smith, who prepared illustrative material, Mr. Ray W. Albright, who carried out the probit analysis on a Bendix G-15 electronic computer, and Mrs. Maxine U. Thibert, who assisted with the illustrations, typed and edited this report, is acknowledged.

This report covers a portion of the work being presented by Mr. Edward G. Damon in a dissertation to be submitted to the University of New Mexico in partial fulfillment of the requirements for the Ph. D. degree.

Experiments described herein were conducted according to the Principles of Laboratory Animal Care as promulgated by the National Society for Medical Research.

## TABLE OF CONTENTS

	<u>Page</u>
Foreword . . . . .	i
Abstract . . . . .	ii
Acknowledgments . . . . .	iii
Introduction . . . . .	1
Methods . . . . .	1
Shock Tube . . . . .	1
Pressure-Time Measurements . . . . .	3
Animal Exposure . . . . .	3
Series I . . . . .	5
Controls . . . . .	5
Series II . . . . .	5
Controls . . . . .	5
Probit Analysis . . . . .	8
Results . . . . .	8
Discussion . . . . .	12

## LIST OF TABLES

No. 1	Results of Probit Analysis of the Series I Data . . . . .	10
No. 2	Comparison of LD <sub>50</sub> Values . . . . .	12
No. 3	Results of Probit Analysis of the Series II Data . . . . .	13

## LIST OF FIGURES

No. 1	Shock-Tube Layout and Pressure-Time Oscillograms . . . . .	2
No. 2	Calibration Curve - 12-Inch Shock Tube . . . . .	4
No. 3 (a-c)	Overall Pressure-Time Profiles for Series I . . . . .	6
No. 3 (d-e)	Overall Pressure-Time Profiles for Series I . . . . .	7
No. 4	Mortality Curves for Mice . . . . .	9
No. 5	Tolerance of Mice to Air Blast as Related to the Initial Pressure . . . . .	11

# THE EFFECTS OF AMBIENT PRESSURE ON THE TOLERANCE OF MICE TO AIR BLAST

Edward G. Damon, Donald R. Richmond  
and Clayton S. White

## INTRODUCTION

Although the relationships between biological response to air blast and various parameters of the pressure wave have been investigated in recent years,<sup>1-4</sup> very little is known about the effects of ambient pressure on mammalian tolerance to overpressure. Not only was attention called to this fact several years ago,<sup>5</sup> but for theoretical and commonsense reasons, it was predicted that the ambient pressure existing at the time an animal was loaded with a pressure pulse would be a significant parameter influencing biological response. Without question, it is important to know whether or not this speculation has validity, and if so, the magnitude of the effect, because human exposures to detonation-produced variations in the environment can and do occur at a variety of ambient pressures such as those existing at different elevations of terrain, at whatever levels above and below the earth's surface are available to man and at duty stations inside different manned vehicles and pressurized spaces wherever they may be.

The present investigation was undertaken to develop shock-tube and related techniques for exposing animals to air blast at different ambient pressures and to explore the tolerance of mice to "sharp"-rising overpressures of "long" duration as related to pre-shot ambient pressures ranging from a fraction of an atmosphere to several atmospheres.


## METHODS

### Shock Tube


A conventional, circular shock tube, 19 ft 6 in. long and 12 in. in diameter, was modified and used to expose mice to air blast at different ambient pressures. The tube had a wall thickness of 3/8 in., and as shown diagrammatically in Figure 1, was divided by a frangible diaphragm into a compression chamber 2 ft 6 in. long and a 17-ft expansion chamber. The latter was closed with an end-plate on which animal cages were mounted.

Appropriate pipes and valves, to allow pre- and post-shot control of the

# SHOC



-2-



# SHOCK TUBE LAYOUT






Figure 1

pressure inside the shock tube, were fitted to the expansion and compression chambers and multiple layers of Dupont Mylar plastic were employed as a diaphragm. Since experience proved that the Mylar sheets exhibited a consistent bursting pressure when tested on 12-in. tubes, different exposure pressures were achieved by varying the total thickness of the plastic and allowing each diaphragm to rupture spontaneously as the compression chamber was progressively pressurized.

### Pressure-Time Measurements

On every test, the shock pressures were measured with piezoelectric gauges mounted side-on in the wall of the tube 6 in. upstream from the end-plate (Figure 1). Occasionally, gauges were also located on the end-plate to record the pressure-time wave form at the position of the animals. The piezoelectric transducers contained sensors of Lead Metaniobate (Model ST-2, Susquehanna Instruments, Bel Air, Maryland). Each signal from a pressure transducer was passed through a cathode follower and was displayed and photographically recorded on a cathode-ray oscilloscope. Details of the system and its calibration already have been reported.<sup>4,6</sup> Typical pressure-time oscillograms obtained with the gauge mounted side-on in the wall of the tube are presented in Figure 1.

The overpressure in the expansion chamber before and after each blast was measured by a Bourdon-type dial pressure gauge (Heise Bourdon Tube Co., Newton, Connecticut). A mercury manometer gave the pressure levels when the expansion chamber was partly evacuated. The time required to increase or decrease the pressure in the expansion chamber was carefully measured with a stopwatch and also checked on oscillograms obtained with Quartz piezoelectric transducers (Model PZ-4, Kistler Instrument Corporation, North Tonawanda, New York). The oscilloscopes were triggered so that the time to increase or release the pressure was recorded.

Figure 2 presents a comparison of the empirical, shock-tube calibration curve with theoretical data. The results indicate that the measured performance of the current hardware was within 10 per cent of that predicted by the theoretical relationships. Since this result is consistent with experience reported elsewhere<sup>7,8</sup> as characteristic of air-driven, conventional shock tubes, it indicates that the methods used to measure shock pressures were reliable at either reduced or elevated initial ambient pressures.

### Animal Exposure

In all, 672 female mice of the Webster strain were employed. Their mean body weight was 19.7 g (standard error of the mean and range were  $\pm 0.84$  and 16 - 24 g, respectively). Except where otherwise mentioned, three animals were exposed per shot. Each animal was oriented right-side-on to the incident shock in an individual, cylindrical, wire-mesh cage mounted against the end-plate. The diameter of the wire from which the cages were constructed was 1/16 in. and the inside diameter of the squares of the mesh was 1/4 inch. The cages were arranged 2 in. apart, one above the other. Since the end-plate of the tube was oriented normal to the incident shock wave, the

CALIBRATION CURVE  
12-inch Shock Tube




Fig. 2. Comparison of the calibration curve for the 12-in. shock tube with the theoretical curve for shock strength as a function of the starting pressure ratio (Bleakney, 1949).<sup>7</sup>

animals were subjected to the incident and the reflected shock almost simultaneously. Consequently, the air-blast dose was taken to be the maximal overpressure in the reflected shock. The duration of the positive phase of the primary blast wave was 16 - 20 msec, which is much longer than the "critical duration" for mice.<sup>4</sup> Following the first positive wave, the animals were subjected to a series of decreasing secondary pressure pulses resulting from the reflection of the shock wave from one end of the tube to the other. Pressure-time record "a" in Figure 1 is a typical oscillogram showing these multiple reflections.

### Series I

Two hundred and seventy mice were exposed in groups to three levels of reflected shock pressures while at initial pressures of 7, 12, 18, 24, and 42 psia. Immediately after the blast, the pressure in the expansion chamber was quickly adjusted to the respective pre-shot level and then held for one hour before it was returned to ambient level.

The five overall pressure-time profiles for Series I experiments are illustrated in Figures 3a - 3e. Indicated are the times required to increase or decrease the pressure on the animals before and after the blast. For instance, Figure 3c shows that 25 seconds ( $t_1$ ) were required to increase the pressure from the atmospheric ambient ( $P_0$ ) of 12 psia to the initial pressure ( $P_i$ ) of 18 psia in the expansion section. It was held for 78 seconds ( $t_2$ ) before the blast. The duration of the blast wave itself was 0.016 seconds ( $t_3$ ). After the blast, the pressure stayed at  $P_b$  (27 psia) for 2 seconds ( $t_4$ ) before it could be reduced to the pre-shot level in 18 seconds ( $t_5$ ). At the end of the 1-hour hold ( $t_6$ ), the pressure was returned to ambient in 15 seconds ( $t_7$ ).

### Controls

Except for exposure to blast overpressures, 16 control animals were subjected to the pressure-time sequence illustrated in Figure 3e.

### Series II

Two hundred and eighty-five mice were exposed to air blast at initial pressures of 7, 18, 30, 36, and 42 psia following the general procedures used in Series I animals, except they were returned to ambient immediately after blast exposure. The rates of pressure changes previous to and following the blast were kept similar to those in the Series I studies, except for the absence of the 1-hour hold period ( $t_6$ ).

### Controls

Fifteen Series II control animals were handled as Series I controls except they were not held for an hour at the pre-shot pressure ( $P_i$ ) of 42 psia.






Fig. 3 (a-c). Overall pressure-time profiles for Series I.





Fig. 3 (d-e). Overall pressure-time profiles for Series I.

## Probit Analysis

Probit analysis was applied to the one-hour mortality data obtained from both experimental series.<sup>9</sup> Thus, the results presented refer to lethality within one hour following the blast.

In Series I, the total number of animals which were dead when first observed at the end of the 1-hour hold were recorded for the one-hour mortality. Of these, the number which exhibited signs of rigor mortis were also recorded. Since some of the animals could have died during the two minutes required for removing them from the tube following decompression, probit analysis was applied to both the total one-hour mortality data and the mortality data based on only those which exhibited signs of rigor mortis. Since there was no significant difference in the LD<sub>50</sub> values computed from the two sets of data, only the results of the analysis<sup>50</sup> of the total one-hour mortality data for Series I are presented.

## RESULTS

### Series I

Probit mortality curves relating the percentage dead in probit units to the log reflected pressure are presented in Figure 4 for the mice exposed at the five initial pressure levels in Series I. The probit regression lines were adjusted to an average slope since statistical tests revealed them to be essentially parallel at the 95-per cent fiducial limits.<sup>9</sup> The LD<sub>50</sub> reflected shock pressures with their 95-per cent confidence limits and the probit regression equations' constants are listed in Table 1 along with the associated number of animals. As indicated in Table 1, the reflected pressure required for 50-per cent lethality rose as the initial pressures were increased. The LD<sub>50</sub> pressures were 20.3, 31.0, 44.5, 55.3, and 91.8 psig when mice were exposed at initial pressures of 7, 12, 18, 24, and 42 psia, respectively. Each LD<sub>50</sub> value differed significantly from the others at the 95-per cent confidence level. Actually, the LD<sub>50</sub> values increased linearly with increasing initial pressures. A Bendix G-15 computer was programmed to fit a regression of the form,  $\log y = a + b \log x$ , to the data. Figure 5 presents the regression and a log-log plot of the data.

Table 2 compares the LD<sub>50</sub> values in terms of reflected overpressure (psig) and atmospheres (atm) of the initial pressure ( $\Delta P/P_i$ ). As noted, the LD<sub>50</sub> pressure ratio ranged from 2.90 to 2.19 for initial pressures of 7 to 42 psia, respectively. Thus, in terms of atm of the initial pressure, biological tolerance decreased somewhat with increases in initial pressure.

### Series II

The results of the probit analysis of the one-hour mortality data from the series in which the animals were returned to ambient immediately after

EFFECT OF INITIAL PRESSURE ON MOUSE RESPONSE TO AIR BLAST




Fig. 4. Probit regression lines relating the percent mortality in probit units to the log of the reflected shock pressures for mice subjected to air blast at different initial air pressures.

**TABLE 1**  
**RESULTS OF PROBIT ANALYSIS OF THE SERIES I DATA**

Initial Pressure, psia	Number of Animals	LD50-1-hour Reflected Pressure ( $\Delta P$ ), psig	Probit Equation Constants	
			intercept, a	slope, b
7	60	20.3 (19.0-21.5)*	-14.481	14.889**
12	45	31.0 (29.3-33.3)	-17.254	14.889
18	48	44.5 (41.9-47.4)	-19.543	14.889
24	60	55.3 (52.4-58.3)	-20.948	14.889
42	57	91.8 (86.1-98.3)	-24.225	14.889
<b>Total</b>	<b>270</b>			

\*Numbers in parentheses are the 95-percent confidence limits.  
\*\*Standard deviation of the slope constant, b -  $\pm 2.154$ .

TOLERANCE OF MICE TO AIR BLAST AS RELATED  
TO THE INITIAL PRESSURE




Fig. 5. LD50-1-hour overpressure as a function of the initial pressure at exposure. Regression equations:  
 Series I,  $\log (\text{LD}_{50}) = 0.590 + 0.842 \log (P_i)$ ;  
 Series II,  $\log (\text{LD}_{50}) = 0.832 + 0.611 \log (P_i)$ .

TABLE 2  
COMPARISON OF LD<sub>50</sub> VALUES

Initial Pressure, P <sub>i</sub> , psia	LD <sub>50</sub> -1-Hour Overpressure	
	ΔP, psig	atm*(ΔP/P <sub>i</sub> )
7	20.3	2.90
12	31.2	2.60
18	44.5	2.47
24	55.3	2.30
42	91.8	2.19
	Average	2.49

\*Atmospheres of the initial pressure.

blast exposure are presented in Table 3. The LD<sub>50</sub> reflected shock pressures were 22.7, 37.9, 53.6, 61.3, and 68.4 psig for initial pressures of 7, 18, 30, 36, and 42 psia, respectively. As illustrated in Figure 5, the LD<sub>50</sub> values were below those of Series I at initial pressures greater than ambient and above them for initial pressures less than ambient.

#### Controls

Results of control experiments revealed that the most rigorous combinations of decompression or compression, hold, and release of pressure (without the blast) encountered in this study, by themselves, produced neither deaths nor noticeable injury in mice. For instance, groups of animals were compressed to 67 psia in 225 seconds, held at that level for 2 minutes, and then returned to 42 psia and held for one hour, after which the pressure was reduced to 12 psia in 34 seconds. In addition, mice were compressed to 67 psia in 225 seconds, held there for 2 minutes, and then returned to 12 psia in 56 seconds.

#### DISCUSSION

This study, designed to explore the significance of ambient pressure on blast tolerance, shows an unequivocal increase in resistance to over-

TABLE 3  
RESULTS OF PROBIT ANALYSIS OF THE SERIES II DATA

Initial Pressure, psia	Number of Animals	LD50-1-Hour Reflected Pressure ( $\Delta P$ ), psig	Probit Equation Constants	
			intercept, a	slope, b
7	45	22.7 (21.0-24.6)*	-18.805	17.554**
18	69	37.9 (35.2-41.2)	-22.717	17.554
30	45	53.6 (49.4-58.7)	-25.359	17.554
36	57	61.3 (55.7-67.2)	-26.379	17.554
42	69	68.4 (64.2-73.2)	-27.211	17.554
Total	285			

\* Numbers in parentheses are the 95-percent confidence limits.

\*\* Standard deviation of the slope constant, b is  $\pm 2.946$ .

pressure in both the Series I and Series II experimental groups compared with controls. In terms of the magnitude of the overpressure of the reflected shock (psig), the Series I mice — those held for one hour at the pre-shot initial pressure before being returned to the Albuquerque ambient pressure — showed a fourfold increase in tolerance to be associated with a sixfold increase in the pre-shot ambient pressure. Series II animals — those returned to the Albuquerque ambient pressure immediately after exposure to blast — only exhibited a threefold increase in tolerance associated with the same sixfold increase in pre-shot ambient pressure.

The differences between the Series I and II data — shown clearly in Figure 5 — are of considerable interest and deserve several comments. First, in the experiments involving pre-shot pressures less than the Albuquerque ambient, blast tolerance was higher in the Series II than in the Series I mice. This means that mortality due to blast can be reduced by promptly pressurizing the animal after exposure to blast, as was the case for the Series II animals. This experience, consistent with the findings of Clemenson<sup>10</sup> and Benzinger<sup>11</sup> who demonstrated experimentally that early pressurization following a severe blast injury was beneficial and effective in reducing mortality, was not unanticipated since arterial air emboli, entering the circulation from the injured lung and known to be a prominent cause of early lethality in blasted animals, would be expected to decrease in size with pressurization and therefore become less hazardous to the animal.

Secondly and in contrast to the above results, the present study showed that blast tolerance of the Series I was greater than that of the Series II animals in all experiments involving pre-shot pressures above the Albuquerque ambient. These data mean that decompression carried out immediately after blast exposure, as was done in the Series II groups, increases lethality. One probable explanation is that blast-produced arterial emboli grow in size and therefore become a greater challenge to the animal. Another possibility is that more arterial emboli are produced by the decompression, a likely sequence of events should air trapping in the distal airways occur as a consequence of intra-bronchial hemorrhage, a not uncommon finding in blast-injured lungs.

Third, the Series I experiments in which the animals were held for one hour at the pre-shot ambient before being returned to laboratory pressures, no doubt are a more valid indication of the true variation in blast tolerance due to ambient pressure changes than are the Series II data. This seems so because (a) air emboli during the hold period have time to produce their biological effects, to decrease in size or to disappear from the circulation and (b) most individuals injured by blast are likely to be treated and held at the ambient pressure existing in the environment in which they were exposed. However, important exceptions to this statement are not improbable. For example, air evacuation of blast casualties could involve hazardous pressure changes. The post-exposure course of blast injuries occurring aloft in aircraft and during pressurized mining and tunneling operations could be worsened, improved or remain unchanged depending upon what pressure variations

occurred. Since control of the post-exposure pressure is not always impossible even in some emergencies, those who treat blast casualties should know two results of the present study; namely, (a) that the rate and range of decompression tolerated without demonstrable effect by control mice proved hazardous to blast-injured animals and (b) the rate and range of compression, producing no effect in controls when applied to experimental animals immediately after blast exposure, reduced mortality significantly.

Fourth, lethality-time data, limited in the Series I animals mostly to crude observation of body temperature and the presence or absence of rigor, would aid further analysis of the differences noted between Series I and II mice. Such information will be forthcoming in future experiments since the end-plate of the shock tube has been fitted with an observation window.

Fifth, though the Series I data, as noted above, seem more applicable than the Series II results to most blast situations in "real life," it is well to consider the validity of the Series I findings further. In this regard, there are at least three matters of interest. The first is whether or not exposing an animal to a series of "sharp"-rising pressure pulses as exemplified in the lower left portion of Figure 1 bears any similarity to an exposure involving only a single "sharp"-rising pressure as might occur near a detonation in the open or in an open-ended or vented shock tube. Though these two situations, a single versus a repetitive pressure pulse, do seem different on the surface, the  $P_{50}$  figure of 31.0 psi (29.3 - 33.3), referable to an ambient pressure of 12 psi found in the present study, is not significantly different from those reported for mice in previous investigations with "long"-duration overpressures carried out at Albuquerque altitude; 2, 4, 12, 13 namely,  $29.8 \pm 0.8$  psi for repetitive shock-tube pulses of 6 - 8 sec duration;  $30.7 \pm 0.6$  psi for single shock-tube pulses of 400 msec duration;  $29.0 \pm 0.6$  psi for single shock-tube pulses of 3 - 4 msec duration;  $26.0 \pm 0.4$  psi for single high-explosive pulses of 2.1 msec duration; and  $29.9 \pm 1.1$  psi for single high-explosive pulses of 1.3 msec duration. These data give power to the argument that it is the initial "sharp"-rising portion of a repetitive pulse of decreasing pressure that is definitive in producing lethality and not the second and subsequent oscillations, which indeed, seem to have little detectable effect.

Then, there is the question of possible differences in biological effect when using the shock-tube procedures described compared with actual free-field exposures to blast at various ambient pressures. For example, the technique developed represents an attempt to simulate a "real-life" blast situation in the laboratory, but the pressure-time variations in the shock tube — particularly over the immediate post-shot period — were hardly constant for the various experiments, represent departures from the ideal, and embody the potential for introducing variables into the experimental situation. Whether or not these may be significant, and if so, eliminated by improving the technique employed is not clear at the present time. However, appropriate work is under way in the laboratory and free-field experiments at different ambient pressures to check the shock tube data are being planned.

Also, since the  $P_{50}$  figures determined for the different ambient pressures represent an equal challenge to the animal — that is, the overpressures of 20.3, 31.0, 44.5, 55.3 and 91.8 psi above their respective ambients of 7, 12, 18, 24, and 42 psi are biologically equivalent — one searches for a constant parameter which, if approximately the same for each experimental group, might indicate consistency in the data, aid in their interpretation and increase confidence in the overall findings. That the pressure ratio,  $\Delta P/P_i$ , where  $P_i$  is the pre-shot ambient pressure and therefore the pressure inside the air-containing cavities of the body and  $\Delta P$  is the blast overpressure and therefore the external pressure loading an animal, might be such a parameter is supported by the common-sense view that blast effects are sure to be importantly related to the magnitudes of the internal and external pressures, by the work of Haber and Clamann on the physics of rapid decompression,<sup>14</sup> by the findings of Luft and Bancroft<sup>15</sup> in biological studies of decompression and by White et al.<sup>5</sup> in blast studies wherein pulmonary lesions in dogs were correlated to the pressure ratio for nuclear blast waves inside shelters that rose in "steps" or in a "saw-toothed" manner. In this regard, the results of the present study are encouraging.

The pressure ratios shown in Table 2 are not constant, but ranged from 2.9 to 2.2 when the pre-exposure ambient pressure was varied from 7 to 42 psi. Blast tolerance, expressed this way, decreased by a factor of 1.3 or near 25 per cent when there was a sixfold increase in ambient pressure. If one uses the average pressure ratio of 2.5 shown in Table 2, it is possible to say that the  $P_{50}$  ratios only varied about 16 and 12 per cent above and below the average, respectively.

While experimental variations of these magnitudes are frequently noted in biological studies, the consistent and apparently not random decrease in the  $P_{50}$  pressure ratios with progressive elevation of the ambient pressures found in the Series I experiments not only stimulates one to search further for analytical understanding of the observed data to improve the grasp of etiologic mechanisms, but prompts a cautious approach to drawing general conclusions applicable to all mammalian species including man on the basis of the experiments reported here on mice.

Fortunately, it is now possible to say that similar studies have been completed using rats and guinea pigs. Preliminary analyses of the data show some randomness in the pressure-ratio figures and, in general, similar trends in blast tolerance with variations in the ambient pressure. Whether or not the pressure-ratio associated with such experiments is indeed a constant, with the differences noted indicating only "normal" experimental error and chance variations, cannot be stated now. But if results from future experiments with other species indicate that the  $LD_{50}$ 's can be expressed as multiples of the initial pressure, biological blast scaling as a function of ambient pressure will become a relatively simple matter. For example, man's tolerance ( $LD_{50}$ -24-hours) to "sharp"-rising overpressures of 400-msec duration has been calculated to be 50 psig from extrapolation of an interspecies correlation involving six different mammals.<sup>2</sup> Since the data

were compiled at an ambient pressure of 12.0 psia, the overpressure – normalized to the initial pressure – would be 4.2 atm. Consequently, to obtain the LD<sub>50</sub> for "long"-duration air blasts for different ambient pressures, one may tentatively multiply the ambient pressure of interest by 4.2. Thus at sea level (14.7 psia), the calculated LD<sub>50</sub> for man would be 62 psig; at 26,400 ft (5.2 psia) it would be 22 psig. It is well to emphasize the tentative and uncertain nature of these procedures, and it is no doubt premature to dwell on this topic further. Let it suffice to say that full understanding of biological blast scaling must await the results of future work.

Be this as it may, it is currently quite clear that the ambient pressure is indeed a physical parameter of major importance in specifying blast effects. Consequently, recording the local barometric pressure now needs to be considered a requirement in all quantitative investigations of blast tolerance.

## REFERENCES

1. Fisher, R. B., P. L. Krohn and S. Zuckerman, "The Relationship Between Body Size and the Lethal Effects of Blast," Ministry of Home Security Report R. C. 284, Oxford University, Oxford, England, December 10, 1941.
2. Richmond, D. R., V. R. Clare, V. C. Goldizen, D. E. Pratt, R. T. Sanchez and C. S. White, "Biological Effects of Overpressure. II. A Shock Tube Utilized to Produce Sharp-Rising Overpressures of 400 Milliseconds Duration and Its Employment in Biomedical Experiments," Technical Progress Report No. DASA 1246, Defense Atomic Support Agency, Department of Defense, Washington 25, D. C., April 7, 1961. Also in Aerospace Med., 32: 997-1008, 1961.
3. Richmond, D. R. and C. S. White, "A Tentative Estimation of Man's Tolerance to Overpressures from Air Blast," in Proceedings of the Symposium on Effectiveness Analysis Techniques for Non-Nuclear Warheads against Surface Targets, October 30-31, 1962, Vol. 1, pp. L to L-34, U. S. Naval Weapons Laboratory, Dahlgren, Virginia, Technical Progress Report No. DASA 1335, Defense Atomic Support Agency, Department of Defense, Washington 25, D. C., November 7, 1962.
4. Richmond, D. R., V. C. Goldizen, V. R. Clare and C. S. White, "The Overpressure-Duration Relationship and Lethality in Small Animals," Technical Progress Report No. DASA 1325, Defense Atomic Support Agency, Department of Defense, Washington 25, D. C., September 10, 1962.
5. White, C. S., T. L. Chiffelle, D. R. Richmond, W. H. Lockyear, I. G. Bowen, V. C. Goldizen, H. W. Merideth, D. E. Kilgore, B. B. Longwell, J. T. Parker, F. Sherping and M. E. Cribb, "The Biological Effects of Pressure Phenomena Occurring Inside Protective Shelters Following Nuclear Detonation," USAEC Civil Effects Test Group Report, WT-1179, Office of Technical Services, Department of Commerce, Washington 25, D. C., October 28, 1957.
6. Granath, B. A. and G. A. Coulter, "BRL Shock Tube Piezo-Electric Blast Gages," BRL Technical Note No. 1478, Ballistic Research Laboratories, Aberdeen Proving Ground, Md., August 1962.
7. Bleakney, W., D. K. Weimer and C. H. Fletcher, "Shock Tube: A Facility for Investigations in Fluid Dynamics," Rev. Sci. Instrum., 20: 807-815, 1949.
8. Lampson, C. W., "Resume of the Theory of Plane Shock and Adiabatic Waves with Applications to the Theory of the Shock Tube," BRL Technical Note No. 139, Ballistic Research Laboratories, Aberdeen Proving Ground, Md., March 1950.

9. Finney, D. J., Probit Analysis. A Statistical Treatment of the Sigmoid Response Curve, (2nd edition), Cambridge University Press, Cambridge, England, 1952.
10. Clemedson, Carl-Johan and H. Hultman, "Air Embolism and the Cause of Death in Blast Injury," Milit. Surg., 114: 424-437, 1954.
11. Benzinger, T., "Physiological Effects of Blast in Air and Water," Chap. XIV-B, German Aviation Medicine, World War II, Vol. II, pp. 1225-1259, U. S. Government Printing Office, Washington 25, D. C., 1950.
12. Richmond, D. R., V. C. Goldizen, V. R. Clare, D. E. Pratt, F. Sherping, R. T. Sanchez, C. C. Fischer and C. S. White, "Biologic Response to Overpressure. II' Mortality in Small Animals Exposed in a Shock Tube to Sharp-Rising Overpressures of 3 to 4 Msec Duration," Technical Progress Report No. DASA 1242, Defense Atomic Support Agency, Department of Defense, Washington 25, D. C., June 15, 1961. Also in Aerospace Med., 33: 1-27, 1962.
13. Richmond, D. R., R. V. Taborelli, F. Sherping, M. B. Wetherbe, R. T. Sanchez, V. C. Goldizen and C. S. White, "Shock Tube Studies of the Effects of Sharp-Rising, Long-Duration Overpressures on Biological Systems," USAEC Technical Report, TID-6056, Office of Technical Services, Department of Commerce, Washington 25, D. C., March 10, 1959.
14. Luft, U. C. and R. W. Bancroft, "Transthoracic Pressure in Man During Rapid Decompression," J. Aviat. Med., 27: 208-220, 1956.
15. Haber, F. and H. G. Clamann, "Physics and Engineering of Rapid Decompression: A General Theory of Rapid Decompression," Report 3, Project 21-1201-0008, USAF School of Aviation Medicine, Randolph AFB, Texas, 1953.

## DISTRIBUTION

### ARMY AGENCIES

Deputy Chief of Staff for Military Operations, Department of the Army,  
ATTN: Dir of SW&R, Washington, D. C. 20301 (1 copy)

Chief of Research & Development, Department of the Army, ATTN: Atomic  
Division, Washington, D. C. 20301 (1 copy)

Deputy Chief of Staff Intelligence, Department of the Army, Washington,  
D. C. (1 copy)

Chief of Engineers, Department of the Army, ATTN: ENGNB, Washington,  
D. C. 20315 (1 copy)

Chief of Engineers, Department of the Army, ATTN: ENGEB, Washington,  
D. C. 20315 (1 copy)

Chief of Engineers, Department of the Army, ATTN: ENGTB, Washington,  
D. C. 20315 (1 copy)

Chief of Ordnance, Department of the Army, ATTN: ORDTN, Washington, D. C.,  
20315 (2 copies)

Chief Signal Officer, Comb Dev & Ops Div, ATTN: SIGCO-4, Washington, D. C.,  
20330 (1 copy)

Chief of Transportation, Office of Planning & Intel, Department of the  
Army, Washington, D. C., 20315 (1 copy)

The Surgeon General, Department of the Army, ATTN: MEDNE, Washington,  
D. C. 20360 (2 copies)

Commander-in-Chief, U.S. Army, Europe, APO 403, ATTN: OPOT Div, Weapons  
Branch, New York, New York (one copy)

Commanding General, U. S. Continental Army Command, Fort Monroe, Virginia,  
23351 (3 copies)

Director of Special Weapons, Development Office, HQ CONARC, ATTN:  
Capt Chester I. Peterson, Fort Bliss, Texas 79906 (1 copy)

President, U. S. Army Artillery Board, Fort Sill, Oklahoma 73503 (1 copy)

President, U. S. Army Aviation Board, ATTN: AIBG-DG, Fort Rucker, Alabama  
36362 (1 copy)

Commandant, U. S. Army C&GS College, ATTN: Archives, Fort Leavenworth,  
Kansas 66027 (1 copy)

Commandant, U. S. Army Air Defense School, ATTN: Command & Staff Dept,  
Fort Bliss, Texas 79906 (1 copy)

Commandant, U. S. Army Armored School, Fort Knox, Kentucky 40120 (1 copy)

Commandant, U. S. Army Arty & Missile School, ATTN: Combat Dev Dept, Fort  
Sill, Oklahoma 73503 (1 copy)

Commandant, U. S. Army Infantry School, ATTN: C.D.S., Fort Benning,  
Georgia 31905 (1 copy)

Commandant, Quartermaster School, U. S. Army, ATTN: Ch, QM Library,  
Fort Lee, Virginia 23801 (1 copy)

Commanding General, Chemical Corps Training Command, Fort McClellan,  
Alabama 36205 (1 copy)

Commandant, U. S. Army Chemical Corps CBR, Weapons School, Dugway Proving  
Ground, Dugway, Utah 84022 (1 copy)

Commandant, U. S. Army Signal School, Fort Monmouth, New Jersey 07703 (1 copy)

(Cont'd on page 2)

Commandant, U. S. Army Transport School, ATTN: Security and Information  
 Office, Fort Eustis, Virginia 79906 (1 copy)  
 Commanding General, The Engineer Center, ATTN: Asst Cndt Engineer  
 School, Fort Belvoir, Virginia 22060 (1 copy)  
 Commanding General, Army Medical Service School, Brooke Army Medical  
 Center, Fort Sam Houston, Texas 78234 (1 copy)  
 Commanding Officer, 9th Hospital Center, APO 180, ATTN: CO, US Army  
 Nuclear Medicine Research Det, Europe, New York, New York (1 copy)  
 Director, Armed Forces Institute of Pathology, Walter Reed Army Medical  
 Center, Washington, D. C. 20315 (1 copy)  
 Commanding Officer, Army Medical Research Laboratory, Fort Knox,  
 Kentucky 40120 (1 copy)  
 Commandant, Walter Reed Army Institute of Research, Walter Reed Army Medical  
 Center, Washington, D. C. 20315 (1 copy)  
 Commanding General, QM R&D Comd, QM R&D Center, ATTN: CBR Liaison Officer,  
 Natick, Massachusetts (1 copy)  
 Commanding General, QM Research & Engineering Command, USA, Natick,  
 Massachusetts (For reports from Opn HARDTACK only) (1 copy)  
 Director, U. S. Army Research and Development Laboratory, ATTN: Chief,  
 Technical Support Branch, Fort Belvoir, Virginia 22060 (1 copy)  
 Director, Waterways Experiment Station, U. S. Army Corp of Engineers, P. O.  
 Box 631, ATTN: Library, Vicksburg, Mississippi (1 copy)  
 Director, U. S. Army Ballistic Research Laboratories, Aberdeen Proving  
 Ground, Maryland 21005 (2 copies)  
 Chief, Army Ballistic Missile Agency, ATTN: ORDAB-HT, Redstone Arsenal,  
 Alabama 35808 (1 copy)  
 Commanding General, Army Electronic Proving Ground, ATTN: Tech Library,  
 Fort Huachuca, Arizona 85613 (1 copy)  
 Director, Research Analysis Corporation, Waverly Bldg, 6935 Arlington  
 Road, Bethesda 14, Maryland (1 copy)  
 Commanding General, Army Materiel Command, Washington, D. C. 20315 (2 copies)  
 Commanding Officer, U. S. Army Nuclear Defense Laboratory, ATTN: Librarian,  
 Edgewood Arsenal, Maryland 21040 (1 copy)  
 Commanding General, Army Materiel Command, ATTN: EOD Division, Washington,  
 D. C. 20315 (1 copy)

#### NAVY AGENCIES

Chief of Naval Operations, Navy Department, ATTN: OPO3EG, Washington,  
 D. C. 20350 (1 copy)  
 Chief of Naval Operation, Navy Department, ATTN: OP-75, Washington,  
 D. C. 20350 (1 copy)  
 Chief of Naval Operations, Navy Department, ATTN: OP-922G2, Washington,  
 D. C. 20350 (1 copy)  
 Chief of Naval Operation, Navy Department, ATTN: OP-91, Washington,  
 D. C. 20350 (1 copy)  
 (Cont'd on page 3)

Chief of Naval Personnel, Navy Department, Washington, D. C. 20350 (1 copy)  
 Chief of Naval Research, Navy Department, ATTN: Code 811, Washington,  
 D. C. 20390 (2 copies)  
 Chief, Bureau of Medicine and Surgery, Navy Department, ATTN: Special  
 Weapons Def Division, Washington, D. C. 20390 (1 copy)  
 Chief, Bureau of Ships, Navy Department, ATTN: Code 423, Washington,  
 D. C. 20360 (1 copy)  
 Chief, Bureau of Yards and Docks, Navy Department, ATTN: D-440,  
 Washington, D. C. 20370 (1 copy)  
 Director, U. S. Naval Research Laboratory, ATTN: Mrs. Katherine H.  
 Cass, Washington, D. C. 20390 (1 copy)  
 Commander, U. S. Naval Ordnance Laboratory, White Oak, Silver Spring,  
 Maryland 20390 (2 copies)  
 Commander, U. S. Naval Material Laboratory (Code 900), New York Naval  
 Shipyard, Naval Base, Brooklyn 1, New York (1 copy)  
 Commanding Officer, U. S. Naval Mine Defense Lab, Panama City, Florida (1 copy)  
 Commanding Officer and Director, U. S. Naval Radiological Defense Laboratory,  
 ATTN: Tech Info Div, San Francisco 24, California (4 copies)  
 Commanding Officer, U. S. Naval Schools Command, U. S. Naval Station,  
 Treasure Island, San Francisco, California (1 copy)  
 Superintendent, U. S. Naval Postgraduate School, Monterey, California (1 copy)  
 Commanding Officer- Nuclear Weapons Training Center, Atlantic, U. S. Naval  
 Base, ATTN: Nuclear Warfare Department, Norfolk 11, Virginia (1 copy)  
 Commanding Officer, Nuclear Weapons Training Center, Pacific, U. S. Naval  
 Air Station, North Island, San Diego 35, California (1 copy)  
 Commanding Officer, U. S. Naval Damage Control Training Center, Naval Base,  
 ATTN: ABC Defense Course, Philadelphia 12, Pennsylvania (1 copy)  
 Commanding Officer, U. S. Naval Air Development Center, ATTN: NAS,  
 Librarian, Johnsville, Pennsylvania (1 copy)  
 Commanding Officer, U. S. Naval Medical Research Institute, National Naval  
 Medical Center, Bethesda, Maryland 20011 (1 copy)  
 Officer in Charge, U. S. Naval Supply Research and Development Facility,  
 Naval Supply Center, Bayonne, New Jersey (1 copy)  
 Commandant, U. S. Marine Corps, ATTN: Code A03H, Washington, D. C. 20380  
 (1 copy)

#### AIR FORCE AGENCIES

HQ USAF (AFTAC) Wash DC 20330 (1 copy)  
 HQ USAF, ATTN: Operations Analysis Office, Vice Chief of Staff, Wash 25 DC  
 (1 copy)  
 HQ USAF (AFNIN3), ATTN: ACS/1 (AFOIN-3VI), Wash DC 20330 (2 copies)  
 Assistant Chief of Staff, Intelligence, HQ USAF, APO 633, ATTN: Directorate  
 of Air Targets, New York, New York (1 copy)  
 HQ USAF (AFRDD), ATTN: Guidance & Weapons Division, Wash DC 20330 (1 copy)  
 TAC, ATTN: Doc Security Branch, Langley AFB VA 23365 (1 copy)  
 The Surgeon General, HQ USAF, ATTN: Bio-Def Pre Med Div, Wash DC 20330 (1 copy)  
 (Cont'd on page 4)

TAC, ATTN: Doc Security Branch, Langley AFB Va 23365 (1 copy)  
 ADC, ATTN: Assistant for Atomic Energy, ADLDC-A, Ent AFB Colo 80912 (1 copy)  
 AFSC, Andrews AFB, ATTN: RDRWA, Wash DC 20331  
 DDC(TIOL), AF Unit Post Office, ATTN: WDSOT, Los Angeles, Calif 90045 (1 copy)  
 Commander-in-Chief, Pacific Air Forces, APO 953, ATTN: PFCIE-MB, Base  
 Recovery, San Francisco, Calif (1 copy)  
 AFRL, L G Hanscom Fld, ATTN: CRQST-2, Bedford Mass 01731 (2 copies)  
 AFSWC, ATTN: Tech Info & Intel Div, Kirtland AFB NMex 87117 (5 copies)  
 AUL, Maxwell AFB Ala 36112 (2 copies)  
 Lowry AFB Colo, ATTN: Dept of Sp Wpns Tng, 80230 (1 copy)  
 USAFSAM, Brooks AFB Tex 78235 (2 copies)  
 Commander, 1009th Sp Wpns Squadron, USAF, Wash DC (1 copy)  
 ASD, ATTN: WCOSI, Wright-Patterson AFB Ohio 45433 (2 copies)  
 Director, USAF Project RAND, VIA: US Air Force Liaison Office, The RAND  
 Corporation, 1700 Main Street, Santa Monica, California (2 copies)  
 Commander, Air Defense Systems Integration Division, L G Hanscom Fld,  
 ATTN: SIDE-S, Bedford Mass 01731 (1 copy)  
 Command, Air Technical Intelligence Center, USAF, ATTN: AFCIN-421A,  
 Library, Wright-Patterson AFB Ohio 45433 (1 copy)

#### OTHER AGENCIES

Director of Defense Research and Engineering, ATTN: Tech Library,  
 Washington, D. C. 20301 (1 copy)  
 Director, Weapons Systems Evaluation Group, Room 1E880, Washington,  
 D. C. 20301 (1 copy)  
 U. S. Documents Officer, Office of the United States National Military  
 Representative-SHAPE APO 55, New York, New York (1 copy)  
 Director, Defense Atomic Support Agency, ATTN: Document Library,  
 Washington, D. C. 20301 (4 copies\*) \*Reduce to 3 cys for all FWE Rpts.  
 Commander, Field Command, Defense Atomic Support Agency, Albuquerque,  
 New Mexico 87115 (1 copy)  
 Commander, Field Command, Defense Atomic Support Agency, ATTN: FCWT,  
 Albuquerque, New Mexico 87115 (2 copies)  
 Commander, Field Command, Defense Atomic Support Agency, ATTN: FCTG,  
 Albuquerque, New Mexico 87115 (1 copy)  
 Director, National Aeronautics & Space Administration, 1520 H Street, N. W.,  
 Washington, D. C. 20546 (1 copy)  
 SAC ATTN: OAWS, Offutt AFB Nebr 68113 (1 copy)  
 Commandant, U. S. Coast Guard, 1300 E. Street, N. W., ATTN: OIN,  
 Washington, D. C. (1 copy)  
 (Cont'd on page 5)

## OTHER DISTRIBUTION

U. S. Atomic Energy Commission, Washington, D. C. 20545, ATTN:  
Chief, Civil Effects Branch, Division of Biology and Medicine (450 copies)  
Director, U. S. Army Ballistic Research Laboratories, ATTN: Terminal Ballistics,  
Mr. Robert O. Clark, Physicist (1 copy); Mr. William J. Taylor, Physicist (1 copy)  
Aberdeen Proving Ground, Maryland 20115

Airborne Instruments Laboratory, Department of Medicine and Biological  
Physics, Deer Park, Long Island, New York, ATTN: Mr. W. J. Carberry (1 copy)

AFSWC, ATTN: WIRS, Kirtland AFB NMex 87117 (1 copy)

AFSC, ATTN: SCGB, Andrews AFB Wash DC 20331 (1 copy)

AiResearch Manufacturing Company, 9851-9951 Sepulveda Blvd, Los Angeles 25,  
California, ATTN: Mr. Frederick H. Green, Assistant Chief, Preliminary  
Design (1 copy); Dr. Names N. Waggoner, Medical Director (1 copy)

AeResearch Manufacturing Company, Sky Harbor Airport, 402 South 36th Street,  
Phoenix, Arizona, ATTN: Delano Debaryshe (1 copy); Leighton S. King (1 copy)

American Airlines, Inc., La Guardia Airport Station, Flushing 71, New York,  
ATTN: Medical Director (1 copy)

USAFSAM, ATTN: Maj Gen Theodore C. Bedwell, Jr., Commandant (1 copy);  
Col Paul A. Campbell, Chief, Space Medicine (1 copy); Dr. Hubertus  
Strughold, Advisor for Research & Professor of Space Medicine (1 copy)

Boeing Company, Aero Space Division, 7755 E Marginal Way, Seattle 24, Washington  
ATTN: Dr. Thrift G. Hanks, Director of Health & Safety (1 copy); Dr. Romney  
H. Lowry, Manager, Space Medicine Branch (1 copy); Dr. F. Werner, Jr,  
Space Medicine Section (1 copy)

Chance Vought Astronautics, P. O. Box 5907, Dallas 22, Texas, ATTN: Life  
Sciences Section (5 copies)

Commanding Officer, U. S. Army Chemical Research & Development Laboratories,  
ATTN: Dr. Fred W. Stemler (1 copy); Dr. R. S. Anderson (1 copy),  
Edgewood Arsenal, Maryland 21040

Civil Aeromedical Research Institute, Oklahoma City, Oklahoma, ATTN:  
Director of Research (1 copy)

Convair Division, General Dynamics Corp, Fort Worth, Texas, ATTN:  
Mr. H. A. Bodely (1 copy); Mr. Schreiber (1 copy)

Convair, General Dynamics Corporation, Mail Zone 1-713, P. O. Box 1950,  
San Diego 12, California, ATTN: Dr. R. C. Armstrong, Chief Physician;  
Dr. J. C. Clark, Assistant to Vice-President Engineering; Mr. James  
Dempsey; Dr. L. L. Lowry, Chief Staff Systems Evaluation Group; Mr. M.  
H. Thiel, Design Specialist; Dr. R. A. Nau (Mail Zone 6-104);  
Mr. W. F. Rector, III (Mail Zone 580-40), P.O. Box 1128; Mr. R. C. Sebald,  
Vice-President Engineering, Convair General Offices, (8 copies)

Commander, Field Command, Defense Atomic Support Agency, Sandia Base,  
ATTN: Col F. Neumann, Surgeon, Albuquerque, New Mexico 87115 (1 copy)

Department of National Defense, Defense Research Board, Suffield Experimental  
Station, Ralston, Alberta, Canada, ATTN: Physiology Section (2 copies)

The Dikewood Corporation, 4805 Menaul Blvd., N. E., Albuquerque, New  
Mexico (1 copy)

Douglas Aircraft Company, Inc., El Segundo Division, El Segundo, California,  
ATTN: Mr. Harvey Glassner (1 copy); Dr. E. B. Konecci (1 copy)

(Cont'd on page 6)

Federal Aviation Agency, 1711 New York Avenue N. W., Washington, D. C. 20553, ATTN: Civil Air Surgeon (1 copy)

Goodyear Aircraft Corporation, Department 475, Plant H, 1210 Massillon Road, Akron 15, Ohio, ATTN: Dr. A. J. Cacioppo (1 copy)

Harvard School of Public Health, Harvard University, 695 Huntington Avenue, Boston 15, Massachusetts, ATTN: Dr. Ross A. McFarland, Associate Professor of Industrial Hygiene (1 copy)

Mr. Kenneth Kaplan, Physicist, Broadview Research Corporation, 1811 Trousdale Drive, Burlingame, California, (1 copy)

Lockheed Aircraft Company, Suite 302, First National Bank Bldg, Burbank, California, ATTN: Dr. Charles Barron (1 copy)

Lockheed Missile and Space Company, Space Physics Department (53-23), Sunnyvale, California, ATTN: Scientific Assistant to Director of Research (3 copies)

Lockheed Missile and Space Company, 1122 Jagels Road, Palo Alto, California, ATTN: Missile Systems Director (1 copy)

Lovelace Foundation for Medical Education and Research, 4800 Gibson Blvd, S.E., Albuquerque, New Mexico, ATTN: Dr. Clayton S. White, Director of Research (50 copies)

The Martin Company, Denver, Colorado, ATTN: Chief, Space Medicine (1 copy)

McDonnell Aircraft Company, Lambert Field, Saint Louis, Missouri, ATTN: Chief Airborne Equipment Systems Engineer (1 copy)

National Aeronautics and Space Administration, 1520 H. Street, N. W., Washington D. C. 20546, ATTN: Director, Life Sciences Program (1 copy)

Naval Medical Research Institute, Bethesda, Maryland 20315, ATTN: Dr. David E. Goldman, MSC, Commander (1 copy)

Chief, Bureau of Medicine & Surgery, Navy Department, ATTN: Director, Submarine Medical Division, Washington, D. C. 20390 (1 copy)

North American Aviation, International Airport, Los Angeles 45, California, ATTN: Scott Crossfield; Dr. Toby Freedman, Flight Surgeon; Mr. Fred A. Payne, Manager Space Planning, Development Planning; Mr. Harrison A. Storms (4 copies)

Director of Defense Research & Engineering, ATTN: Chief, Medical Services Division, Office of Science, Washington, D. C. 20301 (1 copy)

The Ohio State University, 410 West 10th Avenue, Columbus 10, Ohio, ATTN: Dr. William F. Ashe, Chairman, Department of Preventive Medicine; Dean Richard L. Meiling (2 copies)

Director, The RAND Corporation, 1700 Main Street, Santa Monica, California, ATTN: Dr. H. H. Mitchell, Physics Division; Dr. Harold L. Brode (2 copies)

Republic Aviation Corporation, Applied Research & Development, Farmingdale, Long Island, New York, ATTN: Life Sciences Division (3 copies)

President, Sandia Corporation, Sandia Base, Albuquerque, New Mexico 87115, ATTN: Director of Research (5 copies)

System Development Corporation, Santa Monica, California, ATTN: Dr. C. J. Roach (1 copy)

United Aircraft Company, Denver, Colorado, ATTN: Medical Director (1 copy)

(Cont'd on page 7)

Laboratory of Nuclear Medicine & Radiation Biology, School of Medicine,  
University of California, Los Angeles, 900 Veteran Avenue, Los Angeles 24,  
California, ATTN: Dr. G. M. McDonnel, Associate Professor; Dr. Benedict  
Cassen (2 copies)

University of Illinois, Chicago Professional Colleges, 840 Wood Street,  
Chicago 12, Illinois, ATTN: Director, Aeromedical Laboratory (1 copy)

University of Kentucky, School of Medicine, Lexington, Kentucky, ATTN:  
Dr. Loren D. Carlson, Professor of Physiology & Biophysics (1 copy)

University of New Mexico, Albuquerque, New Mexico, ATTN: Library (1 copy)

U. S. Naval Ordnance Laboratory, ATTN: Capt Richard H. Lee, MSC; Mr. James  
F. Moulton, White Oak, Maryland 20390 (2 copies)

U. S. Naval School of Aviation Medicine, U. S. Naval Aviation Medical  
Center, ATTN: Director of Research, Pensacola, Florida (1 copy)

Dr. Shields Warren, Cancer Research Institute, New England Deaconess Hospital,  
194 Pilgrim Road, Boston 15, Massachusetts (1 copy)

ASD, ATTN: Commanding Officer; Chief, Bioacoustics Laboratory, Wright-  
Patterson AFB Ohio 45433 (2 copies)

Dr. Eugene Zwoyer, Director, Shock Tube Laboratory, P. O. Box 188, University  
Station, Albuquerque, New Mexico (1 copy)

DDC, Cameron Stn, Alexandria, Va 22314 (20 copies)

Western Development Laboratories, Philco Corporation, Palo Alto, California,  
ATTN: Chief, Bioastronautics (1 copy)