


AD 715512

A NOTE ON A CONJECTURE OF L. J. MORDELL

BY
MICHAEL A. MALCOLM

STAN-CS-70-184
NOVEMBER 1970



COMPUTER SCIENCE DEPARTMENT
School of Humanities and Sciences
STANFORD UNIVERSITY

Reproduced by
NATIONAL TECHNICAL
INFORMATION SERVICE
Springfield, Va 22151



STANFORD UNIVERSITY
LIBRARY
300 LATHAM DRIVE
STANFORD, CALIF. 94305

A NOTE ON A CONJECTURE OF L. J. MORDELL

by

Michael A. Malcolm

Abstract: A computer proof is described for a previously unsolved problem concerning the inequality

$$\sum_{i=1}^n x_i / (x_{i+1} + x_{i+2}) \geq \frac{n}{2} .$$

A NOTE ON A CONJECTURE OF L. J. MORDELL

by

Michael A. Malcolm ^{*}/

The function

$$S_n(\underline{x}) = \sum_{i=1}^n \frac{x_i}{x_{i+1} + x_{i+2}}, \quad (\text{subscripts mod } n) \quad (1)$$

where $x_i \geq 0$, $x_i + x_{i+1} > 0$, $i = 1, 2, \dots \pmod{n}$, has been studied by Shapiro, Mordell, and others (see [1-2, 6-11]). Let

$$\lambda(n) = \inf_{\underline{x}} S_n(\underline{x}) \quad (2)$$

Then

$$\lambda(n) \leq n/2 \quad (3)$$

H. S. Shapiro [10] suggested the verification of

$$\lambda(n) = n/2 \quad (4)$$

For $n \leq 6$, several authors (see [6]) proved the validity of (4).

Mordell [6] conjectured that (4) is false for all $n \geq 7$, but later [7] stated that (4) is true for $n = 7$. M. J. Lighthill (see [9]) and

A. Zulauf [11] proved that (4) is false for $n = 2k$, $k \geq 7$.

D. Z. Djokovic [2] proved that (4) is true for $n = 8$. P. H. Diananda [1] proved that (4) is false for $n = 2k+1$, $k \geq 13$. In the same paper he

^{*} This work was sponsored in part by the Office of Naval Research (NR 044-211) and by the National Science Foundation (GJ 408).

proved (i) that if (4) is false for some $n = m$, where m is odd, then (4) is false for all $n \geq m$, and (ii) that if (4) is true for some $n = m$, where m is even, then (4) is true for all $n \leq m$. Recently, P. Nowosad [8] proved (4) is true for $n = 10$, and thus (4) is true for all $n \leq 10$.

Therefore, the question remains open for $n = 12$ and for odd n ranging from 11 to 25. In this note I will describe a computer proof that (4) is false for $n = 25$.

The problem can be viewed as a multivariate constrained minimization of S_n . The constraints $x_i \geq 0$, $i = 1, 2, \dots, n$, can be removed by the simple change of variables

$$x_i = \theta_i^2, \quad i = 1, 2, \dots, n.$$

If the second constraint is violated, i.e., $x_i + x_{i+1} = 0$ for some i , the function (1) is undefined. Thus, the function

$$s_n(\theta) = \sum_{i=1}^n \frac{\theta_i^2}{\theta_{i+1}^2 + \theta_{i+2}^2},$$

where the indices are taken mod n , can be minimized using a standard method. For the case $n = 25$, this was done using a subroutine written by J. Alan George [3] on an IBM 360/67. The resulting values for x_i were:

$x_1^* = 5.11133$	$x_{13}^* = 2.47011$
$x_2^* = 0$	$x_{14}^* = 3.15375$
$x_3^* = 6.70077$	$x_{15}^* = 1.66622$
$x_4^* = 1.10052$	$x_{16}^* = 3.058$
$x_5^* = 6.70241$	$x_{17}^* = 0.980738$
$x_6^* = 2.57379$	$x_{18}^* = 3.12582$
$x_7^* = 5.91561$	$x_{19}^* = 0.400648$
$x_8^* = 3.33613$	$x_{20}^* = 3.44328$
$x_9^* = 4.60951$	$x_{21}^* = 0$
$x_{10}^* = 3.47149$	$x_{22}^* = 4.07589$
$x_{11}^* = 3.43693$	$x_{23}^* = 0$
$x_{12}^* = 3.33360$	$x_{24}^* = 4.8248$
	$x_{25}^* = 0$

To prove that $S_{25}(x^*)$ is less than 12.5, the calculation (1) was programmed in a language [4] in which the calculations are carried out in interval arithmetic (see Moore [5]). The inaccuracies due to number conversion and roundoff are automatically accounted for by the language translator. Thus, assuming the translator is properly programmed, the resulting interval is guaranteed to contain the true result. The program was run on the IBM 360/67 at Stanford University, giving

$$12.49847 < S_{25}(x^*) < 12.49851 .$$

Computer time, including program compilations, amounted to 12 sec., costing exactly \$1. Programming time, including typing at a terminal, amounted to about four hours.

This method has been applied to the cases $n = 2^3$ and $n = 21$.
In each case, an \underline{x} could not be found such that $S_n(\underline{x}) < n/2$.

Bibliography

- [1] Diananda, P. H., On a Cyclic Sum, Proc. Glasgow Math. Assoc. 6 (1963), 11-13.
- [2] Djokovic, D. Z., Sur une inégalité, Proc. Glasgow Math. Assoc. 6 (1963), 1-10.
- [3] George, J. Alan, FMINL - Local Minimum of a Real Scalar Function of Several Variables Using a Gradient Search Method, Computer Science Department Program Library, Stanford University, 1968.
- [4] Malcolm, M. A. and Katzman, J., RAT - Range Arithmetic Translator, (unpublished), 1969.
- [5] Moore, R. E., Interval Analysis, Prentice-Hall: Englewood Cliffs, N. J., 1966.
- [6] Mordell, L. J., On the Inequality $\sum_{r=1}^n x_r / (x_{r+1} + x_{r+2}) \geq \frac{n}{2}$ and Some Others, Abh. Math. Sem., Univ. of Hamburg 22 (1958), 221-240.
- [7] _____, Note on the Inequality $\sum_{r=1}^n x_r / (x_{r+1} + x_{r+2}) \geq \frac{n}{2}$, J. London Math. Soc. 37 (1962), 176-178.
- [8] Nowosad, Pedro, Isoperimetric Eigenvalue Problems in Algebras, Comm. Pure Appl. Math., 21 (1968), 401-465.
- [9] Rankin, R. A., An Inequality, Math. Gaz. 42 (1958), 39-40.
- [10] Shapiro, H. S., Problem 4603, Amer. Math. Monthly 61 (1954), 571.
- [11] Zulauf, Von A., Note on a Conjecture of L. J. Mordell, Abh. Math. Sem., Univ. of Hamburg 22 (1968), 240-241.

UNCLASSIFIED

Security Classification

DOCUMENT CONTROL DATA - R & D

Security classification of title, body of abstract and indexing annotation must be entered when the overall report is classified.

1. ORIGINATING ACTIVITY (Corporate author) Stanford University		2a. REPORT SECURITY CLASSIFICATION Unclassified	
		2b. GROUP	
3. REPORT TITLE A Note on a Conjecture of L. J. Mordell			
4. DESCRIPTIVE NOTES (Type of report and inclusive dates)			
5. AUTHOR(S) (First name, middle initial, last name) Michael A. Malcolm			
6. REPORT DATE October 1970		7a. TOTAL NO OF PAGES 5	7b. NO OF REFS 11
8a. CONTRACT OR GRANT NO NR 044-211		9a. ORIGINATOR'S REPORT NUMBER(S) 1	
b. PROJECT NO. NSF (GJ 408)			
c.		9b. OTHER REPORT NO(S) (Any other numbers that may be assigned this report)	
d.		None	
10. DISTRIBUTION STATEMENT Distribution of this document is unlimited			
11. SUPPLEMENTARY NOTES		12. SPONSORING MILITARY ACTIVITY Office of Naval Research National Science Foundation	
13. ABSTRACT A computer proof is described for a previously unsolved problem concerning a conjecture of L. J. Mordell.			

UNCLASSIFIED

Security Classification

14 KEY WORDS	LINK A		LINK B		LINK C	
	ROLE	WT	ROLE	WT	ROLE	WT
Number theory Interval arithmetic						