AD 740846

ARL 71-0321
DECEMBER 1971

Aerospace Research Laboratories

ON WEAK CONVERGENCE OF EMPIRICAL PROCESSES FOR RANDOM NUMBER OF INDEPENDENT STOCHASTIC VECTORS

PRANAB KUMAR SEN

UNIVERSITY OF NORTH CAROLINA
CHAPEL HILL, NORTH CAROLINA

CONTRACT NO. F33615-71-C-1927
PROJECT NO. 7071

Reproduced by
NATIONAL TECHNICAL
INFORMATION SERVICE
Springfield, Va. 22151

Approved for public release; distribution unlimited.

D D C
APR 12 1972
C

AIR FORCE SYSTEMS COMMAND
United States Air Force

18

NOTICES

When Government drawings, specifications, or other data are used for any purpose other than in connection with a definitely related Government procurement operation, the United States Government thereby incurs no responsibility nor any obligation whatsoever; and the fact that the Government may have formulated, furnished, or in any way supplied the said drawings, specifications, or other data, is not to be regarded by implication or otherwise as in any manner licensing the holder or any other person or corporation, or conveying any rights or permission to manufacture, use, or sell any patented invention that may in any way be related thereto.

Agencies of the Department of Defense, qualified contractors and other government agencies may obtain copies from the

Defense Documentation Center
Cameron Station
Alexandria, Virginia 22314

This document has been released to the

CLEARINGHOUSE
U. S. Department of Commerce
Springfield, Virginia 22151

for the sale to the public.

ACCESSION for
CFSTI WHITE SECTION
DDC BUFF SECTION
UNANNOUNCED
JUSTIFICATION
BY
DISTRIBUTION/AVAILABILITY CODES
DIST. AVAIL. and/or SPECIAL
A

Copies of ARL Technical Documentary Reports should not be returned to Aerospace Research Laboratories unless return is required by security considerations, contractual obligations or notices on a specified document.

AIR FORCE 6-1-72/500

Unclassified

Security Classification

DOCUMENT CONTROL DATA - R & D

(Security classification of title, body of abstract and indexing annotation must be entered when the overall report is classified)

1. ORIGINATING ACTIVITY (Corporate author) University of North Carolina Department of Biostatistics Chapel Hill, North Carolina 27514	2a. REPORT SECURITY CLASSIFICATION Unclassified 2b. GROUP M-2

3. REPORT TITLE

On Weak Convergence of Empirical Processes for Random Number of Independent Stochastic Vectors

4. DESCRIPTIVE NOTES (Type of report and inclusive dates)

Scientific Interim

5. AUTHOR(S) (First name, middle initial, last name)

Pranab Kumar Sen

6. REPORT DATE December 1971	7a. TOTAL NO. OF PAGES 15	7b. NO. OF REFS 7
8a. CONTRACT OR GRANT NO F 33615-71-C-1927 b. PROJECT NO 7071-00-12 c. DoD Element 61102 F d. DoD Subelement 681304	9a. ORIGINATOR'S REPORT NUMBER(S) 9b. OTHER REPORT NO(S) (Any other numbers that may be assigned this report) ARL 71-0321	

10. DISTRIBUTION STATEMENT

Approved for public release; distribution unlimited.

11. SUPPLEMENTARY NOTES TECH OTHER	12. SPONSORING MILITARY ACTIVITY Aerospace Research Laboratories(LB) Wright-Patterson AFB, Ohio 45433

13. ABSTRACT

By the use of a semi-martingale property of the Kolmogorov supremum, the results of Pyke [Proc. Cambridge Phil. Soc. 64 (1968), 155-160] on the weak convergence of the empirical process with random sample size are simplified and extended to the case of p(>1)-dimensional stochastic vectors.

DD FORM 1473

Unclassified

Security Classification

Unclassified
Security Classification

14 KEY WORDS	LINK A ROLE	LINK A WT	LINK B ROLE	LINK B WT	LINK C ROLE	LINK C WT
Weak Convergence						
Empirical Processes						
Stochastic Vectors						
Gaussian Processes						

Unclassified
Security Classification

*U.S. Government Printing Office: 1972 — 759-053/402

ARL 71-0321

ON WEAK CONVERGENCE OF EMPIRICAL PROCESSES FOR RANDOM NUMBER OF INDEPENDENT STOCHASTIC VECTORS

PRANAB KUMAR SEN

UNIVERSITY OF NORTH CAROLINA
CHAPEL HILL, NORTH CAROLINA

DECEMBER 1971

CONTRACT F33615-71-C-1927
PROJECT 7071

Approved for public release; distribution unlimited.

AEROSPACE RESEARCH LABORATORIES
AIR FORCE SYSTEMS COMMAND
UNITED STATES AIR FORCE
WRIGHT-PATTERSON AIR FORCE BASE, OHIO

FOREWORD

This is an interim report of the work done under Contract F 33615-71-C-1927 by the University of North Carolina. The work done by Pranab Kumar Sen in this report is sponsored by the Aerospace Research Laboratories under the above contract; it was accomplished on Project 7071, "Research in Applied Mathematics" and is technically monitored by P. R. Krishnaiah of the Aerospace Research Laboratories.

ABSTRACT

By the use of a semi-martingale property of the Kolmogorov supremum, the results of Pyke [Proc. Cambridge Phil. Soc. 64 (1968), 155-160] on the weak convergence of the empirical process with random sample size are simplified and extended to the case of p(>1)-dimensional stochastic vectors.

TABLE OF CONTENTS

Section	Pages
1. Introduction	1
2. Some Preliminary Results	2
3. The Proof of Theorem 1	7
References	10

1. Introduction. Consider a sequence $\{\underset{\sim}{X}_i=(X_{i1},\ldots,X_{ip})', i\geq 1\}$ of independent and identically distributed stochastic $p(\geq 1)$-vectors, defined on a probability space $(\Omega,\mathcal{A},P)$, with each $\underset{\sim}{X}_i$ having a continuous distribution function (df) $F(\underset{\sim}{x})$, $\underset{\sim}{x}\epsilon R^p$, the p-dimensional Euclidean space. We denote the marginal df of X_{ij} by $F_{[j]}$, let $Y_{ij} = F_{[j]}(X_{ij})$, $j=1,\ldots,p$, $\underset{\sim}{Y}_i = (Y_{i1},\ldots,Y_{ip})'$, $i\geq 1$, $\underset{\sim}{t}=(t_1,\ldots,t_p)'$, and define

$$G(\underset{\sim}{t}) = P\{Y_{ij}\leq t_j,\ j=1,\ldots,p\},\ \underset{\sim}{t}\epsilon E^p,$$

where $E^p = \{\underset{\sim}{t}:\ 0\leq t_j\leq 1,\ j=1,\ldots,p\}$. Then, the empirical df for $\underset{\sim}{Y}_1,\ldots,\underset{\sim}{Y}_n$ is defined by

$$G_n(\underset{\sim}{t}) = n^{-1}\sum_{i=1}^{n} c(\underset{\sim}{t}-\underset{\sim}{Y}_i),\ \underset{\sim}{t}\epsilon E^p, \tag{1.2}$$

where $c(\underset{\sim}{u})=1$ iff $u_j\geq 0$, $j=1,\ldots,p$; otherwise, $c(\underset{\sim}{u})=0$. Consider then the empirical process

$$W_n(\underset{\sim}{t}) = n^{\frac{1}{2}}[G_n(\underset{\sim}{t})-G(\underset{\sim}{t})],\ \underset{\sim}{t}\epsilon E^p, \tag{1.3}$$

and denote by

$$W_n = \{W_n(\underset{\sim}{t}):\ \underset{\sim}{t}\epsilon E^p\}. \tag{1.4}$$

For $p=1$, it is well-known that W_n weakly converges to a Brownian motion $W^o = \{W^o(t):\ 0\leq t\leq 1\}$. For $p\geq 1$, on the space $D^p[0,1]$ of all real functions on E^p with no discontinuities of the second kind, W_n converges in distribution (in the (extended) Skorokhod J_1-topology) to an appropriate Gaussian function,

say, $W = \{W(\underset{\sim}{t}): \ \underset{\sim}{t} \in E^p\}$, where $E[W(\underset{\sim}{t})]=0$, and

$$(1.5) \qquad E[W(\underset{\sim}{s})W(\underset{\sim}{t})] = G(\underset{\sim}{t} \wedge \underset{\sim}{s}) - G(\underset{\sim}{t})G(\underset{\sim}{s}), \ \underset{\sim}{t}, \ \underset{\sim}{s} \in E^p,$$

and $\underset{\sim}{t} \wedge \underset{\sim}{s} = (t_1 \wedge s_1, \ldots, t_p \wedge s_p)$, where $a \wedge b = \min(a,b)$; we refer to Neuhaus (1971) who also reviews the earlier literature.

Let now $\{N_\nu, \ \nu \geq 1\}$ be a sequence of positive integer-valued random variables, such that

$$(1.6) \qquad \nu^{-1} N_\nu \to \xi, \text{ in probability, as } \nu \to \infty,$$

where ξ is a positive random variable defined on the same probability space $(\Omega, \mathcal{A}, P)$.

For p=1, Pyke (1968) has shown that under (1.6), W_{N_ν} converges in law to W^o; his result is extended here to the general multivariate case.

<u>Theorem 1.</u> <u>Under</u> (1.6), <u>for every</u> $p \geq 1$,

$$W_{N_\nu} \xrightarrow{\mathcal{B}} W, \ \underline{\text{in the Skorokhod } J_1\text{-typology on}} \ D^p[0,1].$$

The proof is outlined in section 3. Whereas, Pyke's arguments rely heavily on the properties of an equivalently defined Poisson process (which may become quite complicated for p>1), our approach is based on a simple semi-martingale property of the Kolmogorov supremum, which is considered first in section 2.

2. Some preliminary results.

For two real valued functions $Z(\underset{\sim}{t})$ and $Z^*(\underset{\sim}{t})$, defined on E^p, we let

$$(2.1) \qquad \rho(Z,Z^*) = \text{Sup}\{|Z(\underset{\sim}{t}) - Z^*(\underset{\sim}{t})|: \ \underset{\sim}{t} \in E^p\},$$

and for every $n \geq 1$, let

(2.2) $$W_n^+ = \mathrm{Sup}\{W_n(\underset{\sim}{t}):\ \underset{\sim}{t}\in E^p\},\ W_n^- = \mathrm{Sup}\{-W_n(\underset{\sim}{t}):\ \underset{\sim}{t}\in E^p\};$$

(2.3) $$W_n^* = \max\{W_n^+, W_n^-\} = \rho(W_n, 0).$$

Let $\mathcal{F}_n$ be the σ-field generated by $\{X_1,\ldots,X_n\}$, so that $\mathcal{F}_n$ is $\uparrow$ in n (≥ 1). Then, we have the following

<u>Lemma 2.1.</u> $\{n^{\frac{1}{2}}W_n^+,\ \mathcal{F}_n;\ n\geq 1\}$ <u>and</u> $\{n^{\frac{1}{2}}W_n^-,\ \mathcal{F}_n;\ n\geq 1\}$ <u>are both non-negative semi-martingale sequences.</u>

<u>Proof.</u> We only prove the result for W_n^+, as the other follows similarly. Note that W_n^+ is, by definition, non-negative [as $W_n(\underset{\sim}{t})=0$ for $\underset{\sim}{t}=\underset{\sim}{0}$ or $\underset{\sim}{t}=\underset{\sim}{1}$]. Let $\underset{\sim}{t}_n^o(\in E^p)$ be a point such that

(2.4) $$W_n^+ = W_n(\underset{\sim}{t}_n^o);\ \underset{\sim}{t}_n^o \text{ need not be unique.}$$

Then, by definition,

(2.5) $$(n+1)^{\frac{1}{2}}W_{n+1}^+ = (n+1)^{\frac{1}{2}} \sup_{\underset{\sim}{t}\in E^p} W_{n+1}(\underset{\sim}{t}) \geq (n+1)^{\frac{1}{2}}W_{n+1}(\underset{\sim}{t}_n^o),$$

so that, by (1.2), (1.3) and (2.5), for every $n\geq 1$,

(2.6) $$E\{(n+1)^{\frac{1}{2}}W_{n+1}^+ \mid \mathcal{F}_n\} \geq E\{(n+1)^{\frac{1}{2}}W_{n+1}(\underset{\sim}{t}_n^o) \mid \mathcal{F}_n\}$$

$$= \sum_{i=1}^{n+1} E\{[c(\underset{\sim}{t}_n^o-\underset{\sim}{Y}_i) - G(\underset{\sim}{t}_n^o)] \mid \mathcal{F}_n\}$$

$$= \sum_{i=1}^{n} [c(\underset{\sim}{t}_n^o-\underset{\sim}{Y}_i) - G(\underset{\sim}{t}_n^o)] + E\{[c(\underset{\sim}{t}_n^o-\underset{\sim}{Y}_{n+1}) - G(\underset{\sim}{t}_n^o)] \mid \mathcal{F}_n\}$$

$$= n^{\frac{1}{2}}W_n^+ + 0 = n^{\frac{1}{2}}W_n^+,$$

as, given $\mathcal{F}_n$, $c(\underset{\sim}{t}_n^o - \underset{\sim}{Y}_{n+1})$ assumes the values 1 and 0 with respective conditional probabilities $G(\underset{\sim}{t}_n^o)$ and $1-G(\underset{\sim}{t}_n^o)$. Q.E.D.

Lemma 2.2. For every $n \geq 1$, there exist two positive constants c_0 and c_1, independent of n, such that

$$E\{(W_n^+)^2\} \leq c_0/c_1 \text{ and } E\{(W_n^-)^2\} \leq c_0/c_1. \tag{2.7}$$

Proof. By partial integration,

$$E\{(W_n^+)^2\} = 2\int_0^\infty xP\{W_n^+ > x\}dx, \tag{2.8}$$

where by Theorem 1 of Kiefer and Wolfowitz (1958), for all $n \geq 1$,

$$P\{W_n^+ > x\} < c_0 \exp\{-c_1 x^2\} \text{ for all } x \geq 0. \tag{2.9}$$

Consequently, by (2.8) and (2.9), $E\{(W_n^+)^2\} \leq c_0/c_1$. The other result follows similarly.

Lemma 2.3. For every $\varepsilon > 0$, there exists a positive $K_\varepsilon (<\infty)$, such that for every $n \geq 1$,

$$P\{\max_{1 \leq k \leq n} (k/n)^{\frac{1}{2}} \rho(W_k, 0) > K_\varepsilon\} < \varepsilon. \tag{2.10}$$

Proof. By (2.3), for every $\varepsilon > 0$,

$$\begin{aligned} &P\{\max_{1 \leq k \leq n} (k/n)^{\frac{1}{2}} \rho(W_k, 0) > K_\varepsilon\} \\ &\leq P\{\max_{1 \leq k \leq n} k^{\frac{1}{2}} W_k^+ > n^{\frac{1}{2}} K_\varepsilon\} + P\{\max_{1 \leq k \leq n} k^{\frac{1}{2}} W_k^- > n^{\frac{1}{2}} K_\varepsilon\}, \end{aligned} \tag{2.11}$$

and hence, by Lemma 2.1 along with the Kolmogorov inequality for semi-martingales

[viz., Feller (1966, p. 235)], the right and side of (2.11) is bounded above by

$$(2.12) \qquad (nK_\varepsilon^2)^{-1}[nE\{(W_n^+)^2\} + nE\{(W_n^-)^2\}]$$

$$= [E\{(W_n^+)^2\} + E\{(W_n^-)^2\}]/K_\varepsilon^2$$

$$\leq 2c_0/c_1K_\varepsilon^2, \text{ by Lemma 2.2.}$$

The proof then follows by selecting $K_\varepsilon > [2c_0/c_1\varepsilon]^{\frac{1}{2}}$. Q.E.D.

<u>Lemma 2.4</u>. (<u>Uniform continuity in probability</u>). <u>For every</u> $\varepsilon>0$ <u>and</u> $\eta>0$, <u>there exists a</u> $\delta>0$ <u>and an</u> $n_0(\varepsilon,\eta)$, <u>such that for</u> $n\geq n_0(\varepsilon,\eta)$,

$$(2.13) \qquad P\{\max_{k:\ |k-n|<\delta n} \rho(W_k,W_n) > \varepsilon\} < \eta.$$

<u>Proof</u>. Proceeding as in the proof of Theorem 2.1 of Pyke (1968), namely, as in his (2.7) through (2.10), we are only to show that as $n\to\infty$,

$$(2.14) \qquad \max_{1\leq k\leq n} (k/n)^{\frac{1}{2}} \rho(W_k,0) = O_p(1),$$

$$(2.15) \qquad \rho(W_n,0) = \text{Sup}\{|W_n(\underset{\sim}{t})|:\ \underset{\sim}{t}\in E^p\} = O_p(1).$$

Now, (2.14) has already been proved in Lemma 2.3, while by Theorem 3.1 of Neuhaus (1971) along with his treatment on the weak convergence of W_n to W, it follows that for every $\varepsilon>0$, there exists a positive $M_\varepsilon(<\infty)$, such that

$$(2.16) \qquad \lim_{n\to\infty} P\{\rho(W_n,0) > M_\varepsilon\}$$

$$= P\{\rho(W,0) > M_\varepsilon\} < \varepsilon';\quad 0<\varepsilon'<\varepsilon,$$

which completes the proof of the lemma.

We now show that $\{W_n\}$ is a mixing sequence in the sense of Rényi (1958). This follows by defining

$$(2.17) \qquad W_n'(\underset{\sim}{t}) = n^{-1}\{\textstyle\sum_{i=k_n}^{n} [c(\underset{\sim}{t}-Y_i)-G(\underset{\sim}{t})]\}, \quad \underset{\sim}{t}\in R^p,$$

where $k_n \to \infty$ but $n^{-\frac{1}{2}} k_n \to 0$ as $n\to\infty$, and noting that

$$(2.18) \qquad \rho(W_n, W_n') \leq n^{-\frac{1}{2}} k_n \to 0 \quad \text{as} \quad n\to\infty.$$

Consequently, proceeding as in the proof of Lemma 3 of Blum, Hanson and Rosenblatt (1963), we obtain from Lemma 2.4, the following.

<u>Lemma 2.5.</u> <u>If</u> $A\in\mathcal{A}$, <u>then for every</u> $\varepsilon>0$ <u>and</u> $\eta>0$, <u>there exists a</u> $\delta>0$, <u>such that as</u> $n\to\infty$,

$$(2.19) \qquad P\{\max_{k:\ |k-n|<\delta n} \rho(W_k, W_n) > \varepsilon | A\} < \eta.$$

Let us now define

$$(2.20) \qquad \omega_\delta(W_n) = \operatorname{Sup}\{|W_n(\underset{\sim}{t})-W_n(\underset{\sim}{t}')|: \ |\underset{\sim}{t}-\underset{\sim}{t}'|<\delta\}.$$

Then, from the results of section 5 of Neuhaus (1971), for every $\varepsilon>0$ and $\eta>0$, there exists a $\delta>0$, such that

$$(2.21) \qquad \lim_{n\to\infty} P\{\omega_\delta(W_n) > \varepsilon\} < \eta.$$

Hence, again using (2.18) and Rényi's (1958) idea of mixing sequence of sets, we have for $A\in\mathcal{A}$,

$$(2.22) \qquad \lim_{n\to\infty} P\{\omega_\delta(W_n) > \varepsilon | A\} < \eta.$$

3. The proof of Theorem 1. Let $[s]$ denote the largest integer $\leq s$. Then for $\varepsilon>0$,

$$(3.1)\qquad P\{\rho(W_{N_\nu}, W_{[\nu\xi]}) > \varepsilon\}$$

$$\leq P\{|\nu^{-1}N_\nu-\delta|\geq\delta'\} + P\{\rho(W_{N_\nu}, W_{[\nu\xi]}) > \varepsilon,\ |\nu^{-1}N_\nu-\xi|<\delta'\}$$

$$\leq P\{|\nu^{-1}N_\nu-\xi|\geq\delta'\} + P\{\max_{k:\ |k-[\nu\xi]|<\delta'\nu} \rho(W_k, W_{[\nu\xi]})>\varepsilon\},\quad \delta'>0.$$

Thus, if $\xi=c$, a positive constant, with probability one [the case treated in Pyke (1968)], it readily follows from (1.6) and (2.19) that the right hand side of (3.1) can be bounded by $\eta(>0)$ by a proper choice of $\delta'>0$. The proof of the theorem then follows by noting that by the results of Neuhaus (1971), as $\nu\to\infty$,

$$(3.2)\qquad W_{[\nu c]} \xrightarrow{\mathcal{D}} W, \text{ in the Skorokhod } J_1\text{-typology on } D^p[0,1].$$

So, in the sequel, we consider the general case of ξ having an arbitrary distribution on $(0,\infty)$. For every $\eta>0$, there exists an $a_o = a_o(\eta)$, such that

$$(3.3)\qquad P\{\xi \leq a_o(\eta)\} < \frac{1}{4}\eta.$$

Consider then a countable set of events

$$(3.4)\qquad A_h = \{\xi:\ a_o(\eta)+h\delta'<\xi\leq a_o(\eta)+(h+1)\delta'\},\quad h=0,1,\ldots,$$

and let $a_h = a_h(\delta',\eta) = a_o(\eta)+(h+\frac{1}{2})\delta'$, $h=0,1,\ldots$ Then, the right hand side of (3.1) is bounded above by

$$(3.5)\qquad P\{|\nu^{-1}N_\nu-\xi|\geq\delta'\} + P\{\xi \leq a_o(\eta)\} +$$

$$\sum_{h=0}^{\infty} P\{\max_{k:\ |k-[\nu\xi]|<\nu\delta'} \rho(W_k,W_{[\nu\xi]})>\varepsilon|A_h\}P(A_h)$$

$$\leq P\{|\nu^{-1}N_\nu-\xi|\geq\delta'\} + P\{\xi \leq a_o(\eta)\} +$$

$$\sum_{h=1}^{\infty} P\{\max_{k:\ |k-[\nu a_h]|<\frac{3}{2}\delta'\nu} \rho(W_k,W_{[\nu a_h]})>\varepsilon|A_h\}P(A_h).$$

Now, by (1.6) and (3.3), the first two terms on the right hand side of (3.5) are bounded by $\eta/4$ by proper choice of $\delta'>0$, while by (2.19), the last term can also be bounded by $\eta/2$, by proper choice of $\delta'(>0)$, as $\nu a_h\to\infty$ with $\nu\to\infty$, for every $h\geq 0$. Consequently, as $\nu\to\infty$,

$$(3.6)\qquad \rho(W_N,W_{[\nu\xi]}) \xrightarrow{P} 0.$$

Thus, it suffices to show that as $\nu\to\infty$,

$$(3.7)\qquad W_{[\nu\xi]} \xrightarrow{\mathcal{D}} W, \text{ in the Skorokhod } J_1\text{-topology on } D^p[0,1].$$

Now, (3.2) implies the convergence of the finite dimensional distributions of $\{W_\nu\}$ to those of W, while (2.13) implies that for any $\underline{t}\in E^p$, $\{W_\nu(\underline{t}): |\nu-n|<\delta n\}$ satisfy the "uniform continuity in probability" condition; these two conditions, in accordance with Theorem 1 of Mogyorodi (1965), imply the convergence of the finite dimensional distributions of $\{W_{[\nu\xi]}\}$ to those of W. So, to complete the proof of the theorem, we require to establish the 'tightness' property of $\{W_{[\nu\xi]}\}$ when $\nu\to\infty$. By (1.7) and (3.5) of Neuhaus (1971), it suffices to show that for every $\varepsilon>0$ and $\eta>0$, there exists a positive δ, such that as $\nu\to\infty$,

$$(3.8)\qquad P\{\omega_\delta(W_{[\nu\xi]}) \geq \varepsilon\} < \eta.$$

To show this, we note that for every $\varepsilon>0$, $\delta'>0$,

$$(3.9) \qquad P\{\omega_{\delta'}(W_{[\nu\xi]}) \geq \varepsilon\}$$

$$\leq P\{\xi \leq a_o(\eta)\} + P\{\omega_{\delta'}(W_{[\nu\xi]}) \geq \varepsilon,\ \xi > a_o(\eta)\}$$

$$= P\{\xi \leq a_o(\eta)\} + \sum_{h=0}^{\infty} P\{\omega_{\delta'}(W_{[\nu\xi]}) \geq \varepsilon | A_h\} P(A_h)$$

$$\leq P\{\xi \leq a_o(\eta)\} + \sum_{h=0}^{\infty} P\{\omega_{\delta'}(W_{[\nu a_h]}) \geq \tfrac{1}{3}\varepsilon | A_h\} P(A_h)$$

$$+ \sum_{h=0}^{\infty} P\{\rho(W_{[\nu\xi]}, W_{[\nu a_h]}) \geq \tfrac{1}{3}\varepsilon | A_h\} P(A_h)$$

$$\leq P\{\xi \leq a_o(\eta) + \sum_{h=0}^{\infty} P\{\omega_{\delta'}(W_{[\nu a_h]}) \geq \tfrac{1}{3}\varepsilon | A_h\} P(A_h)$$

$$+ \sum_{h=0}^{\infty} P\{\max_{k:\ |k-[\nu a_h]|<\frac{1}{2}\delta'\nu} \rho(W_k, W_{[\nu a_h]}) \geq \tfrac{1}{3}\varepsilon | A_h\} P(A_h),$$

which, by (3.3), (2.22) and (2.19), can be made smaller than $\eta(>0)$ by a proper choice of $\delta'(>0)$. Q.E.D.

REFERENCES

[1] BLUM, J. R., HANSON, D. L., AND ROSENBLATT, J. I. (1963). On the central limit theorem for the sum of a random number of independent random variables. Zeit. Wahrsch Verw. Geb. 1, 389-393.

[2] FELLER, W. (1966). An introduction to Probability Theory and its Applications, Vol 2. John Wiley, New York.

[3] KIEFER, J., AND WOLFOWITZ., J. (1958). On the deviations of the empiric distribution function of vector chance variables. Trans. Amer. Math. Soc. 87, 173-186.

[4] MOGYORODI, J. (1965). Limit distributions for sequences of random variables with random indices. Trans. 4th Prague Confer. Infor. Th. Statist. Dec. fn. Random. Proc. 463-470.

[5] NEUHAUS, G. (1971). On weak convergence of stochastic processes with multidimensional time parameter. Ann. Math. Statist. 42, 1285-1295.

[6] PYKE, R. (1968). The weak convergence of the empirical process with random sample size. Proc. Cambridge Phil. Soc. 64, 155-160.

[7] RÉNYI, A. (1958). On mixing sequences of sets. Acta Math. Acad. Sc. Hungar. 9, 215-228.