

AD 740975

DDC
REFILED
MAY 4 1972
REGITLD
B

Reproduced by
NATIONAL TECHNICAL
INFORMATION SERVICE
Springfield, Va. 22151

UNCLASSIFIED

Security Classification

DOCUMENT CONTROL DATA - R & D

Security classification of title, body of abstract and index... must be entered when the overall report is classified

1. ORIGINATING ACTIVITY (Corporate author) NAVAL MEDICAL RESEARCH INSTITUTE NATIONAL NAVAL MEDICAL CENTER BETHESDA, MARYLAND 20014	2a. REPORT SECURITY CLASSIFICATION UNCLASSIFIED
	2b. GROUP

3. REPORT TITLE
STUDIES ON THE MECHANISM OF FASTING HYPERBILIRUBINEMIA

4. DESCRIPTIVE NOTES (Type of report and inclusive dates)
Medical research progress report

5. AUTHOR(S) (First name, middle initial, last name)
Joseph R. BLOOMER, M.D., Peter V. BARRETT, M.D., F. Lee RODKEY, Ph.D., and Nathaniel I. BERLIN, M.D., Ph.D.

6. REPORT DATE 1971	7a. TOTAL NO OF PAGES 9	7b. NO OF REFS 30
-------------------------------	-----------------------------------	-----------------------------

8a. CONTRACT OR GRANT NO. b. PROJECT NO. c. d.	9a. ORIGINATOR'S REPORT NUMBER(S) M4306-02-4030, Report No. 4
	9b. OTHER REPORT NO(S) (Any other numbers that may be assigned this report)

10. DISTRIBUTION STATEMENT
THIS DOCUMENT HAS BEEN APPROVED FOR PUBLIC RELEASE AND SALE; ITS DISTRIBUTION IS UNLIMITED.

11. SUPPLEMENTARY NOTES GASTROENTEROLOGY Vol. 61, No. 4, Part 1	12. SPONSORING MILITARY ACTIVITY BUREAU OF MEDICINE AND SURGERY (NAVY) WASHINGTON, D. C.
---	--

13. ABSTRACT The total plasma bilirubin concentration (Br) increased by 23 to 334% in 12 individuals (10 healthy volunteers and 2 patients with Gilbert's syndrome) who fasted for 1 to 3 days. Eighty-six per cent of the rise in Br was due to an increase in plasma unconjugated bilirubin. The percentage increase in Br did not correlate with the magnitude of the base line concentration. Studies with bilirubin-³H in 5 subjects showed that the hepatic clearance of bilirubin from the plasma (C_{Br}) was reduced by 28 to 54% during fasting, accounting for the increase in Br. The ratio of the plasma bilirubin turnover during fasting to that in base line state was 0.99 ± 0.16 (mean ± SD) for these five individuals, indicating that increased plasma bilirubin turnover did not contribute to the rise in Br. In 5 additional subjects the mean change in carbon monoxide production with fasting was only +24%, whereas Br increased by 168%. Further studies were done to determine why C_{Br} decreases with fasting. The plasma disappearance rate of indocyanine green in 5 subjects after a 48-hr fast was unchanged from base line. Nine homozygous Gunn rats had a 41 ± 10% (mean ± SE) increase in Br over control values with 48 hr of fasting. Bilirubin-³H clearance studies in 2 homozygous Gunn rats demonstrated 43 and 44% decreases in the whole body clearance of bilirubin during fasting, with return to control rates on refeeding. Multicompartmental analysis of the human bilirubin clearance data showed that the ratio of the hepatic bilirubin pool to plasma bilirubin pool decreased during fasting. These results are most consistent with the hypothesis that C_{Br} decreases with fasting because of reduced hepatic ability to extract bilirubin from the plasma.

UNCLASSIFIED

Security Classification

14. KEY WORDS	LINK A		LINK B		LINK C	
	ROLE	WT	ROLE	WT	ROLE	WT
Fasting Hyperbilirubinemia Carbon monoxide production Bilirubin glucuronyl transferase						

DD FORM 1473

NOV 65

(BACK)

3/73 3102-014-0000

Security Classification

4-3100

STUDIES ON THE MECHANISM OF FASTING HYPERBILIRUBINEMIA

JOSEPH R. BLOOMER, M.D., PETER V. BARRETT, M.D., F. LEE RODKEY, Ph.D.,
AND NATHANIEL I. BERLIN, M.D., Ph.D.

*National Cancer Institute, National Institutes of Health; Naval Medical Research Institute,
Bethesda, Maryland; and Harbor General Hospital, Torrance, California*

The total plasma bilirubin concentration (Br) increased by 23 to 334% in 12 individuals (10 healthy volunteers and 2 patients with Gilbert's syndrome) who fasted for 1 to 3 days. Eighty-six per cent of the rise in Br was due to an increase in plasma unconjugated bilirubin. The percentage increase in Br did not correlate with the magnitude of the base line concentration. Studies with bilirubin-³H in 5 subjects showed that the hepatic clearance of bilirubin from the plasma (C_{br}) was reduced by 28 to 54% during fasting, accounting for the increase in Br. The ratio of the plasma bilirubin turnover during fasting to that in base line state was 0.99 ± 0.16 (mean \pm SD) for these five individuals, indicating that increased plasma bilirubin turnover did not contribute to the rise in Br. In 5 additional subjects the mean change in carbon monoxide production with fasting was only +24%, whereas Br increased by 188%. Further studies were done to determine why C_{br} decreases with fasting. The plasma disappearance rate of indocyanine green in 5 subjects after a 48-hr fast was unchanged from base line. Nine homozygous Gunn rats had a $41 \pm 10\%$ (mean \pm SE) increase in Br over control values with 48 hr of fasting. Bilirubin-³H clearance studies in 2 homozygous Gunn rats demonstrated 43 and 44% decreases in the whole body clearance of bilirubin during fasting, with return to control rates on refeeding. Multicompartmental analysis of the human bilirubin clearance data showed that the ratio of the hepatic bilirubin pool to plasma bilirubin pool decreased during fasting. These results are most consistent with the hypothesis that C_{br} decreases with fasting because of reduced hepatic ability to extract bilirubin from the plasma.

The effect of caloric intake on the serum bilirubin concentration was noted as early as 1906 by Gilbert and Herscher.¹ They observed that the serum bilirubin concen-

tration was higher when individuals were fasting. More recently, Barrett² and Felscher et al.³ have shown that caloric restriction for 24 to 48 hr causes a significant in-

Received March 12, 1971. Accepted May 27, 1971.

Part of this work was presented at the annual meeting of the American Federation for Clinical Research, May 1971, in Atlantic City, New Jersey.

Address requests for reprints to: Dr. Nathaniel I. Berlin, National Cancer Institute, Metabolism Branch, Building 10, Room 4N117, National Institutes of Health, Bethesda, Maryland 20014.

This investigation was supported in part by United

States Public Health Service Grant AM-12306 and CMC RR-00425 from the National Institutes of Health (Dr. Barrett) and by the Bureau of Medicine and Surgery, Research Task M4306.02.40-99BAK9 (Dr. Rodkey). The opinions or assertions contained herein are those of the authors and are not to be construed as official or reflecting the views of the Navy Department or the naval service at large.

Dr. Bloomer's present address is: Liver Unit, Yale

crease in the serum bilirubin concentration of both normal individuals and patients with a variety of types of hepatic dysfunction.

This investigation was designed to determine the mechanism responsible for the hyperbilirubinemia of fasting. The results indicate that hepatic clearance of bilirubin from the plasma decreases during fasting, causing the increase in the plasma bilirubin concentration. Additional studies were done to determine why hepatic bilirubin clearance is altered by fasting.

Methods

Plasma bilirubin concentrations in human subjects were measured by the method of Weber and Schalm.⁴ Plasma bilirubin concentrations in homozygous Gunn rats were measured by the method of Malloy and Evelyn⁵ on 0.05 to 0.15 ml of blood collected from the tail vein in heparinized micro-blood collecting tubes (Arthur H. Thomas Co., Philadelphia, Pa.).

The plasma disappearance rate of bilirubin-³H in human subjects was determined following intravenous administration of 15 to 30 μ c of bilirubin-³H in 0.5 mg of bilirubin,⁶ using the methods of Barrett et al.⁷ and Berk et al.⁸ The following indices of bilirubin metabolism were calculated from parameters of the plasma disappearance curve of bilirubin-³H and the plasma concentration of unconjugated bilirubin (Br):⁹ (1) C_{10} , the volume of plasma cleared of unconjugated bilirubin each minute by the liver; (2) BRT, the plasma bilirubin turnover in milligrams per day; and (3) the percentage retention of labeled bilirubin in the plasma at 4 hr.

$$C_{10} \text{ (ml per min)} = \frac{\text{initial distribution volume of bilirubin-}^3\text{H} \times 1}{\text{area under bilirubin-}^3\text{H plasma disappearance curve}} \quad (1)$$

$$\text{BRT (mg per day)} = C_{10} \text{ (ml per min)} \times \frac{\text{Br}}{100} \text{ (mg per ml)} \times 1400 \text{ (min per day)} \quad (2)$$

New Haven Hospital, New Haven, Connecticut 06511.

The authors thank Mr. John Vergalla, Harold Collison, and John O'Neal for their assistance in performing these studies, and Miss Teri Mills for preparation of the manuscript.

The bilirubin-³H clearance data were also analyzed by multicompartamental analysis in terms of a three-pool model previously described by Berk et al.⁸ In this model the plasma pool of unconjugated bilirubin is considered to exchange with a hepatic bilirubin pool and extravascular extrahepatic bilirubin pool. The pool sizes and fractional transfer rates between pools were determined.

Labeled bilirubin clearance studies in homozygous Gunn rats were done with bilirubin-³H dissolved in male Sprague-Dawley rat plasma. The solution (0.5 to 1.0 ml), containing 0.1 to 0.2 μ c of bilirubin, was injected into a tail vein. Plasma samples for radioactivity were obtained from blood collected at the end of the tail in heparinized micro-blood collecting tubes.

The rate of carbon monoxide production in human subjects was measured in a closed re-breathing system. The methods of Collison et al.⁹ and Rodkey et al.¹⁰ were used.

The plasma disappearance rate of indocyanine green (ICG) in human subjects was determined by the method of Cherrick et al.¹¹ following intravenous administration (6.5 mg per kg of body weight).

The water content of rat liver was determined as the difference between the wet weight and the dry weight of tissue. The protein content was measured on trichloroacetic acid precipitates of liver homogenates by the method of Lowry et al.¹² Bilirubin glucuronyl transferase activity in liver homogenates from male Sprague-Dawley rats was measured by the method of Black et al.¹³

Subjects

Twelve individuals participated in these studies. All individuals except H. G. and K. S. were healthy volunteers. H. G. had Gilbert's syndrome and an undefined congenital hemolytic anemia (red cell life span measured with tritiated diisopropylfluorophosphate was 30 days). K. S. had Gilbert's syndrome. Informed consent was obtained from each individual.

During periods of fasting both human subjects and rats underwent total caloric restriction, with only water intake allowed. During base line studies the human subjects were on an *ad libitum* diet, and rats were fed unlimited Purina rat chow.

Results

The plasma bilirubin concentration of each individual rose with caloric restriction (tables 1 to 3). The percentage increase in the bilirubin concentration did not correlate

with the magnitude of the base line value ($r = 0.19$). Eighty-six per cent (range: 69 to 93%) of the increase in the total plasma bilirubin concentration was due to an increase in the unconjugated fraction. The greatest increase occurred during the first 24 to 34 hr of fasting, with a smaller increase over the subsequent 24 hr of fasting. The plasma bilirubin concentration in each individual returned to the base line value within 24 to 48 hr of refeeding.

The results of bilirubin- ^3H clearance studies in 5 subjects in the base line state as compared with fasting are shown in table 1. The fasting bilirubin- ^3H clearance study was started after 40 to 57 hr of total caloric restriction. Caloric restriction was continued for the duration of the study (an additional 24 to 30 hr). In four of the individuals the plasma bilirubin concentration had reached a new constant level by the time the fasting bilirubin- ^3H clearance study was started (the ratios of the plasma bilirubin concentrations at the end of the study to those at the beginning were 0.89, 0.99, 1.00, and 1.09). This indicates that a new steady state with respect to the plasma bilirubin concentration was present during the fasting bilirubin- ^3H clearance study. The plasma bilirubin concentration in H.

G. was constant over the first 15 hr of the fasting bilirubin- ^3H clearance study, but then increased by 28% during the last 9 hr of the study. However, the measurement of hepatic bilirubin clearance was well determined in this subject (uncertainty in the measurement was $\pm 5\%$).

The difference between the plasma disappearance of bilirubin- ^3H in the base line state compared with that during fasting is shown in figure 1. Indices of bilirubin metabolism obtained from parameters of the curves are given in table 1. In each individual the hepatic clearance of unconjugated bilirubin decreased during fasting, and the plasma retention of labeled bilirubin at 4 hr increased. The average uncertainty in the clearance measurement for the 5 subjects was $\pm 8\%$, both during base line state and fasting.* The decreases in hepatic bilirubin clearance (by 28 to 54%) accounted for the increases in the plasma unconjugated bilirubin concentration. The ratio of the plasma bilirubin turnover during fasting to that during base line was 0.99 ± 0.16 (mean \pm SD) for the 5 subjects, indicating that an increase in plasma bilirubin turnover did not contribute to the increase in the plasma unconjugated bilirubin concentration. The increase in

TABLE 1. Bilirubin- ^3H clearance data*

Subject	Study	Plasma bilirubin concentration ^b		Hepatic bilirubin clearance	Plasma 4 hr bilirubin- ^3H retention	Plasma bilirubin turnover
		Conjugated	Unconjugated			
		mg/100 ml		ml/min		ml/day
C. G.	Base line	0.02 \pm 0.01	0.39 \pm 0.06	48	4	270
	Fasting	0.08 \pm 0.02	0.86 \pm 0.09	22	8	275
L. S.	Base line	0.08 \pm 0.01	0.81 \pm 0.06	28	11	330
	Fasting	0.24 \pm 0.01	1.51 \pm 0.06	18	17	391
B. C.	Base line	0.05 \pm 0.01	0.97 \pm 0.05	29	9	397
	Fasting	0.14 \pm 0.03	1.17 \pm 0.05	21	15	350
K. S.	Base line	0.07 \pm 0.02	1.30 \pm 0.03	15	21	285
	Fasting	0.13 \pm 0.03	1.69 \pm 0.12	9	37	223
H. G.	Base line	0.24 \pm 0.01	8.79 \pm 0.33	7	36	659
	Fasting	0.66 \pm 0.07	15.47 \pm 0.62	4	42	936

* Indices of bilirubin metabolism calculated from parameters of the bilirubin- ^3H disappearance curve and the plasma concentration of unconjugated bilirubin are shown. The fasting studies in K. S. and H. G. were started after 40 hr of total caloric restriction. The fasting studies in the other three individuals were started after 57 hr of total caloric restriction.

^b The plasma concentration listed for both conjugated bilirubin and unconjugated bilirubin is the mean \pm SD of seven to eight samples obtained during the study.

plasma conjugated bilirubin during fasting, which forms a small percentage of the total increase, may also be due to decreased hepatic clearance, although this could not be studied by the present method.

The rate of carbon monoxide production was measured in 5 additional subjects during fasting (table 2). Carbon monoxide and bilirubin are both produced by catabolism of heme compounds.^{14,15} An increase in bilirubin production resulting from increased heme degradation will be accompanied by an equimolar increase in carbon monoxide production,¹⁶⁻¹⁸ although the converse may not be true.¹⁷ Since the plasma unconjugated bilirubin concentration is directly proportional to the plasma bilirubin turnover,¹⁹ carbon monoxide production must change by at least the same percentage as the plasma bilirubin concentration during fasting, if augmented heme degradation alone is to account for the increase. This was not observed. The mean increase in the plasma bilirubin con-

centration in the 5 subjects was 188%, whereas the mean change in the carbon monoxide production rate was only +24%. The coefficients of variation seen on several

TABLE 3. Indocyanine green (ICG) plasma clearance*

Subject	Study	Total plasma bilirubin concentration ^b	ICG plasma half life
		mg/100 ml	min
C. G.	Base line	0.40	3.0
	Fasting	1.12	2.8
M. C.	Base line	0.55	3.2
	Fasting	0.78	3.6
B. C.	Base line	0.81	3.6
	Fasting	1.37	3.6
J. B.	Base line	0.73	3.7
	Fasting	0.99	3.6
R. H.	Base line	0.53	4.0
	Fasting	1.34	4.3
Mean ± SD	Base line	0.62 ± 0.16	3.5 ± 0.4
	Fasting	1.10 ± 0.26	3.6 ± 0.5

*The plasma clearance of ICG during fasting is compared with control clearance. The base line study was done following an 8-hr overnight fast (midnight to 8 AM). Each subject was then fasted an additional 48 hr and the fasting study was performed. There was no difference in the plasma half-life of ICG during fasting compared with base line.

^bThe total plasma bilirubin concentration is the mean of at least two samples obtained during the study.

TABLE 2. Carbon monoxide (CO) production*

Subject	Study	Total plasma bilirubin concentration ^b	CO space	CO production
		mg/100 ml	ml	ml/day
J. V.	Base line	0.79	1024	12.2
	After 24-hr fast	0.53	1048	14.6
R. H.	Base line	2.01	1013	11.2
	After 48-hr fast	0.63	1305	20.4
P. B. ^c	Base line	1.34	1315	23.5
	After 48-hr fast	0.79	1274	2.8
F. R.	Base line	0.74	1096	16.3
	After 24-hr fast	0.85	824	9.2
	After 48-hr fast	0.90	794	10.5
J. R. B.	Base line	0.59	771	16.9
	After 24-hr fast	2.05	1093	13.1
	After 48-hr fast	2.56	1078	18.9
			1111	19.0

*The carbon monoxide production rate during fasting is compared with that during *ad libitum* caloric intake. A closed rebreathing system was used to measure the CO production rate. All studies except those in P. B. were performed on consecutive days.

^bThe total plasma bilirubin concentration is the mean of at least two samples obtained during the study.

^cA base line plasma bilirubin concentration was not established.




FIG. 1. The plasma disappearance of bilirubin-³H in subject L. S. while fasting is compared with that during *ad libitum* caloric intake (base line study). Individual data points are shown as a fraction of the extrapolated value at zero time. The solid lines represent computer fits to the data. The plasma disappearance rate of bilirubin-³H decreased with fasting.

replicate determinations of the base line carbon monoxide production rate in two individuals were $\pm 9\%$ and $\pm 16\%$. These results support the hypothesis that a reduction in hepatic bilirubin clearance is mainly responsible for the hyperbilirubinemia of fasting.

The plasma disappearance rate of ICG was determined in 5 subjects in the base line state and during fasting (table 3). Since ICG is extracted from the plasma by the liver with an efficiency of 63 to 88%, measurement of its plasma disappearance rate has been used to estimate hepatic blood flow indirectly.^{11, 20, 21} In the individuals studied there was no difference in the plasma clearance of ICG in the fasting state compared with control values. These results indicate that a decrease in hepatic blood flow does not occur during fasting.

The total plasma bilirubin concentration in each of 9 homozygous Gunn rats with unconjugated hyperbilirubinemia increased during fasting. The base line concentration in these rats was 10.0 ± 3.5 mg per 100 ml (mean \pm SD). During 48 hr of fasting, the mean plasma bilirubin concentration increased by 41% (range, 12 to 88%) of the




FIG. 2. The plasma clearance of bilirubin-³H in homozygous Gunn rats decreases during fasting, returning to control values on refeeding. Individual data points are shown as a fraction of the extrapolated value at zero time. The least squares fits to the bilirubin-³H clearance data during *ad libitum* caloric intake and fasting are shown by the solid lines and dashed lines respectively.

TABLE 4. Rat liver composition changes during fasting*

Basic composition						
Group	Body weight			Liver weight	Grams of H ₂ O per g of liver	Grams of protein per g of liver
	Base line	Fasted	Refed			
1	264 \pm 20	\bar{x}		10.2 \pm 1.6	0.68 \pm 0.004	0.20 \pm 0.002
2	261 \pm 11	226 \pm 7		7.8 \pm 0.7	0.69 \pm 0.007	0.22 \pm 0.004
3	262 \pm 10	222 \pm 9	249 \pm 12	10.2 \pm 0.8	0.68 \pm 0.004	0.19 \pm 0.005
Bilirubin glucuronid transferase						
Group	Body weight		Liver weight	Milligrams per g of liver per hr	Milligrams per g of whole liver per hr	Milligrams per g of protein per hr
	Base line	Fasted				
4	96 \pm 3	\bar{x}	3.0 \pm 0.1	2.4 \pm 0.03	7.3 \pm 0.3	11.7 \pm 0.3
5	66 \pm 1	50 \pm 2	1.8 \pm 0.1	2.7 \pm 0.2	4.9 \pm 0.6	11.5 \pm 0.7
6	174 \pm 6		6.5 \pm 0.7	2.5 \pm 0.1	15.9 \pm 1.2	
7	173 \pm 6	151 \pm 3	4.0 \pm 0.2	2.8 \pm 0.2	11.2 \pm 1.0	

* Seven groups of 4 rats each were used in these studies. Values listed are the mean \pm SE for the entire group. Groups 1, 4, and 6 were control groups. Groups 2, 5, and 7 were studied after 48 hr of total caloric restriction. Group 3 was fasted for 48 hr, then refed for 48 hr before study.

control value. In 6 rats which were fasted an additional 24 hr there was a further increase (mean increase at 72 hr was 55% of control value). The total body weight in these rats decreased by an average of 11% during this period of fasting, whereas the mean hematocrit rose slightly from 40% to 41%.

Labeled bilirubin clearance studies in 2 homozygous Gunn rats are shown in figure 2. Rat 1 was a 240-g female with a total plasma bilirubin concentration of 15.7 mg per 100 ml. Rat 2 was a 196-g female with a bilirubin concentration of 14.6 mg per 100 ml. The fractional turnover of the total accessible bilirubin pool, determined from the terminal slope of the plasma disappearance curve,²² was 0.397 per day in rat 1 and 0.401 per day in rat 2 during normal caloric intake. The rate decreased to 0.226 per day in rat 1 and to 0.223 per day in rat 2 during fasting, with return to control rates on refeeding. These results in an animal which lacks bilirubin glucuronyl transferase indicate that a decrease in the rate of bilirubin conjugation is unlikely to explain the increase in the plasma unconjugated bilirubin concentration which occurs during fasting.

In male Sprague-Dawley rats a decrease in both total body weight (by 13 to 24%) and liver weight (by 24 to 30%) occurred during

48 hr of fasting (see table 4). The concentration of protein in the liver rose approximately 15% during fasting ($P < 0.01$ by Student's *t*-test). The water content (grams of H₂O per gram of liver (wet weight)) did not change. Identical changes in hepatic protein and water content occurred in Gunn rats during fasting. The changes in bilirubin glucuronyl transferase activity in male Sprague-Dawley rat livers were the same as for total protein, and activity per gram of protein did not change.

The bilirubin-³H clearance data from human subjects were subjected to multi-compartmental analysis in an attempt to define the specific step in hepatic bilirubin excretion which is altered by fasting.⁸ The model used (see fig. 3) considers the plasma pool of unconjugated bilirubin (M_1) to exchange with a hepatic pool (M_2) and ex-



Fig. 3. Three-compartment model for the metabolism of unconjugated bilirubin. Values for the λ 's, which are the fractional transfer rates between compartments, and for the pool sizes M calculated from the plasma disappearance curve of bilirubin-³H and the plasma concentration of unconjugated bilirubin.

TABLE 5. Multicompartmental analysis of bilirubin-³H clearance studies*

Subject	State	λ_{12}	λ_{21}	λ_{23}	M_2/M_1	M_3/M_1
C. G.	Base line	0.032	0.015	0.015	1.86	3.50
	Fasting	0.019	0.016	0.016	0.59	2.67
L. S.	Base line	0.015	0.008	0.009	0.50	2.46
	Fasting	0.012	0.016	0.013	0.40	1.86
B. C.	Base line	0.011	0.001	0.005	1.67	1.68
	Fasting	0.012	0.012	0.014	0.45	1.74
K. S.	Base line	0.016	0.006	0.006	0.82	1.19
	Fasting	0.005	0.004	0.004	0.63	1.20
H. G.	Base line	0.006	0.009	0.006	0.42	1.00
	Fasting	0.005	0.014	0.006	0.27	0.85

* Results of multicompartmental analysis of bilirubin-³H clearance data in human subjects are shown. λ_{12} = the fractional transfer rate of unconjugated bilirubin (FTR) from plasma pool to hepatic pool. λ_{21} = FTR from hepatic pool to plasma pool. λ_{23} = irreversible fractional removal rate of unconjugated bilirubin from hepatic pool (= conjugation). M_2/M_1 = ratio of hepatic pool of unconjugated bilirubin (milligrams) to plasma pool (milligrams). M_3/M_1 = ratio of extravascular extrahepatic pool of unconjugated bilirubin (milligrams) to plasma pool (milligrams).

travascular extrahepatic pool (M_3) of unconjugated bilirubin. The fractional transfer rates between pools are depicted on figure 3 as λ 's. In each individual a decrease in the ratio $M_2:M_1$ occurred during fasting, indicating that the ability of the liver to extract bilirubin from the plasma was reduced by fasting (table 5). This decrease in the ratio of pool sizes was associated with a decrease in the fractional transfer rate of bilirubin from plasma to liver (λ_{21}) and/or an increase in the fractional transfer rate from liver to plasma (λ_{12}). The fractional rate which represents conjugation, λ_{c2} , decreased by 33% in subject K. S. during fasting but was unchanged or increased in the other 4 subjects. The ratio $M_2:M_1$ was unchanged in 2 subjects during fasting compared with base line, and slightly decreased in the other 3 subjects.

Discussion

The hyperbilirubinemia of fasting may be due to either a decrease in hepatic clearance of bilirubin from the plasma or an increase in plasma bilirubin turnover resulting from increased bilirubin production (or a combination of both). In this investigation evidence has been presented which indicates that a reduction in hepatic bilirubin clearance during fasting is principally responsible for the increase in the plasma bilirubin concentration.

Measurements of the rate of carbon monoxide production in human individuals during fasting (table 2), while supporting the hypothesis that reduced hepatic bilirubin clearance is mainly responsible for the hyperbilirubinemia of fasting, suggest that an increased rate of heme degradation may also occur. Bakken et al.²² have in fact shown in rats that the activity of hepatic heme oxygenase, the enzyme responsible for the conversion of heme to bilirubin and carbon monoxide, is increased by fasting and by hormones released during hypoglycemia. The production of carbon monoxide-¹⁴C from heme labeled with glycine-2-¹⁴C is also increased.²² The increased rate of carbon monoxide production observed in our subjects during fasting

may thus indicate an increased rate of hepatic heme degradation. If this also reflects increased hepatic bilirubin production, the portion of hepatic bilirubin production which circulates in the plasma prior to excretion in the bile would contribute to the hyperbilirubinemia of fasting.²⁴ It is necessary to point out, however, that an increase in the rate of carbon monoxide production does not always mean that there is an increase in bilirubin production. Landaw et al.¹⁷ showed in rats with experimental porphyria that hepatic heme may be degraded along nonbilirubin pathways. Likewise, Schacter et al.²⁵ have described a mechanism for the degradation of heme and hemoprotein which results in the formation of carbon monoxide but variable amounts of bilirubin.

The physiological alteration which causes the decrease in hepatic bilirubin clearance during fasting seems to be independent of the base line clearance rate and plasma bilirubin concentration. In 2 subjects with Gilbert's syndrome (K. S. and H. G. in table 1), base line hepatic bilirubin clearance was well below the normal range previously described.⁸ Three other individuals (C. G., L. S., and B. C.) had values for clearance which were within the normal range. Subject L. S. had a plasma bilirubin concentration which was only 10% of that for H. G., and his base line hepatic bilirubin clearance was 3-fold greater. Yet both individuals had nearly identical percentage changes in both the plasma bilirubin concentration and hepatic bilirubin clearance with fasting. Multicompartmental analysis of the bilirubin clearance data demonstrated that the ratio of the hepatic bilirubin pool to plasma bilirubin pool decreased in each individual during fasting (table 5).

It is unlikely that hepatic bilirubin clearance decreases with fasting because of a decrease in hepatic blood flow. In order to explain the magnitude of the increase in the plasma bilirubin concentration, a marked reduction in hepatic blood flow would be necessary. This would cause a decrease in the plasma clearance of ICG, since the plasma disappearance rate of ICG offers an indirect measurement of he-

patic blood flow which correlates very well with direct measurements.^{11, 20, 21} Such a decrease was not observed in five individuals (table 3).

A decrease in the activity of hepatic bilirubin glucuronyl transferase is also unlikely to explain the hyperbilirubinemia of fasting. The demonstration that the plasma bilirubin concentration increases with fasting in Gunn rats which lack this enzyme, direct measurement of bilirubin glucuronyl transferase activity in homogenates of male Sprague-Dawley rat liver (table 4), and multicompartmental analysis of human bilirubin clearance data (table 5) indicate that a decrease in the activity of this enzyme does not explain the increase in the plasma bilirubin concentration during fasting.

Two hepatic cytoplasmic protein fractions which bind bilirubin and other organic anions have been described.²⁴ However, the turnover rates of these proteins (half-times of 19 days and 42 hr)²⁷ are too long to account for the rapidity of rise in the plasma bilirubin concentration during fasting, even if synthesis of the proteins stopped with fasting.

Diminished excretion of bilirubin into bile during fasting could cause the increase in plasma conjugated bilirubin. However, this would not explain the increase in plasma unconjugated bilirubin, since only conjugated bilirubin is excreted into bile.

A mechanism to explain fasting hyperbilirubinemia which is consistent with all the data presented is that a material forms (or increases) during fasting which inhibits hepatic bilirubin clearance. Alternatively, a material which facilitates hepatic bilirubin clearance could be depleted. Several compounds which may influence bilirubin metabolism change rapidly with fasting. For example, the plasma glucagon level increases to a peak by 3 days of fasting, whereas the plasma insulin and glucose levels decrease over the same time period.²⁸ This is thought to reflect a period of accelerated hepatic gluconeogenesis. If hepatic uptake of bilirubin is energy-dependent, such changes might be important. Free fatty acids, which have been shown to

affect the binding of bilirubin to albumin, increase markedly within 24 hr of fasting and return rapidly to base line values on refeeding.²⁹ Although the plasma concentration of free fatty acids attained during fasting is not sufficient to cause displacement of bilirubin from albumin *in vitro*,³⁰ the free fatty acids might compete effectively with bilirubin for the hepatic cytoplasmic binding proteins, since the liver actively metabolizes free fatty acids during fasting.

Many other changes occur with fasting, and further work will be required to define the reason why hepatic bilirubin clearance decreases. Such investigation should add to the knowledge of bilirubin metabolism.

REFERENCES

1. Gilbert A, Herscher M: Sur les variations de la cholemie physiologique. *Presse Med* 14:209-211, 1906
2. Barrett P: Hyperbilirubinemia of fasting. *JAMA* (in press)
3. Felsher BF, Rickard D, Redeker AG: The reciprocal relation between caloric intake and the degree of hyperbilirubinemia in Gilbert's syndrome. *New Eng J Med* 283:170-172, 1970
4. Weber AP, Schalm L: Quantitative separation and determination of bilirubin and conjugated bilirubin in human serum. *Clin Chim Acta* 7:80-10, 1962
5. Malloy H, Evelyn K: The determination of bilirubin with the photoelectric colorimeter. *J Biol Chem* 119:481-490, 1937
6. Howe RB, Berk PD, Bloomer JR, et al: Preparation and properties of specifically labeled radiochemically stable ³H-bilirubin. *J Lab Clin Med* 75:499-502, 1970
7. Barrett PVD, Berk PD, Menken M, et al: Bilirubin turnover studies in normal and pathologic states using bilirubin-¹⁴C. *Ann Intern Med* 68:355-377, 1968
8. Berk PD, Howe RB, Bloomer JR, et al: Studies of bilirubin kinetics in normal adults. *J Clin Invest* 48:2176-2180, 1969
9. Collison HA, Rodkey FL, O'Neal JD: Determination of carbon monoxide in blood by gas chromatography. *Clin Chem* 14:162-171, 1968
10. Rodkey FL, O'Neal JD, Collison HA: Oxygen and carbon monoxide equilibria of human adult hemoglobin at atmospheric and elevated pressure. *Blood* 33:57-65, 1969
11. Cherrick GR, Stein SW, Leevy CM, et al: Indol-

- cyanine green: observations on its physical properties, plasma decay and hepatic extraction. *J Clin Invest* 39:592-600, 1960
12. Lowry OH, Rosebrough NJ, Farr AL, et al: Protein measurement with the Folin phenol reagent. *J Biol Chem* 193:265-275, 1951
 13. Black M, Billing BH, Heirwegh KPM: Determination of bilirubin UDP-glucuronyl transferase activity in needle-biopsy specimens of human liver. *Clin Chim Acta* 29:27-35, 1970
 14. Sjöstrand T: The formation of carbon monoxide by the decomposition of haemoglobin in vivo. *Acta Physiol Scand* 26:338-344, 1952
 15. Ludwig GD, Blakemore WS, Drabkin DL: Production of carbon monoxide by heme oxidation. *J Invest* 36:912, 1957
 16. Engel R, Berk PD, Rodkey FL, et al: Estimation of heme turnover (HT) and erythrocyte survival (RBCLS) in man from clearance of bilirubin (BR)-¹⁴C and from carbon monoxide production (COP). *Clin Res* 17:325, 1969
 17. Landaw SA, Callahan EW Jr, Schmid R: Catabolism of heme in vivo: comparison of the simultaneous production of bilirubin and carbon monoxide. *J Clin Invest* 49:914-925, 1970
 18. Coburn RF, Williams WJ, Kahn SB: Endogenous carbon monoxide production in patients with hemolytic anemia. *J Clin Invest* 45:460-468, 1966
 19. Berk PD, Bloomer JR, Howe RB, and Berlin NI: Constitutional hepatic dysfunction (Gilbert's syndrome). A new definition based on kinetic studies with unconjugated radiobilirubin. *Amer J Med* 49:296-305, 1970
 20. Reemtsma K, Hattinger GC, DeGraff AC, et al: The estimation of hepatic blood flow using indocyanine green. *Surg Gynec Obstet* 110:353-356, 1960
 21. Caesar J, Shaldon S, Chiardussi L, et al: The use of indocyanine green in the measurement of hepatic blood flow and as a test of hepatic function. *Clin Sci* 21:43-57, 1961
 22. Schmid R, Hammaker L: Metabolism and disposition of C¹⁴-bilirubin in congenital nonhemolytic jaundice. *J Clin Invest* 42:1726-1734, 1963
 23. Bakken AF, Thalen MM, Schmid R: Hormonal control of hepatic heme catabolism. *J Clin Invest* 50:50a, 1971
 24. Jones EA, Bloomer JR, Berk PD, et al: The measurement of hepatic synthesis of bilirubin and its delivery to plasma in man. *J Clin Invest* 50:50a, 1971
 25. Schacter BA, Meyer UA, Hildebrandt AG, et al: Microsomal lipid peroxidation: a new mechanism for hemoprotein catabolism. *Clin Res* 19:429, 1971
 26. Levi AJ, Gatmaitan Z, Arias IM: Two hepatic cytoplasmic protein fractions, Y and Z, and their possible role in the hepatic uptake of bilirubin, sulfobromophthalein, and other anions. *J Clin Invest* 48:2156-2167, 1969
 27. Levi AJ, Gatmaitan Z, Arias IM: Study of the rates of turnover and development of two hepatic cytoplasmic proteins (Y and Z) which bind bilirubin and sulfobromophthalein (BSP). *Gastroenterology* 56:401, 1969
 28. Marliss EB, Aoki TT, Unger RH, et al: Glucagon levels and metabolic effects in fasting man. *J Clin Invest* 49:2256-2270, 1970
 29. Morse WJ, Mahabir R: Changes in glucose tolerances and plasma free fatty acids after fasting in obesity. *Diabetes* 13:286-290, 1964
 30. Starinsky R, Shafrir E: Displacement of albumin-bound bilirubin by free fatty acids: Implications for neonatal hyperbilirubinemia. *Clin Chim Acta* 29:311-315, 1970