

AD-A080 824

FOREIGN TECHNOLOGY DIV WRIGHT-PATTERSON AFB OH
MOMENTUM AND ENERGY RELAXATION BY AN ELECTRON IN A CRYSTAL III.--ETC(U)
MAR 79 P A KAZLAUSKAS, I B LEVINSON
FTD-ID(RS)T-0372-79

F/G 20/8

UNCLASSIFIED

NL

1 - 1
A 10-1



END
DATE
FILMED
3 80
301


PHOTOGRAPH THIS SHEET

ADA U 80824

DTIC ACCESSION NUMBER



LEVEL



INVENTORY

FTD-ID(RS) T-0372-79

DOCUMENT IDENTIFICATION

DISTRIBUTION STATEMENT A

Approved for public release;
Distribution Unlimited

DISTRIBUTION STATEMENT

ACCESSION FOR

NTIS GRA&I

DTIC TAB

UNANNOUNCED

JUSTIFICATION

BY

DISTRIBUTION /


AVAILABILITY CODES

DIST

AVAIL AND/OR SPECIAL

A

DISTRIBUTION STAMP



DATE ACCESSIONED

DATE RECEIVED IN DTIC

PHOTOGRAPH THIS SHEET AND RETURN TO DTIC-DDA-2

ADA 080824

FOREIGN TECHNOLOGY DIVISION



MOMENTUM AND ENERGY RELAXATION BY AN ELECTRON IN A CRYSTAL
III. COMPOUND SCATTERING BY OPTICAL PHONONS AT LOW TEMPERATURES

By

P. A. Kazlauskas, I. B. Levinson, G. E. Mazhuolite



Approved for public release;
distribution unlimited.

79 08 20 223

EDITED TRANSLATION

FTD-ID(RS)T-0372-79

21 March 1979

MICROFICHE NR: *FTD-79-C-000421*MOMENTUM AND ENERGY RELAXATION BY AN ELECTRON
IN A CRYSTAL III. COMPOUND SCATTERING BY OPTICAL
PHONONS AT LOW TEMPERATURES

By: P. A. Kazlauskas, I. B. Levinson, G. E. Mazhuolite

English pages: 10

Source: Litovskiy Fizicheskiy Sbornik, Vol. 6,
Nr. 3, 1966, pp. 377-381

Country of Origin: USSR

Translated by: Gale M. Weisenbarger

Requester: AFWL/IN

Approved for public release; distribution unlimited.

THIS TRANSLATION IS A RENDITION OF THE ORIGINAL FOREIGN TEXT WITHOUT ANY ANALYTICAL OR EDITORIAL COMMENT. STATEMENTS OR THEORIES ADVOCATED OR IMPLIED ARE THOSE OF THE SOURCE AND DO NOT NECESSARILY REFLECT THE POSITION OR OPINION OF THE FOREIGN TECHNOLOGY DIVISION.

PREPARED BY:

TRANSLATION DIVISION
FOREIGN TECHNOLOGY DIVISION
WP-AFB, OHIO.

U. S. BOARD ON GEOGRAPHIC NAMES TRANSLITERATION SYSTEM

Block	Italic	Transliteration	Block	Italic	Transliteration
А а	<i>А а</i>	A, a	Р р	<i>Р р</i>	R, r
Б б	<i>Б б</i>	B, b	С с	<i>С с</i>	S, s
В в	<i>В в</i>	V, v	Т т	<i>Т т</i>	T, t
Г г	<i>Г г</i>	G, g	У у	<i>У у</i>	U, u
Д д	<i>Д д</i>	D, d	Ф ф	<i>Ф ф</i>	F, f
Е е	<i>Е е</i>	Ye, ye; E, e*	Х х	<i>Х х</i>	Kh, kh
Ж ж	<i>Ж ж</i>	Zh, zh	Ц ц	<i>Ц ц</i>	Ts, ts
З э	<i>З э</i>	Z, z	Ч ч	<i>Ч ч</i>	Ch, ch
И и	<i>И и</i>	I, i	Ш ш	<i>Ш ш</i>	Sh, sh
Й й	<i>Й й</i>	Y, y	Щ щ	<i>Щ щ</i>	Shch, shch
К к	<i>К к</i>	K, k	Ъ ъ	<i>Ъ ъ</i>	"
Л л	<i>Л л</i>	L, l	Ы ы	<i>Ы ы</i>	Y, y
М м	<i>М м</i>	M, m	Ь ь	<i>Ь ь</i>	'
Н н	<i>Н н</i>	N, n	Э э	<i>Э э</i>	E, e
О о	<i>О о</i>	O, o	Ю ю	<i>Ю ю</i>	Yu, yu
П п	<i>П п</i>	P, p	Я я	<i>Я я</i>	Ya, ya

*ye initially, after vowels, and after ъ, ь; e elsewhere.
When written as ë in Russian, transliterate as yë or ë.

RUSSIAN AND ENGLISH TRIGONOMETRIC FUNCTIONS

Russian	English	Russian	English	Russian	English
sin	sin	sh	sinh	arc sh	sinh ⁻¹
cos	cos	ch	cosh	arc ch	cosh ⁻¹
tg	tan	th	tanh	arc th	tanh ⁻¹
ctg	cot	cth	coth	arc cth	coth ⁻¹
sec	sec	sch	sech	arc sch	sech ⁻¹
cosec	csc	csch	csch	arc csch	csch ⁻¹
		Russian	English		
		rot	curl		
		lg	log		

0372, gw

MOMENTUM AND ENERGY RELAXATION BY AN ELECTRON IN A CRYSTAL

III. COMPOUND SCATTERING BY OPTICAL PHONONS AT LOW TEMPERATURES

P. A. Kazlauskas, I. B. Levinson, G. E. Mazhuolite

(Received 10 Dec. 1965)

The coefficients of the Fokker-Planck equation are calculated during low-temperature scattering by optical phonons for the random law of dispersion and for electron energies up to the energy of an optical phonon.

1. Introduction. The probability of compound scattering.

Under conditions when the lattice temperature kT is much less than the maximum frequency of optical phonons $\hbar\omega_0$, the probability of emission of a phonon is much greater than the probability of absorption of the phonon. Therefore absorption is accompanied by "instantaneous re-emission" and these two sequential processes may be examined as a single act of compound scattering [1, 2]. Thanks to the small dispersion of optical phonons the compound scattering is almost elastic and precisely this circumstance makes it possible to convert the integral collision term of the kinetic equation from Boltzmann's form to the differential form of Fokker-Planck. While completing this procedure B. Davydov and N. Shmushkevich [2] assumed that in a considerable area of energies $\epsilon \ll \hbar\omega_0$, the anisotropy of the distribution function is also small, i.e.,

$$(1.1) \quad f_{(p)} = f_0(\epsilon) + f_1(\epsilon) \cos \vartheta, \quad f_1(\epsilon) \ll f_0(\epsilon).$$

Here p is the pulse of the electron, ϑ is its angle with the direction of the electrical field. These assumptions in fact are not significant; in [3] it was shown that the concept of compound scattering is limited only by the condition $\epsilon \ll \hbar\omega_0$, when the scattering begins necessarily from absorption and that it may be described by the probability of compound scattering:

$$(1.2) \quad \hat{W}(p, p') = \int (dp'') \epsilon_0^+(p'') W^-(p, p'') W^+(p'', p').$$

Here p and p' are the initial and final states with energies ϵ , $\epsilon' < \omega$, p'' is an intermediate state with an energy of $\epsilon' > \omega$. The lifetime in this intermediate state (due to emission) is

$$(1.3) \quad \frac{1}{\tau_0^+(p'')} = \int (dp') W^+(p'', p').$$

Factor W describes the absorption transition from p to p'' , and $W^+ \tau_0^+$ is the conditional probability that emission of the phonon will lead to the transition from p'' to the final point p' which is of interest to us.

Integrating (1.2) with respect to p' it is easy to check that the lifetime of state p relative to compound scattering $\frac{1}{\tau_0}(p)$ coincides with the lifetime relative to initial scattering $\tau_0^+(p)$; this is physically clear because $1/\tau_0$ is contributed to equally by all processes of phonon absorption independently of what happens with this electron following absorption. However, for other characteristic times such a coincidence does not occur.

Subsequently we shall use an isotropic model [4, 5]. Expanding the compound and regular probability through Legendre polynomials [2.7] I, for the momentums we find

$$(1.4) \quad \hat{W}_I(\epsilon, \epsilon') = \int_{\omega_0}^{\infty} d\epsilon'' \epsilon_0^+(\epsilon'') W_I^-(\epsilon, \epsilon'') W_I^+(\epsilon'', \epsilon').$$

It is not difficult to check that if the initial probability W satisfies the principle of detailed scattering, then the latter is automatically satisfied by the compound probability \hat{W} . Let us note one more obvious circumstance: concepts of compound scattering and its probability (1.2) refer not only to polarization scattering by optical phonons PO, examined in (2), but also to differential scattering DO.

2. Characteristics of compound scattering.

Below we calculate values which describe compound scattering of an electron. The sign $\hat{}$, which indicates the compound nature of the scattering, is omitted for brevity.

We introduce the following designations:

minimum and maximum momentums of absorbed phonons

$$(2.1) \quad \begin{aligned} q_1^-(\epsilon) &= p(\epsilon + \hbar\omega_0) - p(\epsilon), \\ q_2^-(\epsilon) &= p(\epsilon + \hbar\omega_0) + p(\epsilon), \end{aligned}$$

momentums of matrix elements of interaction

$$(2.2) \quad B_n(\epsilon) = \int_{q_1^-(\epsilon)}^{q_n^-(\epsilon)} dq q^n B(q),$$

auxiliary function

$$(2.3) \quad \Lambda(\epsilon) = \ln \frac{q_n^-(\epsilon)}{q_1^-(\epsilon)},$$

number of phonons (2.4)

$$N_0 = \exp \left\{ - \frac{\hbar\omega_0}{kT} \right\},$$

characteristic momentum and velocity

$$(2.5) \quad p_0 = p(\hbar\omega_0), \quad v_0 = v(\hbar\omega_0).$$

All other unstipulated designations follow I and II.

During calculation of the lifetime τ_0 and the longitudinal relaxation time of momentum $\tau_{||} = \tau$ the dispersion of optical phonons may be disregarded, i. e., compound scattering may be considered elastic. Using the probability of transition in the form (2.4) I, calculating the momentums of compound probability with (2.9) I and

(1.4) and substituting everything in (3.2) I and (3.5) I, we find the lifetime:

$$(2.6) \quad \frac{1}{\tau_0(\varepsilon)} = \frac{2\pi}{\hbar} N_0 2\pi \frac{\rho(\varepsilon + \hbar\omega_0)}{\rho(\varepsilon) \nu(\varepsilon + \hbar\omega_0)} B_1(\varepsilon),$$

and the relaxation time:

$$(2.7) \quad \frac{1}{\tau(\varepsilon)} = \frac{1}{\tau_0(\varepsilon)} \left[1 - \left(\text{cth } \Lambda(\varepsilon) - \frac{1}{2\rho(\varepsilon) \rho(\varepsilon + \hbar\omega_0)} \frac{B_2(\varepsilon)}{B_1(\varepsilon)} \right)^2 \right].$$

During calculation of the coefficient of energy diffusion D one should take into account the dispersion of optical phonons which is accepted in the following form:

$$(2.8) \quad \hbar\omega(q) = \hbar\omega_0 - \frac{q^2}{2M}.$$

where M is the order of mass of the nucleus. Considering the dispersion term in the law of preservation of energy and expanding the corresponding δ -functions in the probabilities of transitions W^+ and W^- , entering into (1.2), to the second order with respect to M^{-1} inclusively, we obtain

$$(2.9) \quad D(\varepsilon) = \frac{1}{\tau_0(\varepsilon)} \frac{1}{4M^2} \left[\frac{B_2(\varepsilon)}{B_1(\varepsilon)} - \frac{B_2^2(\varepsilon)}{B_1^2(\varepsilon)} \right].$$

Compound scattering is elastic everywhere and satisfies the principle of detailed equilibrium; therefore the power of energy losses Q can be calculated by direct differentiation using (2.9) and (1.32) II.

Using the matrix elements $B(q)$ from I and calculating momentum $B_n(\epsilon)$, it is possible to find the following expressions for specific mechanisms of interaction:

DO

$$\frac{1}{\tau_0(\epsilon)} = \frac{1}{\tau(\epsilon)} = \frac{2\pi}{\hbar} B_0 N_0 4\pi \frac{p^2(\epsilon + \hbar\omega_0)}{v(\epsilon + \hbar\omega_0)}, \quad (2.10)$$

$$D(\epsilon) = \frac{1}{\tau_0(\epsilon)} \frac{p^2(\epsilon) p^2(\epsilon + \hbar\omega_0)}{3M^2}, \quad (2.11)$$

PO

$$\frac{1}{\tau_0(\epsilon)} = \frac{2\pi}{\hbar} B_0 N_0 2\pi \frac{p(\epsilon + \hbar\omega_0)}{p(\epsilon)v(\epsilon + \hbar\omega_0)} \Lambda(\epsilon), \quad (2.12)$$

$$\frac{1}{\tau(\epsilon)} = \frac{1}{\tau_0(\epsilon)} \left[1 - \left(\text{cth } \Lambda(\epsilon) - \frac{1}{\Lambda(\epsilon)} \right)^2 \right], \quad (2.13)$$

$$D(\epsilon) = \frac{1}{\tau_0(\epsilon)} \frac{p^2(\epsilon) p^2(\epsilon + \hbar\omega_0)}{M^2} \frac{1}{\Lambda(\epsilon)} \left(\text{cth } \Lambda(\epsilon) - \frac{1}{\Lambda(\epsilon)} \right). \quad (2.14)$$

For a parabolic zone the results are simplified somewhat and

$$(2.15) \quad \Lambda(\epsilon) = 2 \text{Arsh} \left(\frac{\epsilon}{\hbar\omega_0} \right)^{\frac{1}{2}}.$$

The relaxation time τ for a parabolic zone and the mechanism PO were calculated using methods of resonance scattering [6]. Disregarding of

the virtual transitions leads, in [6], to a formula different from (2.13) by the supplementary factor 2/3 in the right part, which, as was pointed out in [7], is the result of error in integration.

3. Low energies.

The obtained formulas are simplified significantly for low energies $\varepsilon \ll \hbar\omega_0$. Let us note that in this case

$$(3.1) \quad \Lambda(\varepsilon) = 2 \frac{p(\varepsilon)}{p_0} \ll 1,$$

$$(3.2) \quad \operatorname{cth} \Lambda(\varepsilon) - \frac{1}{\Lambda(\varepsilon)} = \frac{2}{3} \frac{p(\varepsilon)}{p_0} \ll 1.$$

Using these formulas we find that for low energies

$$(3.3) \quad \frac{1}{\tau(\varepsilon)} = \frac{1}{\tau_0(\varepsilon)} = \frac{2\pi}{\hbar} B_0 N_0 4\pi \frac{p_0^4}{v_0} \quad (DO),$$

$$(3.4) \quad = \frac{2\pi}{\hbar} B_0 N_0 4\pi \frac{1}{v_0} \quad (PO),$$

$$(3.5) \quad D(\varepsilon) = \frac{1}{\tau_0(\varepsilon)} \frac{1}{3} \frac{p^2(\varepsilon) p_0^2}{M^2}.$$

Times τ and τ_0 in this case turn out to be independent of ε for any law of dispersion. With the parabolic law of dispersion these results coincide with those obtained in [2]. In this case it is also

convenient to introduce the relaxation time $\tilde{\tau}$ and fluctuation time $\bar{\tau}$ of energy

$$(3.7) \quad \frac{1}{\tilde{\tau}(\epsilon)} = \frac{1}{\tau_0} \frac{4}{3} \frac{\hbar\omega_0}{kT} \left(\frac{m}{M}\right)^2, \quad \epsilon^* = \frac{3}{2} kT$$

$$\frac{1}{\bar{\tau}(\epsilon)} = \frac{1}{\tau_0} \frac{4}{3} \frac{\hbar\omega_0}{\epsilon} \left(\frac{m}{M}\right)^2.$$

Here ϵ^* is the energy with which $Q(\epsilon)$ changes sign. From (3.6) it is evident that inelasticity of compound scattering by optical phonons is determined by the parameter

$$(3.8) \quad \bar{\delta} = \frac{4}{3} \frac{\hbar\omega_0}{kT} \left(\frac{m}{M}\right)^2.$$

Let us note additionally that the use of formulas obtained for $\epsilon \ll \hbar\omega_0$ in the entire interval up to $\epsilon = \hbar\omega_0$, gives correct results with respect to the order of magnitude.

Institute of Physics and Mathematics, Academy of Sciences Lithuanian SSR.

LITERATURE

1. H. Frolich, N. F. Mott, Proc. Roy. Soc., 171, 496 (1939).
2. Б. Давыдов, Н. Шмушкевич, ЖЭТФ, 10, 1043 (1940).

3. И. Б. Левинсон, Г. Э. Мажуолите, ЖЭТФ, 50, 1048 (1966).
4. П. А. Казлаускас, И. Б. Левинсон, Liet. fiz. rinkinys, 6, 33 (1966) (цитируется как I).
5. П. А. Казлаускас, И. Б. Левинсон, Liet. fiz. rinkinys, 6, 233 (1966) (цитируется как II).
6. S. Zieman, E. R. A. Report, L/T 236 (1950).
7. H. Frolih, Adv. in Physics, 3, 325 (1954).

Summary

When lattice temperature kT and electron energy ε is lower than optical phonon energy $\hbar\omega_0$, the absorption of a phonon is followed by its immediate re-emission. These two scattering acts may be treated as a compound scattering, whose probability \mathcal{W} is given by (1.2) in terms of ordinary scattering probability W .

General formulae are: (2.6) — life time, (2.7) — momentum relaxation time (longitudinal), (2.9) — energy diffusion coefficient; for special mechanisms: (2.10) and (2.11) — deformation scattering, (2.12) — (2.14) — polarization scattering.

For energies $\varepsilon \ll \hbar\omega_0$ (3.1) — (3.5) are valid. At such energies and parabolic band the energy relaxation and fluctuation times, given by (3.6) and (3.7) respectively may be introduced. The elasticity of compound scattering is given by (3.8).

DISTRIBUTION LIST

DISTRIBUTION DIRECT TO RECIPIENT

<u>ORGANIZATION</u>	<u>MICROFICHE</u>	<u>ORGANIZATION</u>	<u>MICROFICHE</u>
A205 DMATC	1	E053 AF/INAKA	1
A210 DMAAC	2	E017 AF/RDXTR-W	1
B344 DIA/RDS-3C	9	E403 AFSC/INA	1
C043 USAMIA	1	E404 AEDC	1
C509 BALLISTIC RES LABS	1	E408 AFWL	1
C510 AIR MOBILITY R&D LAB/FIO	1	E410 ADTC	1
C513 PICATINNY ARSENAL	1	FTD	
C535 AVIATION SYS COMD	1	CCN	1
C591 FSTC	5	ASD/FTD/NIIS	3
C619 MIA REDSTONE	1	NIA/PHS	1
D008 NISC	1	NIIS	2
H300 USAICE (USAREUR)	1		
P005 DOE	1		
P050 CIA/CRB/ADD/SD	1		
NAVORDSTA (50L)	1		
NASA/KSI	1		
AFIT/LD	1		
ILL/Code L-389	1		
NSA/1213/TDL	2		