

AD-A081 660

ILLINOIS UNIV AT URBANA-CHAMPAIGN DEPT OF CHEMISTRY

F/G 7/8

THE ANOMALOUS METALLOPORPHYRIN AND CHLOROPHYLL A ACTIVATED CHEN--ETC(U)

FEB 80 S P SCHMIDT, G B SCHUSTER

N00014-76-C-0745

UNCLASSIFIED TR-24

NL

| or |
AD
A081 660



END
DATE
FILMED
4-80
DTIC

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

20. (cont.)

role of the central metal atom in eliciting this unusual behavior was evaluated by determining the catalytic rate constants and the initial chemiluminescence intensity of several non-metallated as well as differently metallated porphyrins. These results suggest that formation of a ground state complex between \downarrow and certain metalloporphyrins is the cause of the unusual catalysis. Spectroscopic evidence of a ground state complex between zinc tetraphenyl porphyrin (ZnTPP) and tetramethyldioxetane, a model for \downarrow , has been obtained.

Accession For	
NTIS GSA&I	<input checked="" type="checkbox"/>
DOC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By _____	
Distribution/	
Availability Codes	
Dist	Avail and/or special
A	

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

OFFICE OF NAVAL RESEARCH

Contract ¹⁵ N0014-76-C-0745

Task No. NR-051-616

⁹ TECHNICAL REPORT NO. N0014-76-C-0745-24

¹⁴ TR-24

The Anomalous Metalloporphyrin and Chlorophyll a Activated
Chemiluminescence of Dimethyldioxetanone. Chemically
Initiated Electron-Exchange Luminescence.

by

Steven P. Schmidt and Gary B. Schuster

Prepared for Publication

in

Journal of the American Chemical Society

School of Chemical Sciences

University of Illinois

Urbana, Illinois 61801

Feb ~~1976~~ 1980

12/21/

Reproduction in whole or in part is permitted for
any purpose of the United States Government

Approved for Public Release; Distribution Unlimited.

400-1

MIT

The chemistry of the dioxetane ring system has been under active investigation for the past decade.² In a recent report³ we described our discovery of the operation of the chemically initiated electron-exchange luminescence (CIEEL) mechanism⁴ in the chemiluminescence of dimethyldioxetanone (λ). The involvement of this mechanism was indicated primarily by the correlation of the excited state generating effectiveness of a wide range of electron donors with the one electron oxidation potential (E_{ox}) of the donor. In that report we noted that zinc tetraphenylporphyrin (ZnTPP) is an apparent exception to this correlation. The initial chemiluminescent light intensity from ZnTPP is about 100 times greater than is predicted simply from its one electron oxidation potential. Moreover, the observed rate constant for catalysis of the decomposition of λ by ZnTPP is much larger than is predicted by its oxidation potential.

Unusual chemiluminescent phenomena have been reported previously for metalloporphyrins. Linschitz observed emission from the thermolysis of decalin hydroperoxide in the presence of ZnTPP or chlorophyll.⁵ He noted that the free-base porphyrin, unlike ZnTPP, has no catalytic effect on the rate of reaction of decalin hydroperoxide. Moreover, the free-base reaction is not chemiluminescent. More recently, McCapra reported chemiluminescence from the reaction of a variety of peroxides with ZnTPP.⁶ Again the metal is an absolute requirement.



In this report we describe our investigation of the interaction of ZnTPP and other metallo- and free-base porphyrins with dioxetanone λ . The results of this investigation show conclusively for the first time that the unusual behavior of ZnTPP is consistent with the CIEEL mechanism and that this behavior is due to the formation of a ground state complex between the peroxide and the metalloporphyrin.

Results and Discussion

In our previous report we established that the mechanism for electron donor (activator, ACT) catalyzed chemiluminescence of dioxetanone λ proceeds by

a rate limiting electron transfer from ACT to the peroxide generating eventually acetone radical anion and \cdot ACT radical cation. Annihilation of these oppositely charged radical ions forms the excited state of ACT which we detect by its characteristic luminescence. This mechanism, and the appropriate rate constants, are shown in Scheme 1.

Scheme 1



Our results indicate that the kinetics of the activator catalyzed reaction conform to:

$$k_{\text{obs}} = k_1 + k_{\text{cat}} [\text{ACT}] \quad (4)$$

where k_{obs} is the rate constant for the observed net first order consumption of dioxetanone λ , k_1 is the rate constant for reaction of λ in the absence of ACT, and k_{cat} is rate constant for the bimolecular reaction of λ and ACT. It is the bimolecular interaction that is primarily responsible for the chemiluminescence.

The ratio of initial chemiluminescence intensity (I_0) to the fluorescence efficiency of ACT (ϕ_f) can be related to the magnitude of k_{cat} , and the initial concentrations of λ and ACT according to:


$$I_0/\phi_f = k_{cat} [1]_0 [ACT] \quad (5)$$

Thus under conditions where the concentrations of λ and ACT are constant during the time required for the experiment, I_0/ϕ_f is directly proportional to k_{cat} . We have measured I_0/ϕ_f under these conditions for a wide variety of ACT. In general, for amine and aromatic hydrocarbon ACT their E_{ox} accurately predict the magnitude of I_0/ϕ_f . The exceptions to this correlation are some metalloporphyrins, Figure 1. The measured value of k_{cat} for a variety of ACT are presented in Table I. It should be noted that the exceptionally large value of I_0/ϕ_f for the zinc and magnesium porphyrins are consistent with the independently measured values for k_{cat} .


In order to determine if the unusual catalytic ability of the zinc and magnesium porphyrins is due to the metal ion or to the porphyrin ring system we investigated the behavior of some free-base porphyrins. In particular, we determined I_0/ϕ_f and k_{cat} for tetraanisylporphyrin (H_2TAP). As can be seen from Figure 1 and Table I, H_2TAP does not behave unusually. That, is its E_{ox} is a reliable predictor of both I_0/ϕ_f and k_{cat} . Similarly, tetraphenylporphyrin (H_2TPP) does not appear to catalyze the decomposition of λ or its chemiluminescence beyond that which would be predicted by its E_{ox} . Evidently, the metal ion plays a crucial role in causing the unusually efficient catalysis by MgTPP and ZnTPP.

To probe further the mechanism of this metal ion catalysis we examined the effect of changing the metal ion on I_0/ϕ_f and k_{cat} for tetraphenyl- and tetraanisylporphyrins. Cyclic voltammetric measurements have revealed that for

zinc, cadmium, and magnesium porphyrins the first oxidation is ligand centered.⁷ On the other hand, for silver and cobalt porphyrin the first oxidation is metal centered.^{7,8} Interestingly, zinc, cadmium, cobalt, and magnesium porphyrin (including chlorophyll a) exhibit the unusually effective catalysis, but AgTPP does not. The magnitude of k_{cat} for AgTPP is accurately predicted by its E_{ox} , Table I. Evidently, the identity of the metal ion matters greatly for the catalysis. Compare, for example, AgTPP ($E_{\text{ox}} = 0.59 \text{ V vs SCE}$) which exhibits a k_{cat} of $1.1 \text{ M}^{-1} \text{ s}^{-1}$ and MgTPP ($E_{\text{ox}} = 0.60 \text{ V vs SCE}$) whose k_{cat} is nearly 10,000



Free-Base Porphyrins

 H_2TPP , Ar=Ph H_2TAP , Ar=(*p*-OCH₃)Ph

Metalloporphyrins

MTPP, Ar=Ph

MTAP, Ar=(*p*-OCH₃)Ph

times larger. But, the catalysis can occur with metalloporphyrins that undergo either metal- or ligand-centered oxidation. Apparently, it is some feature other than the site of oxidation of the metalloporphyrin that engenders the remarkable chemiluminescent catalysis.

It is quite well known that certain metalloporphyrins form stable complexes with nitrogen bases such as pyridine.⁹ Moreover, it is known that free-base porphyrins do not form these complexes.¹⁰ Quite recently the results of a thermodynamic study of the complex formation of ZnTPP with several neutral donors was reported.¹¹ In this study the equilibrium constant for the complex was

estimated by the shift of the maximum of the porphyrin Soret absorption band. For metalloporphyrins it has been observed that the maximum of the Soret band shifts to longer wavelengths upon complex formation. For simple ester donors the shift in cyclohexane solution is observed to be ca. 5 nm, for alkyl ethers ca. 6 nm. We investigated the possibility that the formation of a ground state complex between dioxetanone λ and certain metalloporphyrins is the cause of the unusual catalysis.

Addition of ZnTPP or MgTPP to cyclohexane solutions of dioxetanone λ causes such a rapid reaction that it is not possible for us to measure the absorption spectrum of the mixture before most of the peroxide has reacted. To circumvent this problem we used tetramethyldioxetane (λ) as a model for λ . The reactions of λ are not particularly sensitive to metalloporphyrin catalysis, presumably because its reduction potential is far more negative than that of λ . In cyclohexane solution the maximum of the Soret band for ZnTPP is at 416.0 nm. In the presence of 0.36 M λ this maximum shifts to 417.2 nm. A similar result is obtained with MgTPP. Interestingly, AgTPP, which does not readily expand its coordination sphere,¹² and for which no special catalysis is observed, does not give evidence of complex formation with dioxetane λ . This is in line with our observation that even pyridine does not shift the absorption maximum or oxidation potential of this metalloporphyrin. Evidently the ability of the metalloporphyrin to form detectable complexes with dioxetane λ predicts reliably the unusual catalytic behavior.

The magnitude of k_{cat} is related to several of the rate constants specified in Scheme 1. Our previously reported data indicates that $k_{30} \gg k_{\text{-ACT}}$. Thus simple analysis of the kinetics show that:

$$k_{\text{cat}} = K_{12} k_{\text{ACT}} \quad (6)$$

where K_{12} (k_{12}/k_{21}) is the equilibrium constant for complex formation.¹³ For the amine and aromatic hydrocarbon activators K_{12} is evidently independent of the structure of ACT and probably depends, as in the Weller model,¹⁴ only on diffusion. However, ZnTPP, MgTPP, CoTPP, CdTPP, etc. form ground state complexes with peroxide \downarrow and the magnitude of k_{cat} is therefore the product of K_{12} for the complex and k_{ACT} . Indeed, if we assume that complexation does not affect k_{ACT} then we can estimate the magnitude of K_{12} from the measured value of k_{cat} and an interpolated (from the E_{ox} vs. I_0/ϕ_f plot) value for k_{ACT} . The K_{12} of ZnTPP, is thereby estimated to be 100 M^{-1} , a value consistent with equilibrium constants reported by others for complexes of metalloporphyrins with various oxygen donors.¹¹

The electron transfer reaction of chlorophyll a (Chla) continue to be of general interest due to their intimate involvement in the initiating steps of photosynthesis. In connection with their study of the electrochemistry of Chla, Saji and Bard reported their observation of electrogenerated chemiluminescence of Chla in the presence of oxygen.¹⁵ In our system Chla behaves much like the synthetic magnesium and zinc porphyrins in eliciting efficient chemiluminescence from \downarrow by the electron-exchange pathway. The magnitude of the Chla catalytic rate constant is comparable to that of the magnesium tetraphenylporphyrins (Table I), suggesting that ground state complex formation with peroxide \downarrow is important for Chla as well. It is likely therefore that other Chla - Peroxide systems might also be chemiluminescent, the result of an analogous complexation - electron transfer sequence. As noted by others,¹⁵ for example, weak chemiluminescence has been observed from the reaction of O_2 with photoreduced chlorophyll.¹⁶

While the magnesium and zinc porphyrins are among the most effective activators of chemiluminescence for \downarrow which we have found, they are quite ineffective in catalyzing chemiluminescence with diphenylperoxide (DPP). This might be surprising initially in light of the established involvement of the CIEEL pathway for the chemiluminescence of DPP in the presence of aromatic hydrocarbons.⁴ In

fact, bimolecular rate constants for the catalyzed decomposition of DPP are generally several orders of magnitude larger than for λ under comparable conditions, a consequence of the more facile reduction of DPP and diacyl peroxides in general than λ and peroxyesters in general.³ The metalloporphyrin, however, is consumed in the reaction with DPP. We suspect, therefore, that two electron reduction of DPP is occurring. The second oxidation potentials of these metalloporphyrins are relatively low,¹⁷ in comparison to those of the typical aromatic hydrocarbons which catalyze light formation with DPP. The aryloxy radical intermediate produced from the one electron reduction and O-O bond cleavage of DPP is more easily reduced than the alkoxy radical derived from the one electron reduction and O-O bond cleavage of λ .¹⁵ Relative rates of decarboxylation may also contribute to the different behavior of λ and DPP. In any event, further reduction of DPP radical anion effectively removes this species from the light-generating path.

Finally, we would like to point out the remarkable sensitivity of the metalloporphyrin activated chemiluminescence as an analytical technique. The unusually large values of k_{cat} for the complex forming metalloporphyrins, and the emissive properties of ZnTPP and MgTPP conspire to make possible the detection of incredibly small amounts of the metal ions. For example, we have been able to detect unambiguously ca. 1 ppb zinc in solutions of PdTPP and ZnTPP using dioxetanone λ .

Conclusion

The chemical reactions of dimethyldioxetanone with a variety of fluorescent electron donors have revealed new features of the CIEEL mechanism. In this report we have described our observation of a ground state complex between this peroxide and several metalloporphyrins. The complex evidently involves an interaction between the metal ion and the peroxide. In all of the cases, and only in those cases, for which complex

formation is detected we observe an increase in both I_0/ϕ_f and k_{cat} . This finding indicates that the complex is on the reaction path leading to eventual light generation. This remarkably enhanced chemiluminescence catalysis should prove valuable in the detection of trace quantities of these metal ions and in increasing the yield of light when the CIEEL path is in competition with a rapid unimolecular decomposition of the key chemiluminescent intermediate.

Experimental Section

General. All chemiluminescence measurements were made by the photon counting technique as previously described.³ Photoexcited fluorescence spectra were obtained on a Farrand Mark V spectrofluorometer, or by photon counting for direct comparison with chemiluminescence emission spectra. UV-VIS absorption spectra were recorded on a Beckman Acta MVI, or Cary 14 instrument. ¹H NMR spectra were obtained on a Varian HR-220 instrument. Solvents used in the chemiluminescence experiments were purified as previously described.³

Cyclic Voltammetry. Cyclic voltammetry was done in argon saturated dichloromethane solution at (23±2)°C with 0.1 M tetra-n-butyl-ammonium perchlorate as supporting electrolyte. A three-electrode system, consisting of platinum working and counter electrodes, and saturated calomel reference electrode (SCE), was used. Scans were made typically at 0.5 V/s. The E_{ox} values were measured as that potential lying midway between the oxidation and reduction peak for a given couple.

Peroxides. Dimethyldioxetanone,^{3,18} tetramethyldioxetane,¹⁹ and diphenylperoxide^{4b} were prepared and purified as described elsewhere.

Porphyrins. The porphyrin free bases H₂TPP and H₂TAP were kindly provided by Prof. Larry Faulkner. They were purified by treatment with dichlorodicyanobenzoquinone (DDQ) in refluxing chloroform, as described by Barnett et al.²⁰ The porphyrins were shown to be chlorin free by visible absorption spectroscopy.²¹

ZnTPP (Sigma) was also purified by treatment with DDQ.

Other metalloporphyrins were prepared by reaction of the appropriate divalent metal salt with the porphyrin free base in refluxing N,N-dimethylformamide according to the general procedure of Adler et al.²² The magnesium porphyrins and CdTPP were recrystallized from benzene-hexane. ZnTAP was purified by dry

column chromatography on Al_2O_3 , eluting with chloroform, followed by several recrystallizations from chloroform-methanol. CoTPP and AgTPP were purified by dry column chromatography on Florisil, eluting with chloroform, followed by recrystallization from chloroform-methanol.

All porphyrins gave satisfactory elemental analyses, and had visible absorption spectra consistent with literature data.^{21,23}

Chlorophyll a (Sigma) was used as received.

Acknowledgment. This research was supported by the Office of Naval Research and the National Science Foundation. S.P.S. is a University of Illinois Fellow, 1976-1980.

References and Notes

- (1) Fellow of the Alfred P. Sloan Foundation 1977-79.
- (2) K. A. Horn, J.-y. Koo, S. P. Schmidt, and G. B. Schuster, Mol. Photochem., 9, 1 (1978-79); P. D. Bartlett and M. E. Landis, in "Singlet Oxygen", H. H. Wasserman and R. W. Murray, Ed.; Academic Press; New York, 1979; W. Adam, Adv. Heterocycl. Chem., 21, 437 (1977); T. Wilson, Int. Rev. Sci.: Phys. Chem., Ser. Two, 9, 265 (1976).
- (3) S. P. Schmidt and G. B. Schuster, J. Am. Chem. Soc., 102, 0000 (1980).
- (4) (a) G. B. Schuster, Acc. Chem. Res., 12, 366 (1979). (b) J.-y. Koo and G. B. Schuster, J. Am. Chem. Soc., 100, 4496 (1978).
- (5) H. Linschitz in "Light and Life," McElroy and Glass, Eds.; Johns Hopkins Press, 1961, p 173.
- (6) F. McCapra and P. D. Leeson, J. Chem. Soc., Chem. Commun., 114 (1979).
- (7) (a) D. G. Davis in "The Porphyrins", D. Dolphin, Ed.; Academic Press; New York, 1978, Vol V, Chapter 4. (b) R. H. Felton, ibid, Chapter 3. (c) A. Wolberg, Isr. J. Chem., 12, 1031 (1974).
- (8) (a) A. Antipas, D. Dolphin, M. Gouterman, and E. C. Johnson, J. Am. Chem. Soc., 100, 7705 (1978). (b) F. A. Walker, D. Beroiz, and K. M. Kadish, J. Am. Chem. Soc., 98, 3484 (1976). (c) L. A. Truxillo and D. G. Davis, Anal. Chem., 47, 2260 (1975). (d) J. Manassen, Isr. J. Chem., 12, 1059 (1974).
- (9) P. Hambright, Coord. Chem. Rev., 6, 247 (1971); C. B. Storm, A. H. Corwin, R. R. Arellano, M. Martz, and R. Weintraub, J. Am. Chem. Soc., 88, 2525 (1966); C. H. Kirksey, P. Hambright, and C. D. Storm, Inorg. Chem., 8, 2141 (1969).
- (10) J. R. Miller and G. D. Dorough, J. Am. Chem. Soc., 74, 3977 (1952).
- (11) G. C. Vogel and J. R. Stahlbush, Inorg. Chem., 16, 950 (1977).
- (12) The equilibrium constant for pyridine complexation of MgTPP is at least 4×10^4 times greater than that for AgTPP.¹⁰

- (13) G. B. Schuster, J. Am. Chem. Soc., 101, 5851 (1979).
- (14) D. Rehm and A. Weller, Isr. J. Chem., 8, 259 (1970).
- (15) T. Saji and A. J. Bard, J. Am. Chem. Soc., 99, 2235 (1977).
- (16) F. F. Litvin, J. A. Vladimirov, and A. A. Kransnovskii, Usp. Fiz. Nauk, 71, 149 (1960).
- (17) L. Ebersson, Acta. Chem. Scand., 17, 2004 (1963).
- (18) W. Adam and J.-C. Liu, J. Am. Chem. Soc., 94, 2894 (1972); W. Adam, A. Alzerreca, J.-C. Liu and F. Yany, ibid., 99, 5768 (1977).
- (19) K. R. Kopecky, J. E. Kilby, C. Mumford, P. A. Lockwood, and J.-Y. Ding, Can. J. Chem., 53, 1103 (1975).
- (20) G. H. Barnett, M. F. Hudson, and K. M. Smith, J. Chem. Soc., Perkin I, 1401 (1975).
- (21) G. M. Badger, R. A. Jones, and R. L. Laslett, Aust. J. Chem., 17, 1028 (1964).
- (22) A. D. Adler, F. R. Longo, F. Kampas, and J. Kim, J. Inorg. Nucl. Chem., 32, 2443 (1970).
- (23) G. D. Dorough, J. R. Miller, and F. M. Huenneckens, J. Am. Chem. Soc., 73, 4315 (1951).

Table I. Oxidation Potentials of Activators and Rate Constants for Their Catalysis of the Reaction of Dioxetanone \downarrow .

Activator	E_{ox} , V ^a	k_{cat} , M ⁻¹ s ⁻¹ ^b
Chlorophyll a	0.52 ^c	(1.5±0.1)×10 ³
MgTAP	0.54	(2.6±0.2)×10 ⁴
MgTPP	0.60	(1.0±0.1)×10 ⁴
AgTPP	0.59 ^{d,e}	1.1±0.2
CdTPP	0.64 ^f	(2.7±0.2)×10 ³
ZnTAP	0.66 ^g	(1.3±0.1)×10 ³
ZnTPP	0.73	(7.0±0.2)×10 ²
CoTPP	0.77 ^d	>10 ⁵ ^h
H ₂ TAP	0.80 ^g	0.67±0.03
Rubrene	0.82	0.44±0.02

^aThis work, measured in CH₂Cl₂, unless otherwise noted. For details see Experimental Section.

^bMeasured in CH₂Cl₂ solution at 24.5°C. All errors are standard deviations.

^cA. Standianda, Naturwissenschaften, 50, 731 (1963); Z. Phys. Chem., 229, 257 (1965).

^dMetal centered oxidation M^{II}TPP → M^{III}TPP.

^e0.60 V in pyridine.

^fA. Standianda and G. Biebl, Z. Phys. Chem. (Frankfurt am Main) [N.S.] 52, 254 (1967).


^gRef. 7c.

^hEstimated rate constant.

Caption for Figure

Relationship between the initial chemiluminescence intensity in dichloromethane at 24.5°C, corrected for fluorescence efficiency (ϕ_f) and photomultiplier tube and monochromator response, and the one electron oxidation potential of the activator: DPA = 9,10-diphenylanthracene, BPEA = 9,10-bis(phenylethynyl)anthracene.

Figure 1



TECHNICAL REPORT DISTRIBUTION LIST, GEN

	<u>No.</u> <u>Copies</u>		<u>No.</u> <u>Copies</u>
Office of Naval Research Attn: Code 472 800 North Quincy Street Arlington, Virginia 22217	2	U.S. Army Research Office Attn: CRD-AA-IP P.O. Box 1211 Research Triangle Park, N.C. 27709	1
ONR Branch Office Attn: Dr. George Sandoz 536 S. Clark Street Chicago, Illinois 60605	1	Naval Ocean Systems Center Attn: Mr. Joe McCartney San Diego, California 92152	1
ONR Branch Office Attn: Scientific Dept. 715 Broadway New York, New York 10003	1	Naval Weapons Center Attn: Dr. A. B. Amster, Chemistry Division China Lake, California 93555	1
ONR Branch Office 1030 East Green Street Pasadena, California 91106	1	Naval Civil Engineering Laboratory Attn: Dr. R. W. Drisko Port Hueneme, California 93401	1
ONR Branch Office Attn: Dr. L. H. Peebles Building 114, Section D 666 Summer Street Boston, Massachusetts 02210	1	Department of Physics & Chemistry Naval Postgraduate School Monterey, California 93940	1
Director, Naval Research Laboratory Attn: Code 6100 Washington, D.C. 20390	1	Dr. A. L. Slafkosky Scientific Advisor Commandant of the Marine Corps (Code RD-1) Washington, D.C. 20380	1
The Assistant Secretary of the Navy (R,E&S) Department of the Navy Room 4E736, Pentagon Washington, D.C. 20350	1	Office of Naval Research Attn: Dr. Richard S. Miller 800 N. Quincy Street Arlington, Virginia 22217	1
Commander, Naval Air Systems Command Attn: Code 310C (H. Rosenwasser) Department of the Navy Washington, D.C. 20360	1	Naval Ship Research and Development Center Attn: Dr. G. Bosmajian, Applied Chemistry Division Annapolis, Maryland 21401	1
Defense Documentation Center Building 5, Cameron Station Alexandria, Virginia 22314	12	Naval Ocean Systems Center Attn: Dr. S. Yamamoto, Marine Sciences Division San Diego, California 91232	1
Dr. Fred Saalfeld Chemistry Division Naval Research Laboratory Washington, D.C. 20375	1	Mr. John Boyle Materials Branch Naval Ship Engineering Center Philadelphia, Pennsylvania 19112	1

TECHNICAL REPORT DISTRIBUTION LIST, GEN

No.
Copies

Dr. Rudolph J. Marcus
Office of Naval Research
Scientific Liaison Group
American Embassy
APO San Francisco 96503

1

Mr. James Kelley
DTNSRDC Code 2803
Annapolis, Maryland 21402

1

TECHNICAL REPORT DISTRIBUTION LIST, 051A

	<u>No.</u> <u>Copies</u>		<u>No.</u> <u>Copies</u>
Dr. M. A. El-Sayed Department of Chemistry University of California, Los Angeles Los Angeles, California 90024	1	Dr. M. Rauhut Chemical Research Division American Cyanamid Company Bound Brook, New Jersey 08805	1
Dr. M. W. Windsor Department of Chemistry Washington State University Pullman, Washington 99163	1	Dr. J. I. Zink Department of Chemistry University of California, Los Angeles Los Angeles, California 90024	1
Dr. E. R. Bernstein Department of Chemistry Colorado State University Fort Collins, Colorado 80521	1	Dr. B. Schechtman IBM San Jose Research Center 5600 Cottle Road San Jose, California 95143	1
Dr. C. A. Heller Naval Weapons Center Code 6059 China Lake, California 93555	1	Dr. John Cooper Code 6130 Naval Research Laboratory Washington, D.C. 20375	1
Dr. J. R. MacDonald Chemistry Division Naval Research Laboratory Code 6110 Washington, D.C. 20375	1		
Dr. G. B. Schuster Chemistry Department University of Illinois Urbana, Illinois 61801	1		
Dr. E. M. Eyring Department of Chemistry University of Utah Salt Lake City, Utah	1		
Dr. A. Adamson Department of Chemistry University of Southern California Los Angeles, California 90007	1		
Dr. M. S. Wrighton Department of Chemistry Massachusetts Institute of Technology Cambridge, Massachusetts 02139	1		