

AD A 082057

AMRL-TR-79-81

125

125



SOUND-EVOKED VISUAL FIELD SHIFTS: INTERACTION WITH FIVE CLASSES OF STIMULATION

*D. E. PARKER
W. L. GULLEDGE
W. PEREZ
R. POSTON*

*DEPARTMENT OF PSYCHOLOGY
MIAMI UNIVERSITY
OXFORD, OHIO 45056*

JANUARY 1980

SDTIC
ELECTE
MAR 20 1980
A D

Approved for public release; distribution unlimited.

AIR FORCE AEROSPACE MEDICAL RESEARCH LABORATORY
AEROSPACE MEDICAL DIVISION
AIR FORCE SYSTEMS COMMAND
WRIGHT-PATTERSON AIR FORCE BASE, OHIO 45433

DDC FILE COPY

80 3 17 015

NOTICES

When US Government drawings, specifications, or other data are used for any purpose other than a definitely related Government procurement operation, the Government thereby incurs no responsibility nor any obligation whatsoever, and the fact that the Government may have formulated, furnished, or in any way supplied the said drawings, specifications, or other data, is not to be regarded by implication or otherwise, as in any manner licensing the holder or any other person or corporation, or conveying any rights or permission to manufacture, use, or sell any patented invention that may in any way be related thereto.

Please do not request copies of this report from Air Force Aerospace Medical Research Laboratory. Additional copies may be purchased from:

National Technical Information Service
5285 Port Royal Road
Springfield, Virginia 22161

Federal Government agencies and their contractors registered with Defense Documentation Center should direct requests for copies of this report to:

Defense Documentation Center
Cameron Station
Alexandria, Virginia 22314

TECHNICAL REVIEW AND APPROVAL

AMRL-TR-79-81

The voluntary informed consent of the subjects used in this research was obtained as required by Air Force Regulation 169-3.

This report has been reviewed by the Office of Public Affairs (PA) and is releasable to the National Technical Information Service (NTIS). At NTIS, it will be available to the general public, including foreign nations.

This technical report has been reviewed and is approved for publication.

FOR THE COMMANDER



HENNING E. VON GERKE
Director
Biodynamics and Bioengineering Division
Air Force Aerospace Medical Research Laboratory

12 REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM	
18 1. REPORT NUMBER AMRL-TR-79-81	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER	
6 4. TITLE (and Subtitle) SOUND-EVOKED VISUAL FIELD SHIFTS: INTERACTION WITH FIVE CLASSES OF STIMULATION.		9 5. TYPE OF REPORT & PERIOD COVERED Final Report.	
7. AUTHOR(s) 10 D. E. Parker W. Perez W. L. Gullledge R. Poston		15 8. CONTRACT OR GRANT NUMBER(s) AF 33 615-73-C-5029	
9. PERFORMING ORGANIZATION NAME AND ADDRESS Miami University Oxford, Ohio 45056		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS 16 62202F, 17 7231 03-22	
11. CONTROLLING OFFICE NAME AND ADDRESS Air Force Aerospace Medical Research Laboratory, Aerospace Medical Division, Air Force Systems Command, Wright-Patterson Air Force Base, OH 45433		12. REPORT DATE Jan 80	
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office) 1244		13. NUMBER OF PAGES 44	
		15. SECURITY CLASS. (of this report) Unclassified	
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE	
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited			
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)			
18. SUPPLEMENTARY NOTES			
19. KEY WORDS (Continue on reverse side if necessary and identify by block number)			
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) Visual field displacements evoked by acoustical transients in human beings have been described previously, (AMRL-TR-77-81). The influence of several variables on this phenomenon was examined in five studies. Experiment I investigated the interaction between acoustical stimulation and angular acceleration. The results of this experiment support the view that the acoustical transients activate semicircular canal receptors. The influence of head vibration on acoustical			

405-944

cont

Block 20--Continued

transient-evoked visual field displacements was examined in Experiment II. Smaller visual field displacements were reported when head vibration was presented simultaneously with the acoustical stimulation, probably because the head vibration activated the middle ear reflex. No changes in acoustical stimulus-evoked visual field displacements following exposure to a rotating visual field (Experiment III) or as a function of visual target illumination intensity (Experiment IV) were observed. Finally, alcohol ingestion (Experiment V) reduced the magnitudes of the visual field displacement reported by the subjects.



PREFACE

The research described in this report was performed under Contract F 33 615 73 C 5029 between the United States Air Force and Miami University, Oxford, Ohio.

This investigation was conducted in accordance with guidelines established by the Miami University Human Subjects Committee.

We thank our colleagues who have contributed to this effort, particularly H. E. von Gierke and C. S. Harris.

Accession For	
1955 - 1961	
1962 - 1968	
1969 - 1972	
1973 - 1978	
1979 - 1984	
1985 - 1990	
1991 - 1996	
1997 - 2002	
2003 - 2008	
2009 - 2014	
2015 - 2020	
2021 - 2026	
2027 - 2032	
2033 - 2038	
2039 - 2044	
2045 - 2050	
2051 - 2056	
2057 - 2062	
2063 - 2068	
2069 - 2074	
2075 - 2080	
2081 - 2086	
2087 - 2092	
2093 - 2098	
2099 - 2104	
2105 - 2110	
2111 - 2116	
2117 - 2122	
2123 - 2128	
2129 - 2134	
2135 - 2140	
2141 - 2146	
2147 - 2152	
2153 - 2158	
2159 - 2164	
2165 - 2170	
2171 - 2176	
2177 - 2182	
2183 - 2188	
2189 - 2194	
2195 - 2200	
2201 - 2206	
2207 - 2212	
2213 - 2218	
2219 - 2224	
2225 - 2230	
2231 - 2236	
2237 - 2242	
2243 - 2248	
2249 - 2254	
2255 - 2260	
2261 - 2266	
2267 - 2272	
2273 - 2278	
2279 - 2284	
2285 - 2290	
2291 - 2296	
2297 - 2302	
2303 - 2308	
2309 - 2314	
2315 - 2320	
2321 - 2326	
2327 - 2332	
2333 - 2338	
2339 - 2344	
2345 - 2350	
2351 - 2356	
2357 - 2362	
2363 - 2368	
2369 - 2374	
2375 - 2380	
2381 - 2386	
2387 - 2392	
2393 - 2398	
2399 - 2404	
2405 - 2410	
2411 - 2416	
2417 - 2422	
2423 - 2428	
2429 - 2434	
2435 - 2440	
2441 - 2446	
2447 - 2452	
2453 - 2458	
2459 - 2464	
2465 - 2470	
2471 - 2476	
2477 - 2482	
2483 - 2488	
2489 - 2494	
2495 - 2500	
2501 - 2506	
2507 - 2512	
2513 - 2518	
2519 - 2524	
2525 - 2530	
2531 - 2536	
2537 - 2542	
2543 - 2548	
2549 - 2554	
2555 - 2560	
2561 - 2566	
2567 - 2572	
2573 - 2578	
2579 - 2584	
2585 - 2590	
2591 - 2596	
2597 - 2602	
2603 - 2608	
2609 - 2614	
2615 - 2620	
2621 - 2626	
2627 - 2632	
2633 - 2638	
2639 - 2644	
2645 - 2650	
2651 - 2656	
2657 - 2662	
2663 - 2668	
2669 - 2674	
2675 - 2680	
2681 - 2686	
2687 - 2692	
2693 - 2698	
2699 - 2704	
2705 - 2710	
2711 - 2716	
2717 - 2722	
2723 - 2728	
2729 - 2734	
2735 - 2740	
2741 - 2746	
2747 - 2752	
2753 - 2758	
2759 - 2764	
2765 - 2770	
2771 - 2776	
2777 - 2782	
2783 - 2788	
2789 - 2794	
2795 - 2800	
2801 - 2806	
2807 - 2812	
2813 - 2818	
2819 - 2824	
2825 - 2830	
2831 - 2836	
2837 - 2842	
2843 - 2848	
2849 - 2854	
2855 - 2860	
2861 - 2866	
2867 - 2872	
2873 - 2878	
2879 - 2884	
2885 - 2890	
2891 - 2896	
2897 - 2902	
2903 - 2908	
2909 - 2914	
2915 - 2920	
2921 - 2926	
2927 - 2932	
2933 - 2938	
2939 - 2944	
2945 - 2950	
2951 - 2956	
2957 - 2962	
2963 - 2968	
2969 - 2974	
2975 - 2980	
2981 - 2986	
2987 - 2992	
2993 - 2998	
2999 - 3004	
3005 - 3010	
3011 - 3016	
3017 - 3022	
3023 - 3028	
3029 - 3034	
3035 - 3040	
3041 - 3046	
3047 - 3052	
3053 - 3058	
3059 - 3064	
3065 - 3070	
3071 - 3076	
3077 - 3082	
3083 - 3088	
3089 - 3094	
3095 - 3100	
3101 - 3106	
3107 - 3112	
3113 - 3118	
3119 - 3124	
3125 - 3130	
3131 - 3136	
3137 - 3142	
3143 - 3148	
3149 - 3154	
3155 - 3160	
3161 - 3166	
3167 - 3172	
3173 - 3178	
3179 - 3184	
3185 - 3190	
3191 - 3196	
3197 - 3202	
3203 - 3208	
3209 - 3214	
3215 - 3220	
3221 - 3226	
3227 - 3232	
3233 - 3238	
3239 - 3244	
3245 - 3250	
3251 - 3256	
3257 - 3262	
3263 - 3268	
3269 - 3274	
3275 - 3280	
3281 - 3286	
3287 - 3292	
3293 - 3298	
3299 - 3304	
3305 - 3310	
3311 - 3316	
3317 - 3322	
3323 - 3328	
3329 - 3334	
3335 - 3340	
3341 - 3346	
3347 - 3352	
3353 - 3358	
3359 - 3364	
3365 - 3370	
3371 - 3376	
3377 - 3382	
3383 - 3388	
3389 - 3394	
3395 - 3400	
3401 - 3406	
3407 - 3412	
3413 - 3418	
3419 - 3424	
3425 - 3430	
3431 - 3436	
3437 - 3442	
3443 - 3448	
3449 - 3454	
3455 - 3460	
3461 - 3466	
3467 - 3472	
3473 - 3478	
3479 - 3484	
3485 - 3490	
3491 - 3496	
3497 - 3502	
3503 - 3508	
3509 - 3514	
3515 - 3520	
3521 - 3526	
3527 - 3532	
3533 - 3538	
3539 - 3544	
3545 - 3550	
3551 - 3556	
3557 - 3562	
3563 - 3568	
3569 - 3574	
3575 - 3580	
3581 - 3586	
3587 - 3592	
3593 - 3598	
3599 - 3604	
3605 - 3610	
3611 - 3616	
3617 - 3622	
3623 - 3628	
3629 - 3634	
3635 - 3640	
3641 - 3646	
3647 - 3652	
3653 - 3658	
3659 - 3664	
3665 - 3670	
3671 - 3676	
3677 - 3682	
3683 - 3688	
3689 - 3694	
3695 - 3700	
3701 - 3706	
3707 - 3712	
3713 - 3718	
3719 - 3724	
3725 - 3730	
3731 - 3736	
3737 - 3742	
3743 - 3748	
3749 - 3754	
3755 - 3760	
3761 - 3766	
3767 - 3772	
3773 - 3778	
3779 - 3784	
3785 - 3790	
3791 - 3796	
3797 - 3802	
3803 - 3808	
3809 - 3814	
3815 - 3820	
3821 - 3826	
3827 - 3832	
3833 - 3838	
3839 - 3844	
3845 - 3850	
3851 - 3856	
3857 - 3862	
3863 - 3868	
3869 - 3874	
3875 - 3880	
3881 - 3886	
3887 - 3892	
3893 - 3898	
3899 - 3904	
3905 - 3910	
3911 - 3916	
3917 - 3922	
3923 - 3928	
3929 - 3934	
3935 - 3940	
3941 - 3946	
3947 - 3952	
3953 - 3958	
3959 - 3964	
3965 - 3970	
3971 - 3976	
3977 - 3982	
3983 - 3988	
3989 - 3994	
3995 - 4000	
4001 - 4006	
4007 - 4012	
4013 - 4018	
4019 - 4024	
4025 - 4030	
4031 - 4036	
4037 - 4042	
4043 - 4048	
4049 - 4054	
4055 - 4060	
4061 - 4066	
4067 - 4072	
4073 - 4078	
4079 - 4084	
4085 - 4090	
4091 - 4096	
4097 - 4102	
4103 - 4108	
4109 - 4114	
4115 - 4120	
4121 - 4126	
4127 - 4132	
4133 - 4138	
4139 - 4144	
4145 - 4150	
4151 - 4156	
4157 - 4162	
4163 - 4168	
4169 - 4174	
4175 - 4180	
4181 - 4186	
4187 - 4192	
4193 - 4198	
4199 - 4204	
4205 - 4210	
4211 - 4216	
4217 - 4222	
4223 - 4228	
4229 - 4234	
4235 - 4240	
4241 - 4246	
4247 - 4252	
4253 - 4258	
4259 - 4264	
4265 - 4270	
4271 - 4276	
4277 - 4282	
4283 - 4288	
4289 - 4294	
4295 - 4300	
4301 - 4306	
4307 - 4312	
4313 - 4318	
4319 - 4324	
4325 - 4330	
4331 - 4336	
4337 - 4342	
4343 - 4348	
4349 - 4354	
4355 - 4360	
4361 - 4366	
4367 - 4372	
4373 - 4378	
4379 - 4384	
4385 - 4390	
4391 - 4396	
4397 - 4402	
4403 - 4408	
4409 - 4414	
4415 - 4420	
4421 - 4426	
4427 - 4432	
4433 - 4438	
4439 - 4444	
4445 - 4450	
4451 - 4456	
4457 - 4462	
4463 - 4468	
4469 - 4474	
4475 - 4480	
4481 - 4486	
4487 - 4492	
4493 - 4498	
4499 - 4504	
4505 - 4510	
4511 - 4516	
4517 - 4522	
4523 - 4528	
4529 - 4534	
4535 - 4540	
4541 - 4546	
4547 - 4552	
4553 - 4558	
4559 - 4564	
4565 - 4570	
4571 - 4576	
4577 - 4582	
4583 - 4588	
4589 - 4594	
4595 - 4600	
4601 - 4606	
4607 - 4612	
4613 - 4618	
4619 - 4624	
4625 - 4630	
4631 - 4636	
4637 - 4642	
4643 - 4648	
4649 - 4654	
4655 - 4660	
4661 - 4666	
4667 - 4672	
4673 - 4678	
4679 - 4684	
4685 - 4690	
4691 - 4696	
4697 - 4702	
4703 - 4708	
4709 - 4714	
4715 - 4720	
4721 - 4726	
4727 - 4732	
4733 - 4738	
4739 - 4744	
4745 - 4750	
4751 - 4756	
4757 - 4762	
4763 - 4768	
4769 - 4774	
4775 - 4780	
4781 - 4786	
4787 - 4792	
4793 - 4798	
4799 - 4804	
4805 - 4810	
4811 - 4816	
4817 - 4822	
4823 - 4828	
4829 - 4834	
4835 - 4840	
4841 - 4846	
4847 - 4852	
4853 - 4858	
4859 - 4864	
4865 - 4870	
4871 - 4876	
4877 - 4882	
4883 - 4888	
4889 - 4894	
4895 - 4900	
4901 - 4906	
4907 - 4912	
4913 - 4918	
4919 - 4924	
4925 - 4930	
4931 - 4936	
4937 - 4942	
4943 - 4948	
4949 - 4954	
4955 - 4960	
4961 - 4966	
4967 - 4972	
4973 - 4978	
4979 - 4984	
4985 - 4990	
4991 - 4996	
4997 - 5002	
5003 - 5008	
5009 - 5014	
5015 - 5020	
5021 - 5026	
5027 - 5032	
5033 - 5038	
5039 - 5044	
5045 - 5050	
5051 - 5056	
5057 - 5062	
5063 - 5068	
5069 - 5074	
5075 - 5080	
5081 - 5086	
5087 - 5092	
5093 - 5098	
5099 - 5104	
5105 - 5110	
5111 - 5116	

TABLE OF CONTENTS

	<u>Page</u>
I. INTRODUCTION	6
Experiment I--Angular Acceleration	7
Experiment II--Head Vibration	7
Experiment III--Prior Visual Field Rotation	8
Experiment IV--Target Illumination	9
Experiment V--Alcohol Ingestion	9
II. METHOD	10
Experiment I--Angular Acceleration	10
Experiment I-A	10
Experiment I-B	13
Experiment II--Head Vibration	14
Experiment III--Prior Visual Field Rotation	15
Experiment III-A	15
Experiment III-B	17
Experiment IV--Target Illumination	18
Experiment V--Alcohol Ingestion	19
III. RESULTS	21
Experiment I--Angular Acceleration	21
Experiment I-A	21
Experiment I-B	23
Experiment II--Head Vibration	25

TABLE OF CONTENTS (CONTINUED)

	<u>Page</u>
Experiment III--Prior Visual Field Rotation . . .	25
Experiment IV--Target Illumination	28
Experiment V--Alcohol Ingestion	28
IV. DISCUSSION	32
Experiment I--Angular Acceleration	32
Experiment II--Head Vibration	33
Experiment III--Prior Visual Field Exposure . . .	34
Experiment IV--Target Illumination	35
Experiment V--Alcohol Ingestion	35
V. CONCLUSIONS	37
REFERENCES	38

LIST OF FIGURES

	<u>Page</u>
1. Geometric Means of Perceived Visual Field Displacements as a Function of Rotation Rate of Centrifuge Prior to Acoustical Stimulation	22
2. Means of Z-score Transformed Perceived Visual Field Displacement Reports as a Function of Centrifuge Rotation Rate	24
3. Geometric Means of Reported Visual Field Displacements as a Function of Ear Stimulated	26
4. Geometric Means of Visual Field Displacement Reports as a Function of Target Brightness	29
5. Geometric Means of Perceived Visual Field Displacement Reports as a Function of Time After Ingestion of Orange Juice or Orange Juice Plus Alcohol	30

LIST OF TABLES

	<u>Page</u>
1. Reports of Target Motion Evoked by Acoustical Transients Following Rotation at 5, 10, and 15 rpm (%)	21
2. Magnitude Estimations--Experiment I-A	23
3. Geometric Means--Experiment III	27
4. Repeated Measures Analyses of Variance for Experiment III	28

SECTION I

INTRODUCTION

In previous studies, personnel at Miami University and Wright-Patterson Air Force Base have investigated the effects of pressure transients, infrasound, and intense audiofrequency sound on (a) the response of the ossicular chain, (b) pressure changes in inner ear fluids, (c) the response of the vestibular nerve, (d) head and eye movements, and (e) perceived visual field displacements. Guinea pigs, monkeys, and human subjects have been examined in these investigations (refs 13-20). These studies and others (refs 7-9) suggest that sound and pressure stimuli activate the receptors of the semicircular canals to produce head movements, eye movements, and perceived displacements of the visual field.

In the study on visual field displacements evoked from human beings by acoustical transients (ref 16), we found that approximately one-half of the 133 subjects who were exposed to the acoustical stimuli at 125 dB SPL reported visual field displacements. Stimuli in the 500- to 1000-Hz frequency range at repetition rates of about 1/sec resulted in the largest proportion of responses.

The present series of experiments examined the contribution of other variables to visual field displacements evoked by acoustical transients. The specific variables investigated included angular acceleration (Experiment I), mastoid vibration (Experiment II), prior exposure to a rotating visual field (Experiment III), intensity of target illumination (Experiment IV), and alcohol ingestion (Experiment V).

EXPERIMENT I--ANGULAR ACCELERATION

The results of previous investigations supported the hypothesis that pressure transients (step and ramp functions of pressure change) at the tympanic membrane and intense audiofrequency transients affect the receptors of the semicircular canals. The observations that led to this conclusion included the following: (a) pressure transients evoke nystagmus in guinea pigs and monkeys; (b) intense audiofrequency sound evokes nystagmus in human beings; (c) the nystagmus response is lost in animals following eighth nerve section; (d) vestibular nerve responses can be evoked by pressure transients; (e) nystagmus is retained following damage to the otolith organs by centrifugation; (f) the nystagmus response is lost following damage to the semicircular canals.

Experiment I examined the hypothesis that prior exposure to angular acceleration would alter the reports of visual field displacement evoked by acoustical transients. After a sudden angular deceleration, the cupulae of the semicircular canals in the plane of rotation are initially displaced and then gradually return to their resting position. If, as suggested by previous research, acoustical transients result in cupula displacement, the effects of angular deceleration should interact with the effects of acoustical stimulation. Whether the visual field shift is enhanced or suppressed depends on the direction of the angular acceleration and/or the ear stimulated. Given apparatus constraints, it was more convenient to maintain the direction of angular deceleration constant and to vary the ear which received the acoustical stimuli. The hypothesis was that the magnitude of sound-evoked visual field shifts after angular deceleration would differ for right ear and left ear stimulation.

EXPERIMENT II--HEAD VIBRATION

Lackner and Graybiel (ref 12) recently reported that visual and postural illusions, nystagmus, and motion sickness were produced by head vibration. These effects were obtained by

stimulation at various head loci with the clearest responses associated with stimulation around the ear. They suggest that vibration affects the semicircular canals; however, no explanations are offered for the causative biomechanical processes. We and others (C. S. Harris, personal communication) have attempted to repeat the observations reported by Lackner and Graybiel without success. Although some visual field displacement may be associated with vibration onset, this displacement appears to be related to a "startle" reaction rather than to vestibular stimulation. Visual illusions were not reported by our subjects during or after head vibration.

Perhaps the behavioral techniques that we employed were not sufficiently sensitive to duplicate the results of Lackner and Graybiel. If they are correct in assuming that vibration stimulates the semicircular canals and if we are correct in assuming that intense acoustical transients also stimulate the semicircular canals, then either enhancement or suppression of visual field shifts should be associated with simultaneous presentation of vibration and acoustical transients, depending upon the relationship between the ear stimulated and the vibration. Regardless of the direction of the effect obtained with the right ear, the effect obtained with the left ear should be opposite. Experiment II was designed to evaluate this suggestion.

EXPERIMENT III--PRIOR VISUAL FIELD ROTATION

Several investigators have examined the influence of rotating visual environments on eye movement responses (optomotor responses) (ref 11) as well as on subjective perception (ref 3). The purpose of Experiment III was to determine whether prior exposure to a rotating visual field comprised of alternating black and white stripes would influence the incidence and magnitude of visual field displacements evoked by acoustical transients.

EXPERIMENT IV--TARGET ILLUMINATION

Change in visual responses evoked by vestibular stimulation as a function of the illumination intensity of the visual target has been examined in several laboratories (refs 1, 2, 5, 6). Experiment IV was performed to examine the hypothesis that the magnitude of the visual field shift elicited by intense acoustical transients would be reduced when the target was brightly illuminated. This hypothesis was derived from the observation that nystagmus is suppressed by visual field illumination (ref 6).

EXPERIMENT V--ALCOHOL INGESTION

Numerous investigators have examined the influence of alcohol ingestion on responses evoked by vestibular stimulation (e.g., refs 2, 5, 10, 21). Heifer (ref 10) and Schroeder (ref 21) have reported that alcohol depresses vestibular nystagmus when the observations are obtained in darkness, whereas nystagmus responses are enhanced if the recordings are obtained in light. The former observation has been accounted for by the general depression of vestibular responses produced by alcohol and the latter observation has been related to the depressive action of alcohol on the visual fixation mechanism.

Following the results of Schroeder (ref 21) and previous experiments undertaken at Miami University employing guinea pigs and monkeys (ref 17), it was hypothesized that smaller visual field shifts would be reported by human subjects following ingestion of alcohol. Experiment V was designed to evaluate this hypothesis.

SECTION II

METHOD

A. EXPERIMENT I--ANGULAR ACCELERATION

Two experiments were performed on the interaction between acoustical stimulation and angular acceleration. In Experiment I-A changes in the magnitudes of perceived visual field displacements were examined following angular deceleration at three levels, and in Experiment I-B both magnitude and direction effects were studied.

Experiment I-A

Twelve graduate students were paid to participate in Experiment I-A. These twelve were selected from a larger group of volunteers because they exhibited hearing losses less than 20 dB (500- to 8000-Hz) in an audiometric examination and initially reported visual field displacements upon exposure to the acoustical transients presented during the first part of the experiment. All subjects were given a second audiometric examination after completing the experiment to ensure that no permanent threshold shifts were produced by the sound exposures at any of the audiometric test frequencies. Audiometry was performed with a Grason-Stadler tracking audiometer (model 1703).

Tone bursts (pulses) were formed by passing an oscillator (Hewlett-Packard model 200 CD) signal through an electronic switch (Grason-Stadler model 1287), controlled by two timers (Grason-Stadler model 1216), to an attenuator set (Hewlett-Packard 350D) and a 40-W McIntosh amplifier. The signal from the amplifier activated a TDH 49 earphone which was attached to an MX-41/AR cushion. Sound-level calibrations were made using a General Radio octave-band noise analyser (model 1558EP) and a General Radio coupler (model 1560-P83). Absolute sound-pressure-level calibration was obtained with a General Radio piston

phone (model 1562-A).

The visual target was a black cross formed by a horizontal line and a vertical line (both 7.6×7.6 cm) drawn on a piece of white paper (21.6×27.9 cm). The target was placed, at approximately eye level, 150 cm directly in front of the seated subject, whose head was positioned in a U-shaped chin rest. A Naren spotlight (model N-103) illuminated the target, and the brightness at the subjects' eye was 100 mL.

Angular acceleration was obtained with a centrifuge. The centrifuge consisted of a 122-cm radius circular platform which was driven through a reduction gear by a 10-horsepower, variable-speed DC motor. A servo regulator maintained the centrifuge rotation rate at the level determined by the experimenter. The subject was restrained in a padded aircraft chair during centrifugation.

After the audiometric examination all of the subjects were shown the basic apparatus and told that the experiment involved exposure to loud sounds and rotation for a brief period. The subjects were then read the following instructions.

"I would like you to look at the cross on the wall while I present a series of loud beeps through the earphone. I would like to know if you see any changes in the cross while the sound is on. Other subjects have reported a variety of changes; some subjects report no changes at all. I would like to have you tell me about anything that you see."

In the first part of the experiment, the subjects were exposed to six stimulus trains (10 tone bursts in each train) presented to either the left or right ear and following rotation at 30, 60, or 90 deg/sec. Acoustical stimulus characteristics were held constant: frequency--500 Hz, intensity--125 dB SPL,

repetition rate--0.9/sec, duration--100 msec, and onset/offset time--1 msec.

Before the sound presentation, the centrifuge was slowly accelerated until a constant angular velocity of 30, 60, or 90 deg/sec was reached. The angular velocity was maintained for a period of 2 min after which the centrifuge was stopped as rapidly as possible (0.3 sec for 30 deg/sec, 0.85 for 60 deg/sec, and 1.3 for 90 deg/sec). The subject was then oriented toward the target and the earphone was placed over the right or left ear. After the subject's head was in the head holder, a series of 10 tone bursts was presented. A period of 0.7-0.9 sec elapsed between termination of rotation and the start of the tone burst presentation.

A trial consisted of exposure to the series of ten tone bursts to either the right or left ear following rotation at a particular rate. One trial was completed for each of the six experimental conditions (two ears by three rotation rates). The order of trials was counterbalanced and varied across subjects. A 2-min rest interval intervened between trials.

At the end of each trial the subject was asked to report any changes in the target associated with the acoustical stimuli. Only reports of target motion constituted a positive response; other reports (e.g., changes in color, line width, shadowing, and so on) were recorded as negative responses for the purposes of this experiment.

Those subjects who reported target motion on at least one of the six trials during the first part of the experiment proceeded to the second part. In this second part, the subjects were asked to estimate the amount of perceived target motion correlated with each of the six conditions. The subjects were read the following instructions.

"You have reported motion of the cross associated with the beeps. Now I would like you to estimate the amount of motion associated with beeps after rotation at different rates. I will present a series of beeps. Use any number you wish to indicate the amount of motion produced by the beeps. Then I will present another series of beeps after rotation at a different rate. If the amount of motion that you see is twice as large for the second series of beeps as for the first series, use a number twice as large as you used for the first series. If the amount of motion is half as great, use a number half as large. You may use any numbers that you wish--whole numbers, decimals, fractions--to indicate the amount of motion that you see. We ask only that you try to be consistent in the way you use the numbers."

The subjects assigned positive numbers based on their perception of target motion magnitude produced by the various stimuli. A modulus was not employed. As in the first part of the session, the subjects were exposed to six trains of tone bursts; the ten tone bursts that comprised a train were essentially identical (phase was not controlled).

Experiment I-B

Four Miami University graduate students were paid to complete Experiment I-B. All had participated in previous experiments and were highly practiced on the task.

The subjects were selected because they reported lateral visual shifts in one direction when stimulated in the left ear and shifts in the opposite direction when stimulated in the right ear. Two subjects reported maximum displacements at 500 Hz and the remaining two reported maximum displacements at 1000 Hz. Other acoustical stimulus characteristics were the same as in Experiment I-A.

The subjects completed four trials per day on eight consecutive days. A trial proceeded in the following manner: (a) four tone bursts were delivered to either the right or left ear (pretest); (b) the subject was centrifuged for 2 min at 30 or 60 deg/sec in the clockwise direction; (c) the centrifuge was abruptly stopped; and (d) four tone bursts were immediately presented (posttest). After the tone burst presentation the subject reported the magnitude of visual field shift, as previously, and the direction of the displacement.

The trials were divided into two blocks according to the rotation rate; that is, two trials were completed at one of the rotation rates during which the right (left) and then the left (right) ear was stimulated. After a 15-min rest, two more trials were completed employing the second centrifugation rate. A 5-min rest interval was allowed between trials. Each day the subject received pretest and posttest tone bursts presented to each ear coupled with each centrifugation rate. The order in which the trials were presented was counterbalanced across subjects.

B. EXPERIMENT II--HEAD VIBRATION

Seven Miami University graduate and undergraduate students served as subjects for pay. The subjects met the audiometric criteria employed in Experiment I. All seven subjects were pretested and reported lateral visual field displacements when exposed to acoustical stimuli of 125 dB at either 500 or 1000 Hz.

The apparatus employed for acoustical stimulation and the visual target was the same as previously.

Head vibration was obtained with a Sears Vibrator Massage unit (model 753-2259B). The vibrator exhibited spectral peaks at 120 Hz (reference—0 dB), 60 Hz (-9 dB), 180 Hz (-8 dB), 240 Hz (-9 dB), and 300 Hz (-12 dB). The plastic vibrator tip was held against the subject's right mastoid bone with a force of 300 gr. The force at the subject's skull was determined with a

Grass force/displacement transducer (model FT10), a Grass oscillograph (model 7), and a Tektronik storage oscilloscope (type 564).

The subjects were asked to estimate the magnitude of target motion produced by acoustical transients under each of eight conditions (with or without vibration, left or right ear, 500 or 1000 Hz). The instructions to the subjects were similar to those used in the second part of Experiment I.

Six of the seven subjects completed at least eight experimental sessions and one subject completed six sessions. Each session consisted of eight trials under each of the eight conditions indicated above. A trial consisted of presentation of four tone bursts (repetition rate--0.9/sec, duration--100 msec, onset/offset time--1 msec, intensity--125 dB SPL, frequency--500 or 1000 Hz). At the end of each group of four tone bursts, the subjects estimated the magnitude of visual field displacement. The order of presentation of the eight conditions within a session was counterbalanced within subjects across the eight sessions. At least 2 hr elapsed between sessions.

C. EXPERIMENT III--PRIOR VISUAL FIELD ROTATION

Experiment III-A

Six graduate students who received payment for their participation served as subjects in this experiment. The subjects were selected from a larger group of volunteers by means of a screening procedure. This screening was essentially the same as used in the first part of Experiment I-A except that rotation was not employed. Only subjects who reported visual field displacements to either 500 or 1000 Hz stimuli presented to either the left or the right ear participated in Experiment III. As before, all subjects were audiometrically tested to insure that their hearing did not deviate from audiometric zero by more than 20 dB and were retested after the experiment to ensure that no

permanent damage had occurred.

The acoustical stimulation apparatus was the same as that used in Experiment I. The visual target was also the same with the exception that the viewing distance was 122 cm rather than 150 cm.

Visual field rotation was produced by a drum 92 cm high and 92 cm in diameter. Its interior was lined with white illustration board with black vertical stripes 2 cm wide separated by 5 cm. The subjects' heads were restrained in a U-shaped chin rest with their eyes 33 cm below the top of the drum interior and at a viewing distance of 36 cm from the drum walls. The interior of the drum was illuminated by a 65-W incandescent bulb mounted overhead. For the fixation condition, the subjects fixated upon a 1-cm black paper disk, mounted at eye level 33 cm from the eyes on an L-shaped wire arm extending from the chin rest. For the scanning condition, the subjects tracked four 1-cm copper rivet heads mounted at eye level every 90 deg on the interior surface of the drum.

Those subjects who consistently reported target motion during screening were then assigned to one of two groups: left ear or right ear. Selection was determined by which ear when stimulated produced the most consistent reports of target motion. If no preference existed, subjects were randomly assigned to groups.

The subjects' instructions were essentially the same as those used in the second part of Experiment I.

Each subject participated in 16 sessions held on consecutive weekdays. For each session the subject received two acoustic stimulus trains, a 5-min exposure to the rotating drum, and then two more acoustic stimulus trains. Each stimulus train consisted of 20 tone bursts, initially at an

intensity of 80 dB and rising in steps until reaching 125 dB on the tenth and all succeeding tone bursts. The tone bursts had a duration of 100 msec, an onset/offset time of 1 msec, and an interstimulus interval of 0.9 sec.

During each session each subject received one 1000-Hz stimulus train and one 500-Hz train before drum exposure. Train frequency order was counterbalanced within and across sessions. Subjects were then immediately seated in the drum with their heads in the chin rest. They were then read the following instructions.

"I would like you to fixate upon the black dot in front of you" (fixation condition) or "without moving your head, follow with your eyes the copper rivets as they pass through your visual field. The drum will spin for 5 min after which I would like you to return to the target viewing chair as quickly as possible. Close your eyes please."

The subjects' eyes remained closed for 15 sec to allow the drum to reach a rotational speed of 12 rpm either toward the ear which received auditory stimulation (homolateral spin) or away from the stimulated ear (contralateral spin). Spin direction was counterbalanced across sessions. After 5 min of drum viewing, the subjects returned to the visual target-viewing chair where they again received auditory stimulus trains of 1000 and 500 Hz and gave magnitude estimates of perceived motion.

Experiment III-B

Four subjects from Experiment III-A participated in Experiment III-B. Experiment III-B was identical in methodology to III-A except on the following points. During drum stimulation subjects always followed the rivets with their eyes and received either homolateral spin or a control condition of no spin. Under the control condition subjects were still required

to close their eyes for 15 sec before drum exposure. Also, in an attempt to equate attentional states, subjects were required to mentally calculate and orally respond to multiplication problems presented auditorily for 60 sec before and after drum exposure. During those calculations the target sheet was covered by a plain sheet of white paper of the same size.

D. EXPERIMENT IV--TARGET ILLUMINATION

A total of 50 students enrolled in introductory psychology courses participated in Experiment IV either to satisfy an experiment participation course requirement or for pay. All subjects received an audiometric examination as in the previous experiments.

Nineteen subjects reported lateral visual field displacements during a screening procedure that was analogous to the first part of Experiment I-A.

The auditory stimulus apparatus and characteristics as well as the visual target were the same as those used in the previous experiments. The visual target was illuminated with the Naren spotlight which was powered through a variac. Light intensity was modified by varying the variac settings. Five light intensity settings (0.8, 2.3, 8.4, 36, and 110 m^L at the subject's eye), which were found to produce approximately equal steps of perceived illumination change during preliminary testing, were employed in this experiment.

The subjects were asked to estimate the magnitude of visual field displacement during exposure to acoustical stimulation (10 tone bursts) under each of the five target illumination levels, following instructions analogous to those employed in Experiment I-B. Only one frequency (the one which yielded the most consistent responses during screening) was used. The subjects reported visual displacements of the target at each of the five illumination levels. Order of illumination levels was

randomly varied across subjects.

E. EXPERIMENT V--ALCOHOL INGESTION

Six Miami University students who had participated in an earlier experiment were paid to serve as subjects. All subjects reported target motion upon presentation of acoustical transients. They were given an audiometric examination to ensure that their hearing thresholds did not deviate more than 20 dB from audiometric zero at all frequencies examined. Also, a second audiometric examination was given either immediately or one week following exposure to ensure that their ears had not suffered any permanent damage.

At the onset of the experiment the subjects were weighed to determine alcohol dosage. Weights ranged from 50.9 kg to 103.6 kg. One cc of 100-proof Vodka was given for each kg of body weight. Each alcohol dose was mixed with 480 cc of orange juice.

Subjects were run on four consecutive days. On two of the days subjects received the Vodka-orange juice drink and on the other two they were given plain orange juice. The order in which subjects received the two types of drink was counter-balanced. Three subjects received the alcoholic drink on their first and last session and the other three on their second and third session.

Subjects were tested four times each day. The first test session was performed before ingesting either the alcoholic or nonalcoholic drink. Following this session, the subjects were allowed 30 min to drink either type of beverage and were subsequently tested. Subjects were tested again 30 and 90 min later. Therefore, on each day the subjects were given a pretest and three posttests extended over a two-hr session.

During each testing session, subjects were exposed to four stimulus trains comprised of four tone bursts in each train.

Two tone bursts were delivered to each ear at 500 and 1000 Hz at an intensity of 125 dB. Each testing session began with the tone bursts being presented to the left ear at 500 Hz and 1000 Hz respectively. The subjects assigned positive numbers based on their perception of target motion magnitude. A modulus was not employed.

SECTION III

RESULTS

A. EXPERIMENT I--ANGULAR ACCELERATION


Experiment I-A

No differences between ears in the incidence or magnitude of visual field shifts evoked by acoustical transients were found after $-\alpha_2$ (right yaw) angular acceleration/deceleration. The results that lead to this conclusion are illustrated in Tables 1 and 2 and Fig. 1. Table 1 summarizes the percentage of reports of target motion following rotation at various rates when the acoustical transients were presented to either the left or right ears. No clear pattern emerged from the results for the first part of the experiment. Table 2 presents the magnitude estimation data obtained from the twelve subjects who completed the experiment. The geometric means of the magnitude estimations, calculated across subjects, as a function of ear stimulated and rotation rate are illustrated in Fig. 1. The data from the experiment were analyzed by an analysis of variance (after log or z-score transformation). Two-group differences were evaluated employing the Wilcoxon matched-pairs signed-ranks test. In no case, including the difference between ears following 5 rpm rotation, did the results of the various statistical tests approach significance.

TABLE 1
REPORTS OF TARGET MOTION EVOKED BY ACOUSTICAL
TRANSIENTS FOLLOWING ROTATION AT 5, 10, AND
15 rpm (%)

	rpm		
	5	10	15
Left ear	67	83	75
Right ear	67	50	83

PERCEIVED VISUAL FIELD DISPLACEMENT - GEOMETRIC MEAN



ROTATION RATE - deg/sec

Figure 1. Geometric means of perceived visual field displacements as a function of rotation rate of centrifuge prior to acoustical stimulation. No differences were obtained when the right ear was stimulated versus when the left ear was stimulated.

TABLE 2
MAGNITUDE ESTIMATIONS--EXPERIMENT I-A

Subject	5 rpm		10 rpm		15 rpm	
	Left ear	Right ear	Left ear	Right ear	Left ear	Right ear
1	0	0	10	10	20	25
2	25	17	20	10	25	10
3	2	2.3	2	3	1.5	3
4	10	3	10	7	5	5
5	2	2	8	1	5	3
6	6	4	4	2	1	4
7	0.5	0	0.5	0	1	0
8	70	20	25	100	10	15
9	1	0	1	3	5	4
10	40	60	10	9	15	10
11	10	15	15	20	5	20
12	3	25	0	1	0.5	0

Experiment I-B

Two subjects reported visual field displacements in the direction of the stimulated ear, and the remaining two subjects reported displacements away from the stimulated ear.

The results from Experiment I-B are illustrated in Fig. 2. The data were normalized within subjects and averaged across subjects. The ordinate of Fig. 2 indicates the average normalized perceived visual field displacement prior to rotation minus the average normalized displacement following rotation. The figure indicates that the reports of visual field displacement were diminished following rotation at 60 deg/sec (relative to 30 deg/sec) when the subjects reported that the pretest acoustical stimulus produced a leftward displacement of the visual field. The opposite pattern of results was obtained when




Figure 2. Means of Z-score transformed perceived visual field displacement reports as a function of centrifuge rotation rate. Each data point represents the difference between the reports before and after centrifugation. When the acoustical stimuli produced a leftward visual field shift, centrifugation reduced the apparent visual field displacement; when acoustical stimuli produced a rightward visual field shift, centrifugation enhanced the apparent visual field displacement.

the subjects reported rightward displacement following acoustical stimulation during the pretest.

An analysis of variance calculated from the data illustrated in Fig. 2 yielded an interaction term (rotation rate by direction of displacement during the pretest) which approached significance ($F = 3.88$; $df = 1, 3$; $p = 0.14$). The fact that this F ratio did not reach the normal statistical significance level can be attributed primarily to the limited number of degrees of freedom. If more subjects who exhibited the appropriate pretest pattern of visual field displacements could have been located, we are confident that statistically significant results would have been obtained. However, less than 10 percent of the population appears to exhibit the required response pattern.

B. EXPERIMENT II--HEAD VIBRATION

Subjects reported that the magnitudes of the visual field shifts evoked by the acoustical transients were reduced by nearly 50% when the transients were presented with simultaneous head vibration (see Fig. 3). A three-way analysis of variance calculated from the log-transformed magnitude estimation reports and averaged across subjects indicated a statistically significant difference between the magnitude estimations with the vibration present and the vibration absent ($F = 11.94$; $df = 1, 6$; $p = 0.01$). The interaction between ear stimulated and vibration was not significant ($F = 0.06$; $df = 1, 6$; $p = 0.82$), indicating that the effects of vibration were the same whether the acoustical stimulus was applied to the contralateral (left) or homolateral (right) ear.

C. EXPERIMENT III--PRIOR VISUAL FIELD ROTATION

The results of Experiment III are summarized in Table 3, which contains the geometric means of the magnitude estimation observations. A repeated measure analysis of variance is summarized in Table 4. As indicated in Table 3, the magnitude estimations were lower after exposure to the rotating visual

**PERCEIVED VISUAL FIELD DISPLACEMENT -
GEOMETRIC MEAN**




Figure 3. Geometric means of reported visual field displacements as a function of ear stimulated. Simultaneous presentation of vibration and acoustical stimulation reduced the apparent visual field displacement.

field in Experiment III-A; the difference between magnitude estimations before and after drum exposure was significant at the 0.05 probability level (see Table 4). Neither of the other two main effects (drum rotation direction and scan pattern during rotation) reached statistical significance. Analysis of the results from Experiment III-B indicated that neither the magnitude estimation differences before and after drum exposure nor the differences associated with drum rotation (spin vs. no spin) reached statistical significance.

TABLE 3
GEOMETRIC MEANS--EXPERIMENT III

	<u>Experiment III-A</u>			
	<u>Before condition</u>		<u>After condition</u>	
	Homo direction	Contra direction	Homo direction	Contra direction
1000 Hz				
Fix eye movement	5.00	4.47	4.19	2.86
Scan eye movement	3.98	4.15	2.17	2.55
500 Hz				
Fix eye movement	5.71	3.47	3.56	2.89
Scan eye movement	3.54	3.70	1.93	2.88
	<u>Experiment III-B</u>			
	<u>Before condition</u>		<u>After condition</u>	
1000 Hz				
Spin stimulation	3.72		2.79	
No spin stimulation	3.78		3.85	
500 Hz				
Spin stimulation	2.29		2.30	
No spin stimulation	3.15		3.47	

TABLE 4
 REPEATED MEASURES ANALYSES OF VARIANCE FOR EXPERIMENT III


Source	F	df	P
Experiment III-A--1000 Hz data			
Before/after	6.41	1, 5	0.05
Homolateral/contralateral spin	0.03	1, 5	0.87
Fixation/scan	3.46	1, 5	0.12
Experiment III-A--500 Hz data			
Before/after	6.56	1, 5	0.05
Homolateral/contralateral spin	0.03	1, 5	0.86
Fixation/scan	0.97	1, 5	0.37
Experiment III-B--1000 Hz data			
Before/after	0.65	1, 3	0.48
Spin/no spin	0.81	1, 3	0.43
Experiment III-B--500 Hz data			
Before/after	0.36	1, 3	0.59
Spin/no spin	3.08	1, 3	0.18

D. EXPERIMENT IV--TARGET ILLUMINATION

Of the 19 subjects who completed the observations in Experiment IV, five subjects received stimulation at 1000 Hz and 14 subjects were exposed to the acoustical transients at 500 Hz. The magnitude estimation responses from the 19 subjects were combined and the geometric means are presented in Fig. 4. The trend of the data suggests that the higher the level of illumination, the larger the magnitude estimation response; however, the results of a randomized blocks analysis of variance on normalized (within subjects) data failed to provide statistical support for the trend ($F = 1.06$; $df = 4, 68$; $p = 0.38$).

E. EXPERIMENT V--ALCOHOL INGESTION

Alcohol ingestion resulted in diminished magnitude estimation reports. Fig. 5 illustrates that the geometric means for




TARGET BRIGHTNESS - mL

Figure 4. Geometric means of visual field displacement reports as a function of target brightness. The tendency for larger visual field displacements to correlate with target brightness was not statistically significant.

PERCEIVED VISUAL FIELD DISPLACEMENT -

Figure 5. Geometric means of perceived visual field displacement reports as a function of time after ingestion of orange juice or orange juice plus alcohol. Alcohol suppressed visual field displacements.



the magnitude estimations following alcohol consumption were approximately one-half as large as those obtained when the subjects drank orange juice alone. The magnitude estimations following alcohol consumption reached a minimum immediately following ingestion and then gradually returned toward the no-alcohol level, whereas magnitude estimations obtained during the no-alcohol tests exhibited a gradual decline across time within a session. An analysis of variance on subject means (collapsed across days, ears, and frequencies) revealed statistically significant effects associated with the presentation of alcohol ($F = 8.68$; $df = 1, 5$; $p < 0.05$), time of observation ($F = 3.61$; $df = 3, 15$; $p < 0.05$), and the interaction between alcohol and time ($F = 7.31$; $df = 3, 15$; $p < 0.05$).

SECTION IV

DISCUSSION

A. EXPERIMENT I--ANGULAR ACCELERATION

The results of Experiment I-A failed to support the hypothesis that the incidence and magnitude of visual field shifts would differ following $-\alpha_2$ angular acceleration/deceleration when the acoustical stimuli were presented to the left or right ear. This result was also contrary to the impressions reported by several subjects who participated in the experiment.

The direction of the visual field displacements reported by the subjects in Experiment I-A differed across subjects. This observation provided an interpretation of the failure to obtain statistically significant results in Experiment I-A and the hypothesis for Experiment I-B. While the angular acceleration stimuli in Experiment I-A may have had strong effects in particular individuals, these effects varied across subjects and disappeared when the group data were evaluated. This variability may be due to the specific pattern of inner ear fluid displacement produced by the acoustical transients in particular individuals. In order to reliably predict the effect of angular acceleration, we would have to know the specific pattern of inner ear displacement produced by the acoustical transients in different persons, as indicated by the direction of perceived cross motion elicited by stimulation of a particular ear.

The observations obtained in Experiment I-B (Fig. 2) support the view that angular deceleration can modify reports of visual field displacements. These observations can be described in terms of the reafference interpretation of the visual field displacement phenomenon described previously (ref. 16). Because the spin table rotated in the counterclockwise direction, the initial direction of angular deceleration-induced eye motion was

toward the subject's left. According to the reafference interpretation, visual field displacement is reported when there is a mismatch between the eye movement expected on the basis of the vestibular signal and the actually received signal concerning the eye movement. To the extent that this mismatch is large, the reported visual field shift would be large. Figure 2 illustrates that the magnitudes of visual field displacement following angular deceleration were reduced when subjects reported leftward visual field displacement prior to rotation. This would be expected from the reafference interpretation because the angular deceleration-induced tendency of the eye to move to the subject's left would reduce the mismatch between the expected and actually received eye movement signals. On the other hand, when the subjects reported a rightward visual field displacement prior to rotation, the angular deceleration resulted in slightly larger reports of visual field displacement. Following the reafference interpretation, the deceleration-induced tendency of the eye to drift toward the subject's left would increase the mismatch between the expected rightward movement from the vestibular receptors and the actually received eye movement signals.

The observation that angular deceleration resulted in alteration of visual field displacement reports supports the hypothesis that the acoustical stimuli acted by displacement of semicircular canal structures.

Following the reafference interpretation, it can be suggested that environmental conditions that enhance visual fixation or that result in a real tendency of the eye to move in a direction opposite to that signaled by the vestibular receptors would enhance reports of sound-evoked visual field displacement.

B. EXPERIMENT II--HEAD VIBRATION

The results of Experiment II indicate that vibration acts to attenuate visual field shifts associated with acoustical

stimulation. The mechanism of this attenuation is probably the middle ear reflex. Informal observations indicate that mastoid vibration elicits a middle ear reflex, which would limit the sound energy entering the labyrinth for a given intensity of sound at the tympanic membrane. This interpretation is supported by the observation that the vibration-induced response reduction was equally effective for the homolateral or contralateral ear because the middle ear reflex is bilateral.

In the introduction it was suggested that vibration might enhance or suppress the sound-evoked visual field displacements depending on whether the vibration was homolateral or contralateral to the sound. Clearly, this suggestion was not supported by Experiment II; rather, the suppression, which was apparently mediated by the middle ear reflex, was sufficient to mask any interaction between the two types of stimulation at the semi-circular canals.

An alternative mechanism for the vibration-induced reduction in the visual displacements reported by the subjects can be proposed. The head vibration may have been sufficient to reduce visual acuity by blurring the image of the target cross on the retina. However, previous research on the effects of vibratory frequencies in the range employed in Experiment II suggests that acuity decrements are negligible (ref 4).

C. EXPERIMENT III--PRIOR VISUAL FIELD EXPOSURE

Experiment III-A resulted in a significant reduction in the magnitude of the visual field shifts evoked by acoustical transients after exposure to the rotating visual field. One interpretation of this result was that visual field rotation exposure produced a central "depression" of visual/vestibular reflexes. Alternatively, the observed differences might have resulted from fatigue, loss of attention, or some other "order effect." In order to evaluate the second interpretation, Experiment III-B was performed. If the depression in magnitude

estimation reports were due to drum rotation, a difference between the spin and no spin groups should have been observed. This was not the case. Further, if the drum exposure produced a decrement in the visual field shift response, a before/after difference should have been observed. The fact that the before/after difference was lost in Experiment III-B suggests that the difference found in Experiment III-A was due to loss of arousal because the major procedural difference between Experiments III-A and III-B was the use of mental arithmetic problems. Mental arithmetic has been used in several previous investigations to maintain eye movement reflexes mediated by the vestibular system and apparently this procedure helps to maintain the amplitude of sound-evoked visual field shifts.

D. EXPERIMENT IV--TARGET ILLUMINATION

The hypothesis that target illumination intensity would be inversely related to visual field shift magnitude was not supported. In view of these results, an alternative proposal can be advanced. This alternative is based on the suggestions advanced previously to account for the failure to correlate eye movements with reports of visual field shift (ref 16). As noted in the discussion of Experiment I-B, it was suggested that the perception of visual field shifts following exposure to acoustical transients resulted from a discrepancy between expected and actually received information regarding changes of stimulation to the retina, following a reafference model. Intense target illumination might serve to increase stability of visual fixation, exacerbating the discrepancy, thus resulting in the perception of increased motion.

E. EXPERIMENT V--ALCOHOL INGESTION

The results of Experiment V support the hypothesis that alcohol ingestion would result in smaller reports of perceived visual field displacements following stimulation with high intensity acoustical transients. At least three interpretations can be advanced to account for the results of Experiment V.

First, the reduction in perceived visual field displacement may reflect a general suppression of the neural pathway between the visual and vestibular systems, as suggested by Schroeder (ref 21). Second, the alcohol may have decreased the precision of the eye position control system. If the eye is spontaneously "wandering" it would be difficult for the subjects to detect sound-evoked visual field shifts. This interpretation is congruent with the difficulties in target fixation reported by the subjects following angular deceleration and noted previously. Third, the alcohol may have impaired the subjects' arousal or attention. The results of Experiment II-B support the view that lowered arousal or attention might be associated with lower reports of sound-evoked visual field displacements.

SECTION V

CONCLUSIONS

Intense acoustical transients evoke reports of visual field displacement in human subjects. The nature of the displacement reported varies across subjects and as a function of stimulus characteristics (particularly intensity and frequency). Experimental manipulations that increase the ability of subjects to maintain visual fixation or that tend to evoke eye movements in a direction opposite to that expected on the basis of the acoustical stimulation increase the magnitudes of the visual field displacements reported. Disruption of the visual fixation mechanism (e.g., by alcohol ingestion) or reduction of the effective acoustical stimulus at the labyrinth by vibration tend to reduce the magnitudes of the visual field displacements reported by the subjects.

REFERENCES

1. Benson, A. J., and Guedry, F. E., "Comparison of Tracking Task Performance and Nystagmus During Sinusoidal Oscillation in Yaw and Pitch," Aerospace Med. 42, 593-601, 1971.
2. Collins, W. E., Gilson, R. D., Schroeder, D. J., and Guedry, F. E., "Effects of Alcohol Ingestion on Tracking Performance During Angular Acceleration," J. Appl. Psychol. 55, 559-563, 1971.
3. Dichgans, J., Held, R., Young, L. R., and Brandt, T., "Moving Visual Scenes Influence Apparent Direction of Gravity," Science 178, 1217-1219, 1972.
4. von Gierke, H. E., and Clarke, N. P., "Effects of Vibration and Buffeting in Man," In Aerospace Medicine, H. Randal (Ed.), Williams and Wilkins Co., Baltimore, 1971.
5. Gilson, R. D., Schroeder, D. J., Collins, W. E., and Guedry, F. E., Jr., "Effects of Different Alcohol Dosages and Display Illumination on Tracking Performance During Vestibular Stimulation," Aerospace Med. 43(6), 656-660, 1972.
6. Gilson, R. D., Guedry, F. E., and Benson, A. J., "Influence of Vestibular Stimulation and Display Luminance on the Performance of a Compensatory Track Task," Aerospace Med. 41, 1231-1237, 1970.
7. Harris, C. S., Effects of Acoustical Stimuli on the Vestibular System, Technical Report 71-58, Aerospace Medical Research Laboratory, Wright-Patterson Air Force Base, Ohio, 1971.

8. Harris, C. S., Effects of Increasing Intensity Levels of Intermittent and Continuous 1000 Hz Tones on Human Equilibrium, Technical Report 72-11, Aerospace Medical Research Laboratory, Wright-Patterson Air Force Base, Ohio, 1972. [AD 750840]
9. Harris, C. S., and Sommer, H. C., Human Equilibrium During Acoustical Stimulation by Discrete Frequencies, Technical Report 68-7, Aerospace Medical Research Laboratory, Wright-Patterson Air Force Base, Ohio, 1968. [AD 675172]
10. Heifer, U., "Oculovestibuläre Reaktion und Blutalkoholkonzentration," Hefte zur Unfallheilkunde 99, 192-197, 1969.
11. Howard, I. P., and Templeton, W. B., Human Spatial Orientation, John Wiley and Sons, London, 1966.
12. Lackner, J. R., and Graybiel, A., "Elicitation of Vestibular Side Effects by Regional Vibration of the Head," Aerospace Med. 45, 1267-1272, 1974.
13. Parker, D. E., and von Gierke, H. E., "Vestibular Nerve Response to Pressure Changes in the External Auditory Meatus of the Guinea Pig," Acta Oto-laryng. 71, 456, 1971.
14. Parker, D. E., von Gierke, H. E., and Reschke, M. F., "Studies of Acoustical Stimulation of the Vestibular System," Aerospace Med. 39, 1321, 1968.
15. Parker, D. E., and Reschke, M. F., "Mechanisms of Acoustical Vestibular Stimulation I. Middle Ear Nonlinearity," Minerva Otorinolaringologica 22, 240-249, 1972.

16. Parker, D. E., Tubbs, R. L., and Littlefield, V. M., "Visual-Field Displacements in Human Beings Evoked by Acoustical Transients," J. Acoust. Soc. Amer. 63, 1912-1918, 1978.
17. Parker, D. E., Tubbs, R. L., Ritz, L. A., and Wood, D. L., Effects of Sound on the Vestibular System, Technical Report 75-89, Aerospace Medical Research Laboratory, Wright-Patterson Air Force Base, Ohio, 1976.
18. Reschke, M. F., High-intensity, Audiofrequency Vestibular Stimulation in the Guinea Pig, Unpublished Doctoral Dissertation, Department of Psychology, Miami University, Oxford, Ohio, 1971.
19. Reschke, M. F., Homick, J., Landreth, J., and Parker, D. E., "Acoustical Vestibular Stimulation in Man," Minerva Otorinolaringologica 25, 1975.
20. Reschke, M. F., Parker, D. E., and von Gierke, H. E., "Stimulation of the Vestibular Apparatus in the Guinea Pig by Static Pressure Changes: Head and Eye Movements," J. Acoust. Soc. Amer. 48, 913, 1970.
21. Schroeder, D. J., "Influence of Alcohol on Vestibular Responses to Angular Acceleration," Aerospace Med. 42, 959-970, 1971.