

AD-A085 574

FRANK J SEILER RESEARCH LAB UNITED STATES AIR FORCE A--ETC F/G 7/4
MASS SPECTRAL FRAGMENTATION OF A NOVEL CYCLOOCTANE: 2,4-DIMETHY--ETC(U)
MAY 80 R R MCGUIRE, S A SHACKELFORD

UNCLASSIFIED

ML

1 of 1
5/8 6/8 1/2



END
DATE
FILMED
7-80
DTIC



4
LIVEL # 2



FRANK J. SEILER RESEARCH LABORATORY

**FJSRL TECHNICAL REPORT 80-0012
MAY 1980**

**MASS SPECTRAL FRAGMENTATION OF A NOVEL CYCLOOCTANE:
2,4-DIMETHYL-7,7-DINITRO-1,3,5-TRIOXACYCLOOCTANE**

**RAYMOND R. MCGUIRE¹
SCOTT A. SHACKELFORD**

**DTIC
ELECTRONIC
S JUN 18 1980**

**APPROVED FOR PUBLIC RELEASE;
DISTRIBUTION UNLIMITED.**

PROJECT 2303

**AIR FORCE SYSTEMS COMMAND
UNITED STATES AIR FORCE**

80 6 16 186

ADA 085574



FJSRL-TR-80-0012


This document was prepared by the Energetic Materials Division, Directorate of Chemical Sciences, Frank J. Seiler Research Laboratory, United States Air Force Academy, CO. The research was conducted under Project Work Unit Number 2303-F3-01, Energetic Materials Research, Capt Robert A. Hildreth was the Project Scientist in charge of the work.

When U.S. Government drawings, specifications or other data are used for any purpose other than a definitely related government procurement operation, the government thereby incurs no responsibility nor any obligation whatsoever, and the fact that the government may have formulated, furnished or in any way supplied the said drawings, specifications or other data is not to be regarded by implication or otherwise, as in any manner licensing the holder or any other person or corporation or conveying any rights or permission to manufacture, use or sell any patented invention that may in any way be related thereto.


Inquiries concerning the technical content of this document should be addressed to the Frank J. Seiler Research Laboratory (AFSC) FJSRL/NC, USAF Academy, CO 80840. Phone AC 303 472-2655.

This report has been reviewed by the Commander and is releasable to the National Technical Information Service (NTIS). At NTIS it will be available to the general public, including foreign nations.


This technical report has been reviewed and is approved for publication.



ROBERT A. HILDRETH, Captain, USAF
Project Scientist



KENNETH E. SIEGENTHALER, Lt Colonel, USAF
Director, Chemical Sciences



WILLIAM D. SIURU, JR., Lt Colonel, USAF
Commander

Copies of this report should not be returned unless return is required by security considerations, contractual obligations, or notice on a specific document.

Printed in the United States of America. Qualified requestors may obtain additional copies from the Defense Documentation Center. (All others should apply to: National Technical Information Service
6285 Port Royal Road
Springfield, Virginia 22161)

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER 14 FJSRL-TR-80-1012	2. GOVT ACCESSION NO. ADA085574	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) 6 Mass Spectral Fragmentation of a Novel Cyclooctane 2,4-Dimethyl-7,7-Dinitro-1,3,5-Trioxacyclooctane		5. TYPE OF REPORT & PERIOD COVERED
7. AUTHOR(s) 10 Raymond R. McGuire Scott A. Shackelford		6. PERFORMING ORG. REPORT NUMBER
9. PERFORMING ORGANIZATION NAME AND ADDRESS Frank J. Seiler Research Laboratory (AFSC) FJSRL/NC USAF Academy, CO 80840		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS 16 61102F 2303/F3-01
11. CONTROLLING OFFICE NAME AND ADDRESS F. J. Seiler Research Laboratory (AFSC) FJSRL/NC USAF Academy, CO		12. REPORT DATE 11 May 1980
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		13. NUMBER OF PAGES 17 F3
16. DISTRIBUTION STATEMENT (of this Report) 12 14 Approved for public release; distribution unlimited.		15. SECURITY CLASS. (of this report) UNCLASSIFIED
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report) 9 Technical rept.		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE N/A
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Mass spectra 2,4-dimethyl-7,7-dinitro-1,3,5-trioxacyclooctane geminal polynitroalkanes cyclooctanes trioxane acetals cyclic acetals		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) A novel cyclooctane containing structural features unique to both geminal poly-nitroalkanes and cyclic trioxane acetals afforded an initial electron impact fragmentation analogous to cyclic acetals. Only in the latter stages of the fragmentation pathway did species appear that are characteristic of geminal polynitroaliphatic and nitroalkane molecules.		


UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

319920

SM

SECURITY CLASSIFICATION OF THIS PAGE(When Data Entered)



11

SECURITY CLASSIFICATION OF THIS PAGE(When Data Entered)

PREFACE

This document, FJSRL-TR-80-0012, summarizes detailed mass spectroscopic studies done on the novel cyclooctane: 2,4-dimethyl-7,7-dinitro-1,3,5-trioxacyclooctane. The work described here was done under Work Unit 2303-F3-01 and was not previously published by the authors.

TABLE OF CONTENTS

<u>SECTION</u>	<u>PAGE</u>
Preface	i
List of Schemes	iii
List of Tables	iii
Introduction.	1
Results and Discussion.	1
Experimental	7
Acknowledgements.	7
References	7

LIST OF SCHEMES

	<u>PAGE</u>
Scheme 1	3
Scheme 2	3

LIST OF TABLES

TABLE 1	4
-------------------	---

INTRODUCTION

A novel cyclooctane containing structural features unique to both geminal polynitroalkanes and cyclic trioxane acetals afforded an initial electron impact fragmentation analogous to cyclic acetals. Only in the latter stages of the fragmentation pathway did species appear that are characteristic of geminal polynitroaliphatic and nitroalkane molecules.

The subject compound, 2,4-dimethyl-7,7-dinitro-1,3,5-trioxacyclooctane (I), contains two distinctly different chemical structural features representative of a substituted cyclic trioxane acetal and a geminal polynitroalkane. While separate electron impact studies have been reported for cyclic acetals (e.g., dioxanes and trioxanes)^{2,3} and for nitrated alkane derivatives⁴⁻⁶, the limited nitroalkane mass spectral studies offer no clue to the geminal polynitroalkane structure's electron impact stability. Compound I allows a direct intramolecular comparison of electron impact fragmentation stabilities between a cyclic trioxane acetal structure and the geminal dinitroalkane grouping because of its unique incorporation of both structural features in a single compound.

RESULTS AND DISCUSSION

Neither nitro- nor polynitroalkanes form an appreciable molecular ion⁴⁻⁶. Polynitroalkanes characteristically lose NO_2 and NO ; while nitroalkanes generally release NO_2 and HNO_2 to yield charged alkyl species that undergo further fragmentation. The geminal polynitroalkane, 2,2-dinitropropane, a compound closely resembling the polynitroalkane structure in (I), gives predominantly NO^+ (m/e 30)⁶. A moderately intense ion at m/e 39 (C_3H_3)⁺ is generated by stepwise loss of NO_2 , then NO with oxygen migration to the α carbon to form a charged acetone species, and finally, loss of an OH radical.

An appreciable ion at m/e 43 (CH_3CO^+) was also observed and postulated to form from the charged acetone species by loss of a methyl radical⁶.

Cyclic ethylene acetals likewise do not afford a molecular ion, but lose an alkyl radical to form resonance stabilized oxonium ions². Symmetrical trioxane closely resembles the acetal chemical structure found in (I) and exhibits the loss of one methylene hydrogen atom to form an oxonium ion³. Subsequent fragmentation appears to be dictated by the resultant oxonium ion structure.

The initial fragmentation by (I) follows the same electron impact cleavages observed for in normal alicyclic acetals (Scheme 1)^{2,3}. The spectrum shows no molecular ion at m/e 236 (Table I), and loss of a methyl radical to produce the m/e 221 oxonium ion is heavily favored over a hydrogen atom cleavage to yield the m/e 235 oxonium species^{2,3}. The lesser m/e 235 oxonium fragment appears to form four additional ions at m/e 101, 86, 59, and 43, all by loss of neutral species containing the intact geminal dinitro group. The m/e 86 ion ($\text{C}_4\text{H}_6\text{O}_2$) is one of two mass doublets found and confirmed by high resolution mass spectral analysis.

Fragmentation reactions generated from the more favored m/e 221 oxonium ion produce four additional ions (Scheme 2); however, only one species (m/e 45), characteristic of cyclic acetal fragmentation noted of the m/e 235 ion, is formed by the loss of a neutral geminal dinitroalkane molecule. The three remaining ions produced from the eight-membered m/e 221 cyclic oxonium ion are likely six-membered cyclic geminal dinitroalkyl oxonium derivatives formed from normal cyclic acetal fragmentation mechanisms. The m/e 193 molecule forms by a CO loss²; the m/e 191 oxonium species is generated by the loss of formaldehyde (CH_2O); and the base peak, m/e 177,

TABLE 1. RELATIVE ION ABUNDANCES

<u>m/e</u>	<u>Percent</u>	<u>Formula</u>
236 [M] ⁺	0	C ₇ H ₁₂ N ₂ O ₇
235	0.4	C ₇ H ₁₁ N ₂ O ₇
<u>221</u>	15	C ₆ H ₉ N ₂ O ₆
<u>193</u>	28	C ₅ H ₉ N ₂ O ₆
<u>191</u>	16	C ₅ H ₇ N ₂ O ₆
<u>177</u>	100	C ₄ H ₅ N ₂ O ₆
<u>149</u>	20	C ₃ H ₅ N ₂ O ₅
<u>146</u>	6	C ₅ H ₈ NO ₄
<u>145</u>	7	C ₅ H ₇ NO ₄
115	6	C ₄ H ₅ NO ₃ or C ₅ H ₇ O ₃
<u>102</u>	16	C ₃ H ₄ NO ₃
<u>101</u>	27	C ₄ H ₅ O ₃
<u>86</u> (doublet)	16	C ₃ H ₄ NO ₂ C ₄ H ₆ O ₂
<u>85</u>	12	C ₄ H ₅ O ₂
74	8	C ₃ H ₆ O ₂
73	4	C ₃ H ₅ O ₂
<u>59</u>	9	C ₂ H ₃ O ₂
<u>56</u> (doublet)	9	C ₂ H ₂ NO C ₃ H ₄ O
45	40	CNO ₂
43	49	C ₂ H ₃ O
31	14	CH ₂ OH
30	33	CH ₂ O, NO(?)

Underlined ions confirmed by high resolution mass spectrometry.

forms from the $\underline{m/e}$ 221 oxonium ion splitting out an acetaldehyde molecule (CH_3CHO). Meta stable peaks at $\underline{m/e}$ 168.5 and $\underline{m/e}$ 141.5 establish the 193 and 177 ions respectively as having the common $\underline{m/e}$ 221 precursor.

Only at this point do fragmentation mechanisms characteristic of nitroalkane or geminal dinitroalkanes begin to appear. While the $\underline{m/e}$ 193 oxonium ion splits out acetaldehyde in a characteristic cyclic acetal fragmentation to form $\underline{m/e}$ 149, it also can lose an HNO_2 molecule to yield $\underline{m/e}$ 146⁵. The $\underline{m/e}$ 146 fragment continues by splitting out acetic acid to give the second $\underline{m/e}$ 86 mass doublet ($\text{C}_3\text{H}_4\text{NO}_2$). Loss of NO than affords $\underline{m/e}$ 56 ($\text{C}_3\text{H}_4\text{O}$) which also is one ion comprising another mass doublet found in this mass spectrum. The $\underline{m/e}$ 149 species is associated by a metastable appearing at 115 to the $\underline{m/e}$ 193 precursor ion. The $\underline{m/e}$ 149 species also splits out a neutral HNO_2 molecule to yield $\underline{m/e}$ 102. This molecule can lose formaldehyde to generate the other $\underline{m/e}$ 56 ($\text{C}_2\text{H}_2\text{NO}$)⁺ mass doublet, or it can cleave out mononitroacetylene to eventually provide $\underline{m/e}$ 30 (CH_2O^+) via $\underline{m/e}$ 31 (CH_2OH^+). The $\underline{m/e}$ 30 fragment could also be attributed to initial formation of NO^+ from the original geminal dinitro alicyclic eight and/or six-membered oxonium molecules since 2,2-dinitropropane affords an intense $\underline{m/e}$ 30 (NO^+) peak. However, it should be noted that no other transformations characteristic of 2,2-dinitropropane are observed in Schemes 1 and 2. The remaining peak at $\underline{m/e}$ 191 formed by loss of formaldehyde from $\underline{m/e}$ 221 initiates the only remaining fragmentation pathway that contains typical nitroalkane or geminal polynitroalkane cleavages. Although a seemingly minor process, $\underline{m/e}$ 191 cleaves an NO_2 radical to form $\underline{m/e}$ 145. The $\underline{m/e}$ 145, fragment like $\underline{m/e}$ 102, can liberate mononitroacetylene to yield $\underline{m/e}$ 73 via $\underline{m/e}$ 74; or, it can form two possible $\underline{m/e}$ 115 species by losing neutral CH_2O or by undergoing

another characteristic nitroalkane process and cleaving an NO radical. While the m/e 191 fragment could have formed from m/e 235 by loss of acetaldehyde to provide a species analogous to m/e 177, this pathway is ruled out since all reasonable subsequent reactions would lead to charged species not observed in this spectrum.

In summary, the subject compound, 2,4-dimethyl-7,7-dinitro-1,3,5-trioxacyclooctane (I), represents a molecule of novel chemical structure that contains features common to both alicyclic trioxane acetals and geminal dinitroalkanes. A mass spectrum typical of alicyclic acetals predominates and produces cyclic oxonium ions and low molecular weight CHO molecules as major species. Only relatively late in the fragmentation mechanism are transformations observed that represent pathways characteristic of nitro- and polynitroalkanes. There seems to be a reversal in the fragmentation pattern of the nitroalkane species in that the geminal dinitroalkane ions resulting from (I) parallel mass spectral fragmentations observed for pure mononitroalkanes^{4,5}, while mononitroalkane ions from (I) generally fragment as did polynitroalkanes in a previous study⁶. The results described herein suggest the geminal dinitroalkane group is more stable to electron impact fragmentation than one might suspect (a priori). Its electron impact stability apparently exceeds that exhibited by the cyclic acetal structure as demonstrated by the subject compound whose novel chemical architecture provides a direct intramolecular comparison between these two structural groupings.

EXPERIMENTAL

The mass spectra cited were obtained with a DuPont Instrument 21-491 double focusing mass spectrometer at an ionizing voltage of 78 eV and source temperature around 180°C. The sample was introduced into the mass spectrometer by a direct insertion probe. The compound, 2,4-dimethyl-7,7-dinitro-1,3,5-trioxacyclooctane (mp 54-57°C), gave the following analysis: (Found: C, 35.4; H, 4.9; N, 12.0. $C_7H_{12}N_2O_7$ required C, 35.6; H, 5.1; N, 11.9).

ACKNOWLEDGEMENTS

High resolution mass spectra were supplied by the Air Force Materials Laboratory and Los Alamos Scientific Laboratory through the efforts of L. D. Smithson and E. D. Loughran, respectively. We are indebted to P. Brown (Arizona State University), T. O. Tiernan (Wright State University), and W. D. Closson (SUNY Alb.) for constructive comments.

REFERENCES

1. Present Address: L-402, Lawrence Livermore Laboratory, Livermore, CA 94513.
2. J. T. B. Marshall and D. H. Williams, Tetrahedron **23**, 321 (1967).
3. H. B. Budzikiewicz, C. Djerassi, and D. H. Williams, "Mass Spectrometry of Organic Compounds," Holden-Day, San Francisco, pp. 257-258 (1967).
4. J. Collin, Bull. Soc. Roy. Sci. Liege **194** (1954).
5. R. T. Aplin, M. Fischer, D. Becher, H. Budzikiewicz, and C. Djerassi, J. Am. Chem. Soc. **87**, 4888 (1965).
6. J. T. Larkins, F. E. Saalfield, and L. Kaplan, Org. Mass Spectrum. **2**, 213 (1969).