

AD-A104 262

MASSACHUSETTS INST OF TECH CAMBRIDGE DEPT OF NUTRITI--ETC F/8 6/13  
FUNCTION OF THE PEPTIDE ANTIBIOTIC, GRAMICIDIN S, IN ITS PRODUCE--ETC(U)  
SEP 81 J M PIRET DAA629-78-C-0015

UNCLASSIFIED

ARD-14926.2-L

NL

For  
As  
Agency



END  
DATE  
FILMED  
10-8-81  
DTIC

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

LEVEL II

12

REPORT DOCUMENTATION PAGE

READ INSTRUCTIONS BEFORE COMPLETING FORM

1. REPORT NUMBER 14926.2-L	2. GOVT ACCESSION NO. AD-A104262	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) Function of the Peptide Antibiotic, Gramicidin S, in its Producer, <u>Bacillus Brevis</u> Nagano	5. TYPE OF REPORT & PERIOD COVERED Final Report: 15 Mar 78 - 14 Jun 81	
	6. PERFORMING ORG. REPORT NUMBER	
7. AUTHOR(s) Jacqueline Marie Piret	8. CONTRACT OR GRANT NUMBER(s) DAAG29 78 C 0015	
9. PERFORMING ORGANIZATION NAME AND ADDRESS Massachusetts Institute of Technology Cambridge, MA 02139	10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS	
11. CONTROLLING OFFICE NAME AND ADDRESS U. S. Army Research Office Post Office Box 12211 Research Triangle Park, NC 27709	12. REPORT DATE Sep 81	
	13. NUMBER OF PAGES 6	
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)	15. SECURITY CLASS. (of this report) Unclassified	
	15a. DECLASSIFICATION/DOWNGRADING SCHEDULE	

16. DISTRIBUTION STATEMENT (of this Report)  
Approved for public release; distribution unlimited.

DTIC  
SELECTED  
SEP 17 1981

17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)  
NA

18. SUPPLEMENTARY NOTES  
The view, opinions, and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy, or decision, unless so designated by other documentation.

19. KEY WORDS (Continue on reverse side if necessary and identify by block number)

antibiotics	mutants	ultraviolet radiation
peptides	microbiology	solvents
sporulation	germination	
spores	heat	

20. ABSTRACT (Continue on reverse side if necessary and identify by block number)  
The natural function(s) of the peptide antibiotic, gramicidin S (GS) in its producer, Bacillus brevis Nagano, was investigated. Particular attention was paid to the possible role of GS in the differentiation process: sporulation, spore properties and germination. The GS-producing parental strain and a GS-negative mutant of this strain were compared.

AD A104262

MINI FILE COPY

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

81 3 1 0 0 1

Accession For	
NTIS GRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By _____	
Distribution/	
Availability Codes	
Availability Codes	
Dist _____	
Special _____	
A	

DAAG29-78-C-0015

15 11 14 1271

FUNCTION OF THE PEPTIDE ANTIBIOTIC, GRAMICIDIN S,  
IN ITS PRODUCER, BACILLUS BREVIS NAGANO,

by

JACQUELINE MARIE PIRET 11 24

Submitted to the Department of Nutrition and Food Science  
on March 17, 1981 in partial fulfillment of the  
requirements for the Degree of Doctor of Philosophy in  
Nutritional Biochemistry and Metabolism

ABSTRACT

The natural function(s) of the peptide antibiotic, gramicidin S (GS) in its producer, Bacillus brevis Nagano, was investigated. Particular attention was paid to the possible role of GS in the differentiation process: sporulation, spore properties and germination. The GS-producing parental strain and a GS-negative mutant of this strain were compared.

Sporulation rate and efficiency, sporulation-associated events (except for GS production) were similar in both strains. Mature parental and mutant spores were equally resistant to heat, ultraviolet irradiation and solvents. Germination initiation in both strains was rapid and both responded similarly to a spectrum of germinants tested. Thus the lack of GS synthesis impaired none of these properties.

Outgrowth, however, was affected. In the presence of exogenous GS, spore outgrowth was inhibited. Whereas mutant spores proceeded through outgrowth and entered vegetative growth quickly, parental spores remained in outgrowth several hours longer. Extraction of GS from the parental spores reduced the outgrowth delay. GS apparently acted from the outside on spores.

Outgrowth delay was dependent upon the concentration of GS present. There was a distinct time during outgrowth when mutant spores were no longer sensitive to exogenous GS. Outgrowth inhibition by GS was permanent. Alanine uptake experiments suggested that GS affects the entry of nutrients into the outgrowing spore.

Thesis Advisor: Dr. Arnold L. Demain  
Title: Professor of Industrial Microbiology

Approved for publication  
Distribution Unlimited

1

235

## VI. SUMMARY AND CONCLUSIONS

The role of the peptide antibiotic gramicidin S in its producer Bacillus brevis Nagano was studied. The parent strain and its GS-negative mutant, BI-7, were compared. The study focused on the postulated function of GS in differentiation: sporulation, spore properties and germination. Spores were produced in media supporting low, moderate and high GS synthesis. The extent of sporulation was inversely related to antibiotic production.

Growth and sporulation in parent and BI-7 were followed. The rate, extent and sequence of events of sporulation were similar in both strains in all media used.

The properties of mature spores of the parent and BI-7 were also compared. UV, solvent and heat resistances were determined for spores produced in a range of media. DPA contents were also measured. The absence of GS synthesis by the mutant did not impair the "quality" of its spores.

The response of spores to heat activation and to

germinants was measured by loss of absorbance of spore suspensions. The GS-negative mutant and its parent behaved similarly, regardless of the medium from which they were harvested.

Spore outgrowth, however, was affected by GS. Endogenous GS, associated with parental spores or exogenous GS, added to mutant spores, extended outgrowth by several hours. In the absence of GS or when GS was removed from spores, outgrowth was rapid. All or most of the GS was at or near the spore surface. Outgrowth inhibition by GS was probably permanent. At a distinct time during outgrowth, GS no longer exerted its effect. The uptake of labelled L-alanine by outgrowing spores was inhibited by GS. This inhibition was at a step earlier than the incorporation of L-alanine into protein.

Gramicidin S, made during sporulation by Bacillus brevis Nagano, remains associated with the sporulating, dormant and initiating organism but apparently without affecting these stages of development. During spore outgrowth, however, the antibiotic prolongs this process, inhibiting the onset of vegetative growth. In this GS-sensitive period, nutrient uptake is impaired by GS. The data suggest a mechanism of action for GS whereby it

DAAG29-78-C-0015

interferes in membrane function, such as transport or  
energy metabolism, in outgrowing spores.

PUBLICATIONS

- Demain, A.L. and J.M. Piret. 1977. Speculations on the role(s) of antibiotics in spore-forming microorganisms. In Spores VII. G. Chambliss and J.C. Vary, eds. Amer. Soc. Microbiol. Washington, D.C. p. 17.
- Demain, A.L. and J.M. Piret. 1978. Relationship between antibiotic biosynthesis and sporulation. In Regulation of Secondary Product and Plant Hormone Metabolism. M. Luckner and K. Schreiber, eds. Pergamon Press. New York. p. 183.
- Piret, J.M. and A.L. Demain. 1981. Function of the peptide antibiotic gramicidin S in the producer organism, Bacillus brevis. In Sporulation and Germination. Proc. 8th Intl. Spores Conf. Amer. Soc. Microbiol. (in press)
- Piret, J.M., J. Millet and A.L. Demain. Intracellular protease in Bacillus brevis ATCC 9999. (in preparation)
- Piret, J.M. and A.L. Demain. Function of gramicidin S in its producer, Bacillus brevis Nagano. (in preparation)

PERSONNEL

Professor Arnold L. Demain  
Professor of Industrial Microbiology  
Massachusetts Institute of Technology  
Cambridge, Mass. 02139

Dr. Jacqueline M. Piret  
Laboratory of Industrial Microbiology  
Massachusetts Institute of Technology  
Cambridge, Mass. 02139

Dr. Piret obtained her Ph.D. on this project in June, 1981.