

AD-A125 236

MODULATIONAL STABILITY OF TWO-PHASE SINE-GORDON  
WAVE TRAINS(U) NEW YORK UNIV NY COURANT INST OF  
MATHEMATICAL SCIENCES D W MCLAUGHLIN ET AL. 1982  
AFOSR-TR-83-0010 AFOSR-80-0228

1/1

UNCLASSIFIED

F/G 12/1

NL




END

FILMED

BY

DTIC



MICROCOPY RESOLUTION TEST CHART  
NATIONAL BUREAU OF STANDARDS-1963-A

1

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

AD A125236

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER <b>AFOSR-TR- 83-0010</b>	2. GOVT ACCESSION NO. <b>AD-A125236</b>	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) <b>MODULATIONAL STABILITY OF TWO-PHASE SINE-GORDON WAVETRAINS</b>		5. TYPE OF REPORT & PERIOD COVERED <b>Technical</b>
7. AUTHOR(s) <b>David W. McLaughlin* and M. Gregory Forest** and Nicholas Ercolani**</b>		6. PERFORMING ORG. REPORT NUMBER
9. PERFORMING ORGANIZATION NAME AND ADDRESS <b>Courant Institute of Mathematical Sciences New York University 251 Mercer Street New York, NY 10012</b>		8. CONTRACT OR GRANT NUMBER(s) <b>AFOSR-80-0228</b>
11. CONTROLLING OFFICE NAME AND ADDRESS <b>Mathematical and Information Sciences Directorate Air Force Office of Scientific Research Bolling AFB, DC 20332</b>		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS <b>2304/A4 PE 61102F</b>
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		12. REPORT DATE <b>1982</b>
		13. NUMBER OF PAGES <b>10</b>
		15. SECURITY CLASS. (of this report) <b>Unclassified</b>
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report)  <b>Approved for public release -- distribution unlimited</b>		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)  <b>* David W. McLaughlin: Department of Mathematics and Program in Applied Mathematics, University of Arizona, Tucson, Arizona 85721 ** M. Gregory Forest and Nicholas Ercolani: Dept. of Math, Ohio St. Univ., Columbus, Ohio 43210</b>		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number)		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number)  <b>In this note, we study the modulational stability of real, two-phase sine-Gordon wavetrains. There are three classes of such waves; we find the kink-kink trains are stable, while the breather trains and kink-radiation trains are unstable to modulations.</b>		

DTIC FILE COPY

DTIC ELECTRIC  
MAR 02 1983  
S E D

MODULATIONAL STABILITY OF  
TWO-PHASE SINE-GORDON WAVETRAINS

by

David W. McLaughlin\*  
Department of Mathematics and  
Program in Applied Mathematics  
University of Arizona  
Tucson, Arizona 85721

and

M. Gregory Forest\*\* and Nicholas Ercolani  
Department of Mathematics  
The Ohio State University  
Columbus, Ohio 43210

Accession For	
NTIS GRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By _____	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A	



\*Supported in part by N.S.F. Grant #MCS-79-3533 and in part by AFOSR -80-0228  
~~Grant #MCS-79-3533~~. Address during academic year 1981-82: Courant  
Institute, New York University, 251 Mercer Street, New York, N.Y. 10012.

\*\*Supported in part by N.S.F. Grant # MCS-8002969.

88 02 028 068

Approved for public release;  
distribution unlimited.

## **DISCLAIMER NOTICE**

**THIS DOCUMENT IS BEST QUALITY PRACTICABLE. THE COPY FURNISHED TO DTIC CONTAINED A SIGNIFICANT NUMBER OF PAGES WHICH DO NOT REPRODUCE LEGIBLY.**

I. INTRODUCTION

In this note we study the modulational stability of real, two-phase sine-Gordon wavetrains, <sup>are studied.</sup> There are three classes of such waves; we find the kirk-kink trains are stable, while the breather trains and kink-radiation trains are unstable to modulations.

> These results continue the investigations of Flaschka, Forest, and McLaughlin [1] for the KdV equation and of Forest and McLaughlin [2, 3] for the sinh-Gordon and sine-Gordon equations. In a previous paper [2], the sine-Gordon two-phase modulation theory could only be carried to an intermediate stage. Here we use recent results of Ercolani and Forest [4] to complete this project.

References [1, 2, 3] contain detailed accounts of how inverse spectral theory can be used to prescribe and analyze the modulations of quasi-periodic wavetrains. Here we assume some familiarity with these references. In particular, we assume the sine-Gordon relevant discussions in [2] about squared eigenfunctions, conservation laws, and the  $\bar{\theta}$  - and  $\bar{\mu}$  - representations of two-phase solutions. We also refer to several calculations in [2] which directly apply here. Moreover, we quote two important results from [4].

AIR FORCE OFFICE OF SCIENTIFIC RESEARCH (AFSC)  
NOTICE OF TRANSMITTAL TO DTIC  
This technical report has been reviewed and is approved for public release IAW AFR 190-12. Distribution is unlimited.  
MATHEW J. KERPER  
Chief, Technical Information Division

II. NOTATION AND STATEMENT OF THE PROBLEM

We consider a real sine-Gordon wave which locally appears as a two-phase wavetrain, but which has physical characteristics (such as wave numbers and frequencies) that change slowly over large scales in space and time. Thus, this modulating two-phase wave locally is described by

the " $\vec{\theta}$  - representation",

$$u = u(\theta_1, \theta_2; \vec{E}(X, T)) , \quad (\text{II.1a})$$

where  $\vec{E} = (E_1, \dots, E_4)$ . The local or fast dependence is described by

$$\begin{cases} \frac{\partial}{\partial x} \theta_j = \kappa_j , \\ \frac{\partial}{\partial t} \theta_j = \omega_j ; \end{cases} \quad j = 1, 2 \quad (\text{II.1b})$$

the slow dependence of  $\vec{E}$  on  $X = \epsilon t$  models the modulations in the physical characteristics of the wave. (For example,  $\vec{\kappa}$  and  $\vec{\omega}$  are functions of  $\vec{E}(X, T)$ .)

There are three physically distinct classes of real two-phase sine-Gordon wavetrains, corresponding to the following possible configurations of  $E_1, \dots, E_4$  consistent with real waves (see [2]).

Case 1. Breather Train  $E_1, E_2, E_3 = E_1^*, E_4 = E_2^*$ , all distinct. (II.2a)

Case 2. Kink-Kink Train  $E_1 < E_2 < E_3 < E_4 < 0$ . (II.2b)

Case 3. Kink-Radiation Train  $E_1 < E_2 < 0, E_3 = E_4^*, E_3 \neq E_4$ . (II.2c)




FIGURE 1

The modulational equations for such wavetrains constitute a first-order system of quasi-linear partial differential equations for  $E_j(X, T)$ ,  $j = 1, \dots, 4$ . In [2], it is shown this system admits the following representation on the underlying Riemann surface  $\mathcal{R}$  of the curve  $(E, R(E))$ , where

$$R^2(E) = E \prod_{j=1}^4 (E - E_j) . \quad (\text{II.3})$$

The modulation equations are described by

$$\frac{\partial}{\partial T} \Omega^{(-)} - \frac{\partial}{\partial X} \Omega^{(+)} = 0 , \quad (\text{II.4a})$$

$$\Omega^{(\pm)} = \Omega_{+1} \pm \frac{1}{16} \Omega_{-1} , \quad (\text{II.4b})$$

where  $\Omega_{+1}$ ,  $\Omega_{-1}$  are well-defined differentials on  $\mathcal{R}$  given by

$$\Omega_{+1} = -\frac{1}{2} \left\langle \prod_{j=1}^2 (E - \mu_j) \right\rangle \frac{dE}{R(E)} , \quad (\text{II.5a})$$

$$\Omega_{-1} = \frac{\sqrt{\prod E_k}}{2} \left\langle \prod_{j=1}^2 \left( \frac{E}{\mu_j} - 1 \right) \right\rangle \frac{dE}{ER(E)} . \quad (\text{II.5b})$$

The symbol  $\langle f(u) \rangle$  represents an average over the real isospectral manifold for that class of waves, defined with respect to the  $\vec{\theta}$ -representation by

$$\langle f(u(\theta_1, \theta_2)) \rangle = \frac{1}{(2\pi)^2} \int_0^{2\pi} \int_0^{2\pi} f(u(\theta_1, \theta_2)) d\theta_1 d\theta_2 . \quad (\text{II.6})$$

The average is computed for frozen values of  $\vec{E}(X, T)$ .

### III. DERIVATION OF AN INVARIANT REPRESENTATION

We begin with the system (II.4), (II.5) as derived in [2], and now use recent developments in [4] to compute these averages as products of single integrals. These arguments apply to all three classes of real two-phase waves; we specialize only at the end to deduce the consequences for each class.

Consider some function  $f(u)$  of the modulating two-phase wave  $u(\theta_1, \theta_2)$ . Viewed as complex coordinates,  $\theta_1, \theta_2$  parametrize a complex two-torus; the restriction  $\theta_1 \in [0, 2\pi)$ ,  $\theta_2 \in [0, 2\pi)$  identifies the real isospectral manifold  $\mathcal{M}$ , which is a real two-torus  $\mathbb{T}^2 \cong [2, 4]^2$ ,

$$\mathcal{M} = \mathbb{T}^2 = \theta_1 \times \theta_2 = [0, 2\pi) \times [0, 2\pi). \quad (\text{III.1})$$

Thus, the average over  $\mathcal{M}$  is given as in (II.6),

$$\langle f(u) \rangle = \frac{1}{(2\pi)^2} \int_0^{2\pi} \int_0^{2\pi} f(u(\theta_1, \theta_2)) d\theta_1 d\theta_2. \quad (\text{III.2})$$

Next we change variables from  $\vec{\theta}$  to the " $\vec{\gamma}$ -coordinates" of  $\mathcal{M}$  provided in [4],

$$\mathcal{M} = \mathbb{T}^2 = (\gamma_1\text{-cycle}) \times (\gamma_2\text{-cycle}), \quad (\text{III.3})$$

where  $\vec{\gamma} = (\gamma_1, \gamma_2) \in \mathbb{R} \times \mathbb{R}$  are the following cycles (Figure 2) for the three classes of two-phase waves.




FIGURE 2

Changing variables of integration,

with  $\left| \frac{\partial \vec{\theta}}{\partial \vec{Y}} \right|$  the Jacobian of the change of coordinates, (III.2) becomes

$$\langle f(u) \rangle = \frac{1}{(2\pi)^2} \int \int_{\mathcal{M}} f(u(\vec{Y})) \left| \frac{\partial \vec{\theta}}{\partial \vec{Y}} \right| dY_1 dY_2. \quad (\text{III.4})$$

We write the Jacobian as a product,

$$\left| \frac{\partial \vec{\theta}}{\partial \vec{Y}} \right| = \left| \frac{\partial \vec{\theta}}{\partial \vec{\mu}} \right| \left| \frac{\partial \vec{\mu}}{\partial \vec{Y}} \right|,$$

and analyze each factor. First, by parallel arguments in [1, 2],

$$\left| \frac{\partial \vec{\theta}}{\partial \vec{\mu}} \right| = (2\pi)^2 \frac{\det \rho}{\det M}, \quad (\text{III.5a})$$

where

$$M_{ij} \equiv \int_{\gamma_i\text{-cycle}} \mu^{j-1} \frac{d\mu}{R(\mu)}, \quad \rho_{nj} = \frac{\mu_j^{2-n}}{R(\mu_j)}. \quad (\text{III.5b})$$

Second, it is proven in [4] that the map from  $\vec{Y}$  to  $\vec{\mu}$  variables is diagonal and globally defined, with

$$\left. \begin{aligned} \mu_1(\vec{Y}) &\equiv \mu_1(Y_1), & \mu_2(\vec{Y}) &\equiv \mu_2(Y_2), \\ \left| \frac{\partial \vec{\mu}}{\partial \vec{Y}} \right| &= \frac{\partial \mu_1}{\partial Y_1}(Y_1) \frac{\partial \mu_2}{\partial Y_2}(Y_2) \neq 0. \end{aligned} \right\} \quad (\text{III.6})$$

For  $f(u)$  given by  $(E - \mu_1)(E - \mu_2)$  or  $\left(\frac{E}{\mu_1} - 1\right)\left(\frac{E}{\mu_2} - 1\right)$ ,

these facts now allow us to factor the two-fold integrals (III.4). We compute one average and the other will be obvious. From (III.5), (III.6),

$$\left\langle \left( \frac{E}{\mu_1} - 1 \right) \left( \frac{E}{\mu_2} - 1 \right) \right\rangle.$$

$$= \frac{1}{\det M} \int_{\mathcal{M}} \int (1 - \frac{E}{\mu_1(\gamma_1)}) |p(\vec{\mu}(\vec{\gamma}))| \frac{\partial \mu_1}{\partial \gamma_1}(\gamma_1) \frac{\partial \mu_2}{\partial \gamma_2}(\gamma_2) d\gamma_1 d\gamma_2,$$

and by the nature of the integrands,

$$= \frac{1}{\det M} \int_{\mathcal{M}} \int \det \left[ \left(1 - \frac{E}{\mu_i(\gamma_i)}\right) \frac{\mu_i^{j-1}(\gamma_i)}{R(\mu_i(\gamma_i))} \frac{\partial \mu_i}{\partial \gamma_i}(\gamma_i) d\gamma_i \right];$$

since  $\mathcal{M} = (\gamma_1 - \text{cycle}) \times (\gamma_2 - \text{cycle})$  is a product space, Fubini's theorem gives

$$= \frac{1}{\det M} \det \left[ \int_{\gamma_i - \text{cycle}} \left(1 - \frac{E}{\mu_i(\gamma_i)}\right) \frac{\mu_i^{j-1}(\gamma_i)}{R(\mu_i(\gamma_i))} \frac{\partial \mu_i}{\partial \gamma_i}(\gamma_i) d\gamma_i \right]. \quad (\text{III.7})$$

Now that the two-fold integrals have been factored into single integrals over the  $\gamma_i$ -cycles, we change variables by (III.6) from  $\gamma_i$  to  $\mu_i$ , and make use of the result in [4] that the  $\vec{\mu}$ -cycles are globally homologous to the  $\vec{\gamma}$ -cycles of Figure 2,

$$\mu_i - \text{cycle} \sim \gamma_i - \text{cycle}, \quad i = 1, 2. \quad (\text{III.8})$$

This yields

$$\begin{aligned} & \left\langle \left(\frac{E}{\mu_1} - 1\right) \left(\frac{E}{\mu_2} - 1\right) \right\rangle \neq \det \left[ \int_{\gamma_i - \text{cycle}} \left(1 - \frac{E}{\mu_i}\right) \frac{\mu_i^{j-1}}{R(\mu_i)} d\mu_i \right] \\ & = \frac{\det \left[ \int_{\gamma_i - \text{cycle}} \left(1 - \frac{E}{\mu}\right) \frac{\mu^{j-1}}{R(\mu)} d\mu \right]}{\det M} \\ & = \frac{\det \left[ \int_{\gamma_i - \text{cycle}} \left(1 - \frac{E}{\mu}\right) \frac{\mu^{j-1}}{R(\mu)} d\mu \right]}{\det \left[ \int_{\gamma_i - \text{cycle}} \frac{\mu^{j-1}}{R(\mu)} d\mu \right]}. \quad (\text{III.9b}) \end{aligned}$$

REMARK. The formulas (III.9a, b) are precisely those obtained in [2] for the two-phase sinh-Gordon modulations; the only change is the  $\gamma_1$ -cycles are given by Figure 2 for sine-Gordon waves. Therefore, the remaining analysis in [2] applies verbatim, with the following conclusions.

THEOREM (III.1) (AN INVARIANT REPRESENTATION)

(1) The three classes of real two-phase sine-Gordon wavetrains modulate according to

$$\Omega = \frac{\partial}{\partial T} \Omega^{(-)} - \frac{\partial}{\partial X} \Omega^{(+)} = 0, \quad (\text{III.10a})$$

where

$$\Omega^{(\pm)} = \Omega_{+1} \pm \frac{1}{16} \Omega_{-1}, \quad (\text{III.10b})$$

$$\begin{aligned} \Omega_{+1} &= -\frac{1}{2} \langle (E - \mu_1)(E - \mu_2) \rangle \frac{dE}{R(E)} \\ &= -\frac{1}{2} \left( E^2 - \sum_{j=1}^2 c_j^{(+)} E^{j-1} \right) \frac{dE}{R(E)}, \end{aligned} \quad (\text{III.11a})$$

$$\begin{aligned} \Omega_{-1} &= \frac{\sqrt{\frac{4}{\pi} E_k}}{2} \langle \left( \frac{E}{\mu_1} - 1 \right) \left( \frac{E}{\mu_2} - 1 \right) \rangle \frac{dE}{ER(E)} \\ &= \frac{\sqrt{\frac{4}{\pi} E_k}}{2} \left( 1 - \sum_{j=1}^2 c_j^{(-)} E^j \right) \frac{dE}{ER(E)}. \end{aligned} \quad (\text{III.11b})$$

(2)  $\Omega_{\pm 1}$  are the unique Abelian differentials of the second kind which satisfy the following criteria:

- (i)  $\Omega_{+1}$  is holomorphic except at  $E = \infty$ ,  
 $\Omega_{-1}$  is holomorphic except at  $E = 0$ .

(ii) Near  $E = \xi^{-2} = \infty$ , respectively  $E = \xi^2 = 0$ ,

$$\Omega_{\pm 1} \sim \frac{d\xi}{\xi^2} + (\text{a holomorphic part}).$$

(iii)  $\int_{\gamma_i\text{-cycle}} \Omega_{\pm 1} = 0$ ,  $i = 1, 2$ , where

$$\gamma_1 = \begin{cases} a_1 - b_1 + b_2 \\ b_1 \\ b_1 \end{cases}, \quad \gamma_2 = \begin{cases} a_2 + b_1 - b_2 \\ b_2 \\ a_2 \end{cases} \quad \text{for the } \begin{cases} \text{Breather Train} \\ \text{Kink-Kink Train} \\ \text{Kink-Radiation Train} \end{cases}.$$

(3) Away from the branch points, the expansions of  $\Omega_{\pm 1}$  in local coordinates are given by (III.11a, b).

#### IV. CONSEQUENCES OF THE INVARIANT REPRESENTATION

The three classes of real, two-phase sine-Gordon waves modulate according to

$$\Omega = \frac{\partial}{\partial T} \Omega^{(-)} - \frac{\partial}{\partial X} \Omega^{(+)} = 0. \quad (\text{IV.1})$$

As in [2], expanding (IV.1) near  $E = 0$  and  $E = \infty$  yields the averages of the familiar "polynomial" conservation laws, such as energy and momentum. Likewise, the same arguments in Section (IV.c) of [2] yield a Hamiltonian form of these modulation equations. We do not expand on these points of view here, but rather move on to deduce the predictions for modulational stability of these three classes of waves.

As in [2], we expand (IV.1) in the local coordinate near each branch point  $E = E_k$ ,  $k = 1, 2, 3, 4$ , invoke  $\Omega = 0$ , and the vanishing of the leading order term in these expansions yields

THEOREM (IV.1) (Riemann Invariants for the Modulation Equations)

For all three classes of real waves, the branch points  $\{E_k\}_1^4$  are Riemann invariants, with

$$\frac{\partial}{\partial T} E_k - S^{(k)} \frac{\partial}{\partial X} E_k = 0, \quad k = 1, \dots, 4, \quad (\text{IV.2a})$$

with the characteristic speeds given by

$$S^{(k)} = \frac{\sum_{j=0}^3 D_j^{(+)} E_k^{j-1}}{\sum_{j=0}^3 D_j^{(-)} E_k^{j-1}}, \quad k = 1, \dots, 4, \quad (\text{IV.2b})$$

where  $D_j^{(\pm)}$  are given in terms of the coefficients  $c_j^{(\pm)}$  of  $\Omega_{\pm 1}$

$$D_j^{(\pm)} = \frac{1}{2} [c_j^{(+)} \mp \frac{1}{16} \sqrt{\Pi E_k} c_j^{(-)}], \quad j = 1, 2, \quad (\text{IV.2c})$$

$$D_0^{(\pm)} = \pm \frac{\sqrt{\Pi E_k}}{32}, \quad D_3^{(\pm)} = -\frac{1}{2}.$$

Analysis of the characteristic speeds  $S^{(k)}$  for each class of waves fields.

THEOREM (IV.2) (Modulational Stability of Two-Phase Waves)

(1) Kink-Kink Train. The characteristic speeds  $S^{(k)}$ ,  $k = 1, \dots, 4$  are real; modulational stability is predicted.

(2) Breather Train. The characteristic speeds  $S^{(k)}$ ,  $k = 1, \dots, 4$  are complex, with  $[S^{(3)}]^* = S^{(1)}$ ,  $[S^{(4)}]^* = S^{(2)}$ ; modulational instability is predicted.

(3) Kink-Radiation Train. The characteristic speeds  $S^{(k)}$ ,  $k = 1, \dots, 4$  are complex, with  $[S^{(4)}]^* = S^{(4)}$ ; modulational instability is predicted.

REFERENCES

1. Flaschka, H., Forest, M.G., and McLaughlin, D.W., "Multiphase averaging and the inverse spectral solution of the Korteweg-de Vries Equation," C.P.A.M. 33, 739-784 (1980).
2. Forest, M.G., and McLaughlin, D.W., "Modulations of sinh-Gordon and sine-Gordon wavetrains," to appear Studies in Appl. Math., 1982.
3. Forest, M.G., and McLaughlin, D.W., "Canonical variables for the periodic sine-Gordon equation and a method of averaging," LA-UR 78-3318, Report of the Los Alamos Scientific Laboratory, 1978; also submitted in revised form to J. Math. Phys.
4. Ercolani, N. and Forest, M.G., "On the geometry of real, two-phase solutions of the sine-Gordon equation," in preparation.