

AD-A151 100

PROGRESS REPORT GRANT AFOSR-84-0365(U) FLORIDA UNIV  
GAINESVILLE DEPT OF MATHEMATICS I LASIECKA JAN 85  
AFOSR-TR-85-0172 AFOSR-84-0365


1/1

UNCLASSIFIED

F/G 12/1

NL

				END
				FILMED
				DTIC



MICROCOPY RESOLUTION TEST CHART  
NATIONAL BUREAU OF STANDARDS-1963-A

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE

REPORT DOCUMENTATION PAGE				
1a. REPORT SECURITY CLASSIFICATION <b>UNCLASSIFIED</b>		1b. RESTRICTIVE MARKINGS		
2a. SECURITY CLASSIFICATION AUTHORITY		3. DISTRIBUTION/AVAILABILITY OF REPORT Approved for public release; distribution unlimited.		
2b. DECLASSIFICATION/DOWNGRADING SCHEDULE		4. PERFORMING ORGANIZATION REPORT NUMBER(S)		
4. PERFORMING ORGANIZATION REPORT NUMBER(S)		5. MONITORING ORGANIZATION REPORT NUMBER(S) <b>AFOSR-TR- 85 - 0172</b>		
6a. NAME OF PERFORMING ORGANIZATION University of Florida		6b. OFFICE SYMBOL <i>(If applicable)</i>	7a. NAME OF MONITORING ORGANIZATION Air Force Office of Scientific Research	
6c. ADDRESS (City, State and ZIP Code) Department of Mathematics Gainesville FL 32611		7b. ADDRESS (City, State and ZIP Code) Directorate of Mathematical & Information Sciences, Bolling AFB DC 20332-6448		
8a. NAME OF FUNDING/SPONSORING ORGANIZATION AFOSR		8b. OFFICE SYMBOL <i>(If applicable)</i> NM	9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER AFOSR-84-0365	
8c. ADDRESS (City, State and ZIP Code) Bolling AFB DC 20332-6448		10. SOURCE OF FUNDING NOS.		
		PROGRAM ELEMENT NO. 61102F	PROJECT NO. 2304	TASK NO. A1
		WORK UNIT NO.		
11. TITLE (Include Security Classification) PROGRESS REPORT, GRANT AFOSR-84-0365, 1 SEPTEMBER 1984 - 31 DECEMBER 1984				
12. PERSONAL AUTHOR(S) I. Lasiecka				
13a. TYPE OF REPORT Interim		13b. TIME COVERED FROM 1/9/84 TO 31/12/84	14. DATE OF REPORT (Yr., Mo., Day) JAN 85	15. PAGE COUNT 2
16. SUPPLEMENTARY NOTATION				
17. COSATI CODES		18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)		
FIELD	GROUP	SUB. GR.		
19. ABSTRACT (Continue on reverse if necessary and identify by block number)				
<p>&gt; Results have been obtained for certain second order hyperbolic systems with viscous dumping which imply that one can increase "at will" the margin of stability of arbitrarily finite modes of the damped wave equation (those presumed "dominant") by means of certain type of boundary feedback, while the remaining new modes approach asymptotically the original ones from the left of the vertical axis <math>Re z = -k</math>. Numerical testing of the constructive procedure is in the process of being implemented by a Ph.D. student.</p>				
<b>DTIC FILE COPY</b>		<p style="font-size: 2em; font-weight: bold; margin: 0;">DTIC ELECTE</p> <p style="font-size: 1.5em; font-weight: bold; margin: 0;">MAR 13 1985</p> <p style="font-size: 2em; font-weight: bold; margin: 0;">E</p>		
20. DISTRIBUTION/AVAILABILITY OF ABSTRACT UNCLASSIFIED/UNLIMITED <input checked="" type="checkbox"/> SAME AS RPT. <input type="checkbox"/> DTIC USERS <input type="checkbox"/>		21. ABSTRACT SECURITY CLASSIFICATION UNCLASSIFIED		
22a. NAME OF RESPONSIBLE INDIVIDUAL Dr. Marc Q. Jacobs		22b. TELEPHONE NUMBER <i>(Include Area Code)</i> (202) 767- 4940	22c. OFFICE SYMBOL NM	

Reference is made to the Original Proposal (O.P.) which was submitted to the A.F.O.S.R. on January 1984. The grant's official starting date is September 1, 1984.

Of the two canonical problems for second order hyperbolic dynamics with viscous damping, which were proposed in Section 2 of the O.P., we have so far obtained a theoretical solution of the first (case 1, p 3 of O.P.). Our argument is essentially constructive, and this lends itself naturally to numerical algorithms. Numerical testing is about to begin soon. More precisely: consider the following boundary feedback for the damped wave equation

$$(1) \quad x_{tt}(t, \xi) = \Delta x(t, \xi) - 2kx_t(t, \xi) \quad \text{in } (0, \infty) \times \Omega$$

$$\text{I.C. } x_0 \text{ and } x_1 \quad \text{in } \Omega$$

$$(2) \quad x(t, \sigma) = [\langle x(t, \cdot), w_1 \rangle + \langle x_t(t, \cdot), w_2 \rangle] g(\sigma) \quad \text{in } (0, \infty) \times \Gamma$$

Here  $k > 0$  is the coefficient of viscous damping and the right hand side of (2) is the "interior observation" acting on the boundary.

For  $g = 0$  and 'small'  $k > 0$ , the eigenvalues  $\lambda_m^{+, -}$  of the free system lie on the vertical line

$\text{Re } \lambda = -k$  and it is a classical result that all free solutions decay with upperbound  $e^{-kt}$  in the energy norm for  $t > 0$ .

For  $g \neq 0$ , the following result is proved in [L-T.5], [T.4].

Let a real  $g \in L_2(\Gamma)$  be given, with  $(g_{\text{ext}}, e_m)_{L_2(\Omega)} \neq 0$ ,  $m=1, 2, \dots$  where  $g_{\text{ext}}$  = harmonic extension of  $g$  onto  $\Omega$  and  $\{e_m\}$  are the eigenvectors of the Laplacian  $\Delta$  with zero Dirichlet B.C. Let an arbitrary sequence  $\{\epsilon_m^{+, -}\}$  of distinct, non zero complex numbers, with  $\epsilon_m^+$  = complex conjugate of  $\epsilon_m^-$  be given. It is assumed that, moreover, the sequence  $\{\epsilon_m^{+, -}\}$  satisfies some technical assumptions i.e. that a few infinite sums involving  $\{\lambda_m^{+, -}\}$ ,  $\{\epsilon_m^{+, -}\}$  and  $(g_{\text{ext}}, e_m)$  be finite), which allow finitely many  $\{\epsilon_m^{+, -}\}_{m=1}^M$ ,  $M$  arbitrary, to be preassigned at will, while the sequence  $\{\epsilon_m^{+, -}\}$  approach zero 'sufficiently fast'.

Then, one can construct real vectors  $w_1 \in L_2(\Omega)$   $i = 1, 2$ , such that the feedback system (1)-(2) has the following two desirable properties:

(1) its eigenvalues  $\{\alpha_m^{+, -}\}$  are given precisely by  $\alpha_m^{+, -} = \lambda_m^{+, -} + \epsilon_m^{+, -}$  (so that any finitely many

$\{\alpha_m^{+,-}\}$  can be preassigned at will, while the distance  $|\alpha_m^{+,-} - \lambda_m^{+,-}|$  between new and original eigenvalues goes to zero);

(ii) its corresponding eigenvectors form a Riesz basis on  $L_2(\Omega) \times H^{-1}(\Omega)$ .

In particular, we may require  $\text{Re } \epsilon_m^{+,-} < 0$ , in which case all feedback closed loop solutions preserve the overall decay upper bound  $e^{-kt}$ ,  $t > 0$ , in view of properties (i) - (ii). This result, in particular, implies that one can increase at will the margin of stability of arbitrarily finite modes of the damped wave equation (those presumed 'dominant') by means of the boundary feedback in (2), while the remaining new modes approach asymptotically the original ones from the left of the vertical axis  $\text{Re } \lambda = -k$ .

Numerical testing of the procedure, which is essentially constructive is in the process of being implemented by a Ph.D. student, G. Bartolomeo. Thus, the short term goal of the Project Summary of the Original Proposal has been accomplished.

Accession For	
NTIS GRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By _____	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-1	



AIR FORCE OFFICE OF SCIENTIFIC RESEARCH (AFOSR)  
 NOTICE OF TRANSMITTAL TO DTIC  
 This technical report has been reviewed and is  
 approved for public release in accordance with AFOSR 199-12.  
 Distribution is unlimited.  
 MATTHEW J. KEEPER  
 Chief, Technical Information Division

**END**

**FILMED**

**4-85**

**DTIC**