

AD-A165 187

PROPERTIES OF BATCH MEANS FROM STATIONARY ARMA
(AUTOREGRESSIVE MOVING AVERAGE) TIME SERIES(U) NAVAL
POSTGRADUATE SCHOOL MONTEREY CA K KANG ET AL FEB 86
NP555-86-004


1/1

UNCLASSIFIED

F/G 12/1

NL





MICROCOPY RESOLUTION TEST CHART
NATIONAL BUREAU OF STANDARDS-1963-A

2

NPS55-86-004

NAVAL POSTGRADUATE SCHOOL

Monterey, California

AD-A165 187



FD

DTIC FILE COPY

PROPERTIES OF BATCH MEANS FROM
STATIONARY ARMA TIME SERIES

KEEBOM KANG
BRUCE SCHMEISER

FEBRUARY 1986

Approved for public release; distribution unlimited.

Prepared for:
Naval Postgraduate School
Monterey, CA 93943-5100

00

14

NAVAL POSTGRADUATE SCHOOL
MONTEREY, CALIFORNIA


Rear Admiral R. H. Shumaker
Superintendent

D. A. Schrady
Provost

The work reported herein was supported in part by the Office of Naval Research.

Reproduction of all or part of this report is authorized.


This report was prepared by:


BRUCE SCHMEISER
Adjunct Research Professor

Reviewed by:


ALAN R. WASHBURN
Chairman
Department of Operations Research

Released by:


KNEALE T. MARSHALL
Dean of Information
and Policy Sciences

REPORT DOCUMENTATION PAGE

1a. REPORT SECURITY CLASSIFICATION UNCLASSIFIED		1b. RESTRICTIVE MARKINGS	
2a. SECURITY CLASSIFICATION AUTHORITY		3. DISTRIBUTION/AVAILABILITY OF REPORT Approved for public release; distribution unlimited.	
2b. DECLASSIFICATION/DOWNGRADING SCHEDULE		5. MONITORING ORGANIZATION REPORT NUMBER(S)	
4. PERFORMING ORGANIZATION REPORT NUMBER(S) NPS55-86-004		7a. NAME OF MONITORING ORGANIZATION	
6a. NAME OF PERFORMING ORGANIZATION Naval Postgraduate School	6b. OFFICE SYMBOL (If applicable) Code 55	7b. ADDRESS (City, State, and ZIP Code)	
6c. ADDRESS (City, State, and ZIP Code) Monterey, CA 93943-5000		9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER	
8a. NAME OF FUNDING/SPONSORING ORGANIZATION	8b. OFFICE SYMBOL (If applicable)	10. SOURCE OF FUNDING NUMBERS	
8c. ADDRESS (City, State, and ZIP Code)		PROGRAM ELEMENT NO	PROJECT NO
		TASK NO.	WORK UNIT ACCESSION NO.
11. TITLE (include Security Classification) PROPERTIES OF BATCH MEANS FROM STATIONARY ARMA TIME SERIES			
12. PERSONAL AUTHOR(S) Kang, Keebom (University of Miami) and Schmeiser, Bruce W.			
13a. TYPE OF REPORT Technical	13b. TIME COVERED FROM _____ TO _____	14. DATE OF REPORT (Year, Month, Day) 1986, February	15. PAGE COUNT 10
16. SUPPLEMENTARY NOTATION			
17. COSATI CODES		18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)	
FIELD	GROUP	aggregated time series, batch means, autoregressive, moving average, Monte Carlo, simulation	
19. ABSTRACT (Continue on reverse if necessary and identify by block number) → The batch means process arising from an arbitrary autoregressive moving-average (ARMA) process time series is derived. As side results, the variance and correlation structures of the batch means process as functions of the batch size and parameters of the original process are obtained. Except for the first-order ARMA process, for which a closed-form expression is obtained, the parameters of the batch-means process are determined numerically. <i>Monte Carlo method; simulation</i>			
20. DISTRIBUTION/AVAILABILITY OF ABSTRACT <input type="checkbox"/> UNCLASSIFIED UNLIMITED <input checked="" type="checkbox"/> SAME AS RPT <input type="checkbox"/> DTIC USERS		21. ABSTRACT SECURITY CLASSIFICATION Unclassified	
22a. NAME OF RESPONSIBLE INDIVIDUAL Bruce W. Schmeiser		22b. TELEPHONE (Include Area Code) (408)646-2119	22c. OFFICE SYMBOL Code 55Sc

1. INTRODUCTION

Corresponding to any time series $\{X_i; i = 0, 1, 2, \dots\}$ is the time series $\{\bar{X}_j; j = 0, 1, 2, \dots\}$, where

$$\bar{X}_j = b^{-1} \sum_{h=1}^b X_{(j-1)b+h} \quad (1)$$

is the j^{th} batch mean of size b . The batch-means process, or aggregated time series, is of interest when observations are actually batched (see, e.g., Telser [1967], Amemiya and Wu [1972], and Tiao [1972]), when calculations need to be simplified (see, e.g., Blackman and Tukey [1958, Sec. B. 17]), or when the process mean, $E(X)$, is to be estimated. The third context motivates our work.

Consider estimating $E(X)$ with the average of n observations. $\bar{X} = \sum_{i=1}^n X_i / n$. Using batch means to estimate the variance of the sample mean, $V(\bar{X})$, has long been considered (see, e.g., Conway [1963]). Brillinger (1973) shows that if the values of a process at a distance from each other are only weakly dependent, then the batch means are asymptotically independent and normally distributed. Thus, for large batch size b , $V(\bar{X})$ can be estimated using S^2_k / k , where $k = \lfloor n / b \rfloor$, $S^2_k = (k-1)^{-1} \left[\sum_{j=1}^k \bar{X}_j - k\bar{X}^2 \right]$, and $\lfloor \cdot \rfloor$ denotes the floor function. (Moran [1965] discusses related estimators.) A nominal $100(1-\alpha)$ percent confidence interval on $E(X)$ is then $\bar{X} \pm t_{1-\alpha/2, k-1} S_k k^{-1/2}$, where $t_{1-\alpha/2, k-1}$ is the $1-(\alpha/2)$ quantile of Student's t distribution with $k-1$ degrees of freedom.

The batch means algorithms developed by Mechanic and McKay (1966), Fishman (1978), Law and Carson (1979), Schriber and Andrews (1979), and Adam (1983) empirically calculate measures of batch dependency for various batch sizes b in an attempt to determine a reasonably small value of b that yields batch means that are almost independent and normally distributed. These procedures require substantial calculation; Law and Carson (1979), for example, calculate first-order correlations based on 400 batches. That so many batches are required for accurate estimation of dependency measures is unfortunate, since Schmeiser (1982) shows that, for fixed n , additional batches beyond some small number (ten to thirty) do little to improve the statistical properties of the batch means confidence interval procedures. The results of Section 2 are motivated by the idea that knowledge of the relationship between $\{X_i\}$ and $\{\bar{X}_j\}$ can be used to measure properties of $\{\bar{X}_j\}$ even for small values of k .

A second reason for studying the relationship between $\{X_i\}$ and $\{\bar{X}_j\}$ is to allow more efficient simulation studies of batch-means procedures. Studying the performance of several batch-means procedures in the context of various distributions assumptions for $\{X_i\}$ requires a large computational effort, especially when the large sample sizes required to simulate a system and the large number of replications required for meaningful conclusions are considered. A crude Monte Carlo method is to generate X_1, X_2, \dots, X_n and calculate the batch means $\bar{X}_1, \bar{X}_2, \dots, \bar{X}_{\lfloor n/b \rfloor}$ for all values of b of interest. A computationally more efficient alternative is to derive the properties of $\{\bar{X}_j\}$ from the properties of $\{X_i\}$ and to generate directly the batch means $\{\bar{X}_j\}$, as discussed in Section 4.

A third motivation is that direct insights might result from studying properties of $\{\bar{X}_j\}$ as functions of the parameters of $\{X_i\}$. Particularly interesting is the sensitivity of $\{\bar{X}_j\}$ to the underlying process and to the batch size b , as discussed in Kang (1984, Chapter 5).

Section 2 contains results relating batch-means processes to arbitrary stationary autoregressive moving-average (ARMA) processes. Section 3 considers the special case of the underlying process being ARMA (1,1). Section 4 is a summary containing an algorithm for determining the batch-means process from the underlying process.

2. BATCH MEANS OF STATIONARY ARMA PROCESSES

The ARMA (p, q) process $\{X_i\}$ by definition satisfies

$$\sum_{h=0}^p \phi_h X_{i-h} = \sum_{h=0}^q \theta_h \epsilon_{i-h} \quad (2)$$

where $\phi_0 \equiv 1$, $\theta_0 \equiv 1$, and the error terms ϵ_i are independent with zero mean and variance σ_ϵ^2 . The main result of this paper is Theorem 1, which states that batch-means processes arising from ARMA underlying processes are themselves ARMA and specifies the parameter values.

Theorem 1. *Consider the stationary ARMA (p, q) process of equation (2). Then $\{\bar{X}_j\}$ is the stationary ARMA (p, \bar{q}) process*

$$\sum_{h=0}^p \bar{\phi}_h \bar{X}_{j-h} = \sum_{h=0}^{\bar{q}} \bar{\theta}_h \bar{\epsilon}_{j-h}$$

where $\bar{\phi}_0 \equiv 1$, $\bar{\theta}_0 \equiv 1$, the batch-means error terms $\bar{\epsilon}_j$ are independent random variables with zero mean and variance σ_ϵ^2 , and \bar{q} , $\bar{\phi}_1, \bar{\phi}_2, \dots, \bar{\phi}_p$, and $\bar{\theta}_1, \bar{\theta}_2, \dots, \bar{\theta}_{\bar{q}}$ are functions (of the parameters of the underlying process and the batch size b) given in Lemmas 1, 2, and 4, respectively.

The proof of Theorem 1 requires the following lemmas.

Lemma 1 (Anderson [1979a, p. 155]). *If the underlying process $\{X_i\}$ is a stationary ARMA (p, q) process, then the batch-means process $\{\bar{X}_j\}$ is a stationary ARMA (p, \bar{q}) process, where $\bar{q} = p - \lfloor (p - q) / b \rfloor$.*

Anderson uses the more complicated, but equivalent, expression $\bar{q} = \left\lfloor \frac{q - (p+1)(b-1)}{b} \right\rfloor$.

Lemma 1 has several direct implications, as discussed in the Appendix.

Lemma 2 (Amemiya and Wu [1972]). *The AR parameters $\bar{\phi}_1, \bar{\phi}_2, \dots, \bar{\phi}_p$ of the batch means process $\{\bar{X}_j\}$ are the coefficients of B^1, B^2, \dots, B^p of $\prod_{h=1}^p (1 - \alpha_h^b B)$, respectively, where $\alpha_1, \alpha_2, \dots, \alpha_p$ are the roots of the characteristic equation $\Phi(B) = \sum_{h=0}^p \phi_h B^h = 0$.*

Lemma 3. For any stationary process $\{X_i\}$, the lag- h autocorrelation of the batch-means process $\{\bar{X}_j\}$ is

$$\bar{\rho}_h \equiv \text{Corr}(\bar{X}_j, \bar{X}_{j+h}) = \left[\sum_{i=1}^b i \rho_{(h-1)b+i} + \sum_{i=1}^{b-1} i \rho_{(h-1)b+2b-i} \right] / [bc]$$

where $c = 1 + 2 \sum_{h=1}^{b-1} (1 - (h/b)) \rho_h$, $\rho_h \equiv R_h / R_0$, and $R_h \equiv E[(X_i - E X)(X_{i+h} - E X)]$ for $h = 0, 1, 2, \dots$ and $i = 0, 1, 2, \dots$.

Proof. For any stationary process, the lag- h covariance of the batch means process is

$$\text{Cov}(\bar{X}_j, \bar{X}_{j+h}) = b^{-2} \sum_{i=1}^b \sum_{k=1}^b \text{Cov}(X_{(j-1)b+i}, X_{(j+h-1)b+k}) \quad (3)$$

(see, e.g., Kleijnen [1975, p. 507]), which is a special case of the covariance of linear combinations of random variables as discussed in Box and Jenkins (1976, pp. 28-29). Also, for any stationary process, each batch mean \bar{X}_j has variance

$$\bar{R}_0 \equiv V(\bar{X}_j) = c R_0 / b \quad (4)$$

(see, e.g., Fishman [1973, p. 281]). The definition of correlation

$$\bar{\rho}_h \equiv \text{Cov}(\bar{X}_j, \bar{X}_{j+h}) / [V(\bar{X}_j)V(\bar{X}_{j+h})]^{1/2}$$

and equations (3) and (4) yield

$$\begin{aligned} \bar{\rho}_h &= b^{-2} \sum_{i=1}^b \sum_{k=1}^b \text{Cov}[X_{(j-1)b+i}, X_{(j+h-1)b+k}] / [c R_0 / b] \\ &= \sum_{i=1}^b \sum_{k=1}^b \text{Corr}[X_{(j-1)b+i}, X_{(j+h-1)b+k}] / [bc]. \end{aligned}$$

Counting like terms arising from stationarity yields the result. \square

Lemma 4 (Anderson [1971, p. 237]). Consider a stationary ARMA(p, \bar{q}) process with known AR (autoregressive) parameters $\bar{\phi}_1, \bar{\phi}_2, \dots, \bar{\phi}_p$; variance \bar{R}_0 , and autocorrelation coefficients $\bar{\rho}_1, \bar{\rho}_2, \dots, \bar{\rho}_{\bar{q}}$. Then the MA (moving-average) parameters $\bar{\theta}_1, \bar{\theta}_2, \dots, \bar{\theta}_{\bar{q}}$ are determined.

Theorem 1 can now be proven using the four lemmas.

Proof of Theorem 1.

1. From Lemma 1, the AR and MA orders of $\{\bar{X}_j\}$ are determined: in particular, $\{\bar{X}_j\}$ is an ARMA(p, \bar{q}) process.
2. From Lemma 2, the AR parameters of $\{\bar{X}_j\}$ are determined.
3. The autocorrelations $\rho_1, \rho_2, \dots, \rho_{b(\bar{q}+1)-1}$ of a stationary ARMA process can be calculated using the algorithm of Sweet and Mazaheri (1979).

4. Given the autocorrelations from Step 3, the batch-means variance \bar{R}_0 is determined by equation (4) and the batch-means autocorrelations $\bar{\rho}_1, \bar{\rho}_2, \dots, \bar{\rho}_{\bar{q}}$ are determined by Lemma 3.
5. The MA parameters $\bar{\theta}_1, \bar{\theta}_2, \dots, \bar{\theta}_{\bar{q}}$ are then determined from Lemma 4. \square

The representation of the batch-means process is not unique. The batch-means MA parameters of Lemma 4 require the solution of a polynomial equation of order $2\bar{q}$ to determine $\bar{\theta}_1, \bar{\theta}_2, \dots, \bar{\theta}_{\bar{q}}$. The $2\bar{q}$ roots can be partitioned $2^{\bar{q}}$ ways into two sets $(x_1, x_2, \dots, x_{\bar{q}})$ and $(x_{\bar{q}+1}, x_{\bar{q}+2}, \dots, x_{2\bar{q}})$ such that $x_{\bar{q}+i} = 1/x_i$. All such subsets of size \bar{q} from $(x_1, x_2, \dots, x_{2\bar{q}})$ determine $\bar{\theta}_1, \bar{\theta}_2, \dots, \bar{\theta}_{\bar{q}}$ corresponding to stochastically equivalent processes. But there is a unique subset of size \bar{q} having $|x_i| \leq 1$, and therefore $|x_{\bar{q}+i}| = |1/x_i| \geq 1$, for $i = 1, 2, \dots, \bar{q}$. Thus for the ARMA(p, \bar{q}) process there exist $2^{\bar{q}} - 1$ non-invertible processes corresponding to a unique invertible process.

3. ARMA(1,1) BATCH MEANS

Now consider the special case of stationary first-order ARMA processes

$$X_i + \phi X_{i-1} = \epsilon_i + \theta \epsilon_{i-1} \quad \text{for } i = 1, 2, \dots \quad (5)$$

The low-order moments of $\{X_i\}$ are the zero mean, variance

$$R_0 = \sigma_\epsilon^2 (1 + \theta^2 - 2\phi\theta) / (1 - \phi^2), \quad (6)$$

the lag-one autocorrelation

$$\rho_1 = (1 - \phi\theta)(\theta - \phi) / (1 + \theta^2 - 2\phi\theta), \quad (7)$$

and lag-h autocorrelations

$$\rho_h = (-\phi)\rho_{h-1} = (-\phi)^{h-1}\rho_1 \quad \text{for } h = 2, 3, \dots \quad (8)$$

Closed-form expressions for the parameters of the batch-means process are given in Theorem 2.

Theorem 2. Consider the stationary ARMA(1,1) process of equation (5). The corresponding batch-means process is the stationary ARMA(1,1) process

$$\bar{X}_j + \bar{\phi}\bar{X}_{j-1} = \bar{\epsilon}_j + \bar{\theta}\bar{\epsilon}_{j-1} \quad \text{for } j = 1, 2, \dots \quad (9)$$

where

$$\bar{\phi} = (-1)^{h+1}\phi^h \quad (10)$$

$$\bar{\theta} = \frac{(2\bar{\rho}_1 + \bar{\phi}^2 - 1) \pm \sqrt{(1 - \bar{\phi}^2)(1 - (2\bar{\rho}_1 - \bar{\phi})^2)^{1/2}}}{2(\rho_1 + \bar{\phi})} \quad (11)$$

and

$$\sigma_{\bar{\epsilon}}^2 = \bar{R}_0 (1 - \bar{\phi}^2) / (\bar{\theta}^2 - 2\bar{\phi}\bar{\theta} + 1) \quad (12)$$

where

$$\bar{R}_0 = c R_0 / b \quad (13)$$

$$\bar{\rho}_1 = \rho_1(1 + \bar{\phi})^2 / [bc(1 + \phi)^2] \quad (14)$$

and

$$c = 1 + \{2\rho_1[(-\phi)^b + b(1 + \phi) - 1] / [b(1 + \phi)^2]\} \quad (15)$$

Proof of Theorem 2.

Equations (13), (14), and (15) follow from Lemma 3 and equation (4) via equation (8). Since $\{X_i\}$ is ARMA(1,1), $\{\bar{X}_j\}$ is also ARMA(1,1) by Lemma 1: that is, equation (9) holds. Equation (10) is a special case of Lemma 2. Since $\{\bar{X}_j\}$ is ARMA(1,1), equation (7) yields

$$\bar{\rho}_1 = (1 - \bar{\phi}\bar{\theta})(\bar{\theta} - \bar{\phi}) / (1 + \bar{\theta}^2 - 2\bar{\phi}\bar{\theta}).$$

Solving for $\bar{\theta}$ yields the two roots of equation (11). Since $\{\bar{X}_j\}$ is an ARMA(1,1) process, equation (6) holds with the batch-means parameters:

$$\bar{R}_0 = \sigma_{\bar{\epsilon}}^2(1 + \bar{\theta}^2 - 2\bar{\phi}\bar{\theta}) / (1 - \bar{\phi}^2).$$

Solving for the variance of the batch-means error term, $\sigma_{\bar{\epsilon}}^2$, with either value of $\bar{\theta}$ from equation (11) yields equation (12). (But note that the value of $\sigma_{\bar{\epsilon}}^2$ depends on the choice of $\bar{\theta}$). □

4. SUMMARY

Properties of batch means are studied under the assumption that the underlying process is ARMA(p, q). For ARMA(1,1) processes, closed-form expressions for the corresponding batch-means processes are obtained. A numerical procedure is developed for calculating the parameters of the ARMA batch-means process from the parameters of the underlying process and the batch size b . This procedure is stated concisely here for convenience.

ARMA(p, q) Procedure.

Given parameters $\phi_1, \phi_2, \dots, \phi_p, \theta_1, \theta_2, \dots, \theta_q$, error variance σ_{ϵ}^2 , and batch size b , calculate

1. $\bar{q} = p - [(p - q) / b]$
2. $\bar{\phi}_1, \bar{\phi}_2, \dots, \bar{\phi}_p$ using Lemma 2
3. $R_0, \rho_1, \rho_2, \dots, \rho_{b(\bar{q}-1)}$ from the Yule-Walker equations, probably using the algorithm of Sweet and Mazaheri (1979)
4. $c = 1 - 2 \sum_{h=1}^{b-1} (1 - (h/b)) \rho_h$

5. \bar{R}_0 from equation (4)
6. $\bar{\rho}_1, \bar{\rho}_2, \dots, \bar{\rho}_{\bar{q}}$ using Lemma 3
7. $\bar{\theta}_1, \bar{\theta}_2, \dots, \bar{\theta}_{\bar{q}}$ from Lemma 4
8. $\sigma_{\bar{\epsilon}}^2$ from equation (13) of Anderson (1971, p. 237).

A FORTRAN implementation of the ARMA(p, q) procedure is given in Kang (1984).

When the underlying process is ARMA(1,1), the following closed-form procedure can be used:

ARMA(1,1) Procedure.

Given parameters ϕ, θ , error variance σ_{ϵ}^2 , and batch size b , calculate

1. $\bar{q} = 1$
2. $\bar{\phi}$ from equation (10)
3. R_0 from equation (6), ρ_1 from equation (7)
4. c from equation (15)
5. \bar{R}_0 from equation (13)
6. $\bar{\rho}_1$ from equation (14)
7. $\bar{\theta}$ from equation (11)
8. $\sigma_{\bar{\epsilon}}^2$ from equation (12)

Notice in the ARMA(1,1) special case that calculation in step 3 of all $2b - 1$ autocorrelations of the underlying process is not necessary.

If the underlying error terms ϵ_t are normally distributed, then the batch-means error terms $\bar{\epsilon}_j$ are also normally distributed (see, e.g., Johnson and Kotz [1971, p. 51]). Therefore, generation of random variates directly from the batch-means process is straightforward using equation (9), thereby avoiding the costly computations of aggregating observations from the underlying process. Initialization for steady-state results is straightforward for AR and MA processes, but initialization for ARMA is complicated unless the process is warmed-up by discarding some initial observations (Anderson [1979b]).

REFERENCES

- Adam, N. (1983). "Achieving a Confidence Interval for Parameters Estimated by Simulation," *Management Science* **29**, 856-866.
- Amemiya, T., and Wu, R.Y. (1972). "The Effect of Aggregation on Prediction in the Autoregressive Model," *Journal of the American Statistical Association* **67**, 628-632.
- Anderson, O.D. (1979a). *Time Series Analysis and Forecasting: The Box-Jenkins Approach*, London: Butterworths.
- Anderson, O.D. (1979b). "On Warming-up Time Series Simulations Generated by Box-Jenkins Models," *Journal of the Operational Research Society* **30**, 587-589.
- Anderson, T.W. (1971). *The Statistical Analysis of Time Series*, New York: John Wiley & Sons.
- Blackman, R.B., and Tukey, J.W. (1958). *The Measurement of Power Spectra*, New York: Dover.
- Box, G.E.P., and Jenkins, G.M. (1976). *Time Series Analysis: Forecasting and Control*, San Francisco: Holden Day.
- Brillinger, D.R. (1973). "Estimation of the Mean of a Stationary Time Series by Sampling," *Journal of Applied Probability* **10**, 419-431.
- Conway, R.W. (1963). "Some Tactical Problems in Digital Simulation," *Management Science* **10**, 47-61.
- Fishman, G.S. (1973). *Concepts and Methods in Discrete Event Simulation*, New York: John Wiley & Sons.
- Fishman, G.S. (1978). "Grouping Observations in Digital Simulation," *Management Science* **24**, 510-521.
- Johnson, N.L., and Kotz, S. (1970). *Distributions in Statistics: Continuous Univariate Distributions - 1*, New York: John Wiley & Sons.
- Kang, K. (1984). *Confidence Interval Estimation Via Batch Means and Time Series Modeling*. Ph.D. dissertation, Purdue University, West Lafayette, Indiana.
- Kleijnen, J.P.C. (1975). *Statistical Techniques in Simulation. Part II*. New York: Marcel Dekker.
- Law, A.M., and Carson, J.S. (1979). "A Sequential Procedure for Determining the Length of a Steady State Simulation," *Operations Research* **27**, 1011-1025.
- Mechanic, H., and McKay, W. (1966). "Confidence Intervals for Averages of Dependent Data in Simulation II." Technical Report ASDD 17-202, IBM Corporation, Yorktown Heights, New York.
- Moran, P.A.P. (1975). "The Estimation of Standard Errors in Monte Carlo Simulation Experiments." *Biometrika* **62**, 1-4.

- Schmeiser, B.W. (1982). "Batch Size Effects in the Analysis of Simulation Output." *Operations Research* 30, 556-568.
- Schriber, T.J., and Andrews, R.W. (1979). "Interactive Analysis of Simulation Output by the Method of Batch Means," *Proceedings of the Winter Simulation Conference*, 513-525.
- Sweet, A.L., and Mazaheri, F. (1979). "Computation of the Autocovariances of Stationary ARMA Processes," *Computers and Industrial Engineering* 3, 313-320.
- Telser, L.G. (1967). "Discrete Samples and Moving Sums in Stationary Stochastic Processes." *Journal of the American Statistical Association*, 62, 484-499.
- Tiao, G.C. (1972). "Asymptotic Behaviour of Temporal Aggregates of Time Series." *Biometrika* 59, 525-531.

Accession For	
1982-1981	<input checked="" type="checkbox"/>
1981-1980	<input type="checkbox"/>
1980-1979	<input type="checkbox"/>
1979-1978	
1978-1977	
1977-1976	
1976-1975	
1975-1974	
1974-1973	
1973-1972	
1972-1971	
1971-1970	
1970-1969	
1969-1968	
1968-1967	
1967-1966	
1966-1965	
1965-1964	
1964-1963	
1963-1962	
1962-1961	
1961-1960	
1960-1959	
1959-1958	
1958-1957	
1957-1956	
1956-1955	
1955-1954	
1954-1953	
1953-1952	
1952-1951	
1951-1950	
1950-1949	
1949-1948	
1948-1947	
1947-1946	
1946-1945	
1945-1944	
1944-1943	
1943-1942	
1942-1941	
1941-1940	
1940-1939	
1939-1938	
1938-1937	
1937-1936	
1936-1935	
1935-1934	
1934-1933	
1933-1932	
1932-1931	
1931-1930	
1930-1929	
1929-1928	
1928-1927	
1927-1926	
1926-1925	
1925-1924	
1924-1923	
1923-1922	
1922-1921	
1921-1920	
1920-1919	
1919-1918	
1918-1917	
1917-1916	
1916-1915	
1915-1914	
1914-1913	
1913-1912	
1912-1911	
1911-1910	
1910-1909	
1909-1908	
1908-1907	
1907-1906	
1906-1905	
1905-1904	
1904-1903	
1903-1902	
1902-1901	
1901-1900	
1900-1899	
1899-1898	
1898-1897	
1897-1896	
1896-1895	
1895-1894	
1894-1893	
1893-1892	
1892-1891	
1891-1890	
1890-1889	
1889-1888	
1888-1887	
1887-1886	
1886-1885	
1885-1884	
1884-1883	
1883-1882	
1882-1881	
1881-1880	
1880-1879	
1879-1878	
1878-1877	
1877-1876	
1876-1875	
1875-1874	
1874-1873	
1873-1872	
1872-1871	
1871-1870	
1870-1869	
1869-1868	
1868-1867	
1867-1866	
1866-1865	
1865-1864	
1864-1863	
1863-1862	
1862-1861	
1861-1860	
1860-1859	
1859-1858	
1858-1857	
1857-1856	
1856-1855	
1855-1854	
1854-1853	
1853-1852	
1852-1851	
1851-1850	
1850-1849	
1849-1848	
1848-1847	
1847-1846	
1846-1845	
1845-1844	
1844-1843	
1843-1842	
1842-1841	
1841-1840	
1840-1839	
1839-1838	
1838-1837	
1837-1836	
1836-1835	
1835-1834	
1834-1833	
1833-1832	
1832-1831	
1831-1830	
1830-1829	
1829-1828	
1828-1827	
1827-1826	
1826-1825	
1825-1824	
1824-1823	
1823-1822	
1822-1821	
1821-1820	
1820-1819	
1819-1818	
1818-1817	
1817-1816	
1816-1815	
1815-1814	
1814-1813	
1813-1812	
1812-1811	
1811-1810	
1810-1809	
1809-1808	
1808-1807	
1807-1806	
1806-1805	
1805-1804	
1804-1803	
1803-1802	
1802-1801	
1801-1800	
1800-1799	
1799-1798	
1798-1797	
1797-1796	
1796-1795	
1795-1794	
1794-1793	
1793-1792	
1792-1791	
1791-1790	
1790-1789	
1789-1788	
1788-1787	
1787-1786	
1786-1785	
1785-1784	
1784-1783	
1783-1782	
1782-1781	
1781-1780	
1780-1779	
1779-1778	
1778-1777	
1777-1776	
1776-1775	
1775-1774	
1774-1773	
1773-1772	
1772-1771	
1771-1770	
1770-1769	
1769-1768	
1768-1767	
1767-1766	
1766-1765	
1765-1764	
1764-1763	
1763-1762	
1762-1761	
1761-1760	
1760-1759	
1759-1758	
1758-1757	
1757-1756	
1756-1755	
1755-1754	
1754-1753	
1753-1752	
1752-1751	
1751-1750	
1750-1749	
1749-1748	
1748-1747	
1747-1746	
1746-1745	
1745-1744	
1744-1743	
1743-1742	
1742-1741	
1741-1740	
1740-1739	
1739-1738	
1738-1737	
1737-1736	
1736-1735	
1735-1734	
1734-1733	
1733-1732	
1732-1731	
1731-1730	
1730-1729	
1729-1728	
1728-1727	
1727-1726	
1726-1725	
1725-1724	
1724-1723	
1723-1722	
1722-1721	
1721-1720	
1720-1719	
1719-1718	
1718-1717	
1717-1716	
1716-1715	
1715-1714	
1714-1713	
1713-1712	
1712-1711	
1711-1710	
1710-1709	
1709-1708	
1708-1707	
1707-1706	
1706-1705	
1705-1704	
1704-1703	
1703-1702	
1702-1701	
1701-1700	
1700-1699	
1699-1698	
1698-1697	
1697-1696	
1696-1695	
1695-1694	
1694-1693	
1693-1692	
1692-1691	
1691-1690	
1690-1689	
1689-1688	
1688-1687	
1687-1686	
1686-1685	
1685-1684	
1684-1683	
1683-1682	
1682-1681	
1681-1680	
1680-1679	
1679-1678	
1678-1677	
1677-1676	
1676-1675	
1675-1674	
1674-1673	
1673-1672	
1672-1671	
1671-1670	
1670-1669	
1669-1668	
1668-1667	
1667-1666	
1666-1665	
1665-1664	
1664-1663	
1663-1662	
1662-1661	
1661-1660	
1660-1659	
1659-1658	
1658-1657	
1657-1656	
1656-1655	
1655-1654	
1654-1653	
1653-1652	
1652-1651	
1651-1650	
1650-1649	
1649-1648	
1648-1647	
1647-1646	
1646-1645	
1645-1644	
1644-1643	
1643-1642	
1642-1641	
1641-1640	
1640-1639	
1639-1638	
1638-1637	
1637-1636	
1636-1635	
1635-1634	
1634-1633	
1633-1632	
1632-1631	
1631-1630	
1630-1629	
1629-1628	
1628-1627	
1627-1626	
1626-1625	
1625-1624	
1624-1623	
1623-1622	
1622-1621	
1621-1620	
1620-1619	
1619-1618	
1618-1617	
1617-1616	
1616-1615	
1615-1614	
1614-1613	
1613-1612	
1612-1611	
1611-1610	
1610-1609	
1609-1608	
1608-1607	
1607-1606	
1606-1605	
1605-1604	
1604-1603	
1603-1602	
1602-1601	
1601-1600	
1600-1599	
1599-1598	
1598-1597	
1597-1596	
1596-1595	
1595-1594	
1594-1593	
1593-1592	
1592-1591	
1591-1590	
1590-1589	
1589-1588	
1588-1587	
1587-1586	
1586-1585	
1585-1584	
1584-1583	
1583-1582	
1582-1581	
1581-1580	
1580-1579	
1579-1578	
1578-1577	
1577-1576	
1576-1575	
1575-1574	
1574-1573	
1573-1572	
1572-1571	
1571-1570	
1570-1569	
1569-1568	
1568-1567	
1567-1566	
1566-1565	
1565-1564	
1564-1563	
1563-1562	
1562-1561	
1561-1560	
1560-1559	
1559-1558	
1558-1557	
1557-1556	
1556-1555	
1555-1554	
1554-1553	
1553-1552	
1552-1551	
1551-1550	
1550-1549	
1549-1548	
1548-1547	
1547-1546	
1546-1545	
1545-1544	
1544-1543	
1543-1542	
1542-1541	
1541-1540	
1540-1539	
1539-1538	

APPENDIX

Although Lemma 1 is simple to state compactly, its implications are more clear when cases are considered individually:

If	and	then
$p > q$	$b < p - q$	$q \leq \bar{q} \leq p - 1$
	$b = p - q$	$\bar{q} = p - 1$
	$b > p - q$	$\bar{q} = p$
$p = q$		$\bar{q} = p$
$p < q$	$b < q - p$	$p + 2 \leq \bar{q} \leq q$
	$b = q - p$	$\bar{q} = p + 1$
	$b > q - p$	$\bar{q} = p + 1$

Many results can be stated immediately from examination of these individual cases. Five such results are:

Result 1. *If $\{X_t\}$ is $AR(p)$, then $\{\bar{X}_t\}$ is $ARMA(p, \bar{q})$, as studied by Amemiya and Wu (1972). Additionally, $1 \leq \bar{q} \leq p$.*

Result 2. *If $\{X_t\}$ is $MA(q)$, then $\{\bar{X}_t\}$ is $MA(\bar{q})$, where $1 \leq \bar{q} \leq q$.*

Result 3. *If $\{X_t\}$ is AR or $ARMA$ with batch size satisfying $0 \leq p - q < b$, then $\{\bar{X}_t\}$ is $ARMA(p, p)$.*

Result 4. *If $p \geq q$, then $\lim_{b \rightarrow \infty} \bar{q} = p$.*

Result 5. *If $p < q$, then $\lim_{b \rightarrow \infty} \bar{q} = p + 1$.*

Of course, considering only the order of the batch-means process can be misleading. For example, Results 4 and 5 indicate that large batches lead to MA components of order p or $p + 1$; in particular, a batched $MA(q)$ process converges to an $MA(1)$ process. But large batches are asymptotically independent. The explanation is that $\bar{\theta}_1$ is approaching zero as batch size increases. An implication is that, even for this nicest case of $ARMA$ underlying processes, estimation of the order of the batch-means process is likely to be difficult.

DISTRIBUTION LIST

	NO. OF COPIES
Library (Code 0142) Naval Postgraduate School Monterey, CA 93943-5000	2
Defense Technical Information Center Cameron Station Alexandria, VA 22314	2
Office of Research Administration (Code 012) Naval Postgraduate School Monterey, CA 93943-5000	1
Center for Naval Analyses 2000 Beauregard Street Alexandria, VA 22311	1
Operations Research Center, Room E40-164 Massachusetts Institute of Technology Attn: R. C. Larson and J. F. Shapiro Cambridge, MA 02139	1
Library (Code 55) Naval Postgraduate School Monterey, CA 93943-5000	1
Office of Naval Research Arlington, VA 22217	1
Keebom Kang Department of Industrial Engineering University of Miami Coral Gables, FL 33124	10
Bruce W. Schmeiser Code 55Sc Naval Postgraduate School Monterey, CA 93943-5000	10

DTIC

FILMED

4-86

END