

AD-A173 340


TRANSITION DIPOLE-SOLVENT INTERACTION IN OPTICAL
ELECTRON TRANSFER(U) NEW YORK UNIV NY DEPT OF CHEMISTRY
P DELAHAY ET AL. 30 SEP 86 NYU/DC/TR-12-NEW SERIES-2
N00014-82-K-0113


1/1

UNCLASSIFIED

F/G 7/4

ML





MICROCOPY RESOLUTION TEST CHART
NATIONAL BUREAU OF STANDARDS-1963-A

AD-A173 340

OFFICE OF NAVAL RESEARCH
Contract N00014-82-K-0113
Task No. NR 359-258



TECHNICAL REPORT NO. NYU/DC/TR-12-NEW SERIES-2

TRANSITION DIPOLE-SOLVENT INTERACTION
IN OPTICAL ELECTRON TRANSFER

by

Paul Delahay and Andrew Dzedzic

Prepared for publication in
Accounts of Chemical Research

New York University
Department of Chemistry
New York, NY

September 1986

DTIC
OCT 22 1986
A

Reproduction in whole or in part is permitted for
any purpose of the United States Government

This document has been approved for public release
and sale; its distribution is unlimited

OTIC FILE COPY

406817- SH

6 10 017

REPORT DOCUMENTATION PAGE

1a. REPORT SECURITY CLASSIFICATION NA		1b. RESTRICTIVE MARKINGS NA	
2a. SECURITY CLASSIFICATION AUTHORITY NA		3. DISTRIBUTION / AVAILABILITY OF REPORT This document has been approved for public release and sale; its distribution is unlimited.	
2b. DECLASSIFICATION / DOWNGRADING SCHEDULE NA		4. PERFORMING ORGANIZATION REPORT NUMBER(S) NYU/DC/TR-12-NEW SERIES-2	
4. PERFORMING ORGANIZATION REPORT NUMBER(S) NYU/DC/TR-12-NEW SERIES-2		5. MONITORING ORGANIZATION REPORT NUMBER(S) NA	
6a. NAME OF PERFORMING ORGANIZATION Department of Chemistry New York University	6b. OFFICE SYMBOL (If applicable) NA	7a. NAME OF MONITORING ORGANIZATION ONR	
6c. ADDRESS (City, State, and ZIP Code) New York, NY 10003		7b. ADDRESS (City, State, and ZIP Code) 800 N. Quincy Street Arlington, VA 22217	
8a. NAME OF FUNDING / SPONSORING ORGANIZATION	8b. OFFICE SYMBOL (If applicable) NA	9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER Contract N00014-82-K-0113	
8c. ADDRESS (City, State, and ZIP Code)		10. SOURCE OF FUNDING NUMBERS	
		PROGRAM ELEMENT NO.	PROJECT NO.
		TASK NO.	WORK UNIT ACCESSION NO.
11. TITLE (Include Security Classification) Transition Dipole-Solvent Interaction in Optical Electron Transfer			
12. PERSONAL AUTHOR(S) Paul Delahay and Andrew Dziedzic			
13a. TYPE OF REPORT Technical	13b. TIME COVERED FROM _____ TO _____	14. DATE OF REPORT (Year, Month, Day) 9/30/86	15. PAGE COUNT 30
16. SUPPLEMENTARY NOTATION Prepared for publication in Accounts of Chemical Research			
17. COSATI CODES		18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)	
FIELD	GROUP	SUB-GROUP	
			Dielectric dispersion Dispersion spectra Photoemission
			Solutions Transition dipole VUV spectroscopy
19. ABSTRACT (Continue on reverse if necessary and identify by block number) A new and fundamental feature of photoionization from solutions is presented. The oscillating transition dipole for electronic transitions from bound states in solution to the continuum interacts with the surrounding molecules of the solvent. This interaction is similar to solvation. The transition dipole, oscillating at the frequency of the incident radiation, induces electronic polarization in the solvent. Thus, the minimum energy required for photoionization and the kinetic energy of quasifree electrons generated in solution vary with photon energy. This variation or shift results from dielectric dispersion of the solvent and is displayed in dispersion spectra. Such dispersion shifts are fundamental to all photoionization processes in condensed matter and thus a nonequilibrium electronic contribution to the photoionization energy is required. Before this work, only the contribution from nuclear reorganization was recognized. This phenomenon may have significant effects on photoionization rates and emission laws because of the rich structure of the dispersion spectra. The effect of ionic atmosphere screening on the dispersion shift is also investigated. <i>Keywords:</i>			
20. DISTRIBUTION / AVAILABILITY OF ABSTRACT <input checked="" type="checkbox"/> UNCLASSIFIED/UNLIMITED <input type="checkbox"/> SAME AS RPT. <input type="checkbox"/> DTIC USERS		21. ABSTRACT SECURITY CLASSIFICATION Unclassified	
22a. NAME OF RESPONSIBLE INDIVIDUAL Paul Delahay		22b. TELEPHONE (Include Area Code) 212-598-2811	22c. OFFICE SYMBOL

Transition Dipole-Solvent Interaction in Optical Electron Transfer

PAUL DELAHAY* AND ANDREW DZIEDZIC[†]

Department of Chemistry, New York University, New York, New York 10003

Received

Spectrographic transitions from one bound state to another involve a transition dipole moment. The 1s and 2p states of the hydrogen atom, for instance, have no permanent dipole moment, but the combination of these two states, which is involved in the 1s-2p transition, has a dipole moment. This dipole moment oscillates at or near the frequency of the incident radiation according to the time-dependent perturbation theory of quantum mechanics.¹ Such an oscillating transition dipole is also involved in transitions to the continuum² (photoionization), but in that case there is, of course, ultimate separation of the emitted electron from its parent atomic or molecular system.

If the substance being photoionized is in solution, the oscillating transition dipole interacts with the surrounding solvent molecules in a process which is similar to solvation. Only electronic motion is involved in this interaction to the exclusion of nuclear motion. This process is described microscopically in terms of dipole-dipole interaction and more generally as a multipole-multipole interaction. The oscillating electric field of the oscillating transition dipole in the macroscopic treatment induces electronic polarization of the solvent which is treated as a continuous medium. The transition dipole-solvent interaction causes the minimum energies required for photoionization of substances in solution to vary with the photon energy at which photoionization is observed. Thus, the kinetic energy of quasifree electrons emitted into the liquid upon generation by the photoionization process does not depend on a constant photoionization

energy, as in the gas phase, but on a photoionization energy varying with photon energy. The varying shift in the photoionization energy results from dielectric dispersion, as will be shown in the present account. This dispersion shift (≤ 0.2 eV) was recently discovered by the authors³ and interpreted^{4,5} quantitatively. The dispersion shift affects all photoionization processes in solution and more generally in condensed matter. The dispersion shift is negligible in gas-phase photoionization because the prevailing distances between adjacent molecules are too great to allow significant transition dipole-molecule interaction.

Let us first examine the dispersion shift in terms of a continuous medium model of the solvent. A microscopic description will be given in the theoretical section. The transition dipole oscillates at or near the frequency of the incident radiation, as pointed out above, and consequently the response of the solvent is determined by its dielectric properties at the frequency of the oscillating dipole. The optical dielectric constant of gases, liquids and solids varies with the frequency of the applied electric field on account of dielectric dispersion. Thus, the optical dielectric constant ϵ_{op} of the solvent increases progressively with frequency in the vicinity of an absorption band of the solvent (normal dispersion). This increase of ϵ_{op} for water easily reaches 40 percent of its limiting value of ϵ_{op}^0 in the near-infrared region. The oscillating transition dipole therefore interacts with the solvent at the prevailing value ϵ_{op} rather than the limiting value ϵ_{op}^0 . This effect of dispersion results in a nonequilibrium electronic contribution to the photoionization energy, as will be shown presently.

If dielectric dispersion is negligible, the free energy of photoionization of a substance in solution consists of two contributions⁶:

(i) The difference between the free energies of the initial and final states which can be calculated from thermodynamic data. For example, the initial state in the photoionization of Fe^{2+} ions in aqueous solution is the hydrated Fe^{2+} ion, and the final state consists of the hydrated Fe^{3+} ion and a quasifree electron in water.

(ii) The free energy of reorganization of the nuclear coordinates of the solute and solvent molecules near the solute. Nuclear reorganization occurs after removal of the emitted electron from the solute because the solvent initially retains the nuclear configuration it had prior to photoionization. For instance, the nuclear configuration of the solvent around a Fe^{2+} ion in aqueous solution is initially unchanged after photoionization to a Fe^{3+} ion. The solvent around the Fe^{3+} ion subsequently reorganizes to the final nuclear configuration of a solvated Fe^{3+} ion. The free energy R thus corresponds to a nonequilibrium nuclear contribution to the free energy of photoionization.

The energy of transition dipole-solvent interaction under conditions of negligible dielectric dispersion of the solvent is included in the difference between the free energies of the initial and final states. It does not matter in that case how the transition occurs from the point of view of energetics. There is, of course, a transition dipole-solvent interaction in the absence of dispersion, but this interaction prevails under conditions in which the solvent has the optical dielectric constant ϵ_{op}^0 , and no explicit consideration of transition dipole-solvent interaction is necessary. If dielectric dispersion is not negligible at the prevailing photon energy, the transition dipole interacts with a solvent characterized by the optical dielectric constant ϵ_{op} rather than its limiting value ϵ_{op}^0 . The difference W between the energies for transition dipole-solvent interaction at

ϵ_{op} and ϵ_{op}^0 therefore must be added to the two previously considered contributions to the photoionization energy. This energy consequently consists of three contributions: (i) the thermodynamic contribution determined by the initial and final states; (ii) the nonequilibrium nuclear contribution R; and (iii) the nonequilibrium electronic contribution W.

The foregoing considerations are fundamental to the understanding of photoionization in solutions. Their validity is amply documented as far as the thermodynamic and nuclear reorganization contributions are concerned.⁶ Experimental evidence for the third contribution, the dispersion shift W, is presented in the next section.

Experimental Evidence of the Dispersion Shift

The observation of how the dispersion shift W varies with photon energy requires a precise determination of the rate of photoionization in solution as a function of photon energy. This experimental requirement is satisfied in the study of photoelectron emission by liquids and solutions.^{6,7} The basic experiment is as follows:

The surface of the liquid or solution is irradiated at normal incidence with photons of sufficient energy to cause emission of electrons into the vapor phase above the liquid. Electrons emitted into the gas phase are collected by a grid electrode parallel to the surface of the liquid, and the emission current is measured. The emission yield, Y at the photon energy E, is defined as the number of electrons collected per incident photon. An emission spectrum is obtained by plotting Y as a function of E. Quantum theory^{8,9} of the emission process predicts and experiment confirms^{6,7} that emission follows a quadratic law,

$$Y = K(E - E_t)^2.$$

(1)



APP

The quantity K is independent of E and its explicit form is unimportant for our purpose. The quantity E_t is the threshold energy for photoelectron emission by the liquid or solution into the gas phase. Equation 1 holds for photon energies E higher than E_t by at least a few tenths of an electronvolt. The exponent is greater than 2 near the threshold, but only the range of energies over which the quadratic law applies concerns us.

The threshold energy E_t can be identified to a good approximation (to within 0.1 eV) with the photoionization free energy.^{6,7} Thus,

$$E_t = \Delta G_{th} + R + W, \quad (2)$$

where ΔG_{th} is the difference between the free energies of the initial and final states of the photoemission process, R is the free energy of nuclear reorganization, and W is the dispersion shift. For instance, the hydrated Fe^{2+} ion is the initial state in photoelectron emission from aqueous solutions of Fe^{2+} ions, and the final state consists of the hydrated Fe^{3+} ion and an electron in the gas phase. The terms ΔG_{th} and R do not depend on E , as noted in the introductory remarks, and consequently the dependence of E_t on E is determined solely by the variations of W with photon energy. If W is nearly independent of E , one deduces from eq 2 that Y exhibits a parabolic dependence on E . Indeed, this is very nearly the case for photoelectron emission by liquid water^{6,7} ($E_t = 10.06$ eV) in agreement with predictions from the dielectric properties of liquid water near 10 eV. In general, fine structure is observed in the emission spectra, that is, in the plots of Y against E (Figure 1). This is the case for emission by aqueous³ and nonaqueous⁵ solutions of all substances examined thus far (anions, cations, molecules). This fine structure and its quantitative interpretation constitute the evidence for the dispersion shift W .

The departure from the quadratic law resulting from the dispersion shift can be displayed conveniently by direct application of eq 1. It follows from this equation that a plot of $Y^{1/2}$ against E is linear if E_t is independent of E . This linear relationship is obeyed in the range of validity of the quadratic emission law, that is, for E higher than E_t by at least a few tenths of an electronvolt. One deduces from eq 1,

$$dY^{1/2}/dE = K(1 - dE_t/dE) \quad (3)$$

or in view of the equality $dE_t/dE = dW/dE$ (see above),

$$dY^{1/2}/dE = K(1 - dW/dE) \quad (4)$$

Variations of E_t with E therefore are detected readily from a plot (Figure 2) of $dY^{1/2}/dE$ against E . Such plots are termed dispersion spectra.

Dispersion spectra such as those of Figure 2 are determined essentially by the solvent and in a minor way by the nature and concentration of the substance being photoionized. This conclusion is valid for figure 2 in which the four dispersion spectra of very different inorganic anions are strikingly similar. This essential property of dispersion spectra was established in a series of experiments covering the photoelectron emission by aqueous solution of 17 inorganic anions,³ inorganic cations¹⁰ and various molecules.¹⁰ The evidence also includes a study of photoelectron emission by various organic liquids.⁵ It was also shown that dispersion spectra do not result from experimental artifacts having their origin in the source of vacuum ultraviolet radiation or the monochromator and optical system.⁵ Attenuation of the photon flux by absorption by the solvent was also ruled out as a possible cause of the results displayed in dispersion spectra.^{3,4} Finally, it should be noted that dispersion spectra are not very sensitive to the choice of exponent in the emission law of eq 1. If one assumes, for instance, that the yield Y is proportional to $(E - E_t)^{5/2}$, the resulting dispersion

spectrum, $dY^{2/5}/dE$ against E , is quite similar to the plot of $dY^{1/2}/dE$ against E for the same substance, and both dispersion spectra exhibit the same characteristic features (e.g., extrema) at nearly the same photon energies.

Theory of the Dispersion Shift

A theory of the dispersion shift must account for the rather rich structure of experimental dispersion spectra (Figure 2). Furthermore, this structure must be accounted for primarily on the basis of the dielectric properties of the solvent in the prevailing range of photon energies. These requirements follow directly from the observation that dispersion spectra are determined primarily by the nature of the solvent.

Two limiting cases may be considered in the calculation of the transition dipole-solvent interaction energy:

(i) One assumes that the photoelectron produced by photoionization is "ejected" so far from its parent (ion or molecule) that the electrostatic interaction between the parent species and the ejected electron is negligible. Thus, one assumes complete separation of the ejected electron from its parent. The energy for the interaction between the oscillating transition dipole and the solvent is then calculated by considering only the change in the parent field from initial to final state. No attempt is made to follow the time evolution of the transition dipole-solvent interaction. Indeed, the matrix element for absorption refers only to the initial and final states, and photon absorption is interpreted in terms of an annihilation operator acting in occupation space.¹¹

(ii) One assumes that the transition dipole is represented by a point dipole whose magnitude oscillates at or near the frequency of radiation. Energy for interaction with the solvent is then calculated for a given, plausible magnitude of the point dipole.

The first approach is akin to the treatment of ionic solvation with the restriction that only electronic motion is taken into account to the exclusion of nuclear motion in the solvent. The second approach is related directly to the calculation of the free energy of solvation of a dipole with the same restriction about nuclear motion as for the first approach. The dielectric properties of the solvent in both approaches are taken at the frequency of the incident radiation (cf. introductory remarks). Detailed theoretical calculations of the enthalpy of ionic solvation have been made,¹²⁻¹⁴ and a fairly standard model is available¹³ for such calculations for inorganic cations. The different orientations of solvent molecules around inorganic anions and cations in aqueous solution¹⁵ can readily be taken into account.⁴ This model yields solvation enthalpies within a few percent of the enthalpies deduced from experimental data for inorganic ions in aqueous solution. In contrast, the solvation of a point dipole is treated¹⁵ for a cruder model (continuous medium) than ionic solvation. It turns out that the ionic and dipole solvation models yield essentially the same functional dependence on photon energy for dispersion spectra of aqueous solutions in the 7 to 10 eV range of photon energies.⁵ The dispersion shifts calculated from the ionic solvation model are too high by a factor of 2 or 3 whereas the values of W from the dipole solvation model are too low by at least an order of magnitude. This is to be expected from models corresponding to limiting cases. Only the dependence of W on photon energy, as reflected in plots of $-dW/dE$ against E (eq 4), will be discussed in the present account. The ionic solvation model, which is fully satisfactory for this purpose, is retained solely here.

In the case of ions, the volume around the ion being photoionized is divided into two regions. (i) The inner-sphere shell consists of N solvent

molecules around the photoionized ion. The solvent molecules are treated as point dipoles at a distance of $r_c + r_w$ from the ionic charge, r_c and r_w being the crystallographic radii of the central ion and water, respectively. (ii) The outer-sphere region outside the sphere of radius $a = r_c + 2r_w$ is treated as a continuous medium. The continuous medium model is used exclusively for the photoionization of electrically neutral species. The inner-sphere shell is treated microscopically in terms of the polarizability α of the solvent whereas the outer-sphere region is given a macroscopic treatment based on the optical dielectric constant ϵ_{op} of the solvent. The quantities α and ϵ_{op} are related by the Lorenz-Lorentz equation.¹⁷ The polarizability α and dielectric constant ϵ_{op} are real quantities for transparent solvents and are complex for absorbing solvents. The case of transparent solvents is treated first.

The electric field of the solute induces a dipole in each of the N solvent molecules in the inner-sphere shell. The induced moment is proportional to the solvent polarizability. These induced dipoles experience different interactions which can be calculated by using a multipole expansion of the central field: charge-induced dipole, dipole-induced dipole, induced dipole-induced dipole and quadrupole-induced dipole interactions. Solute-solvent and solvent-solvent London dispersion and Born repulsion must also be taken into account. All these energies of interaction depend directly or indirectly on the solvent polarizability.

The change in the ionic valence of the solute resulting from photoionization causes a change in the energies of the interactions in the inner-sphere shell. If dielectric dispersion is negligible, the net change of interaction energies in the inner-sphere shell is included in the difference of energies between the initial and final states of the photoionization

process. Actually, the solvent polarizability α at the radiation frequency prevails in photoionization instead of the value α^0 in the absence of radiation (cf. introductory remarks). The contribution ΔP_{in} for the inner-sphere shell to the dispersion shift W therefore is equal to the difference between the net interaction energies calculated for the polarizabilities α and α^0 . The detailed calculation is given in ref. 4.

The contribution ΔP_{out} from the outer-sphere region to the dispersion shift W is equal to the difference between the free energies of electronic polarization of the medium for the dielectric constants ϵ_{op} at the radiation frequency and ϵ_{op}^0 without radiation. The free energy difference ΔP_{out} is calculated from the Born model of ionic solution. Thus,

$$\Delta P_{out} = (e^2/2a)[(\epsilon_{op}^0)^{-1} - \epsilon_{op}^{-1}], \quad (5)$$

where e is the electronic charge, and the radii a was defined above.

The optical dielectric constant at photon energies at which the solvent absorbs radiation is a complex quantity¹⁷ $\epsilon_1 - i\epsilon_2$ in which the real part ϵ_1 pertains to polarization of the medium and the imaginary part ϵ_2 corresponds to the dissipation of energy in the medium. The polarizability α is also a complex quantity.¹⁷ The dispersion shift W consists now of a real part $P = \Delta P_{in} + \Delta P_{out}$ for polarization of the solvent and an imaginary part L for losses or dissipation of energy in the medium. The real P of W is calculated for the same model of inner- and outer-sphere regions as for transparent solvents by generalization of the equations applied to the latter case.⁴ For instance, the result,

$$\Delta P_{out} = (e^2/2a)[(\epsilon_{op}^0)^{-1} - \epsilon_1/(\epsilon_1^2 + \epsilon_2^2)], \quad (6)$$

for an absorbing solvent reduces to eq 5 for $\epsilon_2 = 0$ (transparent medium).

The calculation of the imaginary part L of W poses a problem which is not encountered in ionic solvation. The latter deals with a static problem of

polarization which is treated by considering the initial and final states of the solvation process. The calculation of L pertains to a dynamic problem dealing with the energy dissipated as heat in the medium between the initial and final states of the photoionization process. The problem is related to the treatment of the dissipation of energy in a dielectric having a significant ionic and/or electronic conductivity. The calculation for absorbing solvents is attacked classically by introducing the concept of an optical conductivity $\omega\epsilon_2/4\pi$ at the photon energy $h\omega$. This interpretation holds even if the dielectric (solvent) does not exhibit any conductivity from free charges,¹⁸ and the imaginary part ϵ_2 of the optical dielectric constant arises solely from bound-bound transitions of the solvent. The loss of energy is calculated in ref. 4 for the inner-sphere shell (L_{in}) and the outer-sphere region (L_{out}). The quantity L_{out} , for instance, is obtained from the conduction current and the corresponding ohmic loss which results from the optical conductivity and the change of the field around the photoionized species from the initial to the final state. One obtains⁴

$$L_{out} = (e^2/a)\epsilon_2/(\epsilon_1^2 + \epsilon_2^2). \quad (7)$$

The loss L_{in} is calculated much in the same way as ΔP_{in} for transparent solvents by summing the negative imaginary parts of the complex components of ΔP_{in} . Details are given in ref. 4.

The dispersion shift is

$$W = (P^2 + L^2)^{1/2}, \quad (8)$$

where $P = \Delta P_{in} + \Delta P_{out}$ and $L = L_{in} + L_{out}$ are the real and imaginary parts of W , respectively, as noted above. Comparison with experimental dispersion spectra by application of eq 4 requires the derivative dW/dE , namely

$$dW/dE = [P/(P^2 + L^2)^{1/2}]dP/dE + [L/(P^2 + L^2)^{1/2}]dL/dE. \quad (9)$$

Variations of $-dW/dE$ and its two components (eq 9) with photon energy are displayed in Figure 3 for liquid water.⁴ These curves were computed by using the literature values of ϵ_1 and ϵ_2 obtained by reflectance spectroscopy of liquid water. The latter exhibits (up to 10 eV) two absorption bands¹⁹ having their maxima at ca. 8.2 and 10.0 eV, respectively. Figure 3 shows that the term in the derivative dL/dE (eq 9) pertaining to the energy loss in the solvent approaches zero at or near the photon energies of the absorption maxima. Conversely, the term in the derivative of dP/dE (eq 9) corresponding to polarization of the solvent reaches a maximum at or near the photon energy of the absorption maximum. The relative importance of these two contributions to dW/dE determines the effect of the ionic strength of the solution on the shape of dispersion spectra. This point is discussed in the next section.

Inner-outer Sphere Splitting from Screening by the Ionic Atmosphere

The curve representing the variations of $-dW/dE$ with photon energy in Figure 3 resembles the dispersion spectra of Figure 2, but a more detailed comparison of experiment and theory requires the consideration of electrostatic screening by the ionic atmosphere. This screening was not taken into account in the previous calculation. One may assume to a first approximation that the ionic atmosphere around the ion being photoionized lowers the amplitude of the electric field in the outer-sphere region but not in the inner-sphere shell. Screening in the inner-sphere is introduced later. Thus, ΔP_{out} and L_{out} decrease with increasing ionic strength whereas ΔP_{in} and L_{in} are not affected, at least to a first approximation. The relative importance of the contributions from the inner-sphere and outer-sphere regions therefore varies with ionic strengths, and the shape of the dispersion spectrum changes accordingly. This change is shown in Figure 4

in which screening is characterized by the factor S . One has $s = 1$ in the absence of screening and $S = 0$ for complete screening of the outer-sphere regions ($\Delta P_{\text{out}} = 0$, $L_{\text{out}} = 0$). A calculation of S based on the Debye-Hückel theory is given in ref. 4. The factor $S = -0.1$ in Figure 4 is empirical and is introduced to account for partial screening of the inner-sphere shell as well as possible counterion penetration of the first hydration sphere for high ionic strength solutions.

The change in the shape of the dispersion spectra of Figure 4 becomes quite pronounced for $S \leq 0.2$. There is a change in shape because the terms in dP/dE and dL/dE of eq 9 peak at different photon energies (Figure 3). The photon energies at the extrema of the dispersion spectra for $S = 1$ and $S = 0$ are listed in Table I. A maximum in dispersion spectra is observed between two minima in the range of normal dispersion, and conversely a minimum between two maxima appears in the range of anomalous dispersion. This pattern of extrema was referred to as inner-outer sphere splitting in ref. 4.

The photon energies at the extrema of experimental dispersion spectra from rather extensive data are compared in Table I with the corresponding theoretical values. The excellent agreement between experiment and theory give strong support to the present interpretation of the dispersion shift and its calculation. The shapes of calculated and experimental curves representing dispersion spectra also agree quite well (Figure 5) for dilute (0.25 M NaCNS) and concentrated (2 M $\text{Na}_2\text{S}_2\text{O}_3$) solutions. A divalent anion was selected for the concentrated solution to achieve high ionic strength and insure extensive screening by the ionic atmosphere. The experimental results of Figure 5 and the change in the shape of dispersion spectra upon increase of the ionic strength confirm the theoretical predictions of Figure 4.

Conclusion

A fundamental and novel feature of photoionization in solution is discussed in the present account. Thus, dielectric dispersion of the solvent and interaction between transition dipoles and the solvent result in a nonequilibrium electronic contribution (dispersion shift) to the photoionization energy of substances in solution. Only the nonequilibrium contribution from nuclear reorganization to the photoionization energy had been recognized and studied prior to the work discussed here. The dispersion shift should affect all photoionization processes in solution and, more generally, in condensed matter. The effect of this shift is readily observed under conditions of significant dielectric dispersion. Photoionization rates and emission laws therefore can be affected significantly because of the rather rich structure of dispersion spectra. Screening by the ionic atmosphere can also be investigated in a novel way on the basis of dispersion spectra.

The present account was prepared with support from the Office of Naval Research.

[‡]Present address: The Perkin-Elmer Corp., 50 Danbury Road, Wilton, Conn. 06897.

- (1) J. D. Macomber, "The Dynamics of Spectroscopic Transitions," Wiley, New York, 1976, pp. 95-108.
- (2) E. Merzbacher, "Quantum Mechanics," Wiley, New York, 1961, pp. 459-460.
- (3) P. Delahay and A. Dziedzic, J. Chem. Phys., 80, 5381 (1984).
- (4) P. Delahay and A. Dziedzic, J. Chem. Phys., 84, 936 (1986).
- (5) P. Delahay and A. Dziedzic, Chem. Phys. Lett., in press.

- (6) P. Delahay in "Electron Spectroscopy: Theory, Techniques and Applications," vol. 5, C. R. Brundle and A. D. Baker, eds., Academic Press, London, 1984, pp. 123-196.
- (7) P. Delahay, Acc. Chem. Res., 15, 40 (1982).
- (8) A. M. Brodsky and A. V. Tsarevsky, J. Chem. Soc., Faraday Trans. 2, 72, 1781 (1976).
- (9) A. M. Brodsky, J. Phys. Chem., 84, 1856 (1980).
- (10) P. Delahay and A. Dziedzic, J. Chem. Phys., 81, 3678 (1984).
- (11) J. J. Sakurai, "Advanced Quantum Mechanics," Addison-Wesley, Reading, Mass., 1967, p. 37.
- (12) A. D. Buckingham, Discuss. Faraday Soc., 24, 151 (1957).
- (13) W. E. Morf and W. Simon, Helv. Chim. Acta, 54, 794 (1971).
- (14) B. E. Conway, "Ionic Hydration in Chemistry and Biophysics," Elsevier, Amsterdam, 1981, pp. 312-341.
- (15) J. E. Enderby, Ann. Rev. Phys. Chem., 34, 155 (1983).
- (16) J. G. Kirkwood, J. Chem. Phys., 2, 351 (1934)
- (17) C. J. F. Böttcher and P. Bordewijk, "Theory of Electric Polarization," 2nd ed., Elsevier, Amsterdam, 1978, Vol. II, pp. 23-27, 292-301, 395-404.
- (18) J. D. Axe in "Solid State Chemistry and Physics," vol. 1, P. F. Waller, ed., Marcel Dekker, New York, 1983, p. 423.
- (19) J. M. Heller, Jr., R. N. Hamm, R. D. Birkhoff and L. R. Painter, J. Chem. Phys., 60, 3483 (1974).

Table I

Photon Energies at the Extrema of Calculated and Experimental
Dispersion Spectra of Aqueous Solutions⁴

Dispersion	Extremum	Dominant contribution	Calculated photon energy, eV	Experimental photon energy, eV
normal (< 7.75 eV)	min	inner	7.24	no datum
	max	mixed	7.30	7.30
	min	outer	7.42	7.41
anomalous (7.75 to 8.70 eV)	max	inner	8.24	8.16 ± 0.06
	min	mixed	8.58	8.42 ± 0.06
	max	outer	8.68	8.65 ± 0.03
normal (8.70 to 9.70 eV)	min	inner	9.18	9.11 ± 0.03
	max	mixed	9.38	9.37 ± 0.06
	min	outer	9.63	9.63 ± 0.12
anomalous (> 9.7 eV)	max	inner	9.96	9.86 ± 0.04
	min	mixed	10.11	10.18
	max	outer	10.32	10.35

Data for 0.5 M VCl_2 and 1 M CrCl_2 (7.30 and 7.41 eV) and 0.05 M $\text{K}_4\text{Fe}(\text{CN})_6$ (7.41 eV) from ref 10. Average values with standard deviation ($8.16 \leq E \leq 10$ eV) for 1 M solutions of 17 inorganic anions from ref 3. Extrema at 10.18 and 10.35 eV obtained in ref 4 with 0.2 M KPF_6 .

List of Captions

Figure 1. Variations of the yield with photon energy for photoelectron emission by 1 M aqueous solutions of different anions. Sodium salts except for potassium carbonate. The yield is defined as the number of collected electrons per incident photon. Relative scales of ordinates A to E: 3.0, 5.8, 2.2, 1.5, 1.0 (weakest signal).³

Figure 2. Dispersion spectra of 1 M aqueous solutions of different anions. Sodium salts except for potassium carbonate. Relative scale of ordinates from A to E: 1.9, 1.4, 1.1, 1.0 (weakest signal).³

Figure 3. Variations of $-dW/dE$ with photon energy (top) and dependence of the two contributions of $-dW/dE$ (eq 9) on photon energy (bottom).⁴

Figure 4. Variations of $-dW/dE$ with photon energy for different values of the screening factor S ($S = 0$ for complete screening of the outer-sphere region, $S = -0.1$ for partial screening of the inner-sphere shell). Complete data for the calculations in ref 4. Ordinates at the minimum at 7.42 eV from $S = -0.1$ to $S = 1.0$: -0.164, -0.196, -0.299, -0.263, -0.297, -0.541. Ordinates at the maximum (photon energy between parentheses) from $S = -0.1$ to $S = 1.0$: 0.151 (8.20 eV), 0.151, (8.24 eV), 0.162 (8.35 eV), 0.181 (8.41 eV), 0.216 (8.68 eV), 0.509 (8.68 eV).

Figure 5. Comparison of the experimental dispersion spectra ($dY^{1/2}/dE$ against E) of 0.25 M NaCNS (bottom) and 2 M $Na_2S_2O_3$ (top). Calculated dispersion spectra ($-dW/dE$ against E) for the same data as Figure 4 and a screening factor $S = 0.3$ (bottom) and -0.1 (top). The dispersion spectra for 2 M NaCNS and 0.25 M $Na_2S_2O_3$ (not shown) are intermediate between the curves shown here.




FIG. 1






FIG. 2



PHOTON ENERGY (eV)

FIG. 3




FIG 4




FIG. 5

TECHNICAL REPORT DISTRIBUTION LIST, GEN

	<u>No. Copies</u>		<u>No. Copies</u>
Office of Naval Research Attn: Code 1113 800 N. Quincy Street Arlington, Virginia 22217-5000	2	Dr. David Young Code 334 NORDA NSTL, Mississippi 39529	1
Dr. Bernard Douda Naval Weapons Support Center Code 50C Crane, Indiana 47522-5050	1	Naval Weapons Center Attn: Dr. Ron Atkins Chemistry Division China Lake, California 93555	1
Naval Civil Engineering Laboratory Attn: Dr. R. W. Drisko, Code L52 Port Hueneme, California 93401	1	Scientific Advisor Commandant of the Marine Corps Code RD-1 Washington, D.C. 20380	1
Defense Technical Information Center Building 5, Cameron Station Alexandria, Virginia 22314	12 high quality	U.S. Army Research Office Attn: CRD-AA-IP P.O. Box 12211 Research Triangle Park, NC 27709	1
DTNSRDC Attn: Dr. H. Singerman Applied Chemistry Division Annapolis, Maryland 21401	1	Mr. John Boyle Materials Branch Naval Ship Engineering Center Philadelphia, Pennsylvania 19112	1
Dr. William Tolles Superintendent Chemistry Division, Code 6100 Naval Research Laboratory Washington, D.C. 20375-5000	1	Naval Ocean Systems Center Attn: Dr. S. Yamamoto Marine Sciences Division San Diego, California 91232	1

ABSTRACTS DISTRIBUTION LIST, 359/627

~~Dr. Paul Delahay
Department of Chemistry
New York University
New York, New York 10003~~

Dr. J. Driscoll
Lockheed Palo Alto Research
Laboratory
3251 Hanover Street
Palo Alto, California 94304

Dr. D. N. Bennion
Department of Chemical Engineering
Brigham Young University
Provo, Utah 84602

Dr. R. A. Marcus
Department of Chemistry
California Institute of Technology
Pasadena, California 91125

Dr. J. J. Auborn
Bell Laboratories
Murray Hill, New Jersey 07974

Dr. Joseph Singer, Code 302-1
NASA-Lewis
21000 Brookpark Road
Cleveland, Ohio 44135

Dr. P. P. Schmidt
Department of Chemistry
Oakland University
Rochester, Michigan 48063

Dr. Roger Belt
Litton Industries Inc.
Airtron Division
Morris Plains, NJ 07950

Dr. Ulrich Stimming
Department of Chemical Engineering
Columbia University
New York, NY 10027

Dr. Manfred Breiter
Institut für Technische Elektrochemie
Technischen Universität Wien
9 Getreidemarkt, 1160 Wien
AUSTRIA

Dr. E. Yeager
Department of Chemistry
Case Western Reserve University
Cleveland, Ohio 44106

Dr. C. E. Mueller
The Electrochemistry Branch
Naval Surface Weapons Center
White Oak Laboratory
Silver Spring, Maryland 20910

Dr. Sam Perone
Chemistry & Materials
Science Department
Lawrence Livermore National Laboratory
Livermore, California 94550

Dr. Royce W. Murray
Department of Chemistry
University of North Carolina
Chapel Hill, North Carolina 27514

Dr. Adam Heller
Bell Laboratories
Murray Hill, New Jersey 07974

Dr. A. B. Ellis
Chemistry Department
University of Wisconsin
Madison, Wisconsin 53706

Dr. Steven Greenbaum
Department of Physics and Astronomy
Hunter College
695 Park Ave.
New York, NY 10021

ABSTRACTS DISTRIBUTION LIST, 359/627

Dr. M. Wrighton
Chemistry Department
Massachusetts Institute
of Technology
Cambridge, Massachusetts 02139

Dr. B. Stanley Pons
Department of Chemistry
University of Utah
Salt Lake City, Utah 84112

Donald E. Mains
Naval Weapons Support Center
Electrochemical Power Sources Division
Crane, Indiana 47522

S. Ruby
DOE (STOR)
Room 5E036 Forrestal Bldg., CE-14
Washington, D.C. 20595

Dr. A. J. Bard
Department of Chemistry
University of Texas
Austin, Texas 78712

Dr. Janet Osteryoung
Department of Chemistry
State University of New York
Buffalo, New York 14214

Dr. Donald W. Ernst
Naval Surface Weapons Center
Code R-33
White Oak Laboratory
Silver Spring, Maryland 20910

Mr. James R. Moden
Naval Underwater Systems Center
Code 3632
Newport, Rhode Island 02840

Dr. Bernard Spielvogel
U.S. Army Research Office
P.O. Box 12211
Research Triangle Park, NC 27709

Dr. Aaron Fletcher
Naval Weapons Center
Code 3852
China Lake, California 93555

Dr. Michael J. Weaver
Department of Chemistry
Purdue University
West Lafayette, Indiana 47907

Dr. R. David Rauh
EIC Laboratories, Inc.
Norwood, Massachusetts 02062

Dr. Aaron Wold
Department of Chemistry
Brown University
Providence, Rhode Island 02192

Dr. Martin Fleischmann
Department of Chemistry
University of Southampton
Southampton SO9 5NH UNITED KINGDOM

Dr. R. A. Osteryoung
Department of Chemistry
State University of New York
Buffalo, New York 14214

Dr. John Wilkes
Air Force Office of Scientific
Research
Bolling AFB
Washington, D.C. 20332

Dr. D. Rolison
Naval Research Laboratory
Code 6171
Washington, D.C. 20375-5000

Dr. D. F. Shriver
Department of Chemistry
Northwestern University
Evanston, Illinois 60201

Dr. Edward M. Eyring
Department of Chemistry
University of Utah
Salt Lake City, UT 84112

Dr. M. M. Nicholson
Electronics Research Center
Rockwell International
3370 Miraloma Avenue
Anaheim, California

ABSTRACTS DISTRIBUTION LIST, 359/627

Dr. Hector D. Abruna
Department of Chemistry
Cornell University
Ithaca, New York 14853

Dr. A. B. P. Lever
Chemistry Department
York University
Downsview, Ontario M3J1P3

Dr. Stanislaw Szpak
Naval Ocean Systems Center
Code 633, Bayside
San Diego, California 95152

Dr. Gregory Farrington
Department of Materials Science
and Engineering
University of Pennsylvania
Philadelphia, Pennsylvania 19104

M. L. Robertson
Manager, Electrochemical
and Power Sources Division
Naval Weapons Support Center
Crane, Indiana 47522

Dr. T. Marks
Department of Chemistry
Northwestern University
Evanston, Illinois 60201

Dr. Micha Tomkiewicz
Department of Physics
Brooklyn College
Brooklyn, New York 11210

Dr. Lesser Blum
Department of Physics
University of Puerto Rico
Rio Piedras, Puerto Rico 00931

Dr. Joseph Gordon, II
IBM Corporation
5600 Cottle Road
San Jose, California 95193

Dr. Nathan Lewis
Department of Chemistry
Stanford University
Stanford, California 94305

Dr. D. H. Whitmore
Department of Materials Science
Northwestern University
Evanston, Illinois 60201

Dr. Alan Bewick
Department of Chemistry
The University of Southampton
Southampton, SO9 5NH UNITED KINGDOM

Dr. E. Anderson
NAVSEA-56233 NC #4
541 Jefferson Davis Highway
Arlington, VA

Dr. Bruce Dunn
Department of Engineering &
Applied Science
University of California
Los Angeles, California 90024

Dr. Elton Cairns
Energy & Environment Division
Lawrence Berkeley Laboratory
University of California
Berkeley, California 94720

Dr. Richard Pollard
Department of Chemical Engineering
University of Houston
Houston, Texas 77004

Dr. M. Philpott
IBM Corporation
5600 Cottle Road
San Jose, California 95193

Dr. Donald Sandstrom
Boeing Aerospace Co.
P.O. Box 3999
Seattle, Washington 98124

Dr. Carl Kannewurf
Department of Electrical Engineering
and Computer Science
Northwestern University
Evanston, Illinois 60201

Dr. Joel Harris
Department of Chemistry
University of Utah
Salt Lake City, Utah 84112

ABSTRACTS DISTRIBUTION LIST, 359/627

Dr. Robert Somoano
Jet Propulsion Laboratory
California Institute of Technology
Pasadena, California 91103

Dr. Johann A. Joebstl
USA Mobility Equipment R&D Command
DRDME-EC
Fort Belvoir, Virginia 22060

Dr. Judith H. Ambrus
NASA Headquarters
M.S. RTS-6
Washington, D.C. 20546

Dr. Albert R. Landgrebe
U.S. Department of Energy
M.S. 6B025 Forrestal Building
Washington, D.C. 20595

Dr. J. J. Brophy
Department of Physics
University of Utah
Salt Lake City, Utah 84112

Dr. Charles Martin
Department of Chemistry
Texas A&M University
College Station, Texas 77843

Dr. H. Tachikawa
Department of Chemistry
Jackson State University
Jackson, Mississippi 39217

Dr. Farrell Lytle
Boeing Engineering and
Construction Engineers
P.O. Box 3707
Seattle, Washington 98124

Dr. Robert Gotscholl
U.S. Department of Energy
MS G-226
Washington, D.C. 20545

Dr. Edward Fletcher
Department of Mechanical Engineering
University of Minnesota
Minneapolis, Minnesota 55455

Dr. John Fontanella
Department of Physics
U.S. Naval Academy
Annapolis, Maryland 21402

Dr. Martha Greenblatt
Department of Chemistry
Rutgers University
New Brunswick, New Jersey 08903

Dr. John Wasson
Syntheco, Inc.
Rte 6 - Industrial Pike Road
Gastonia, North Carolina 28052

Dr. Walter Roth
Department of Physics
State University of New York
Albany, New York 12222

Dr. Anthony Sammells
Eltron Research Inc.
4260 Westbrook Drive, Suite 111
Aurora, Illinois 60505

Dr. C. A. Angell
Department of Chemistry
Purdue University
West Lafayette, Indiana 47907

Dr. Thomas Davis
Polymer Science and Standards
Division
National Bureau of Standards
Washington, D.C. 20234

ABSTRACTS DISTRIBUTION LIST, 359/627

Dr. John Owen
Department of Chemistry and
Applied Chemistry
University of Salford
Salford M5 4WT ENGLAND

Dr. J. O. Thomas
University of Uppsala
Institute of Chemistry
Box 531
S-751 21 Uppsala, Sweden

Dr. Boone Owens
Department of Chemical Engineering
and Materials Science
University of Minnesota
Minneapolis, Minnesota 55455

Dr. O. Stafsudd
Department of Electrical Engineering
University of California
Los Angeles, California 90024

Dr. Menahem Anderman
W.R. Grace & Co.
Columbia, MD 20144

END

12-86

DTIC