

AD-A189 407

BIAS REDUCTION WHEN THERE IS NO UNBIASED ESTIMATE(U)  
FLORIDA STATE UNIV TALLAHASSEE DEPT OF STATISTICS  
H DOSS ET AL. JAN 88 FSU-TR-M777 AFOSR-TR-87-219

1/1


UNCLASSIFIED

DAAL83-86-K-0094

F/G 12/3

NL





UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

2

REPORT DOCUMENTATION PAGE

READ INSTRUCTIONS BEFORE COMPLETING FORM

1. REPORT NUMBER

ARO 23699.17-MA

2. GOVT ACCESSION NO.

N/A

3. RECIPIENT'S CATALOG NUMBER

N/A

DTIC FILE COPY

4. TITLE (and Subtitle)

Bias Reduction When There is No Unbiased Estimate

5. TYPE OF REPORT & PERIOD COVERED

Technical Report

6. PERFORMING ORG. REPORT NUMBER

FSU Technical Report M-777

8. CONTRACT OR GRANT NUMBER(s)

DAAL03-86-K-0094

7. AUTHOR(s)

Hani Doss and Jayaram Sethuraman

9. PERFORMING ORGANIZATION NAME AND ADDRESS

Florida State University  
Department of Statistics  
Tallahassee, FL 32306-3033

10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS

11. CONTROLLING OFFICE NAME AND ADDRESS

U.S. Army Research Office  
Post Office Box 12211  
Research Triangle Park, NC 27709

12. REPORT DATE

13. NUMBER OF PAGES

4

15. SECURITY CLASS. (of this report)

UNCLASSIFIED

15a. DECLASSIFICATION/DOWNGRADING SCHEDULE

16. DISTRIBUTION STATEMENT (of this Report)

for public release; distribution unlimited

DTIC ELECTED  
FEB 02 1988  
S E D

17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)

N/A

18. SUPPLEMENTARY NOTES

19. KEY WORDS (Continue on reverse side if necessary and identify by block number)

bias reduction; jackknife estimate of bias; bootstrap estimate of bias.

20. ABSTRACT (Continue on reverse side if necessary and identify by block number)

Let  $\phi$  be a parameter for which there is no unbiased estimator. This note shows that for an arbitrary sequence of estimators  $T^{(k)}$ , if the biases of  $T^{(k)}$  tend to 0 then their variances must tend to  $\infty$ .

AD-A189 407

## BIAS REDUCTION WHEN THERE IS NO UNBIASED ESTIMATE

by

Hani Doss\*

and

Jayaram Sethuraman†

Department of Statistics  
 Florida State University  
 Tallahassee, FL 32306

FSU Technical Report No. M-777  
 AFOSR Technical Report No. 87-219  
 USARO Technical Report No. D-101



January, 1988

Accession For	
NTIS GRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By _____	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-1	

\* Research supported by the Air Force Office of Scientific Research Grant Number F49620-85-C-0007.

† Research supported by the U. S. Army Research office under Grant Number DAAL03-86-k-0094.

Key words and phrases: bias reduction, jackknife estimate of bias, bootstrap estimate of bias.

AMS 1980 subject classifications. Primary 62F11; secondary 62A99.

# BIAS REDUCTION WHEN THERE IS NO UNBIASED ESTIMATE

by

Hani Doss and Jayaram Sethuraman

Florida State University

## ABSTRACT

Let  $\phi$  be a parameter for which there is no unbiased estimator. This note shows that for an arbitrary sequence of estimators  $T^{(k)}$ , if the biases of  $T^{(k)}$  tend to 0 then their variances must tend to  $\infty$ .

## 1. INTRODUCTION.

Let  $X = (X_1, \dots, X_n)$  have distribution  $P_\theta$ , where the unknown parameter varies in  $\Theta$ . Suppose that we need to estimate a real valued function  $\phi(\theta)$  of the parameter. Let  $\hat{\phi} = \hat{\phi}(X)$  be a biased estimator of  $\phi$ . There exist several procedures for reducing the bias of  $\hat{\phi}$ : jackknifing, bootstrapping (see Efron (1982)), and other procedures based on expansions of  $E_\theta(\hat{\phi})$  (see Cox and Hinkley (1974, Section 8.4)). These procedures may not eliminate the bias completely, and one often hears the following suggestion. Let  $\hat{\phi}^{(1)}$  be obtained from  $\hat{\phi}$  by one of these bias-reduction procedures. If  $\hat{\phi}^{(1)}$  is still biased, repeat the bias-reduction procedure and obtain  $\hat{\phi}^{(2)}, \hat{\phi}^{(3)}$  etc. until a desired amount of reduction in bias is obtained or the bias is removed completely. Such "higher-order bias corrections" are described for instance in the review paper of Miller (1974) in connection with the jackknife. There are examples where no unbiased estimator of  $\phi$  exists but there exists a sequence of estimators  $\hat{\phi}, \hat{\phi}^{(1)}, \hat{\phi}^{(2)}, \dots$  whose biases converge to zero (see Section 2).

The purpose of this note is to show (Theorem 1) that when no unbiased estimator of  $\phi$  exists, then reducing the bias to zero necessarily forces the variance of the estimators to tend to  $\infty$ . This theorem therefore gives qualitative support to the widely held view that bias reduction is by itself not a desirable property, but becomes desirable only if it can be demonstrated that it is accompanied by a reduction in mean squared error.

## 2. MAIN RESULT AND REMARKS.

Let  $(X, S)$  be a measurable space and  $(P_\theta, \theta \in \Theta)$  be a family of probability measures on  $(X, S)$ . Let  $\phi$  be a real valued function defined on  $\Theta$ . The bias of an estimator  $T = T(X)$  is defined by  $\beta_T(\theta) = E_\theta(T(X)) - \phi(\theta)$ , assuming that  $E_\theta(T(X))$  exists.

**THEOREM 1.** Suppose that

$$(A1) \quad P_{\theta_1} \ll P_{\theta_2} \text{ for all } \theta_1, \theta_2 \text{ in } \Theta,$$

$$(A2) \quad \int \left( \frac{dP_{\theta_1}}{dP_{\theta_2}} \right)^2 dP_{\theta_2} < \infty \text{ for all } \theta_1, \theta_2 \text{ in } \Theta,$$

and that  $\{T_k\}_{k=1}^\infty$  is a sequence of estimators for which

$$(1) \quad \beta_{T_k}(\theta) \rightarrow 0 \text{ for all } \theta \text{ in } \Theta.$$

If there does not exist an unbiased estimator of  $\phi$  then

$$(2) \quad \text{Var}_\theta(T_k) \rightarrow \infty \text{ as } k \rightarrow \infty, \text{ for all } \theta \in \Theta.$$

Proof: Suppose that (2) is not true. Then there exists a  $\theta_0$  in  $\Theta$  and a subsequence  $\{k^*\}$  of  $\{k\}$  such that  $\text{Var}_{\theta_0}(T_{k^*})$  is bounded. Now, consider the usual Hilbert space  $H_{\theta_0} = L^2(\mathcal{X}, \mathcal{S}, P_{\theta_0})$  of all functions that are square-integrable with respect to  $P_{\theta_0}$ . Notice that  $\{T_{k^*}\}$  is a norm-bounded set in  $H_{\theta_0}$ . From the sequential weak-compactness of norm-bounded sets, there exists a  $T$  in  $H_{\theta_0}$  and a subsequence  $\{k^{**}\}$  of  $\{k^*\}$  such that  $T_{k^{**}} \rightarrow T$  weakly in  $H_{\theta_0}$  along the subsequence  $\{k^{**}\}$ , i.e.

$$\int T_{k^{**}} f dP_{\theta_0} \rightarrow \int T f dP_{\theta_0} \text{ for every function } f \text{ in } H_{\theta_0}.$$

In particular, setting  $f = dP_{\theta}/dP_{\theta_0}$ , we get

$$E_{\theta}(T_{k^{**}}) \rightarrow E_{\theta}(T),$$

along the subsequence  $\{k^{**}\}$ , for all  $\theta$  in  $\Theta$ . From (1), it now follows that  $E_{\theta}(T) = \phi(\theta)$ , that is  $T$  is unbiased for  $\phi$ , which contradicts one of our assumptions. Hence (2) holds and the proof is complete. ■

There are many examples of situations to which this theorem applies. One class can be obtained from the idea of the following example. Consider the family of Poisson distributions with parameter  $\lambda$  with  $\lambda > 0$ . It is well known that there exists no unbiased estimator of  $1/\lambda$ , and that all polynomials in  $\lambda$  are unbiasedly estimable. From (a slight modification of) the Stone-Weirstrass theorem, there exists a sequence of polynomials in  $\lambda$  which converge to  $1/\lambda$  for each  $\lambda$ . Thus there exists a sequence of estimators which are unbiased for these polynomials in  $\lambda$ , and whose biases in estimating  $1/\lambda$  converge to zero. A simple calculation shows that  $\int \left(\frac{dP_{\lambda_1}}{dP_{\lambda_2}}\right)^2 dP_{\lambda_1} = \exp(\lambda_2 - 2\lambda_1 + \lambda_1^2/\lambda_2)$ . Thus Theorem 1 applies to this case and the variances of these estimators must tend to  $\infty$ .

It may appear that Theorem 1 does not apply to estimates based on the jackknife, since the "delete-one" jackknife can be formed only a finite number of times. However, a situation with an infinite sequence of estimators based on the jackknife arises in the following example, based on an idea of Gaver and Hoel (1970). Suppose that the data consists of a Poisson process  $\{N(t); t \in [0, 1]\}$  with rate  $\lambda$ . In connection with the biased maximum likelihood estimator  $\hat{\phi} = e^{-\lambda N(1)}$  of  $e^{-\lambda}$ , Gaver and Hoel suggest splitting the interval  $[0, 1]$  into  $n$  nonoverlapping intervals each of length  $1/n$ , and letting  $N_i$  be the number of events in the  $i$ th interval. These are independent and identically distributed and one can therefore form the delete-one jackknife as usual. This yields, for each  $n$ , an estimate  $\hat{\phi}_{(n)}$  and they show that as  $n \rightarrow \infty$   $\hat{\phi}_{(n)}$  converges to an estimate  $\hat{\phi}^{(1)}$

which depends on the Poisson process only through the sufficient statistic  $N(1)$ . This procedure can be repeated indefinitely in principle, giving a sequence of estimators  $\{\hat{\phi}^{(k)}\}_{k=1}^{\infty}$ .

## REFERENCES

- Cox, D. R. and Hinkley, D. V. (1974). *Theoretical Statistics*. Chapman and Hall, London.
- Efron, B. (1982). *The Jackknife, the Bootstrap and Other Resampling Plans*. SIAM, Philadelphia.
- Miller, R. G. (1974). The jackknife - a review : *Biometrika* **61**, 1-15.
- Gaver, D. P., Jr. and Hoel, D. G. (1970). Comparison of certain small-sample Poisson probability estimates. *Technometrics* **12**, 835-850.

## REPORT DOCUMENTATION PAGE

1a. REPORT SECURITY CLASSIFICATION UNCLASSIFIED		1b. RESTRICTIVE MARKINGS	
2a. SECURITY CLASSIFICATION AUTHORITY NA		3. DISTRIBUTION/AVAILABILITY OF REPORT Approved for Public Release; Distribution Unlimited	
2b. DECLASSIFICATION/DOWNGRADING SCHEDULE NA			
4. PERFORMING ORGANIZATION REPORT NUMBER(S) FSU Technical Report No. M-777		5. MONITORING ORGANIZATION REPORT NUMBER(S) AFOSR Technical Report No. 87-219	
6a. NAME OF PERFORMING ORGANIZATION Florida State University	6b. OFFICE SYMBOL (If applicable)	7a. NAME OF MONITORING ORGANIZATION AFOSR/NM	
6c. ADDRESS (City, State and ZIP Code) Department of Statistics Tallahassee, FL 32306-3033		7b. ADDRESS (City, State and ZIP Code) Bldg. 410 Bolling AFB, DC 20332-6448	
8a. NAME OF FUNDING/SPONSORING ORGANIZATION AFOSR	8b. OFFICE SYMBOL (If applicable) NM	9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER	
8c. ADDRESS (City, State and ZIP Code) Bldg. 410 Bolling AFB, DC 20332-6448		10. SOURCE OF FUNDING NOS.	
		PROGRAM ELEMENT NO.	PROJECT NO.
		TASK NO.	WORK UNIT NO.
11. TITLE (Include Security Classification) Bias Reduction When There is No Unbiased Estimate			
12. PERSONAL AUTHOR(S) Hani Doss and Jayaram Sethuraman			
13a. TYPE OF REPORT Technical	13b. TIME COVERED FROM _____ TO _____	14. DATE OF REPORT (Yr., Mo., Day)	15. PAGE COUNT 4
16. SUPPLEMENTARY NOTATION			
17. COSATI CODES		18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)	
FIELD	GROUP	SUB. GR.	bias reduction, jackknife estimate of bias, bootstrap estimate of bias.
19. ABSTRACT (Continue on reverse if necessary and identify by block number)			
<p>Let <math>\phi</math> be a parameter for which there is no unbiased estimator. This note shows that for an arbitrary sequence of estimators <math>T^{(k)}</math>, if the biases of <math>T^{(k)}</math> tend to 0 then their variances must tend to <math>\infty</math>.</p>			
20. DISTRIBUTION/AVAILABILITY OF ABSTRACT UNCLASSIFIED/UNLIMITED <input checked="" type="checkbox"/> SAME AS RPT. <input type="checkbox"/> DTIC USERS <input type="checkbox"/>		21. ABSTRACT SECURITY CLASSIFICATION	
22a. NAME OF RESPONSIBLE INDIVIDUAL Frank Proshan	22b. TELEPHONE NUMBER (Include Area Code) (904) 644-1832	22c. OFFICE SYMBOL AFOSR/NM	

END

DATE

FILMED

MARCH

1988

DTIC