

AD-A193 111

A SPECTROSCOPY OF HELIX MELTING(U) PURDUE RESEARCH
FOUNDATION LAFAYETTE IN E W PRONOFKY 11 MAR 87
N00014-86-K-0252


1/1

UNCLASSIFIED

F/G 6/7

NL





SECURITY CLASSIFICATION OF THIS PAGE

DTIC REPORT DOCUMENTATION PAGE

AD-A193 111

DTIC
D

1. RESTRICTIVE MARKINGS NA	
2. DISTRIBUTION AVAILABILITY OF REPORT DISTRIBUTION UNLIMITED	
3. PERFORMING ORGANIZATION REPORT NUMBER Purdue Research Foundation	4. MONITORING ORGANIZATION REPORT NUMBER NA
5a. NAME OF PERFORMING ORGANIZATION Purdue Research Foundation	5b. OFFICE SYMBOL (if applicable) NA
6a. ADDRESS (City, State, and ZIP Code) W. Lafayette, IN 47907	6b. ADDRESS (City, State, and ZIP Code) OSD/SDIO/TA Washington, D.C. 20301-7100
7a. NAME OF FUNDING SPONSORING AGENCY Strategic Defense Initiative Organization	7b. OFFICE SYMBOL (if applicable)
8a. ADDRESS (City, State, and ZIP Code) Washington, D.C. 20301-7100	8b. ADDRESS (City, State, and ZIP Code)
9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER N00014-86-K-0252	
10. SOURCE OF FUNDING NUMBERS	
PROGRAM ELEMENT NO 63222 c	PROJECT NO
TASK NO	WORK UNIT ACCESSION NO

11. TITLE (and De Security Classification)
A Spectroscopy of Helix Melting

12. PERSONAL AUTHOR(S)
Prohofsky, Earl William

13a. TYPE OF REPORT
Annual and Progress

13b. TIME COVERED
FROM 4/87 TO 3/88

14. DATE OF REPORT (Year, Month, Day)
1987-3-11

15. PAGE COUNT
3

16. SUPPLEMENTARY NOTES

17. UNCLASSIFIED

FIELD	GROUP	SUBGROUP
-------	-------	----------

18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)
DNA>HELIX MELTING, Deoxyribonucleic Acid, FREE ELECTRON LASER IRRADIATION

19. ABSTRACT (Continue on reverse if necessary and identify by block number)
We have calculated the far infrared absorption for several models of double helical DNA. These results are in good agreement with experimental observations. We have also fitted our results to observed inelastic neutron scattering data. Again the results are in excellent agreement with observation. We have extended our calculations on helix strand melting to include methylated DNA and many other DNA's of varying base pair sequence. All of the results are useful in leading to more precise predictions of the dynamics of DNA strand separation in a far infrared radiation field. We have also extended our model so as to predict energy flow in the natural process of RNA transcription.

DISTRIBUTION STATEMENT A
Approved for public release;
Distribution Unlimited

20. DISTRIBUTION STATEMENT (See instructions on reverse side)
U

21. DISTRIBUTION STATEMENT (See instructions on reverse side)
OSD/SDIO/TA

Charles L. Houston, III, Captain USAF (202)-693-1556

DATE: 10 June 1988

PROGRESS AND ANNUAL REPORT ON CONTRACT N00014-86-K-0252

PRINCIPAL INVESTIGATOR: E. W. Prohofsky

CONTRACTOR: Purdue Research Foundation

CONTRACT TITLE: A Spectroscopy of Helix Melting

START DATE: 1 April 1986

RESEARCH OBJECTIVE: To investigate the role of far infrared excitations that can be excited by FEL radiation, in the melting of DNA: to develop an understanding of the role of such excitations in transcription and replication.

PROGRESS (Year 2): We have made considerable progress toward a calculation of the amount of energy in a free electron laser beam that would actually be absorbed by DNA. The first step required a careful fit of our calculated spectrum to infrared absorption observations. This has been completed and the fit is detailed in the manuscript "Calculation of Far Infrared Absorption in B poly(dA)·poly(dT) which is enclosed as a technical report. We find that the actual absorption strength does depend on the state of aggregation of the DNA i.e., there are strong interhelical effects. More sophisticated calculations are needed to accurately model the true absorption and we are proceeding with such calculations.

We have been able to fit our basic vibrational mode model to newly published inelastic neutron scattering observations. We find that our model fits the observations very well. We have used the information obtained to improve our model by allowing us to separate charge effects from dielectric constant effects in a more complete manner. The reason for this is the inelastic neutron data is at high enough frequencies to be above the large rapidly changing dielectric response of water. I enclose two manuscripts on this subject as technical reports.

We have made progress on investigating the role of base pair sequence in DNA melting. We have also studied sequence dependant response to far infrared absorption. We have calculated the melting behavior differences for methylated vs. unmethylated DNA and we have calculated the melting behavior for Z conformation DNA. For the methylation case we predict a 7K increase in melting temperature for methylated Z-poly(dGC)·poly(dGC) compared to unmethylated B poly(dGC)·poly(dGC). The experimentally observed increase is 6K. The work on this melting variation is contained in three manuscripts which are enclosed as technical reports.

We have also explored the nonlinearity of the far infrared modes involved in melting and FEL absorption. We found that these modes are likely to display solitary wave narrowing which will be important in more detailed analysis of the effects of FEL irradiation. This work is detailed in a manuscript enclosed as a technical report.

88 4 8 020

WORK PLAN (Year 3): We plan to continue along the lines of progress for the first two years. We will explore further the effect of state of aggregation on the actual far infrared absorption. When this is understood we can predict quantitatively the results of FEL irradiation on DNA. We plan to study absorption for helices in a crystal interacting with other helices. We plan to complete our study of base pair sequence effects on melting and far infrared absorption. Much of this effort has been completed but a manuscript of the comparison across the board will have to wait until we have a better handle on the melting of A-T base pair systems. We have also begun calculations on the transition from small amplitude normal mode behavior to large amplitude solitary wave behavior for these far infrared modes. We will continue such calculations in year three.

All of the efforts listed are key elements in the overall plan to study the absorption of FEL radiation and to predict the dynamic effects of that absorbed radiation. All of the individual elements in that program are progressing as outlined above. The task remaining is to complete the individual calculations and integrate the pieces to the analysis and prediction of the dynamical consequences of FEL radiation.

PUBLICATIONS AND REPORTS (Year 2):

1. L. Young and E. W. Prohofsky, "Calculation of Far Infrared Absorption in B poly(dA)·poly(dT) submitted for publication.
2. V. V. Prabhu, W. K. Schroll, L. L. Van Zandt, and E. W. Prohofsky. "Helical Lattice Vibrational Modes in DNA", to be published in Physical Review Letters.
3. W. K. Schroll, V. V. Prabhu, E. W. Prohofsky and L. L. Van Zandt, "Phonon Interpretation of Inelastic Neutron Scattering in DNA Crystals", submitted for publication.
4. X. M. Hua and E. W. Prohofsky "Vibrational Spectra of Double Helical Molecules with C₂ Symmetry," submitted for publication.
5. X. M. Hua and E. W. Prohofsky, "Normal Mode Calculation for Methylated Z-DNA poly(dG - m⁵dC). (dG - m⁵dC)", submitted for publication.
6. X. M. Hua, Y. Feng and E. W. Prohofsky, "Calculated Melting Temperature of Methylated Z-DNA", submitted for publication.
7. E. W. Prohofsky "Where Solitons May be Hiding in DNA and Their Possible Significance in RNA Transcription", submitted for publication

TRAINING ACTIVITIES: Two postdoctoral persons and five graduate students have had some level of support from this contract.

Women or minorities - 2

Announced		<input checked="" type="checkbox"/>
Justification		<input type="checkbox"/>
By _____		
Distribution/		
Availability Codes		
Dist	Avail and/or Special	
A-1		NSF

END

DATE

FILMED

DTIC

July 88