

AD-A198 719



## Comparison of Three Anesthetics for Chinchilla

By

C.E. Hargett, Jr.\*  
Illa M. Lomba Gautler  
Melvin Carrier, Jr.  
Carol S. Landon  
Irvin W. McConnell\*\*  
James H. Patterson, Jr.

Sensory Research Division

April 1988



- \* Research Foundation, State University of New York at Plattsburgh
- \*\* Schering Research, Safety Evaluation Center, Lafayette, New Jersey

Approved for public release; distribution unlimited.

United States Army Aeromedical Research Laboratory  
Fort Rucker, Alabama 36362-5292

88

REPORT DOCUMENTATION PAGE

1a. REPORT SECURITY CLASSIFICATION <b>Unclassified</b>		1b. RESTRICTIVE MARKINGS	
2a. SECURITY CLASSIFICATION AUTHORITY		3. DISTRIBUTION / AVAILABILITY OF REPORT Approved for public release; distribution unlimited	
2b. DECLASSIFICATION / DOWNGRADING SCHEDULE		4. PERFORMING ORGANIZATION REPORT NUMBER(S) USAARL Report No. 88-4	
4. PERFORMING ORGANIZATION REPORT NUMBER(S) USAARL Report No. 88-4		5. MONITORING ORGANIZATION REPORT NUMBER(S)	
6a. NAME OF PERFORMING ORGANIZATION U.S. Army Aeromedical Research Laboratory	6b. OFFICE SYMBOL (If applicable) SGRD-UAS-AS	7a. NAME OF MONITORING ORGANIZATION U.S. Army Medical Research and Development Command	
6c. ADDRESS (City, State, and ZIP Code) Fort Rucker, AL 36362-5292		7b. ADDRESS (City, State, and ZIP Code) Fort Detrick Frederick, MD 21701-5012	
8a. NAME OF FUNDING / SPONSORING ORGANIZATION	8b. OFFICE SYMBOL (If applicable)	9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER	
8c. ADDRESS (City, State, and ZIP Code)		10. SOURCE OF FUNDING NUMBERS	
		PROGRAM ELEMENT NO. 61102A	PROJECT NO. 3M1611 02BS15
		TASK NO. CB	WORK UNIT ACCESSION NO. 282
11. TITLE (Include Security Classification) Comparison of Three Anesthetics for Chinchilla (U)			
12. PERSONAL AUTHOR(S) C.E. Hargett, Jr., Iffa M. Lomba-Gautier, Melvin Carrier, Jr., Carol S. Landon, Irving W. McConnell, James H. Patterson, Jr.			
13a. TYPE OF REPORT Final	13b. TIME COVERED FROM _____ TO _____	14. DATE OF REPORT (Year, Month, Day) 1988 April	15. PAGE COUNT 19
16. SUPPLEMENTARY NOTATION			
17. COSATI CODES		18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)	
FIELD	GROUP	Chinchilla	
20	01	Acepromazine	
24	07	Anesthesia; (K)	
		Ketamine	
		Xylazine	
19. ABSTRACT (Continue on reverse if necessary and identify by block number) Anesthesia techniques which successfully induce surgical anesthesia in Chinchilla villidera are described and compared. Two injectables, ketamine-acepromazine and ketamine-xylazine, are compared to halothane-nitrous oxide administered by mask only and the same mixture administered by induction chamber until loss of righting reflex and then mask. Forty laboratory-raised adult chinchillas in four groups were used in this study. Subjects were weighed and dosages calculated and administered. All achieved surgical anesthesia with no deaths. Time to loss of righting reflex, time to surgical anesthesia, duration of surgical anesthesia, time from end of surgical anesthesia to standing unaided, and total time from administration of anesthesia to standing unaided are detailed. Findings indicate that the doses evaluated of ketamine-acepromazine, ketamine-xylazine, and halothane-nitrous oxide all provide safe, dependable surgical anesthesia.			
20. DISTRIBUTION / AVAILABILITY OF ABSTRACT <input checked="" type="checkbox"/> UNCLASSIFIED/UNLIMITED <input type="checkbox"/> SAME AS RPT. <input type="checkbox"/> DTIC USERS		21. ABSTRACT SECURITY CLASSIFICATION Unclassified	
22a. NAME OF RESPONSIBLE INDIVIDUAL Chief, Scientific Information Center		22b. TELEPHONE (Include Area Code) (205) 255-6907	22c. OFFICE SYMBOL SGRD-UAX-SI

Notice

Qualified requesters

Qualified requesters may obtain copies from the Defense Technical Information Center (DTIC), Cameron Station, Alexandria, Virginia 22314. Orders will be expedited if placed through the librarian or other person designated to request documents from DTIC.

Change of address

Organizations receiving reports from the US Army Aeromedical Research Laboratory on automatic mailing lists should confirm correct address when corresponding about laboratory reports.

Animal use

In conducting the research described in this report, the investigators adhered to the Guide for Laboratory Animal Facilities and Care, as promulgated by the Committee on the Guide for Laboratory Animal Resources, National Academy of Sciences-National Research Council.


Disposition

Destroy this report when it is no longer needed. Do not return it to the originator.

Disclaimer


The views, opinions, and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy, or decision, unless so designated by other official documentation. Citation of trade names in this report does not constitute an official Department of the Army endorsement or approval of the use of such commercial items.

Reviewed:



BRUCE C. LEIBRECHT, Ph.D.  
LTC, MS  
Director, Sensory Research  
Division

Released for publication:



J. D. LaMOTHE, Ph.D.  
COL, MS  
Chairman, Scientific  
Review Committee



DUDLEY R. PRICE  
Colonel, MC  
Commanding

### Acknowledgments

This work was supported by the U.S. Army Medical Research and Development Command (USAMRDC) under Contract Number DAMD17-86-C-6139.

C.E. Hargett, Jr., an employee of the Research Foundation, State University of New York at Plattsburgh, is assigned full-time at the U.S. Army Aeromedical Research Laboratory (USAARL) as part of the above contract.

Irving W. McConnell, at the time of this research, was a Veterinary Corps officer in the U.S. Army. Dr. McConnell now is at Schering Research, Safety Evaluation Center, Lafayette, NJ.

The views of the authors do not purport to reflect the positions of the Department of the Army or the Department of Defense.

In conducting research using animals, the investigators adhered to the "Guide for the Care and Use of Laboratory Animals," prepared by the Committee on Care and Use of Laboratory Animals of the Institute of Laboratory Animal Resources, National Research Council (NIH Publication 86-23, revised 1985).

The authors wish to acknowledge and thank Ms. Linda S. Barlow for her valuable statistical assistance.

Accession For	
NTIS GRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By _____	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-1	



## Table of contents

	Page no.
Introduction .....	3
Methods and procedures .....	4
Results and discussion .....	7
Conclusion .....	9
References .....	11
Appendix	
A. List of manufacturers .....	12

## List of tables

Table no.		Page no.
1	Time to loss of righting reflex in seconds .....	7
2	Time to surgical anesthesia in seconds .....	7
3	Duration of surgical anesthesia in minutes .....	8
4	Minutes from end of surgical anesthesia to standing unaided .....	9
5	Time from administration of anesthesia to standing unaided .....	9

=====  
This page intentionally left blank  
=====

## Introduction

The chinchilla (*Chinchilla villidera*) is the animal model of choice for certain types of auditory research (Miller, 1970). The Acoustical Sciences Branch, Sensory Research Division of the U.S. Army Aeromedical Research Laboratory (USAARL), Fort Rucker, Alabama, has used the chinchilla as a research animal model since 1977 (Burdick, et al., 1978).

In auditory research, it is necessary to perform surgical procedures such as monauralizations and electrode implants as part of acoustical experimentation. This requires an acceptable method of anesthetizing the animals. Relatively few studies have been reported documenting the use of anesthetics in the chinchilla. Intramuscular injections of ketamine-acepromazine have been used successfully by some research teams (Morgan, et al., 1981). Ketamine is a nonbarbiturate that rapidly produces a cataleptic state of anesthesia characterized as disassociate anesthesia. Acepromazine has a depressant effect on the central nervous system. While anesthesia is rapid with this combination, recovery tends to be slow. In one study, the duration of surgical anesthesia exceeded 40 minutes and required 120 to 300 minutes for complete recovery (Morgan, et al., 1981).

There were no references in the literature that we found to ketamine-xylazine combinations used to anesthetize chinchillas. CI-744 (tiletamine-HCL and zolezapan-HCL) was reported to be effective for chinchilla anesthesia with a mean duration of 202 minutes for five animals given the minimum dosage (22 mg/kg) that provides surgical anesthesia (Schultz and Fowler, 1974). Xylazine is an alternative to acepromazine.

Experience at USAARL has shown that 20 to 25 minutes of surgical anesthesia is sufficient for most procedures. For this reason, a mixture of halothane and nitrous oxide has been used to anesthetize our chinchillas for surgery. Halothane is a halogenated hydrocarbon that produces a potent nonexplosive anesthesia agent (Deutsch, 1978). Inhalation anesthesia has a number of advantages. The agents are eliminated primarily through the lungs, so recovery from anesthesia does not rely upon redistribution within the body and detoxification mechanisms (Lumb and Jones, 1973). Halothane is a "complete anesthetic" capable of producing surgical anesthesia without oxygen deprivation and is nonirritating to the upper and lower respiratory tracts (Deutsch, 1971). Salivary, mucous, and bronchial secretions, as well as laryngospasms and vomiting, are not problems when using halothane anesthesia. Nitrous oxide is used with halothane to provide rapid induction, analgesia, and to complement the more potent anesthetic agent (Lumb and Jones, 1973; Soma, 1971). Inhalation anesthesia requires specialized equipment, but it is the only technique that allows for control

of anesthetic depth and surgical anesthesia duration. The present study was undertaken to compare halothane anesthesia with ketamine in combination with either acepromazine or xylazine. In addition, two methods for inducing halothane anesthesia were compared.

#### Methods and procedures

This study utilized 40 healthy chinchillas of both sexes from the USAARL colony. Individual stainless steel laboratory cages (483mm x 607mm x 203mm) were used as housing for these chinchillas. They were provided with commercial chinchilla ration\* and water ad libitum. They weighed from 375.2 grams to 643.1 grams with a mean weight of 496.1 grams and a median weight of 487.2 grams. Ages ranged from 10 to 48 months. They were assigned randomly to four groups of 10 subjects each. The chinchillas were not deprived of food or water prior to the experiment and were returned to individual cages upon being able to stand unaided.

Each subject was anesthetized to surgical depth by the method of anesthesia for its assigned group. Surgical depth is defined as not exhibiting pedal reflex, or palpebral reflex, and is corroborated by subjective observations evaluating the chinchilla's overall appearance and condition. On each subject, we collected the following data: respiration rates, time to loss of righting reflex, time to loss of palpebral reflex, time to loss of pedal reflex, time at surgical depth, time to return of pedal reflex, and time to standing unaided. Time to head lift from lateral recumbency and both pinna and whisker reflex were taken on some of the injectable subjects. However, it was not possible to acquire these data on the gas anesthesia subjects as the mask is in place providing oxygen flow at the time these reflexes occur. Time at surgical depth for the inhalation anesthesia subjects was limited to 20 minutes for the purposes of this report. This is not to be taken as an upper time limit for this technique, merely the time we find to be optimal for our surgical techniques.

The four methods used to induce surgical anesthesia were:

Group I: Halothane\* and nitrous oxide administered by the mask only.

Group II: Halothane and nitrous oxide administered by the anesthesia chamber until loss of righting reflex, then by the mask.

---

\*See Appendix A.

Group III: Ketamine\*-acepromazine administered by injection.

Group IV: Ketamine-xylazine\* administered by injection.

All times were taken with a stopwatch and recorded as the nearest minute and second, except duration of surgical depth and time to standing unaided for Groups III and IV, which were taken to the nearest minute.

The long duration of the effects of the injections make time rounded to the nearest whole minute more appropriate. Statistical tests of the equivalence of means across the four groups were accomplished using a one-way analysis of variance followed by a posteriori comparisons using Tukey's honestly significant differences (Winer, 1971). A separate analysis was done for each dependent variable.

#### Group I

Ten chinchillas with a mean weight of 522.6 grams were anesthetized using a mask-only system of inhalation anesthesia. The mask system is a nonrebreathing system. It is well-suited for smaller animals such as a chinchilla. The chinchillas were picked up from the transport cage, placed on the surface of the Narkovet\* anesthesia machine and their heads inserted into the mask. They were held with heads in the mask until loss of righting reflex occurred. A 4 percent setting of halothane, with a flow rate of 4 liters per minute of nitrous oxide and 2 liters per minute of oxygen, was used.

When the palpebral reflex was gone, the chinchilla was removed from the mask and Optivet\* chloramphenocol eye ointment was placed in each eye, and the eyes were closed. This kept the eye tissue from drying out. Then, the subject was returned to the mask. The loss of pedal reflex was taken as beginning of surgical depth. The chinchilla was maintained at surgical depth using a 2 1/2 percent halothane setting, with a flow rate of 2 liters per minute of nitrous oxide and 2 liters per minute of oxygen.

At the end of the procedure, the halothane was set at 0 percent (no flow), the nitrous oxide at zero liters per minute (no flow) and the oxygen maintained at 2 liters per minute. This enabled the chinchilla to return to consciousness rapidly, without complications induced by nitrous oxide (Soma, 1971).

## Group II

Ten chinchillas with a mean weight of 459.6 grams were anesthetized using a precharged Ejay\* international anesthesia chamber. It is made of tough, lightweight, transparent plastic. This chamber is supposed to reduce the stress on the subject prior to being anesthetized. The unit was precharged with halothane-nitrous oxide. Then the chinchilla was inserted and the lid quickly closed. The chamber is transparent, so ataxia could be observed.

Gently shaking the unit would cause the chinchilla to topple over at the onset of loss of righting reflex. At the loss of righting reflex, the chinchilla was removed from the unit, tested for palpebral reflex, eyes filled with Optivet ointment, and placed on the mask that is attached to the Narkovet anesthesia machine. At this point, all further details were the same as for subjects in Group I.

## Group III

Ten chinchillas with a mean weight of 513.6 grams were injected intramuscularly in the left gluteal area with a mixture of ketamine hydrochloride (40 mg/kg) and acepromazine (approximately 0.5 mg/kg) using the technique described by Morgan et al., 1981.

Subjects were weighed and individual doses calculated and drawn up into a 1 cc tuberculin syringe. Upon injection, subjects were placed in a 457 mm x 304 mm x 228 mm transparent plastic box for observation.

At loss of righting reflex, subjects were tested for palpebral reflex and then eyes were filled with Optivet ointment and closed to prevent drying. Then, subjects were placed in a recumbent position for further observations.

## Group IV

Ten chinchillas with a mean weight of 488.7 grams were injected in the left gluteal area with a mixture of ketamine hydrochloride (40 mg/kg) and xylazine (2 mg/kg). After calculating the dosage of ketamine to be injected, this dosage was drawn into a 1 cc tuberculin syringe, then cleared of air bubbles. This syringe was inserted into the xylazine vial and the calculated dosage of xylazine drawn on top of the ketamine. Next, air was drawn into the syringe and the two agents well blended. All the air was expelled and then the blend of ketamine-xylazine was injected. Upon injection, subjects were placed in a 457mm x 304 mm x 228 mm transparent plastic box for observation.

## Results and Discussion

The times to loss of righting reflex are shown in Table 1. The analysis of variance revealed significant differences ( $\alpha = .05$ ) in mean time to loss of righting reflex among the four groups. However, a posteriori analysis failed to find any differences between individual groups.

Table 1

### Time to loss of righting reflex in seconds

	<u>Group I</u>	<u>Group II</u>	<u>Group III</u>	<u>Group IV</u>
Median	128.5	117.5	87.5	85.0
Range	(46-170)	(60-205)	(72-135)	(61-124)
Average	124.1	129.4	94.9	86.8
S.D.	34.3	54.9	22.6	20.5

Group I = Mask only  
 Group II = Induction chamber, then mask  
 Group III = Ketamine-acepromazine  
 Group IV = Ketamine-xylazine

Table 2 shows the time to surgical anesthesia in seconds. The time to surgical anesthesia was shortest for the ketamine-xylazine combination; both gas techniques required about the same time. The ketamine-acepromazine combination yielded the longest time to surgical depth. The slowest subject to reach surgical anesthesia (no pedal reflex, no palpebral reflex) in this group took 723 seconds (just over 12 minutes). The analysis of variance showed a significant difference ( $\alpha = .05$ ) between groups. The a posteriori analysis revealed significant differences between Group IV and both Groups I and II, and between Group III and the other groups. No significant difference existed between Groups I and II.

Table 2

### Time to surgical anesthesia in seconds

	<u>Group I</u>	<u>Group II</u>	<u>Group III</u>	<u>Group IV</u>
Median	332.5	298.0	455.5	203.0
Range	(150-402)	(230-445)	(324-723)	(121-311)
Average	308.2	298.8	481.8	197.7
S.D.	72.7	67.6	123.6	453.4

Group I = Mask only  
 Group II = Induction chamber, then mask  
 Group III = Ketamine-acepromazine  
 Group IV = Ketamine-xylazine

The durations of surgical anesthesia are shown in Table 3. The time at surgical anesthesia can best be controlled by use of a gas anesthesia machine. If using injectables, once administered, the rate at which the agent is metabolized is the major factor. Although additional doses can be given to increase time at surgical depth, this method is not as readily controlled as with the gas machine. The average time for Group III (ketamine-acepromazine) to be at surgical anesthesia was 54.4 minutes while for Group IV (ketamine-xylazine) it was 121.8 minutes. Both Groups I and II (gas techniques) began to recover fairly rapidly as soon as there was no flow of both halothane and nitrous oxide and oxygen remained at 2 liters per minute. The analysis of variance showed a significant difference between groups. Significant differences between Group IV and Groups I and II, and Group III were found with the a posteriori analysis. The difference between Groups I and II was not significant.

Table 3

Duration of surgical anesthesia in minutes

	<u>Group I</u>	<u>Group II</u>	<u>Group III</u>	<u>Group IV</u>
Median	24.5	25.0	58.0	114.5
Range	(24-26)	(23-26)	(30-82)	(103-153)
Average	24.6	24.7	54.4	121.8
S.D.	0.7	1.0	16.7	16.8

Group I = Mask only  
 Group II = Induction chamber, then mask  
 Group III = Ketamine-acepromazine  
 Group IV = Ketamine-xylazine

One of the major considerations is the total time from the end of surgical depth anesthesia to complete recovery, determined as the time to standing unaided shown in Table 4. Groups I and II showed very rapid recovery with an average of 8.2 and 9.5 minutes, respectively. As expected, there was no statistical difference between the two gas methods. Group III (ketamine-acepromazine) showed an average time of 81.3 minutes and Group IV (ketamine-xylazine) showed an average time of 55.8 minutes. These are significantly different from each other and from Groups I and II. Rapid recovery is one of the primary advantages of the gas anesthesia method.

Table 4

Minutes from end of surgical anesthesia to standing unaided

	<u>Group I</u>	<u>Group II</u>	<u>Group III</u>	<u>Group IV</u>
Median	7.3	9.2	65.0	51.5
Range	(4.2-16.0)	(7.3-12.3)	(52-135)	(23-97)
Average	8.2	9.5	81.3	55.8
S.D.	3.2	1.7	32.9	26.1

Group I = Mask only  
 Group II = Induction chamber, then mask  
 Group III = Ketamine-acepromazine  
 Group IV = Ketamine-xylazine

Table 5 summarizes the total time from administration of anesthesia to standing unaided, defined as complete recovery. The average for this measure was over 100 minutes longer with injectables than the worst of the two gas methods. The analysis of variance showed a significant difference between groups. The a posteriori analysis revealed differences between Group IV, Groups I and II, and Group III. No significant difference between Groups I and II was found.

Table 5

Time from administration of anesthesia to standing unaided

	<u>Group I</u>	<u>Group II</u>	<u>Group III</u>	<u>Group IV</u>
Median	32.0	34.0	121.5	178.5
Range	(30-47)	(31-39)	(83-216)	(138-231)
Average	33.8	34.4	135.7	180.4
S.D.	4.9	2.6	42.1	32.8

Group I = Mask only  
 Group II = Induction chamber, then mask  
 Group III = Ketamine-acepromazine  
 Group IV = Ketamine-xylazine

## Conclusion

Although each of the four methods of anesthesia will provide a satisfactory depth and duration of surgical anesthesia, there are advantages to each. Our findings with ketamine-acepromazine injected IM are in general agreement with those of Morgan, et al., 1981. They reported an induction period of about 5 minutes. We found a range of about 5-12 minutes with an average of 5.2 minutes. They reported surgical anesthesia lasting from 40-60 minutes. We found a range of 30-82 minutes with an average of 54.4 minutes. They reported

complete recovery in 2-5 hours. We found a range of 1.4-3.6 hours with an average of 2.3 hours from injection until the subject could stand unaided.

The injectables are less controllable once the ketamine-acepromazine or ketamine-xylazine is administered and have a long duration from injection to complete recovery. The ease of administering gas is apparent, but so is the initial expense of procuring the anesthesia unit. The use of the anesthesia chamber allows a smoother induction of anesthesia with minimal stress to the subject, but the subject must be removed from the anesthesia chamber and placed on the mask. This results in loss of gas anesthesia from the chamber. Then the chamber must be precharged again for each subsequent subject. With the ketamine combination injections, time to loss of righting reflex is somewhat shorter. The ketamine-xylazine has the shortest time to loss of righting reflex and to surgical anesthesia, however duration of surgical depth and time to complete recovery are excessive for our needs. On the other hand, time at surgical depth can be controlled with either gas method, minimizing the time at surgical depth.

We conclude that for situations requiring control over length of anesthesia and rapid recovery from anesthesia, either inhalation technique is superior to either ketamine-acepromazine or ketamine-xylazine injections, with the anesthesia chamber minimizing stress on the individual subjects.

Appendix A  
List of Manufacturers

Ralston Purina Company  
Checkerboard Square  
St. Louis, MO 63164  
(Purina Chin Chow)

Halocarbon Laboratories, Incorporated  
82 Berlews Court  
Hackensack, NJ 07601  
(Halothane U.S.P.)

Bristol Laboratories  
Division of Bristol-Myers Company  
Syracuse, NY 13201  
(Ketaset)

CEVA Laboratories, Incorporated  
Overland Park, KS 66212  
(Acepromazine)

Bayvet Division  
Miles Laboratories, Incorporated  
Shawnee, KS 66201  
(Rompum)

North American Drager  
148B Quarry Road  
Telford, PA 18969  
(Narkovet)

Burns-Biotec Laboratories, Incorporated  
8536 K Street  
Omaha, NE 68127  
(Optivet)

Ejay International, Incorporated  
P.O. Box 1835  
Glendora, CA 91740  
(Ejay)

## References

- Burdick, C. K., Patterson, J. H., Jr., Mozo, B. T. and Camp, R. T. 1978. Threshold shifts in chinchillas exposed to octave bands of noise centered at 63 and 1000 Hz for three days. Journal of the Acoustical Society of America. Volume 64:458-466.
- Deutsch, S. 1978. The pharmacodynamics of halothane. The Textbook of Veterinary Medicine. Baltimore, MD: Williams and Wilkins Company.
- Lumb, W. V., and Jones, E. W. 1973. Veterinary Anesthesia. Philadelphia, PA: Lea & Febiger.
- Miller, J. D. 1970. Audibility curve of the chinchilla. Journal of the Acoustical Society of America. Volume 48:513-523.
- Morgan, R. J., Eddy, L. B., Solie, T. N. and Turbes, C. C. 1981. Ketamine-acepromazine as an anesthetic agent for chinchillas (*Chinchilla laniger*). Laboratory Animal (United Kingdom). Volume 15:281-283.
- Schultz, T. A., and Fowler, M. E. 1974. The clinical effects of CI 744 in chinchillas, *Chinchilla villidera (laniger)*. Laboratory Animal Science. Volume 24(5):810-812.
- Soma, L. R. 1971. Systems and techniques for inhalation anesthesia. The Textbook of Veterinary Anesthesia. Baltimore, MD: Williams and Wilkins Company.
- Winer, B. J. 1971. Statistical Principles in Experimental Design. New York, NY: McGraw-Hill Book Company.

Initial distribution

Commander  
U.S. Army Natick Research  
and Development Center  
ATTN: Documents Librarian  
Natick, MA 01760

Naval Submarine Medical  
Research Laboratory  
Medical Library, Naval Sub Base  
Box 900  
Groton, CT 05340

Commander/Director  
U.S. Army Combat Surveillance  
& Target Acquisition Lab  
ATTN: DELCS-D  
Fort Monmouth, NJ 07703-5304

Commander  
10th Medical Laboratory  
ATTN: Audiologist  
APO NEW YORK 09180

Commander  
Naval Air Development Center  
Biophysics Lab  
ATTN: G. Kydd  
Code 60B1  
Warminster, PA 18974

Naval Air Development Center  
Technical Information Division  
Technical Support Detachment  
Warminster, PA 18974

Dr. E. Hendler  
Human Factors Applications, Inc.  
295 West Street Road  
Warminster, PA 18974

Under Secretary of Defense  
for Research and Engineering  
ATTN: Military Assistant  
for Medical and Life Sciences  
Washington, DC 20301

Commander  
U.S. Army Research Institute  
of Environmental Medicine  
Natick, MA 01760

U.S. Army Avionics Research  
and Development Activity  
ATTN: SAVAA-P-TP  
Fort Monmouth, NJ 07703-5401

U.S. Army Research and Development  
Support Activity  
Fort Monmouth, NJ 07703

Chief, Benet Weapons Laboratory  
LCWSL, USA ARRADCOM  
ATTN: DRDAR-LCB-TL  
Watervliet Arsenal, NY 12189

Commander  
Man-Machine Integration System  
Code 602  
Naval Air Development Center  
Warminster, PA 18974

Commander  
Naval Air Development Center  
ATTN: Code 6021 (Mr. Brindle)  
Warminster, PA 18974

Commanding Officer  
Naval Medical Research  
and Development Command  
National Naval Medical Center  
Bethesda, MD 20014

Director  
Army Audiology and Speech Center  
Walter Reed Army Medical Center  
Washington, DC 20307-5001

COL Franklin H. Top, Jr., MD  
Walter Reed Army Institute  
of Research  
Washington, DC 20307-5100

HQ DA (DASG-PSP-0)  
Washington, DC 20310

Naval Research  
Laboratory Library  
Code 1433  
Washington, DC 20375

Harry Diamond Laboratories  
ATTN: Technical Infor-  
mation Branch  
2800 Powder Mill Road  
Adelphi, MD 20783-1197

U.S. Army Materiel Systems  
Analysis Agency  
ATTN: Reports Processing  
Aberdeen proving Ground  
MD 21005-5017

U.S. Army Ordnance Center  
and School Library  
Building 3071  
Aberdeen Proving Ground,  
MD 21005-5201

U.S. Army Environmental Hygiene  
Agency Laboratory  
Building E2100  
Aberdeen Proving Ground,  
MD 21010

Technical Library  
Chemical Research  
and Development Center  
Aberdeen Proving Ground,  
MD 21010-5423

Commander  
U.S. Army Institute  
of Dental Research  
Walter Reed Army Medical Center  
Washington, DC 20307-5300

Naval Air Systems Command  
Technical Air Library 950D  
Rm 278, Jefferson Plaza II  
Department of the Navy  
Washington, DC 20361

Naval Research Laboratory Library  
Shock and Vibration Infor-  
mation Center, Code 5804  
Washington, DC 20375

Director  
U.S. Army Human Engineer-  
ing Laboratory  
ATTN: Technical Library  
Aberdeen Proving Ground,  
MD 21005-5001

Commander  
U.S. Army Test  
and Evaluation Command  
ATTN: AMSTE-AD-H  
Aberdeen Proving Ground,  
MD 21005-5055

Director  
U.S. Army Ballistic  
Research Laboratory  
ATTN: DRXBR-OD-ST Tech Reports  
Aberdeen Proving Ground,  
MD 21005-5066

Commander  
U.S. Army Medical Research  
Institute of Chemical Defense  
ATTN: SGRD-UV-AO  
Aberdeen Proving Ground,  
MD 21010-5425

Commander  
U.S. Army Medical Research  
and Development Command  
ATTN: SGRD-RMS (Ms. Madigan)  
Fort Detrick, Frederick, MD 21701

Commander  
U.S. Army Medical Research  
Institute of Infectious Diseases  
Fort Detrick, Frederick,  
MD 21701

Director, Biological  
Sciences Division  
Office of Naval Research  
600 North Quincy Street  
Arlington, VA 22217

Commander  
U.S. Army Materiel Command  
ATTN: AMCDE-S (CPT Broadwater)  
5001 Eisenhower Avenue  
Alexandria, VA 22333

Commandant  
U.S. Army Aviation  
Logistics School  
ATTN: ATSQ-TDN  
Fort Eustis, VA 23604

U.S. Army Training  
and Doctrine Command  
ATTN: ATCD-ZX  
Fort Monroe, VA 23651

Structures Laboratory Library  
USARTL-AVSCOM  
NASA Langley Research Center  
Mail Stop 266  
Hampton, VA 23665

Naval Aerospace Medical  
Institute Library  
Bldg 1953, Code 102  
Pensacola, FL 32508

Command Surgeon  
U.S. Central Command  
MacDill Air Force Base  
FL 33608

Air University Library  
(AUL/LSE)  
Maxwell AFB, AL 36112

Commander  
U.S. Army Medical Bioengineering  
Research and Development Lab  
ATTN: SGRD-UBZ-I  
Fort Detrick, Frederick,  
MD 21701

Defense Technical  
Information Center  
Cameron Station  
Alexandria, VA 22313

U.S. Army Foreign Science  
and Technology Center  
ATTN: MTZ  
220 7th Street, NE  
Charlottesville, VA 22901-5396

Director,  
Applied Technology Laboratory  
USARTL-AVSCOM  
ATTN: Library, Building 401  
Fort Eustis, VA 23604

U.S. Army Training  
and Doctrine Command  
ATTN: Surgeon  
Fort Monroe, VA 23651-5000

Aviation Medicine Clinic  
TMC #22, SAAF  
Fort Bragg, NC 28305

U.S. Air Force Armament  
Development and Test Center  
Eglin Air Force Base, FL 32542

U.S. Army Missile Command  
Redstone Scientific  
Information Center  
ATTN: Documents Section  
Redstone Arsenal, AL 35898-5241

U.S. Army Research and Technology  
Laboratories (AVSCOM)  
Propulsion Laboratory MS 302-2  
NASA Lewis Research Center  
Cleveland, OH 44135

AFAMRL/HEX  
Wright-Patterson AFB, OH 45433

University of Michigan  
NASA Center of Excellence  
in Man-Systems Research  
ATTN: R. G. Snyder, Director  
Ann Arbor, MI 48109

John A. Dellinger, MS, ATP  
University of Illinois-  
Willard Airport  
Savoy, IL 61874

Project Officer  
Aviation Life Support Equipment  
ATTN: AMCPO-ALSE  
4300 Goodfellow Blvd.  
St. Louis, MO 63120-1798

Commander  
U.S. Army Aviation  
Aviation Systems Command  
ATTN: DRSV-ED  
4300 Goodfellow Blvd  
St. Louis, MO 63120

Commanding Officer  
Naval Biodynamics Laboratory  
P.O. Box 24907  
New Orleans, LA 70189

U.S. Army Field Artillery School  
ATTN: Library  
Snow Hall, Room 14  
Fort Sill, OK 73503

Commander  
U.S. Army Health Services Command  
ATTN: HSOP-SO  
Fort Sam Houston, TX 78234-6000

U.S. Air Force Institute  
of Technology (AFIT/LDEE)  
Building 640, Area B  
Wright-Patterson AFB, OH 45433

Henry L. Taylor  
Director, Institute of Aviation  
University of Illinois-  
Willard Airport  
Savoy, IL 61874

Commander  
U.S. Army Aviation  
Systems Command  
ATTN: DRSV-WS  
4300 Goodfellow Blvd  
St. Louis, MO 63120-1798

Commander  
U.S. Army Aviation  
Systems Command  
ATTN: SGRD-UAX-AL (MAJ Lacy)  
4300 Goodfellow Blvd., Bldg 105  
St. Louis, MO 63120

U.S. Army Aviation  
Systems Command  
Library and Information  
Center Branch  
ATTN: DRSV-DIL  
4300 Goodfellow Blvd  
St. Louis, MO 63120

Federal Aviation Administration  
Civil Aeromedical Institute  
CAMI Library AAC 64D1  
P.O. Box 25082  
Oklahoma City, OK 73125

Commander  
U.S. Army Academy  
of Health Sciences  
ATTN: Library  
Fort Sam Houston, TX 78234

Commander  
U.S. Army Institute  
of Surgical Research  
ATTN: SGRD-USM (Jan Duke)  
Fort Sam Houston, TX 78234-6200

Director of Professional Services  
AFMSC/GSP  
Brooks Air Force Base, TX 78235

U.S. Army Dugway Proving Ground  
Technical Library  
Bldg 5330  
Dugway, UT 84022

U.S. Army Yuma Proving Ground  
Technical Library  
Technical Library  
Yuma, AZ 85364

AFFTC Technical Library  
6520 TESTG/ENXL  
Edwards Air Force Base,  
CAL 93523-5000

Commander  
Code 3431  
Naval Weapons Center  
China Lake, CA 93555

Aeromechanics Laboratory  
U.S. Army Research  
and Technical Labs  
Ames Research Center,  
M/S 215-1  
Moffett Field, CA 94035

Sixth U.S. Army  
ATTN: SMA  
Presidio of San Francisco,  
CA 94129

Commander  
U.S. Army Aeromedical Center  
Fort Rucker, AL 36362

Directorate  
of Combat Developments  
Bldg 507  
Fort Rucker, AL 36362

U.S. Air Force School  
of Aerospace Medicine  
Strughold Aeromedical Library  
Documents Section, USAFSAM/TSK-4  
Brooks Air Force Base, TX 78235

Dr. Diane Damos  
Department of Human Factors  
ISSM, USC  
Los Angeles, CA 90089-0021

U.S. Army White Sands  
Missile Range  
Technical Library Division  
White Sands Missile Range,  
NM 88002

U.S. Army Aviation Engineering  
Flight Activity  
ATTN: SAVTE-M (Tech Lib)  
Stop 217  
Edwards Air Force Base,  
CA 93523-5000

U.S. Army Combat Developments  
Experimental Center  
Technical Information Center  
Bldg 2925  
Fort Ord, CA 93941-5000

Commander  
Letterman Army Institute  
of Research  
ATTN: Medical Research Library  
Presidio of San Francisco,  
CA 94129

Director  
Naval Biosciences Laboratory  
Naval Supply Center, Bldg 844  
Oakland, CA 94625

Commander  
U.S. Army Aviation Center  
and Fort Rucker  
ATTN: ATZQ-CDR  
Fort Rucker, AL 36362

Directorate  
of Training Development  
Bldg 502  
Fort Rucker, AL 36362

Chief  
Army Research Institute  
Field Unit  
Fort Rucker, AL 36362

Commander  
U.S. Army Safety Center  
Fort Rucker, AL 36362

U.S. Army Aircraft Development  
Test Activity  
ATTN: STEBG-MP-QA  
Cairns AAF  
Fort Rucker, AL 36362

Chief  
Defence and Civil Institute  
of Environmental Medicine  
P.O. Box 2000  
ATTN: Director MLSD  
Downsview, Ontario Canada M3M 3B9

Staff Officer, Aerospace Medicine  
RAF Staff, British Embassy  
3100 Massachusetts Avenue, NW  
Washington, DC 20008

Canadian Society  
of Aviation Medicine  
c/o Academy of Medicine, Toronto  
ATTN: Ms. Carment King  
288 Bloor Street West  
Toronto, Canada M5S 1V8

Canadian Forces  
Medical Liaison Officer  
Canadian Defence Liaison Staff  
2450 Massachusetts Avenue, NW  
Washington, DC 20008

Officer Commanding  
School of Operational  
and Aerospace Medicine  
DCIEM P.O. Box 2000  
1133 Sheppard Avenue West  
Downsview, Ontario, Canada M3M 3B9

Chief  
Human Engineering Laboratory  
Field Unit  
Fort Rucker, AL 36362

Commander  
U.S. Army Aviation Center  
and Fort Rucker  
ATTN: ATZQ-T-ATL  
Fort Rucker, AL 36362

President  
U.S. Army Aviation Board  
Cairns AAF  
Fort Rucker, AL 36362

USA Medical Liaison Officer  
U.S. Embassy Box 54  
ATTN: USADO-AMLO  
FPO New York 09509

HQ, Department of the Army  
Office of The Surgeon General  
British Medical Liaison Officer  
DASG-ZX/COL M. Daly  
5109 Leesburg Pike  
Falls Church, VA 22401-3258

Canadian Airline Pilot's  
Association  
MAJ (Retired) J. Soutendam  
1300 Steeles Avenue East  
Brampton, Ontario, Canada L6T 1A2

Commanding Officer  
404 Squadron CFB Greenwood  
Greenwood, NS, Canada B0P 1N0

National Defence Headquarters  
101 Colonel By Drive  
ATTN: DPM  
Ottawa, Ontario, Canada K1A 0K2

Commanding Officer  
Headquarters, RAAF Base  
Point Cook Victoria,  
Australia 3029

Netherlands Army Liaison Office  
Buildingg 602  
Fort Rucker, AL 36362

British Army Liaison Office  
Building 602  
Fort Rucker, AL 36362

Canadian Army Liaison Office  
Building 602  
Fort Rucker, AL 36362

German Army Liaison Office  
Buildingg 602  
Fort Rucker, AL 36362

French Army Liaison Office  
Building 602  
Fort Rucker, AL 36362