

DTIC FILE UFR

USAFSAM-TP-88-10

AD-A201 274

**AN ANNOTATED BIBLIOGRAPHY OF HYPOBARIC DECOMPRESSION SICKNESS RESEARCH CONDUCTED AT THE CREW TECHNOLOGY DIVISION, USAF SCHOOL OF AEROSPACE MEDICINE, BROOKS AFB, TEXAS FROM 1983 TO 1988**

James T. Webb, Ph.D.  
Robert W. Krutz, Jr., Ph.D.  
Gene A. Dixon, M.B.A. (USAFSAM/VNS)

KRUG International  
Technology Services Division  
P.O. Box 790644  
San Antonio, Texas 78279-0644

October 1988

**DTIC**  
**ELECTE**  
DEC 1 2 1988  
**S** **D**  
H

Interim Report for Period 1 June 1983 - 15 October 1988

Approved for public release; distribution is unlimited.

Prepared For  
USAF SCHOOL OF AEROSPACE MEDICINE  
Human Systems Division (AFSC)  
Brooks Air Force Base, TX 78235-5301



88 12 12 050

NOTICES

This interim report was submitted by KRUG International, Technology Services Division, P.O. Box 790644, San Antonio, Texas, under contract F33615-85-C-4503, job order 7930-17-05, with the USAF School of Aerospace Medicine, Human Systems Division, AFSC, Brooks Air Force Base, Texas. Mr. Larry J. Meeker (USAFSAM/VNS) was the Laboratory Project Scientist-in-Charge.

When Government drawings, specifications, or other data are used for any purpose other than in connection with a definitely Government-related procurement, the United States Government incurs no responsibility or any obligation whatsoever. The fact that the Government may have formulated or in any way supplied the said drawings, specifications, or other data, is not to be regarded by implication, or otherwise in any manner construed, as licensing the holder or any other person or corporation; or as conveying any rights or permission to manufacture, use, or sell any patented invention that may in any way be related thereto.

The Office of Public Affairs has reviewed this report, and it is releasable to the National Technical Information Service, where it will be available to the general public, including foreign nationals.


This report has been reviewed and is approved for publication.



LARRY J. MEEKER, B.S.  
Project Scientist



W. CARTER ALEXANDER, Ph.D.  
Supervisor



JEFFREY G. DAVIS, Colonel, USAF, MC  
Commander

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE

REPORT DOCUMENTATION PAGE				Form Approved OMB No. 0704-0188	
1a. REPORT SECURITY CLASSIFICATION Unclassified			1b. RESTRICTIVE MARKINGS		
2a. SECURITY CLASSIFICATION AUTHORITY			3. DISTRIBUTION / AVAILABILITY OF REPORT Approved for public release; distribution is unlimited.		
2b. DECLASSIFICATION / DOWNGRADING SCHEDULE					
4. PERFORMING ORGANIZATION REPORT NUMBER(S)			5. MONITORING ORGANIZATION REPORT NUMBER(S) USAFSAM-TP-88-10		
6a. NAME OF PERFORMING ORGANIZATION KRUG International Technology Services Division		6b. OFFICE SYMBOL (if applicable)	7a. NAME OF MONITORING ORGANIZATION USAF School of Aerospace Medicine (VNS)		
6c. ADDRESS (City, State, and ZIP Code) P.O. Box 790644 San Antonio, TX 78279-0644			7b. ADDRESS (City, State, and ZIP Code) Human Systems Division (AFSC) Brooks AFB, TX 78235-5301		
8a. NAME OF FUNDING / SPONSORING ORGANIZATION		8b. OFFICE SYMBOL (if applicable)	9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER F33615-85-C-4503		
8c. ADDRESS (City, State, and ZIP Code)			10. SOURCE OF FUNDING NUMBERS		
			PROGRAM ELEMENT NO. 62202F	PROJECT NO. 7930	TASK NO. 17
11. TITLE (Include Security Classification) An Annotated Bibliography of Hypobaric Decompression Sickness Research Conducted at the Crew Technology Division, USAF School of Aerospace Medicine, Brooks AFB, Texas from 1983 to 1988					
12. PERSONAL AUTHOR(S) Webb, James T.; Krutz, Robert W., Jr.; and Dixon, Gene A.					
13a. TYPE OF REPORT Interim		13b. TIME COVERED FROM 83/06/01 TO 88/10/15	14. DATE OF REPORT (Year, Month, Day) 1988, October		15. PAGE COUNT 25
16. SUPPLEMENTARY NOTATION					
17. COSATI CODES			18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number) Decompression Sickness, Altitude, Hypobaric, Bubbles, Ultrasound, Doppler, Hematologic, Biochemical, Blood Susceptibility, Review, USAFSAM - (K - )		
FIELD	GROUP	SUB-GROUP			
06	04				
06	10				
19. ABSTRACT (Continue on reverse if necessary and identify by block number) Four major protocols, one of which includes five studies, have been initiated or completed in the 5-year period from 1983 to 1988. The studies have resulted in numerous publications which are listed as the references for this review. The purpose of this review is to provide an accessible summary of these extensive efforts and document the history of their accomplishments. The cross-reference information contained in this review is intended to simplify data accession within both published and data base records. A listing of the abbreviated title, protocol approval information, sponsorship information, computer database (HYPOB) retrieval numbers/titles, dates of exposure, and information about subjects, prebreaths, and exposure parameters for each study is followed by the published abstracts from each publication. <i>Keywords: Bibliographies;</i>					
20. DISTRIBUTION / AVAILABILITY OF ABSTRACT <input checked="" type="checkbox"/> UNCLASSIFIED / UNLIMITED <input type="checkbox"/> SAME AS RPT <input type="checkbox"/> DTIC USERS			21. ABSTRACT SECURITY CLASSIFICATION Unclassified		
22a. NAME OF RESPONSIBLE INDIVIDUAL Larry W. Meeker, S.S.			22b. TELEPHONE (Include Area Code) (512) 536-3811		22c. OFFICE SYMBOL USAFSAM/VNC

DD Form 1473, JUN 86

Previous editions are obsolete.

SECURITY CLASSIFICATION OF THIS PAGE

## TABLE OF CONTENTS

	<u>Page</u>
INTRODUCTION. . . . .	1
METHODS . . . . .	1
CHRONOLOGICAL PRESENTATION OF PROTOCOLS . . . . .	2
Sample Format . . . . .	2
Previous and Current Protocols. . . . .	3
1. Bends Screening Index. . . . .	3
2. Decompression Sickness Protection Using an 8 psia Suit Environment (7.8 psia Male Study) . . . . .	6
3. Decompression Sickness Protection Using an 8 psia Suit Environment (7.8 psia Female Study) . . . . .	7
4. Decompression Sickness Protection Using an 8 psia Suit Environment (Bubble Threshold Study). . . . .	9
5. Decompression Sickness Protection Using an 8 psia Suit Environment (9.5 psia Bubble Threshold Validation). . . . .	10
6. Decompression Sickness Protection Using an 8 psia Suit Environment (8.3 psia Study). . . . .	10
7. Pulmonary Tolerance to 100% Oxygen at 9.5 psia (9.5 psia Oxygen Toxicity Study) . . . . .	12
8. Effect of Carbon Dioxide on Decompression Sickness (5.46 psia, 3% Carbon Dioxide Prebreathe Study). . . . .	13
Studies Planned but not Begun as of October 15, 1988. . . . .	14
Effect of Postbreathing on Altitude DCS . . . . .	14
Effect of Isometric and Isotonic Exercise on Altitude Decompression Sickness . . . . .	14
Effect of Prebreathe with Exercise on Incidence of Altitude Decompression Sickness. . . . .	15
Effect of Rate of Ascent on Incidence of Altitude Decompression Sickness. . . . .	15
Bubble and Bends Correlation Study. . . . .	15
APPENDIX A: SUMMARY OF EXPOSURE PRESSURES AND RESULTS. . . . .	18
APPENDIX B: BIBLIOGRAPHY OF PUBLISHED RESULTS: 1983-1988. . . . .	21

For	<input type="checkbox"/>
	<input type="checkbox"/>
	<input type="checkbox"/>



Available	<input type="checkbox"/>
Available/Notes	<input type="checkbox"/>
Available and/or	<input type="checkbox"/>
Dist Special	<input type="checkbox"/>

A-1

AN ANNOTATED BIBLIOGRAPHY OF HYPOBARIC DECOMPRESSION SICKNESS RESEARCH  
CONDUCTED AT THE CREW TECHNOLOGY DIVISION, USAF SCHOOL OF AEROSPACE MEDICINE,  
BROOKS AFB, TEXAS FROM 1983 TO 1988

## INTRODUCTION

Four major decompression sickness (DCS) protocols, one of which includes five studies, have been initiated or completed in the period from 1983 to 1988. Data from these experiments have been entered on the USAFSAM VAX computer under the various study titles accessible via the menu-driven HYPOB retrieval program. The studies have resulted in numerous publications which are listed as the references for this review. The purpose of this review is to provide an accessible summary of these extensive efforts and document the history of their accomplishments. The cross-reference information contained in this review is intended to simplify data accession within both published and data base records.

The studies are presented in chronological order of their first exposure date. The abbreviated title, protocol approval information, sponsorship information, computer database (HYPOB) retrieval numbers/titles, dates of exposure, and information about subjects, prebreathe, and exposure parameters for each study are followed by the published abstracts from each publication. In some cases, the original protocol approval was used as authority to begin a study and later amendments provided for more efficient acquisition of subjects or extension of protocol parameters.

Appendix A is a table of results by pressure in which percentage bends is reported in two formats. One format indicates the percentage of subjects in the study who had Grade 2 bends during any of the exposures. The other format indicates the percentage of subjects in the study who had Grade 2 bends during the first of the exposures. Presenting both formats should provide some basis of comparison between studies in which there were only one or multiple exposures per subject. Comparisons between studies should be done with the understanding that other experimental parameters were variable in addition to pressure and number of exposures per subject (e.g., duration of exposure, exercise level, and breathing mixture). Appendix B is a current bibliography of published works which used data from any of the studies.

## METHODS

Prebreathe was with 100% O<sub>2</sub> unless otherwise specified. Subjects were monitored for intravenous gas bubbles, graded by the method of Neuman et al.<sup>1</sup> as follows: Grade 0, no bubbles; Grade 1, an occasional bubble signal, with the majority of the cardiac cycles being bubble free; Grade 2, bubbles in many but less than one-half of the cardiac cycles; Grade 3, bubbles in all

<sup>1</sup>Neuman TS, Hall DA, Linaweaver PG, Jr. Gas phase separation during decompression in man: Ultrasound monitoring. Undersea Biomed. Res. 1976;3:121-30.

the cardiac cycles; and Grade 4, numerous bubbles that obscure the heart sounds. Grades 1 and 2 bubbling were classified as "not severe" and Grades 3 and 4 were classified as "severe".

Bends screening exercises (Krutz and Dixon, 1987) consisted of five upward extensions of the arms while holding 5-lb weights and five chair-height knee bends. During doppler monitoring, joint flexion aided in releasing bubbles from extremities and increasing the reliability of precordial bubble detection.

Extravehicular Activity (EVA) exercises as described by Dixon et al. (1986) were similar, both quantitatively and qualitatively, to light workloads expected during actual EVA. The exercise workloads consisted of a different activity at each of three work stations and were designed to emphasize upper body stress. All exercise was performed in synchrony with a 5-s audible signal transmitted to headsets worn by the subjects who rotated sequentially from one station to another every 4 min. The subject at station 1 operated a cycle ergometer set at 0.5 kiloponds (kp) at the rate of 2 revolutions per 5 s, simulating manual closing of the Shuttle bay doors. At station 2, the subject operated a torque wrench while seated, at the rate of one tightening movement per 5 s to simulate operational tool use while weightless. The subject at station 3 pulled a Mini-Gym rope, at a resistance of 7.7 kg, using alternate hands, simulating translation along a cable in the Shuttle bay.

## CHRONOLOGICAL PRESENTATION OF PROTOCOLS

### Sample Format

(Study #) Name of Protocol  
Short Name (SGOR #; SAM ACHE #)  
(Approval authority and date)  
(Addenda with approval authority and dates)  
Sponsorship:  
VAX Title Code #: Name on VAX  
(first - last exposure dates)  
Number of subjects, gender  
    # Exposures/subject  
    # Subjects/Chamber Run  
Prebreathe information  
Exposure information  
    Exercise information

Abstracts

## Previous and Current Protocols

### -- 1. Bends Screening Index

Bends Screening Index (SGOR 83-001; SAM ACHE 82-19)  
(SGP approval, 7Apr83)

(Addendum, blood drawing; SGP approval, 26Apr85; SAM ACHE #85-8)

Sponsorship: This research was supported in part by USAF Contracts  
F33615-81-C-0600 and F33615-85-C-4503 and NASA Contract T-82170.

VAX Title Code 1/: Bends Screening Index  
(May83-Aug87)

Male Subjects

6 Exposures/Subject, > 4 days between exposures  
3 Subjects(max)/Chamber Run

Prebreathe: 1 h

8 h/Exposure @ 4.37 psia (30,000 ft; 14 subjects w/o exercise; 31  
subjects with exercise), 4.89 psia (27,500 ft; 33 subjects), 5.46  
psia (25,000 ft; 29 subjects), 6.08 psia (22,500 ft; 19 subjects);  
breathing 100% oxygen

Bends screening exercises (4 cycles/h)

Abstract - Bends Susceptibility (Adams et al., 1984):

"Introduction. The purpose of this study was to develop a method of producing a broad spectrum of bends symptom onset times to differentiate between bends-prone and -resistant individuals. A pilot study was conducted to determine the optimal altitude to produce this differentiation. Methods. Ten male subjects were exposed to 3 altitudes: 25K, 27.5K, and 30K ft, for a maximum of 8 hrs. Each flight, conducted at least 1 week apart, was preceded by 1 hr of prebreathing 100% O<sub>2</sub>. An exercise regimen--consisting of 5 knee bends and the lifting of two 5-lb weights, 5 times each 15 min--was performed. The subjects were monitored for intravascular bubbles and subjective symptoms. The onset times were recorded. The subjects were removed from the chamber if continuous symptoms were reported, or at the end of 8 hrs. Results. All subjects developed symptoms at 30K ft, with an average onset time of 93.5 min and a range of 42 to 172 min. At 27.5K ft, only 6 subjects developed symptoms. The average onset time was 127 min with a range of 59 to 204 min for those who developed symptoms. Eight subjects reported symptoms at 25K ft; the mean onset time was 133 min, ranging between 82 and 234 min. A high tolerance to decompression stress was exhibited by 3 subjects, and 2 exhibited high susceptibility at all altitudes. Conclusions. The data suggest that, under the stated parameters, an altitude of 27.5 K ft allows the greatest differentiation of bends susceptibility, although more data are necessary for validation."

Abstract - Validity of Ultrasonic Monitoring at Altitude for Bends Detection (Olson et al., (1986):

"Introduction. The ultrasonic precordial bubble detector (UPBD) is an inexpensive, portable device used to detect intravascular bubbles associated with bends. Its value as a device to warn of impending bends is controversial. This paper will present precise, well established numerical estimates of the sensitivity and specificity of

the UPBD based on data from over 100 subjects at various altitudes. Methods. Eighty-nine male and thirty female subjects were taken to simulated altitudes of 16,000 to 30,000 ft after prebreathing 100% oxygen for all but the lowest level flights. At altitude, the subjects performed mild exercise and were monitored with a continuous wave, 5 MHz Doppler bubble detector. They were returned to ground level after 6 to 8 hours or when they developed bends. UPBD sensitivity, defined as: number of bubblers who also bend/number of benders, as calculated from high altitude data where bends was prevalent. Specificity, defined as: Number of bubble-free subjects who also are bends-free/number of bends-free subjects, was determined at low altitudes where there were many non-benders. Results. The results showed that the specificity of the UPBD was 64% at 16,000 ft and 50% at 27,500 ft. The sensitivity was 86% at 30,000 ft and 80% at 27,500 ft. Conclusions. It was concluded that the UPBD, as a warning device for impending bends, is reasonably sensitive and can be expected to predict the majority of cases. It is, however, not very specific and will yield many false positives."

Abstract from The Effects of Exercise on Bubble Formation and Bends Susceptibility at 9,100 m (30,000 ft; 4.3 psia) (Kruz & Dixon, 1987): "This study assessed the value of controlled exercise in a bends susceptibility test. Healthy male subjects were exposed to a pressure altitude of 9,100 m (30,000 ft; 4.3 psia) for a period not exceeding 8 h on two separate days at least 1 week apart. During one exposure, subjects performed five deep knee bends followed by five upward arm extensions with 5-lb weights every 15 min; during the other exposure, they remained sedentary. Exercise and no-exercise altitude exposures were randomized between subjects. A precordial Doppler monitoring technique was used to record venous bubbling at 15-min intervals. Bends was diagnosed from subjective symptoms. Results have shown that controlled exercise decreases time to maximum venous bubbling and increases the incidence of Grade 2 bends compared to no-exercise conditions. Exercise also appears to increase the validity of precordial Doppler monitoring as a method to predict bends onset."

Abstract from Preliminary Findings: Bends Screening Index Study (Smead, 1987): "Objective: To determine the extent of the relationship between intravascular bubbling at altitude and the development of limb bends. Such data could be used to develop a test for bends susceptibility based on the timing and/or intensity of intravascular bubbling. An additional purpose of the investigation was to identify characteristics which might predispose an individual to bends. Methods: Forty-eight male volunteers made a total of 190 man-flights to four selected altitudes (30000, 27500, 25000, and 22500 ft pressure equivalent) in a hypobaric chamber. The subjects' ages ranged from 19 to 48, and all were in good health. Subjects denitrogenated for one hour prior to the flight, and continued breathing 100% oxygen at altitude. Light periodic exercise was performed during the flight, which continued for eight hours unless terminated sooner by the development of grade 2 limb bends (constant moderate pain which resolved on descent). Intravascular bubble grade was determined by ultrasound every 15 minutes: 0 to 4 on the Neuman scale with grades 3

and 4 classified as significant. Results: Altitude (25000 ft and above) did not significantly affect the incidence of limb bends which ranged from 70% to 74% (43% at 22500 ft), but did affect the time of onset, with increased time to bends at each successively lower altitude. Incidence of significant intravascular bubbling varied erratically between 66% and 78% at and above 25000 ft (51% at 22500 ft), but time to onset was similar to the bends pattern. Although the association between bends and bubbling was strong (Chi Square = 9.404,  $p = 0.002$ ), and intravascular bubbling preceded limb bends in 93 of 126 cases, significant bubbling was correctly negative in only 31 of 64 flights without bends (sensitivity = 0.738, specificity = 0.484). Age stood out as the most significant predeterminate for bubble susceptibility ( $t = 3.13$ ,  $p = 0.002$ )."

Abstract from An Evaluation of Precordial Ultrasonic Monitoring to Avoid Bends at Altitude (Olson et al., 1988):

"Several investigators have reported that intravascular bubbles can be detected in decompressed subjects before they develop bends. The altitude exposures were generally of short duration with a limited number of subjects. This important preliminary finding needed to be verified in a larger sampling of long duration altitude exposures. In this experiment, 32 subjects in 82 flights were taken to 27,500 ft simulated altitude for 8 h or until the subject developed mild but steady joint pain (bends). Many subjects took more than one flight. At altitude, the subjects were monitored for circulating bubbles by a team of well-trained, experienced technicians. It was determined that bubbles, clearly audible even to untrained observers, occurred in 77% of the flights in which the subjects developed bends. On the other hand, no bubbles were found in 61% of the flights in which the subjects remained bends free even though the subjects were monitored by more than one experienced technician. Therefore, at 27,500 ft ultrasonic monitoring will miss about 25% of the subjects who developed bends (false negatives) and will incorrectly identify a little less than half of the subjects who do not develop bends as potential benders (false positives)."

## 2. Decompression Sickness Protection Using an 8 psia Suit Environment

7.8 psia Male Study (SGOR 83-010; SAM ACHE #83-14)

1st study within this protocol

(SGP approval, 21Oct83)

Sponsorship: This research was supported in part by USAF Contracts F33615-81-C-0600 and F33615-85-C-4503 and NASA Contract T-82170.

VAX Title Code 18: 7.8 psia Male Study

(Oct83-Jun84)

30 Male Subjects

3 Consecutive Daily Exposures/Subject

3 Subjects(Max)/Chamber Run

Prebreathe: None

6 h/Exposure @ 7.8 psia (16,500 ft); breathing 50% O<sub>2</sub>, 50% N<sub>2</sub>

EVA exercises

Abstract from Decompression Sickness and Intravenous Bubble Formation Using a 7.8 psia Simulated Pressure-Suit Environment (Dixon et al., 1986):

"The purpose of this study was to determine the minimum spacesuit pressure required to prevent decompression sickness (DCS) during operational conditions in a 50% oxygen/50% nitrogen environment. In this study, 30 male volunteer subjects were exposed in groups of three, to three consecutive daily extravehicular activity (EVA) simulations at 7.8 psia (5,031 m altitude equivalent) for a continuous period of 6 h. During each altitude exposure, the subjects participated in similar exercise workloads expected to be experienced by astronauts during a typical EVA scenario. Precordial Doppler monitoring revealed that 73.3% of the subjects had intravenous bubbling during at least 1 d of the 3 d of exposure, with 26.7% remaining bubble-free during the entire study. No correlation was found between either body fat or age and incidence of bubble formation. One case of DCS occurred during the study indicating that 7.8 psia is not sufficient pressure to totally preclude DCS in a 50% oxygen/50% nitrogen environment. The necessary pressure awaits further study."

Abstract - Determining a Bends-Preventing Pressure for a Space Suit (Krutz et al., 1988):

"It is desirable that a pressure suit used for extravehicular activity (EVA) (1) eliminate the threat of decompression sickness (bends) and (2) require no preoxygenation prior to EVA. This paper chronicles the definition of a pressure to prevent bends during EVA without preoxygenation and subsequent studies to test and evaluate this pressure using both male and female subjects with different breathing gas mixtures. METHODS. Initially, a study was conducted at 7.8 psia using a 50:50::O<sub>2</sub>:N<sub>2</sub> breathing gas mixture without prebreathing O<sub>2</sub> and simulated EVA workloads. Since this pressure did not totally eliminate bends, a subsequent study was conducted using step-wise increases in pressure to determine a suit pressure at which both significant intravascular bubbles and bends were eliminated without preoxygenation. The results indicated that 9.5 psia met the aforementioned criteria in male subjects. To validate 9.5 psia,

subsequent studies were conducted using both males and females and 40:60::O<sub>2</sub>:N<sub>2</sub> and 100% O<sub>2</sub> breathing gases. RESULTS AND CONCLUSIONS. No cases of significant bubbling, bends, or any other detrimental physiologic effects were noted during any exposure. It appears then, from a physiologic viewpoint, that a minimum pressure of 9.5 psia should be considered as the standard for EVA from a 14.7 psia space station.

See the abstract of Olson et al. (1986) under Bends Screening Index (1), for additional information about this study.

See the abstract of Krutz et al. (1985) under Bubble Threshold Study (4), for additional information about this study.

### 3. Decompression Sickness Protection Using an 8 psia Suit Environment

7.8 psia Female Study (SGOR 83-010; SAM ACHE #83-14)

2nd study within this protocol

(SGP approval, 21Oct83)

(Addendum 2; use contractor-procured female subjects; ACHE review, 14Aug84; SAM/CC review/approval, 7Sep84; SG approval, 15Mar85; SGP approval, 22Mar85; SAM ACHE #84-15)

Sponsorship: This research was supported in part by USAF Contracts F33615-81-C-0600 and F33615-85-C-4503 and NASA Contract T-82170.

VAX Title Code 20: 7.8 psia Female Study

(Nov83-Jul85)

30 Female Subjects

3 Consecutive Daily Exposures/Subject

3 Subjects(Max)/Chamber Run

Prebreathe: None

6 h/Exposure @ 7.8 psia (16,500 ft); breathing 50% O<sub>2</sub>, 50% N<sub>2</sub>

EVA exercises

Abstract - Female Susceptibility to Decompression Sickness and Bubble Formation Using a Simulated 7.8 psia Suit Environment (Dixon & Krutz, 1986):

Introduction. The purpose of this study was to determine female susceptibility to decompression sickness (DCS) during simulated extravehicular activity (EVA). Methods. Thirty female volunteer subjects were exposed in groups of three to three consecutive daily EVA simulations at 7.8 psia (5031 m altitude equivalent) for a continuous period of 6 h. During each altitude exposure, the subjects breathed a gas mixture of 50% oxygen/50% nitrogen and participated in exercise workloads similar to those expected to be experienced by astronauts during a typical EVA scenario. Results. Precordial Doppler monitoring revealed that 43.3% of the subjects had intravenous bubbling during at least one of the three days of exposure. Seventeen of the thirty subjects (56.7%) were bubble free during the entire study. Five of the thirty subjects (16.7%) developed DCS as a result of exposure to these study conditions and two of these developed delayed symptoms. Three of these five subjects underwent hyperbaric oxygen treatment. Conclusion. Female subjects appear to suffer more delayed DCS symptoms necessitating more hyperbaric oxygen treatment than male subjects under the same experimental conditions. Female

subjects do not appear to bubble as frequently as male subjects when exposed to these study conditions."

Abstract from Decompression Sickness and Bubble Formation in Females Exposed to a Simulated 7.8 psia Suit Environment (Dixon et al., 1988): "The purpose of this study was to measure female susceptibility to decompression sickness (DCS) during simulated extravehicular activity (EVA) at a candidate (7.8 psia) suit pressure. Thirty female volunteer subjects in groups of three, were exposed to three consecutive daily EVA simulations at 7.8 psia (5031 m altitude equivalent) continuously for 6 h. During each altitude exposure, the subjects breathed a gas mixture of 50% oxygen/50% nitrogen, and participated in exercise workloads similar to those expected to be experienced by astronauts during a typical EVA scenario. Precordial Doppler bubble monitoring was accomplished after each cycle of exercise workload simulations. During at least one of the three days of exposure, 43% of the subjects experienced intravenous bubbling. Of the 30 subjects, 17 (57%) did not experience detectable bubbling on any of the three days of exposure and 5 (17%) developed decompression sickness (DCS) during the study. Two cases were delayed, occurring after recompression to ground level; and three subjects required hyperbaric oxygen treatment. The results of this study suggest that female subjects may suffer more delayed DCS symptoms, necessitating hyperbaric oxygen treatment, than their male counterparts under the same experimental conditions. Female subjects did not experience intravenous bubbling as frequently as male subjects when exposed to these study conditions."

See the abstract of Olson et al. (1986) under Bends Screening Index (1), for additional information about this study.

See the abstract of Krutz et al. (1988) under Decompression Sickness Protection Using an 8 psia Suit Environment (2), for additional information about this study.

#### 4. Decompression Sickness Protection Using an 8 psia Suit Environment

Bubble Threshold Study (SGOR 83-010; SAM ACHE #83-14)

3rd study within this protocol

(SGP approval, 21Oct83)

(Addendum; use bubble-prone subjects at higher pressures; ACHE review, 17Aug84; SAM/CC approval, 7Sep84; SAM ACHE #84-19)

(Addendum 3; add more male subjects, alter the purpose, add new altitudes, and use contractor-procured subjects; ACHE review, 14Aug84; SAM/CC review/approval/referral, 7Sep84; SGP approval, 22Mar85; SAM ACHE #84-16)

(Amendment to Addendum 3; blood drawing; SGP approval, 26Apr85; SAM ACHE #85-7)

Sponsorship: This research was supported in part by USAF Contracts F33615-81-C-0600 and F33615-85-C-4503 and NASA Contract T-82170.

VAX Title Code 19: Bubble Threshold Study  
(Apr84-Apr85)

Male Subjects

1 Exposure/Subject at a given pressure

3 Subjects(Max)/Chamber Run

Prebreathe: None

6 h/Exposure @ 8.0 psia (25 subjects), 8.5 psia (10 subjects), 9.0 psia (22 subjects), 9.5 psia (6 subjects), 10.0 psia (9 subjects), 10.5 psia (2 subjects); breathing 50% O<sub>2</sub>, 50% N<sub>2</sub>

EVA exercises

Abstract from Minimum Pressure for a Zero-Prebreathe Pressure Suit (Krutz et al., 1985):

"There are two current approaches to reducing the risk of decompression sickness during repeated extravehicular activity (EVA) without prebreathing 100% oxygen. One approach suggests the use of different pressures in the transfer vehicle, the station, and the suit. The other would use advanced pressure-suit technology to build a suit that will make different pressures unnecessary provided the bends-free suit pressure is reasonable and can be readily determined. Research at the USAF School of Aerospace Medicine since November 1982 has been directed at determining this suit pressure using human subjects at simulated altitudes of 16,500 to 10,000 feet (7.8 to 10 psia). An earlier report of this ongoing research showed that bends is not totally eliminated at 7.8 psia. The present study is a continuation of this effort to define a "bends-free" suit pressure with the initial results suggesting that this pressure begins at around 9.5 psia (from a 14.7 psia station)."

See the abstract of Krutz et al. (1988) under Decompression Sickness Protection Using an 8 psia Suit Environment (2), for additional information about this study.

## 5. Decompression Sickness Protection Using an 8 psia Suit Environment

9.5 psia Bubble Threshold Validation (SGOR 83-010; SAM ACHE #83-14)  
4th study within this protocol  
(Addendum 4; validate 9.5 psia with additional subjects; ACHE review,  
16May85; SAM/CC approval, 12Jul85; SAM ACHE #85-13)

Sponsorship: This research was supported in part by USAF Contracts  
F33615-81-C-0600 and F33615-85-C-4503 and NASA Contract T-82170.

VAX Title Code 21: 9.5 psia Bubble Threshold Validat.  
(Aug86-Nov86)

20 Male & 10 Female Subjects

1 Exposure/Subject

3 Subjects(Max)/Chamber Run

Prebreathe: None

6 h/Exposure @ 9.5 psia (11,500 ft); breathing 40% O<sub>2</sub>, 60% N<sub>2</sub>  
EVA exercises

Abstracted from Evaluation of 9.5 psia as a Suit Pressure for Prolonged  
Extravehicular Activity (Dixon and Krutz 1985):

The present study was undertaken to determine if a pressure of 9.5  
psia would serve to prevent the occurrence of DCS in both males and  
females, without the requirement of prebreathing or the use of stage  
decompression, during a typical simulated Extravehicular Activity  
(EVA) scenario. Four of the twenty (20%) male subjects experienced  
Grade 1 or 2 bubble formation. None of the females exhibited any  
bubble formation. The study conditions of 40% O<sub>2</sub>, 60% N<sub>2</sub> at 9.5 psia  
for 6 h did not result in the development of bends among any of the  
subjects. Thus, this pressure (9.5 psia) appears to protect the  
astronaut from both formation of severe bubbling and development of  
bends symptoms when exposed to these study conditions.

See the abstract of Krutz et al. (1988) under Decompression Sickness  
Protection Using an 8 psia Suit Environment (2), for additional  
information about this study.

## 5. Decompression Sickness Protection Using an 8 psia Suit Environment

8.3 psia Study (SGOR 83-010; SAM ACHE #83-14)

5th study within this protocol

(Addendum 5; test proposed space suit pressure; ACHE review, 29Aug85;  
SAM/CC approval, 13Sep85; SAM ACHE #85-25)

Sponsorship: This research was supported in part by USAF Contracts  
F33615-81-C-0600 and F33615-85-C-4503 and NASA Contract T-82170.

VAX Title Code 24: 8.3 psia Study  
(Aug86-Nov86)

20 Male & 11 Female Subjects

1 Exposure/Subject

3 Subjects(Max)/Chamber Run

Prebreathe: None

6 h/Exposure @ 8.3 psia (15,000 ft); breathing 50% O<sub>2</sub>, 50% N<sub>2</sub>  
EVA exercises

Abstract from Decompression Sickness and Venous Gas Emboli at 8.3 psia (Smead et al., 1986):

"This study sought to determine the bends risk on decompression from sea level to 8.3 psia. On the basis of several prior studies by NASA and the Air Force, this differential was expected to result in a minimal (about 5%) incidence of mild decompression sickness, and may be the pressure of choice for the next generation NASA extravehicular activity (EVA) pressure suit. Thirty-one volunteer subjects, performing light work characteristic of EVA, were exposed to 8.3 psia (4572 m) pressure altitude for six hours in an altitude chamber. Limb bends incidence was 3.2%, and 25.8% of the subjects demonstrated significant intravascular bubbling. Those who bubbled were significantly older than the bubble-free group, but differed in no other aspect. An 8.3 psia advanced pressure suit design was considered insufficient to totally preclude the risk of decompression sickness."

Abstract - Blood Factors and Venous Gas Emboli: Surface to 429 mmHg (8.3 psia) (Webb et al., 1988a; short version of Webb et al., 1987):

"Analyses of 43 parameters were performed on blood obtained from 30 volunteer subjects before and after a 6-h chamber decompression from the surface to 429 mmHg. Eight subjects (5 male, 3 female) were bubble-prone (bubble grades 3 and 4), and 22 (15 male, 7 female) were resistant (bubble grade 0) to forming bubbles as detected with precordial Doppler. Significant ( $P < .05$ ) differences include the following: Higher levels of cholesterol in the bubble-prone males and combined subjects (males and females) than in their resistant counterparts; higher magnesium in the bubble-prone males; shorter pre-exposure prothrombin time in bubble-prone males and combined subjects; increased partial thromboplastin time in bubble-prone females versus the resistant females who showed a decrease during exposure; higher pre-exposure hemoglobin, hematocrit, and red blood cell count in the bubble-prone females; and significant reduction in hemoglobin, red blood cell count, and serum osmolality in the bubble-prone females during the exposure relative to changes in the resistant females. In this study, high cholesterol and hemoconcentration appear to be characteristics of bubble-prone subjects."

See the abstract of Krutz et al. (1988) under Decompression Sickness Protection Using an 8 psia Suit Environment (2), for additional information about this study.

## 7. Pulmonary Tolerance to 100% Oxygen at 9.5 psia

9.5 psia Oxygen Toxicity Study (SGOR 86-007; SAM ACHE #86-05)  
(SG approval, 26Mar87; SGPT approval, 30Mar87)  
(Addendum; change investigators and define test subject physical; SGPT approval, 21Jan88)

Sponsorship: This research was supported in part by USAF Contract F33615-85-C-4503 and NASA Contract T-82170.

VAX Title Code 22: 9.5 psia Oxygen Toxicity Study  
(Sep87-May88)

12 Male & 12 Female Subjects

5 Consecutive Daily Exposures/Subject

3 Subjects(Max)/Chamber Run

Prebreathe: None

8 h/Exposure @ 9.5 psia (11,500 ft); breathing 100% O<sub>2</sub>  
EVA exercises (4 cycles/h)

Abstract from Oxygen Toxicity During 5 8-h Exposures to 100% Oxygen at 9.5 psia (Webb et al., 1988b):

"We conducted a study to determine if oxygen toxicity occurs in a proposed extravehicular activity (EVA) pressure suit environment. Twelve male subjects were exposed to 100% oxygen at 9.5 psia for five consecutive days, 8 h/day while performing moderate exercise. No decompression sickness or venous gas bubbles were detected. Pulmonary function tests, physical exams, blood analyses, arterial oxygen saturation monitoring, and x-rays showed no evidence of oxygen toxicity. These results suggest that a 100% oxygen, 9.5 psia pressure suit environment could avoid both decompression sickness and oxygen toxicity during EVAs of comparable duration and physical activity."

Abstract from Human Tolerance to Five Daily Simulated Eight-hour EVA Exposures to 100% Oxygen at 9.5 psia (Webb et al., In press):

"Extravehicular Activity (EVA) currently involves decompression to 4.3 psia. This degree of decompression carries a significant potential for decompression sickness (DCS) which could be alleviated if a pressure of 9.5 psia could be maintained in the pressure suit. Previous research has not evaluated the potential for oxygen toxicity at 9.5 psia. Twenty-one subjects were exposed to 100% oxygen at 9.5 psia for five consecutive days, 8 h/day while performing moderate exercise to simulate a typical work-week in the proposed pressure suit environment. No DCS or venous gas bubbles were detected. Pulmonary function tests, physical exams, blood analyses, arterial oxygen saturation monitoring, and x-rays showed no evidence of oxygen toxicity under these conditions. These results suggest that a 100% oxygen, 9.5 psia pressure suit environment could avoid both DCS and oxygen toxicity during EVAs of comparable duration and physical activity."

See the abstract of Krutz et al. (1988) under Decompression Sickness Protection Using an 8 psia Suit Environment (2), for additional information about this study.

## 8. Effect of Carbon Dioxide on Decompression Sickness

5.46 psia, 3% Carbon Dioxide Prebreathe Study (SGOR 87-004;  
SAM ACHE #87-8)

(SGPT approval, 9Jul87)

(Addendum; change investigators; SGPT approval, 22Apr88)

(Addendum; change exposure time; SGPT approval, 22Jul88)

(Addendum; finger stick; ACHE review, 31Aug88; SAM/CC approval, 11Oct88)

Sponsorship: This research was supported in part by USAF Contract  
F33615-85-C-4503 and NASA Contract T-82170.

VAX Title Code 23: 5.46 psia, 3% CO<sub>2</sub> Prebreathe Study  
(Nov87-)

30 Male Subjects

4 Exposures/Subject

3 Subjects(Max)/Chamber Run

Prebreathe: 1 h 100% O<sub>2</sub> (2 exposures with 2.9-3.1% CO<sub>2</sub>)

3 or 6 h/Exposure @ 5.46 psia (25,000 ft); breathing 100% O<sub>2</sub>

Bends screening exercises (4 cycles/h)

### Abstract - Audio and Visual Ultrasonic Monitoring of Altitude

Decompression (Baas et al., 1988):

"Of major concern in high altitude and space flight is the development of altitude-induced decompression sickness. Bubbling occurs in a large percentage of subjects who subsequently develop decompression sickness (bends), and therefore monitoring for bubbles is an important tool in evaluating the effectiveness of specific space suit pressurization schedules. This paper details use in our laboratory of two-dimensional ultrasound with the Hewlett Packard Model Number 77020 Ultrasonic Imaging System (Sono 500). This device incorporates both visual and auditory (Doppler) real-time information for the detection of bubbles in subjects exposed to altitude. Ultrasound transducer placement is over the precordium in the acoustic window located between the fourth and sixth intercostal space, with a resulting modified short axis view of the heart. Doppler flow signals are obtained as blood passes from the right atrium through the tricuspid valve. Bubbles, when present, can be seen within the heart and heard in the flow signal. Data collected over an 18 month period of equipment operation will be presented, illustrating the suitability of an integrated auditory and visual ultrasonic system for detection of circulating bubbles.

### Abstract - Effects of a Carbon Dioxide Prebreathe Gas Mixture on the Incidence of Altitude-Induced Decompression Sickness. (Baas, et al., In press):

Introduction. Because of the vasodilative effect of carbon dioxide in some tissues, it is hypothesized that increased perfusion from prebreathing low tensions of carbon dioxide will accelerate the nitrogen washout rate and thereby decrease the probability of bubble formation and subsequent development of decompression sickness (DCS). The purpose of this study is to determine the effect of increased carbon dioxide tension in the prebreathe gas mixture on the incidence of altitude DCS. Methods. Thirty male subjects were exposed to a simulated altitude of 25,000 ft after 1 h prebreathe of (1) 100%

oxygen, i.e., control (2 exposures), or (2) 3% sea level equivalent carbon dioxide (23 torr) with the balance oxygen (2 exposures) for a total of 4 exposures. Subjects remained at 25,000 ft for 3 h or until they experienced persistent DCS symptoms. Results. To date (Sep 88), 78 altitude exposures have been conducted which resulted in 8 cases of DCS. There was no difference in incidence of DCS between the test and control groups. The carbon dioxide prebreathe test group and the control group each experienced 4 cases of DCS (10.3%). These experiments are ongoing. Conclusions. For relatively short prebreathe periods of 1 h and altitude exposures to 25,000 ft, supplementing the prebreathe mixture with 3% carbon dioxide appears to have no effect on the incidence of DCS.

Studies Planned but not Begun as of October 15, 1988

Effect of Postbreathing on Altitude DCS

4.46 psia SAC Study (SGOR 88-003; SAM ACHE #87-30)  
(SGPT approval, 9Nov87; SG approval, 9Nov87)  
(Jan89?-)  
15 Subjects  
    7 Exposures/Subject  
    3 Subjects(Max)/Chamber Run  
Variable pre/postbreathe  
4 h/Exposure @ 4.46 psia; breathing 100% O<sub>2</sub>  
    Minimal exercise

Effect of Isometric and Isotonic Exercise on Altitude Decompression Sickness

6.07 psia Isometric/Isotonic Exercise Study (SGOR 87-007)  
(SGPT approval, 17Aug87)  
(Jan89?-)  
24 Subjects  
    3-9 Exposures/Subject  
    3 Subjects(Max)/Chamber Run  
Prebreathe: None  
4 h/Exposure @ 6.07 psia (22,500'); 100% O<sub>2</sub>  
    Moderate-heavy exercise

Effect of Prebreathe with Exercise on Incidence of Altitude Decompression Sickness

5.46 psia Prebreathe with Exercise Study (SGOR 8-)  
VAX Title Code :  
(Apr89?-)  
30 Subjects  
    3 Exposures/Subject  
    3 Subjects(Max)/Chamber Run  
3 Prebreathe variables: 1) 15 min with heavy exercise  
                          2) 15 min w/o exercise  
                          3) 1 h w/o exercise  
4 h/Exposure @ 5.46 psia (25,000'); 100% O<sub>2</sub>  
    EVA exercise (2 cycles/h)

Effect of Rate of Ascent on Incidence of Altitude Decompression Sickness

5.46 psia Rate of Ascent Study (SGOR 8-)  
VAX Title Code :  
(Jun89?-)  
24 Subjects  
    3 Exposures/Subject  
    3 Subjects(Max)/Chamber Run  
Prebreathe: 1 h  
4 h/Exposure @ 5.46 psia (25,000 ft); 100% O<sub>2</sub>  
    EVA exercise (2 cycles/h)  
    3 Rates of ascent: 1) Standard; 5000'/min  
                          2) Fast; 11,500'/min  
                          3) Rapid; 2-3 s to 25,000'

Bubble and Bends Correlation Study

Bubble/Bends Correlation Study (SGOR 87-002; SAM ACHE # 87-4)  
VAX Title Code :  
(Aug89?-)  
30 Subjects  
    6 Exposures/Subject  
    3 Subjects(Max)/Chamber Run  
Prebreathe: None  
6 h/Exposure @ 5.46-7.8 psia (25,000 ft-16,500 ft); 100% O<sub>2</sub>  
    EVA exercise

APPENDIXES

APPENDIX A

SUMMARY OF EXPOSURE PRESSURES AND RESULTS

PSIA	mmHg	STUDY #	PREBREATHE TIME, MIN	BREATHING % O <sub>2</sub>	SUBJECTS #/GENDER	EXPOSURE: SUBJ, TIME	TOTAL # OF EXPOSURES	EXERCISE TYPE	% SURJ W/HENDS	MEAN MIN TO BEND	% WITH GRADE		% WITH GRADE	% BUBBLES
											NOTE 1	NOTE 2		
4.37	226	30000	60	100	31M	1-3, 8h	43	B	71	81	95	81	3	16
4.37	226	30000	60	100	14M	1-2, 8h	15	C	50	57	151	57	0	43
4.89	252	27500	60	100	33M	1-4, 8h	83	B	76	85	144	91	3	6
5.46	282	25000	60	100	30M	NOTE 8	120	B						
5.46	282	25000	60	100	29M	1-2, 8h	30	B	69	69	151	79	14	7
6.08	314	22500	60	100	19M	1-3, 8h	46	B	47	68	201	68	0	32
7.80	404	16500	0	50	32M	1-3, 6h	94	A	3	3	126	56	19	25
7.80	404	16500	0	50	32F	1-3, 6h	92	A	0	13	183	44	0	56
8.00	412	16000	0	50	25M	1, 6h	25	A	0	0	N/A	44	4	52
8.30	429	15000	0	50	20M	1, 6h	20	A	5	5	261	25	0	75
8.30	429	15000	0	50	11F	1, 6h	11	A	0	0	N/A	27	0	73
8.50	440	14500	0	50	10M	1, 6h	10	A	0	0	N/A	10	20	70
9.00	465	13000	0	50	22M	1-2, 6h	23	A	0	0	N/A	23	14	64
9.50	493	11500	0	50	6M	1, 6h	6	A	0	0	N/A	0	17	83
9.50	493	11500	0	40	20M	1, 6h	20	A	0	0	N/A	0	20	80
9.50	493	11500	0	40	12F	1, 6h	12	A	0	0	N/A	0	0	100
9.50	493	11500	0	100	12F	1-5, 8h	54	A	0	0	N/A	0	0	100
9.50	493	11500	0	100	12M	4-5, 8h	59	A	0	0	N/A	0	0	100
10.00	517	10250	0	50	9M	1, 6h	9	A	0	0	N/A	0	22	78
10.50	543	9000	0	50	2M	1, 6h	2	A	0	0	N/A	0	0	100

- NOTE 1 -- Prebreathe with 100% oxygen
- NOTE 2 -- If <100% oxygen, the rest is nitrogen
- NOTE 3 -- Number of exposures per subject, duration of protocol exposure
- NOTE 4 -- Exercise Types: A - EVA Exercises law Dixon et al. (1986)  
 B - Bends Screening Exercise law Krutz & Dixon (1987)  
 C - Joint flexion only; no exercise
- NOTE 5 -- % of subjects with Grade 2 bends during any exposure
- NOTE 6 -- Mean time (min) to Grade 2 Bends for all cases of bends (not incl. delayed)
- NOTE 7 -- % of subjects with maximum Grade of bubbles indicated on any exposure
- NOTE 8 -- Each subject will experience up to 4 exposures for 3 or 6 h depending upon the protocol modification in effect at that date. Study in progress

Short Name of Each Study

- Study #1: Bends Screening Index  
 Study #2: 7.8 psia Male Study  
 Study #3: 7.8 psia Female Study  
 Study #4: Bubble Threshold Study  
 Study #5: 9.5 psia Bubble Threshold Validat.  
 Study #6: 8.3 psia Study  
 Study #7: 9.5 psia Oxygen Toxicity Study  
 Study #8: 5.46 psia, 3% CO2 Prebreathe Study

## APPENDIX B

### BIBLIOGRAPHY OF PUBLISHED RESULTS: 1983-1988

- Adams JD, Dixon GA, Harvey WT. Bends susceptibility. (Abstract) *Aviat Space Environ Med* 55:452 (1984).
- Baas CL, Olson RM, Dixon GA. Audio and visual ultrasonic monitoring of altitude decompression. Abstract accepted for 26th Annual SAFE Symposium, Las Vegas, NV, 1988.
- Baas CL, Dixon GA, Hill RC. Effect of a carbon dioxide prebreathe gas mixture on the incidence of altitude-induced decompression sickness. (Abstract; in press) Aerospace Medical Association. 1989 Annual Meeting, Washington, DC.
- Dixon GA, Adams JD, Harvey WT. Decompression sickness and intravenous bubble formation using a 7.8 psia simulated pressure-suit environment. *Aviat Space Environ Med* 57:223-228 (1986).
- Dixon GA, Krutz RW. Evaluation of 9.5 psia as a suit pressure for prolonged extravehicular activity. 23rd Annual SAFE Symposium Proceedings, Las Vegas, NV, 1985.
- Dixon GA, Krutz RW. Female susceptibility to decompression sickness and bubble formation using a simulated 7.8 psia suit environment. (Abstract) *Aviat Space Environ Med* 57:511 (1986).
- Dixon GA, Krutz RW Jr, Fischer JR. Decompression sickness and bubble formation in females exposed to a simulated 7.8 psia suit environment. *Aviat Space Environ Med* (In press).
- Inderbitzen RS, DeCarlis JJ Jr. Energy expenditure during simulated EVA workloads. SAE Technical Paper #860921. ICES, San Diego, CA, 1986.
- Krutz RW Jr, Dixon GA. The effects of exercise on bubble formation and bends susceptibility at 9,100 m (30,000 ft; 4.3 psia). *Aviat Space Environ Med* 58:A97-A99 (1987).
- Krutz RW Jr, Dixon GA, Harvey WT. Minimum pressure for a zero-prebreathe pressure suit. SAE Technical Paper #851315. ICES, San Francisco, CA, 1985.
- Krutz RW Jr, Webb JT, Dixon GA. Determining a bends-preventing pressure for a space suit. Abstract accepted for 26th Annual SAFE Symposium, Las Vegas, NV, 1988.
- Olson RM, Krutz RW Jr, Dixon GA, Smead KW. An evaluation of precordial ultrasonic monitoring to avoid bends at altitude. *Aviat Space Environ Med* 59:635-39 (1988).
- Olson RM, Krutz RW, Dixon GA. Validity of ultrasonic monitoring at altitude for bends detection. (Abstract) *Aviat Space Environ Med* 57:511 (1986).

- Smead KW. Preliminary findings: Bends screening index study. Space Life Sciences Symposium Abstracts (unpaginated, unindexed addendum), Washington, DC, 1987.
- Smead KW, Krutz Jr RW, Dixon G, Webb JT. Decompression sickness and venous gas emboli formation at 8.3 psia. 24th Annual SAFE Symposium Proceedings, San Antonio, TX, 1986.
- Webb JT, Olson RM, Dixon GA, Krutz RW, Barnicott PT. Oxygen toxicity during 5 simulated 8-h EVA exposures to 100% oxygen at 9.5 psia. SAE Technical Paper #881071. ICES, San Francisco, CA, 1988.
- Webb JT, Olson RM, Krutz RW Jr, Dixon GA, Barnicott PT. Human tolerance to 100% oxygen at 9.5 psia during five daily simulated eight-hour EVA exposures. Abstract accepted for 26th Annual SAFE Symposium, Las Vegas, NV, 1988. Aviat Space Environ Med (In press).
- Webb JT, Smead KW, Jauchem JR, Barnicott PT. Susceptibility to decompression-induced venous gas emboli: Hematology and biochemistry. Space Life Sciences Symposium Abstracts, Washington, DC, 1987.
- Webb JT, Smead KW, Jauchem JR, Barnicott PT. Blood factors and venous gas emboli: Surface to 429 mmHg (8.3 psi). Undersea Biomed Res 15:107-121 (1988).