

AD-A202 928

TIC
REPORT DOCUMENTATION PAGE

②

2a. SECURITY CLASSIFICATION AUTHORITY EGTE C. 1. 9 1988		1b. RESTRICTIVE MARKINGS DTIC FILE COPY			
2b. DECLASSIFICATION / DOWNGRADING SCHEDULE DCS		3. DISTRIBUTION / AVAILABILITY OF REPORT Approved for public release; Distribution unlimited			
4. PERFORMING ORGANIZATION REPORT NUMBER(S) AFOSR Contract No. F49620-86-C-0010		5. MONITORING ORGANIZATION REPORT NUMBER(S) AFOSR-TR- 88-1265			
6a. NAME OF PERFORMING ORGANIZATION University of Wisconsin	6b. OFFICE SYMBOL (if applicable)	7a. NAME OF MONITORING ORGANIZATION AFOSR/NC			
6c. ADDRESS (City, State, and ZIP Code) Department of Chemistry Madison, WI 53706		7b. ADDRESS (City, State, and ZIP Code) Building 410 Bolling AFB, D. C. 20332-6448			
8a. NAME OF FUNDING / SPONSORING ORGANIZATION AFOSR	8b. OFFICE SYMBOL (if applicable) NC	9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER Contract No. F49620-86-C-0010			
8c. ADDRESS (City, State, and ZIP Code) Building 410 Bolling AFB, D.C. 20332-6448		10. SOURCE OF FUNDING NUMBERS			
		PROGRAM ELEMENT NO. 61102F	PROJECT NO. 2303	TASK NO. B2	WORK UNIT ACCESSION NO.
11. TITLE (Include Security Classification) Chemical Reactions & Properties of Organosilicon Compounds Related to New Materials					
12. PERSONAL AUTHOR(S) Robert West					
13a. TYPE OF REPORT Final Technical	13b. TIME COVERED FROM 10/1/85 TO 9/30/88	14. DATE OF REPORT (Year, Month, Day)	15. PAGE COUNT 14		
16. SUPPLEMENTARY NOTATION					
17. COSATI CODES			18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)		
FIELD	GROUP	SUB-GROUP			
19. ABSTRACT (Continue on reverse if necessary and identify by block number)					
<p>I. Polysilane High Polymers and Oligomers. a) Light-scattering studies of molecular weight and chain conformations were carried out for six polysilane samples. b) Silicon-29 NMR spectroscopy was employed to study configurations of polysilane polymers and oligomers. c) A new class of substances, poly(disilanylene-co-ethynylene) copolymers, were synthesized which show properties indicating σ-π electron conjugation.</p>					
<p>II. Divalent and Multiply-Bonded Silicon Compounds. a) Complexes of silylenes, R_2Si, with various Lewis bases have been observed spectroscopically. b) Reactions of disilenes ($R_2Si=SiR_2$) with nitro compounds, nitroso compounds, azides and oxygen have been studied, leading to previously unknown types of cyclic silicon compounds. c) A new, very stable disilene, 1,2-dimesityl-1,2-bis(1-adamantyl)disilene, was synthesized and investigated. d) The first silandiimines, $RN=Si=NR$, have been made and observed at low temperatures. e) Stable silan-amidines, $RNH=Si(X)=NR$, and silanamide anions, $RN^{\ominus}=Si(X)^{\ominus}=NR^{\ominus}$, have been synthesized and studied.</p>					
20. DISTRIBUTION / AVAILABILITY OF ABSTRACT <input checked="" type="checkbox"/> UNCLASSIFIED/UNLIMITED <input checked="" type="checkbox"/> SAME AS RPT. <input type="checkbox"/> DTIC USERS			21. ABSTRACT SECURITY CLASSIFICATION Unclassified		
22a. NAME OF RESPONSIBLE INDIVIDUAL Dr. Anthony J. Matuszko			22b. TELEPHONE (Include Area Code) (202) 767-4960	22c. OFFICE SYMBOL NC	

DD FORM 1473, 84 MAR

83 APR edition may be used until exhausted.

All other editions are obsolete.

SECURITY CLASSIFICATION OF THIS PAGE

UNCLASSIFIED

Organic silicon compounds (organic)

Final Technical Report



CHEMICAL REACTIONS AND PROPERTIES OF ORGANOSILICON COMPOUNDS RELATED TO NEW MATERIALS

Period Covered: October 1, 1985 - September 30, 1988

In this report will be described the most significant findings from research under AFOSR sponsorship during the three-year period of this Contract. The areas to be covered are: I. Polysilane High Polymers and Oligomers; and II. Divalent and Multiply Bonded Silicon Compounds, including A. Silylenes, $RR'Si$; B. Disilenes, $RR'Si=SiRR'$; and C. Silanediimines, $RN=Si=NR$ and Silanamidines, $RNH-Si(X)=NR$.

I. Polysilane High Polymers and Oligomers

The polysilane polymers are linear polymers in which the "backbone" polymer chain is made up entirely of silicon atoms. They are generally synthesized by sodium condensation of diorganodichlorosilanes. Either homopolymers or copolymers can be made:




The first light-scattering studies of polysilanes were carried out, partially in collaboration with Drs. Robert Miller and Pat Cotts of IBM Company. The results show that the polysilanes have very high molecular weights, up to 7×10^6 in some samples, and have persistence lengths indicating that they behave as slightly stiffened random coils in solution.

Silicon-29 NMR spectroscopy has been used extensively by our group to study the stereochemistry of polysilane polymers. Polysilane homopolymers with the two organic groups on each silicon identical, such as $(n\text{-Hexyl}_2Si)_n$, show only a single sharp line in the ^{29}Si NMR, indicating that conformational effects on the silicon nucleus are rapidly averaged. Polymers with two different

alkyl groups, such as $(n\text{HexylSiCH}_3)_n$, show symmetrical multiplets of 5-7 closely-spaced lines. These spectra are consistent with a random, atactic arrangement of groups along the polysilane chain. Polymers which contain an aryl and an alkyl group, such as $(\text{PhMeSi})_n$, show three distinct multiplet bands, due to different, nonrandom distributions of stereoisomers along the chain.

In order to elucidate the stereochemical arrangement in $(\text{PhMeSi})_n$, the ^{29}Si NMR of model six-membered ring compounds, $\text{ttttt}-(\text{PhMeSi})_6$ and $\text{ttttcc}-(\text{PhMeSi})_6$, have been carefully investigated. The proton and ^{29}Si spectra for both molecules have been correlated by 2-dimensional NMR studies.

NMR investigations have also been used to determine the structure of some cyclic polysilanes. Cyclic permethylpolysilanes undergo aluminum chloride-catalyzed rearrangement to branched cyclic polysilanes with smaller ring size. For example, $(\text{Me}_2\text{Si})_7$ forms compound 1 under these conditions. The rearrangement of $(\text{Me}_2\text{Si})_8$ leads to a compound whose structure was uncertain between the alternatives 2 and 3:



The structure of rearranged $(\text{Me}_2\text{Si})_8$ was solved by NMR and shown to be 2. This experiment and a similar proof of structure for the rearrangement product from $(\text{Me}_2\text{Si})_9$, represent the first 2-dimensional silicon-silicon NMR spectra ever reported.

A new class of polymers has been synthesized: The disilanylene-ethynylene copolymers, 4a-c. The route of synthesis is shown below:



- 1) $2n\text{BuLi}$
- 2) $\text{ClSiR}'_2\text{SiR}'_2\text{Cl}$




4a, R = R' = CH₃

4b, R = R' = n-butyl

4c, R = CH₃, R' = n-butyl

These polymers are unique in that they show electron delocalization involving mixing of σ and π orbitals. Other electron-delocalized polymers involve only π orbitals (polyacetylenes, polyaniline, etc.) or only σ orbitals (polysilanes). Unusual electronic properties for the new polymers 4a-c are anticipated but remain to be explored.

If ethylmagnesium bromide is used in place of n-butyllithium in the second step of the equation shown above, the products are not polymers but instead are eight-membered rings having structure 5a-c. These novel structures are expected to lead to interesting new types of materials through further reactions.





5a, R = R' = CH₃

5b, R = R' = n-butyl

5c, R = CH₃, R' = n-butyl

As models for exploring the electronic properties of polysilanes, some linear oligomeric polysilanes have been synthesized. $\text{Me}(\text{SiMe}_2)_{10}\text{Me}$ and $\text{Me}(\text{SiMe}_2)_{16}\text{Me}$ have been made by the following reactions:




The structure and conformations of these linear oligomers are being studied.

QUALITY
CHECKED
2

Date	
Prepared by Dates	
Dist	No. and/or Serial
A-1	

B. Disilenes, $R_2Si=SiR_2$


The first compound containing a silicon-silicon double bond, tetramesityldisilene (8), was synthesized at the University of Wisconsin in 1981. Under this contract the chemistry of disilenes has been further explored. The disilenes are synthesized by photolysis of trisilanes. The new, highly stable disilene 9 was obtained by this procedure. Compound 9 survives heating to 275°C and is nearly inert to oxygen.




Mes = mesityl; Ad = 1-adamantyl

The activation energy for cis-trans interconversion of 9 was measured by NMR spectroscopy, and found to be 28 kcal mol⁻¹. The crystal structure of 9 has been partially solved by x-ray diffraction.

New reactions of disilenes discovered under the contract include the reaction with nitrobenzene, to give 10; and the reaction with nitrobenzene to give 11, which undergoes further oxidation to 10.



The oxidation of disilenes has been further elucidated. The reaction pathways are illustrated below:




The oxidation products have novel structures with short Si-Si bonds, suggesting possible pi-bonding between the silicons.

The first electrochemical studies of disilenes have been carried out by means of cyclic voltammetry. Oxidation potentials for the disilenes are quite low (+0.5 v) and vary little with substitution at silicon. Reduction potentials are fairly negative, and vary with substituent, indicating that conjugation with aromatic rings influences the energy of the lowest unoccupied molecular orbital in disilenes.


C. Silanediimines, $\text{RN}=\text{Si}=\text{NR}$, and Silanamidines, $\text{RNH}-\text{SiX}=\text{NR}$

Organosilicon compounds in which silicon forms two double bonds to other atoms were completely unknown before the beginning of this contract. Two examples have now been obtained, although both are stable only at low temperatures.


The first silanamidine was obtained by photolysis of bis(trimethylsilyl)silyl diazide, 12, in hydrocarbon matrix at 77K. The first product is the silanimine 13, which can be trapped by chemical reaction. Further photolysis converts 13 to the silanediimine 14.



Photolysis of di-tert-butylsilyl diazide also leads to a silanediimine, but in this case the major product is the silylene, 15:



Recently, we have been attempting to make a stable silanediimine. These experiments have led to the synthesis of the first silanamidines (17). The starting compound is the silanediamine 16, made from tri-tert-butylaniline and silicon tetrahalides.



Treatment of 16 with *n*-butyllithium affords silanamidine 17, while deprotonation of 16 with $\text{KN}(\text{SiMe}_3)_2$ yields the silanamidine anion, 18. These are the first examples of each of these structural types. Loss of halide from 18 should provide the silandiimine; experiments to bring about this reaction are continuing.

COMPLETED PROJECT SUMMARY

1. **TITLE:** Chemical Reactions and Properties of Organosilicon Compounds Related to New Materials
2. **PRINCIPAL INVESTIGATOR:** Robert West
Department of Chemistry
University of Wisconsin
Madison, WI 53706
3. **INCLUSIVE DATES:** October 1, 1985 - September 30, 1988
4. **CONTRACT NOS.:** F49620-86-C-0010; F49620-86-C-0010, Mod. 1
5. **COSTS AND FY SOURCE:** \$222,397, FY85; \$222,658, FY86;
\$241,093, FY87

6. **SENIOR RESEARCH PERSONNEL:**

<u>Name</u>	<u>Present Affiliation</u>
Mary-Ann Pearsall	Drew University, Madison, NJ
Akira Sekiguchi	Tohoku University, Sendai, Japan
Takahisa Iwahara	Kanegafuchi Chem. Industries, Kobe, Japan
Johannes Belzer	University of Wisconsin, Madison, WI
Yuxia Liu	University of Wisconsin, Madison, WI
Li-Ming Huang	University of Wisconsin, Madison, WI
Yoshitaka Hamada	University of Wisconsin, Madison, WI

7. **JUNIOR RESEARCH PERSONNEL:**

Kevin Welsh	Columbia University, New York, NY
Steven S. Zigler	duPont Co., Wilmington, DE
Gregory Jamison	Univ. of North Carolina, Durham, NC
Howard Yokelson	Amoco Co., Naperville, IL
Jim Maxka	Technion, Haifa, Israel*
Gregory Gillette	Univ. of Toulouse, Toulouse, France*
Brian Shepherd	General Electric Co., Schenectady, NY*
Gail Underiner	University of Wisconsin, Madison, WI
Eric Pham	University of Wisconsin, Madison, WI

JUNIOR RESEARCH PERSONNEL (continued)

Alan Fanta	University of Wisconsin, Madison, WI
Tetsuya Asuke	University of Wisconsin, Madison, WI
Tony Millevolte	University of Wisconsin, Madison, WI
Jeff Cavalieri	University of Wisconsin, Madison, WI
Scott Archibald	University of Wisconsin, Madison, WI

*Locations as of January 1, 1989

8. PUBLICATIONS:

1. X-Ray Crystal Structure and Conformational Analysis of Tetradecamethylcycloheptasilane, $(\text{Me}_2\text{Si})_7$, F. Shafiee, J. R. Damewood, Jr., K. J. Haller and R. West, *J. Am. Chem. Soc.*, 107, 6950-6956 (1985).
2. Polysilane High Polymers with Olefinic Side Groups: Synthesis, Properties and Addition of Hydrogen Halides, H. Stuger and R. West, *Macromolecules*, 18, 2349-2352 (1985).
3. Study of the Reaction of Dimethyldichlorosilane with Lithium for Synthesis of $(\text{Me}_2\text{Si})_n$ by Gas Chromatography, S.-M. Chen and R. West, *Huaxue Xuebao*, 43, 365-368 (1985).
4. Metal Halide Catalyzed Rearrangements of Alkylcyclosilanes, T. A. Blinka and R. West, *Organometallics*, 5, 128-133 (1986).
5. Conformational Analysis of Branched Cyclosilanes, T. A. Blinka and R. West, *Organometallics*, 5, 133-139 (1986).
6. The Polysilane High Polymers, R. West, *J. Organomet. Chem.*, 300, 327-346 (1986).
7. IR Transition Moment Directions in Matrix-Isolated Dimethylsilylene and 1-Methylsilene, G. Raabe, H. Vancik, R. West and J. Michl, *J. Am. Chem. Soc.*, 108, 671-677 (1986).
8. Structural and Chemical Properties of 1,3-Cyclodisiloxanes, M. J. Michalczyk, M. J. Fink, K. J. Haller, R. West and J. Michl, *Organometallics*, 5, 531-538 (1986).
9. Cycloaddition of Stable Disilenes to Terminal Acetylenes, D. J. De Young and R. West, *Chem. Lett.*, 6, 883-884 (1986).
10. Polysilanes as Photoinitiators for Vinyl Polymerization, R. West, A. R. Wolff and D. J. Peterson, *J. Radiat. Curing*, 13, 35-40 (1986).
11. Polysilane High Polymers and Their Technological Applications, R. West, *L'Actualite Chimique*, 3, 64-70 (1986).

12. Possible π -Complex Intermediates in the Reaction of Disilenes with Mercury(II) Trifluoroacetate, C. Zybilla and R. West, *J. Chem. Soc., Chem. Commun.*, 857 (1986).
13. Crosslinking of Polysilanes as Silicon Carbide Precursors, R. West, X-H. Zhang, P. I. Djurovich and H. Stuger, in Science of Ceramic Chemical Processing, L. L. Hench and D. R. Ulrich, Eds., Wiley, New York, pp 337-344 (1986).
14. ^{29}Si NMR Observation of an Unprecedented Rearrangement in Tetraaryldisilenes, H. B. Yokelson, J. Maxka, D. A. Siegel and R. West, *J. Am. Chem. Soc.*, 108, 4239-4242 (1986).
15. A Synthon for the Silicon-Silicon Triple Bond, A. Sekiguchi, S. S. Zigler, R. West and J. Michl, *J. Am. Chem. Soc.*, 108, 4241-4242 (1986).
16. Observation of a Silanimine in an Inert Matrix and in Solution at Low Temperature, S. S. Zigler, R. West and J. Michl, *Chem. Lett.*, 1033 (1986).
17. X-Ray Crystal and Molecular Structures of Hexacosamethylcyclo-tetra decasilane $(\text{Me}_2\text{Si})_{13}$ and Dotriacontamethylcyclohexa-decasilane, $(\text{Me}_2\text{Si})_{16}$, F. Shafiee, K. J. Haller and R. West, *J. Am. Chem. Soc.*, 108, 5478-5482 (1986).
18. Trapping of Silylenes by 9,10-Dimethylantracene: 2,3,5,6-Dibenzo-7-silabicyclo[2.2.1]hepta-2,5-dienes, A. Sekiguchi and R. West, *Organometallics*, 5, 1911-1913 (1986).
19. Electrochemical Oxidation of Cyclopolysilanes Using Cyclic Voltammetry, F. Shafiee and R. West, *Silicon, Germanium, Tin & Lead Compds.*, 9, 1-10 (1986).
20. Organosilylenes and Their Dimerization to Disilenes, M. J. Michalczyk, M. J. Fink, D. J. De Young, C. W. Carlson, K. M. Welsh, R. West and J. Michl, *Silicon, Germanium, Tin & Lead Compds.*, 9, 75-80 (1986).
21. Rearrangements in Mass Spectrometry of Cyclosilanes, T. A. Blinka and R. West, *Silicon, Germanium, Tin & Lead Compds.*, 9, 82-85 (1986).
22. Soluble Polysilanes: An Interesting New Class of Radiation Sensitive Materials, R. D. Miller, D. Hofer, G. N. Fickes, C. G. Willson, E. Marinero, P. Trefonas III and R. West, *Polym. Engr. Sci.*, 26, 1129-1134 (1986).
23. Synthesis and Crosslinking of Liquid Polysilane Polymer, Z-H. Zhang and R. West, *Gaofenzi Tongxun*, 4, 257-263 (1986) (China).

24. A Probe for Substituent Hyperconjugative Power: MCD (Magnetic Circular Dichroism) of the Benzene L_b Band, G. W. Weeks, W. Adcock, K. A. Klingensmith, J. W. Waluk, R. West, M. Vasak, J. Downing and J. Michl, *Pure Appl. Chem.*, 58, 1, 39-53 (1986).
25. Photoinitiation of Vinyl Polymerization by Polysilanes, A. R. Wolff and R. West, *Applied Organomet. Chem.*, 1, 7-14 (1987).
26. (7-Silanorbornadienyl)cyclopentadienyldicarbonyl Iron Complexes: An Approach to Iron Substituted Silylenes, A. Marinetti-Magnani and R. West, *Organometallics*, 6, 141-144 (1987).
27. The Generation of Hexamethyl-1,4-Disilabenzene and Its Novel Thermal Chemistry, K. M. Welsh, J. D. Rich, J. Michl and R. West, *J. Organomet. Chem.*, 325, 105-115 (1987).
28. The Bonding in 1,3-Cyclodisiloxanes: ^{29}Si NMR Coupling Constants in Disilenes and 1,3-Cyclodisiloxanes, H. B. Yokelson, A. J. Millevolte, B. R. Adams and R. West, *J. Am. Chem. Soc.*, 109, 4116-4118 (1987).
29. Matrix Isolation of the First Silanediimine, N,N'-bis(trimethylsilyl)silanediimine, S. S. Zigler, K. M. Welsh and R. West, *J. Am. Chem. Soc.*, 109, 4392-4393 (1987).
30. The Addition Reactions of Two Disilenes, D. J. De Young, M. J. Fink, J. Michl and R. West, *Main Group Metal Chem.* 1, 19-43 (1987).
31. Disilaoxiranes: Synthesis and Crystal Structure, H. B. Yokelson, A. J. Millevolte, G. R. Gillette and R. West, *J. Am. Chem. Soc.*, 109, 6865-6866 (1987).
32. The Synthesis and Molecular Structure of a Disilacyclopropanimine, H. B. Yokelson, A. J. Millevolte, K. J. Haller and R. West, *J.C.S. Chem. Comm.*, 21, 1605-1606 (1987).
33. Light Scattering from Solutions of Organosilane Polymers, P. M. Cotts, R. D. Miller, P. T. Trefonas III, R. West and G. Fickes, *Macromolecules*, 20, 1046-1052 (1987).
34. Spectroscopic Observation of Silylene-Ether Complexes, G. R. Gillette, G. H. Noren and R. West, *Organometallics*, 6, 2617-2618 (1987).
35. Chemistry of the Silicon-Silicon Double Bond, R. West, *Angew. Chem.Int. Ed. Eng.*, 26, 1201-1211 (1987).
36. Polysilane High Polymers - An Overview, R. West and J. Maxka in *Inorganic and Organometallic Polymers*, M. Zeldin, K. J. Wynne and H. R. Allcock, Eds., ACS Sym. Series 360, Ch. 2, pp. 6-20 (1988).

37. Cyclic Voltammetric Investigation of Disilenes, B. D. Shepherd and R. West, *Chem. Lett.*, 2, 183-186 (1988).
38. ^{29}Si NMR of Dimethyl and Phenylmethyl Containing Polysilanes, A. R. Wolff, I. Nozue, J. Maxka and R. West, *J. Polym. Sci., Polym. Chem. Ed.*, 26, 701-713 (1988).
39. ^{29}Si NMR of Dialkylpolysilanes, A. R. Wolff, J. Maxka and R. West, *J. Polym. Sci., Polym. Chem. Ed.*, 26, 713-720 (1988).
40. Preparation and Reactions of a Disilyne Synthone, Di(2,3-benzo-7-methyl-1,4,5,6-tetraphenyl-7-silanorbornadien-7-yl), A. Sekiguchi, S. S. Zigler, K. J. Haller and R. West, *Recl. Trav. Chim. Pays-Bas*, 107, 197-202 (1988).
41. Oxidation of Disilenes with Atmospheric Oxygen: A Status Report, R. West, H. B. Yokelson, G. R. Gillette, A. J. Millevolte, in *Silicon Chemistry*, J. Y. Corey, E. R. Corey and P. P. Gaspar, Eds., Ellis Horwood, Ch. 26, pp. 269-281 (1988).
42. A New Route to 1,4-Disilabenzenes and 1,4-Disilabarrelenes, A. Sekiguchi, G. R. Gillette and R. West, *Organometallics*, 7, 1226-1227 (1988).
43. Dodecamethylcyclohexasilane, S.-M. Chen and R. West, *Organomet. Synth.*, 4, 506-507 (1988).
44. E-1,2-Dimesityl-1,2-di(*tert*-butyl)disilene, G. R. Gillette, H. B. Yokelson and R. West, *Organomet. Synth.*, 4, 529-533 (1988).
45. Organosilane High Polymers: Poly(phenylmethylsilylene), P. Trefonas and R. West, *Inorg. Synth.*, 25, 58-62 (1988).
46. Generation of Silanimines by Photolysis of Hindered Azidosilanes, S. S. Zigler, L. M. Johnson and R. West, *J. Organomet. Chem.*, 341, 187 (1988).
47. A New Route to Strained Cyclic Disilanylene-Acetylenes, T. Iwahara and R. West, *J. C. S. Chem. Comm.*, 14, 954-955 (1988).
48. The Photochemistry of Matrix Isolated Di-*tert*-Butyldiazidosilane, K.M. Welsh, J. Michl and R. West, *J. Am. Chem. Soc.*, 110, 6689-6696 (1988).
49. The Reactions of Diorganosilylenes with Carbon Monoxide, M.-A. Pearsall and R. West, *J. Am. Chem. Soc.*, 110, 7228-7229 (1988).

IN PRESS

- A. Formation of Cyclic Silicon-Silicon Systems, R. West in *Inorganic Reactions and Methods*, J. J. Zuckerman, Ed., Verlag Chemie.

- B. Trapping of Silylenes by 9,10-Dimethylanthracene: 2,3,5,6-Dibenzo-7-silabicyclo[2.2.1]hepta-2,5-dienes, A. Sekiguchi and R. West, Organometallics.
- C. Polysilanes, R. West in The Chemistry of Silicon Compounds, S. Patai and Z. Rappoport, Eds., Wiley.
- D. Lewis Base Adducts to Diorganosilylenes, G. R. Gillette, G. H. Noren and R. West, Organometallics.
- E. Synthesis of the Novel Disila-1,2-oxazetidene and Disila-1,2,4-dioxazolidine Ring Systems, G. R. Gillette, J. Maxka and R. West, Angewandte Chemie.

Submitted

- F. High-Yield Synthesis and Crystal Structure of a Novel Benzene Tris-Annulated with Bicyclo[2.2.2]octene, K. Komatsu, Y. Jinbu, G. R. Gillette and R. West, Chem. Letters.
- G. Three-, Four- and Five-Membered Rings from Disilenes, R. West, G. R. Gillette, H. B. Yokelson and A. J. Millevolte, Phosphorus, Sulfur, Silicon and Related Elements.
- H. 1,2-Aryl Rearrangement in Tetraaryldisilenes, H. B. Yokelson, D. A. Siegel, A. J. Millevolte, J. Maxka and R. West, J. Am. Chem. Soc.
- I. Organosilane Polymers: Synthesis and Crosslinking of Formable Polymers Containing Phenylsilylene Units, X-H. Zhang and R. West, Macromolecules.
- J. Conformational Energies and Unperturbed Chain Dimensions of Poly-(phenylmethylsilylene) [-SiPhMe-] and Poly(silastyrene) [-SiPhH-SiH₂-], W. J. Welsh, J. R. Damewood, Jr. and R. West, Macromolecules.
- K. Structures of Two Organosilyl Azides, S. S. Zigler, K. J. Haller, R. West and M. S. Gordon, Organometallics.
- L. Tetramesityldisilene, R. West, G. R. Gillette and H. B. Yokelson, Inorg. Synth.
- M. Use of 2-D INEPT-INADEQUATE ²⁹Si NMR to Determine Structures of Organosilicon Rings, J. Maxka, B. R. Adams and R. West, J. Am. Chem. Soc.

9. ABSTRACT OF OBJECTIVES AND ACCOMPLISHMENTS:

I. Polysilane High Polymers and Oligomers. a) Light-scattering studies of molecular weight and chain conformations were carried out for six polysilane samples. b) Silicon-29 NMR spectroscopy was employed to study configurations of polysilane polymers and oligomers. c) A new class of substances, poly(disilanylene-co-ethynylene) copolymers, were synthesized which show properties indicating σ - π electron conjugation.

II. Divalent and Multiply-Bonded Silicon Compounds. a) Complexes of silylenes, $R_2Si:$, with various Lewis bases have been observed spectroscopically. b) Reactions of disilenes ($R_2Si=SiR_2$) with nitro compounds, nitroso compounds, azides and oxygen have been studied, leading to previously unknown types of cyclic silicon compounds. c) A new, very stable disilene, 1,2-dimesityl-1,2-bis(1-adamantyl)disilene, was synthesized and investigated. d) The first silandiimines, $RN=Si=NR$, have been made and observed at low temperatures. e) Stable silanamidines, $RNH-Si(X)=NR$, and silanamidine anions, $RN^{\ominus}-Si(X)=NR^{\ominus}$, have been synthesized and studied.

Approved for public release
Distribution unlimited

1970-1971
1972-1973
1974-1975
1976-1977
1978-1979
1980-1981
1982-1983
1984-1985
1986-1987
1988-1989
1990-1991
1992-1993
1994-1995
1996-1997
1998-1999
2000-2001
2002-2003
2004-2005
2006-2007
2008-2009
2010-2011
2012-2013
2014-2015
2016-2017
2018-2019
2020-2021
2022-2023
2024-2025