

AD-A204 489 MENTATION PAGE

Form Approved
OMB No. 0704-0188


1a. F		1b. RESTRICTIVE MARKINGS OTIC FILE COPY	
2a. SECURITY CLASSIFICATION AUTHORITY N/A		3. DISTRIBUTION / AVAILABILITY OF REPORT Approved for Public Release; Distribution Unlimited	
2b. DECLASSIFICATION / DOWNGRADING SCHEDULE N/A		5. MONITORING ORGANIZATION REPORT NUMBER(S) AFOSR-TR- 89-0121	
4. PERFORMING ORGANIZATION REPORT NUMBER(S) Technical Report No. 247		7a. NAME OF MONITORING ORGANIZATION AFOSR/NM	
6a. NAME OF PERFORMING ORGANIZATION University of North Carolina Center for Stochastic Processes	6b. OFFICE SYMBOL (If applicable)	7b. ADDRESS (City, State, and ZIP Code) Bldg. 410 Bolling Air Force Base, DC 20332-6448	
6c. ADDRESS (City, State, and ZIP Code) Statistics Department CB #3260, Phillips Hall Chapel Hill, NC 27599-3260		9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER F49620 85C 0144	
8a. NAME OF FUNDING / SPONSORING ORGANIZATION AFOSR	8b. OFFICE SYMBOL (If applicable) NM	10. SOURCE OF FUNDING NUMBERS	
8c. ADDRESS (City, State, and ZIP Code) Bldg. 410 Bolling AFB, DC 20332-6448		PROGRAM ELEMENT NO. 6.1102F	PROJECT NO. 2304
		TASK NO. A5	WORK UNIT ACCESSION NO.
11. TITLE (Include Security Classification) On the rate of convergence in Strassen's functional law of the iterated logarithm			
12. PERSONAL AUTHOR(S) Mijnheer, J.			
13a. TYPE OF REPORT preprint	13b. TIME COVERED FROM <u>Sep 88</u> TO <u>Sep 87</u>	14. DATE OF REPORT (Year, Month, Day) 1988, Sept.	15. PAGE COUNT 20
16. SUPPLEMENTARY NOTATION N/A			
17. COSATI CODES		18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)	
FIELD	GROUP	rate of convergence, functional limit theorem	
XXXXXX	XXXXXXXXXXXXXXX		
19. ABSTRACT (Continue on reverse if necessary and identify by block number)			
<p>→ An improvement of the rate of convergence in the functional law of the iterated logarithm (F.L.I.L) is given. <i>Keywords: Brownian motion, Wiener process, continuous functions. (ker) ←</i></p>			
20. DISTRIBUTION / AVAILABILITY OF ABSTRACT <input checked="" type="checkbox"/> UNCLASSIFIED/UNLIMITED <input type="checkbox"/> SAME AS RPT. <input type="checkbox"/> DTIC USERS		21. ABSTRACT SECURITY CLASSIFICATION UNCLASSIFIED	
22a. NAME OF RESPONSIBLE INDIVIDUAL Major Brian W. Woodruff		22b. TELEPHONE (Include Area Code) (202) 767-5026	22c. OFFICE SYMBOL AFOSR/NM

DTIC ELECTED
21 FEB 1989
S E D

APOSR-TR. 89-0121

CENTER FOR STOCHASTIC PROCESSES

Department of Statistics
University of North Carolina
Chapel Hill, North Carolina



ON THE RATE OF CONVERGENCE IN STRASSEN'S FUNCTIONAL LAW
OF THE ITERATED LOGARITHM

by

Joop Mijneer

Technical Report No. 247

September 1988

89 2 16 073

179. E. Frigola, Remark on the multiple Wiener integral, Mar. 87.
180. E. Frigola, Stochastic filtering solutions for ill-posed linear problems and their extension to measurable transformations, Mar. 87.
181. G. Samorodnitsky, Maxima of symmetric stable processes, Mar. 87.
182. H.L. Hurd, Representation of harmonizable periodically correlated processes and their covariance, Apr. 87. *J. Multivariate Anal.*, to appear.
183. H.L. Hurd, Nonparametric time series analysis for periodically correlated processes, Apr. 87. *IEEE Trans. Information Theory*, to appear.
184. T. Mori and H. Oodaira, Freidlin-Wentzell estimates and the law of the iterated logarithm for a class of stochastic processes related to symmetric statistics, May 87. *Yokohama Math. J.*, to appear.
185. R.F. Serfozo, Point processes, May 87. *Operations Research Handbook on Stochastic Processes*, to appear.
186. Z.D. Bai, W.Q. Liang and W. Vervaat, Strong representation of weak convergence, June 87.
187. O. Kallenberg, Decoupling identities and predictable transformations in exchangeability, June, 87.
188. O. Kallenberg, An elementary approach to the Danielli-Kolmogorov theorem and some related results, June 87. *Math. Nachr.*, to appear.
189. G. Samorodnitsky, Extrema of skewed stable processes, June 87. *Stochastic Proc. Appl.*, to appear.
190. D. Kualart, M. Sanz and M. Zaki, On the relations between increasing functions associated with two-parameter continuous martingales, June 87.
191. F. Avram and M. Taqqu, Weak convergence of sums of moving averages in the α -stable domain of attraction, June 87.
192. M.R. Leadbetter, Harald Cramér (1893-1985), July 87. *ISI Review*, 56, 1988, 89-97.
193. R. LePage, Predicting transforms of stable noise, July 87.
194. R. LePage and B.M. Schreiber, Strategies based on maximizing expected log, July 87.
195. J. Rosinski, Series representations of infinitely divisible random vectors and a generalized shot noise in Banach spaces, July 87. *Ann. Probability*, to appear.
196. J. Szulga, On hypercontractivity of α -stable random variables, $0 < \alpha < 2$, July 87.
197. I. Kuznezova-Shoipo and S.T. Rachev, Explicit solutions of moment problems I, July 87. *Probability Math. Statist.*, 10, 1989, to appear.
198. T. Hsing, On the extreme order statistics for a stationary sequence, July 87. *Stochastic Proc. Appl.*, 29, 1988, 155-169.
199. T. Hsing, Characterization of certain point processes, Aug. 87. *Stochastic Proc. Appl.*, 26, 1987, 297-316.
200. J.P. Nolan, Continuity of symmetric stable processes, Aug. 87. *J. Multivariate Anal.*, 1988, to appear.
201. M. Marques and S. Cambanis, Admissible and singular translates of stable processes, Aug. 88.
202. O. Kallenberg, One-dimensional uniqueness and convergence criteria for exchangeable processes, Aug. 87. *Stochastic Proc. Appl.*, 28, 1988, 159-183.
203. R.J. Adler, S. Cambanis and G. Samorodnitsky, On stable Markov processes, Sept. 87.
204. G. Kallianpur and V. Perez-Abreu, Stochastic evolution equations driven by nuclear space valued martingales, Sept. 87. *Appl. Math. Optimization*, 17, 1988, 237-272.
205. R.L. Smith, Approximations in extreme value theory, Sept. 87.
206. E. Willekens, Estimation of convolution tails, Sept. 87.
207. J. Rosinski, On path properties of certain infinitely divisible processes, Sept. 87. *Stochastic Proc. Appl.*, to appear.
208. A.H. Koreziloglu, Computation of filters by sampling and quantization, Sept. 87.
209. J. Bather, Stopping rules and observed significance levels, Sept. 87. *Sequential Anal.*, to appear.
210. S.T. Rachev and J.E. Yukich, Convolution metrics and rates of convergence in the central limit theorem, Sept. 87. *Ann. Probability*, 1988, to appear.
211. M. Fujisaki, Normed Bellman equation with degenerate diffusion coefficients and its applications to differential equations, Oct. 87.
212. G. Simons, Y.C. Yao and X. Wu, Sequential tests for the drift of a Wiener process with a smooth prior, and the heat equation, Oct. 87.
213. R.L. Smith, Extreme value theory for dependent sequences via the Stein-Chen method of Poisson approximation, Oct. 87.
214. C. Houdré, A vector bimeasure integral with some applications, June 88 (Revised).
215. M.R. Leadbetter, On the exceedance random measures for stationary processes, Nov. 87.
216. M. Marques, A study on Lebesgue decomposition of measures induced by stable processes, Nov. 87 (Dissertation).
217. M.T. Alpuim, High level exceedances in stationary sequences with extremal index, Dec. 87. *Stochastic Proc. Appl.*, to appear.
218. R.F. Serfozo, Poisson functionals of Markov processes and queueing networks, Dec. 87.
219. J. Bather, Stopping rules and ordered families of distributions, Dec. 87. *Sequential Anal.*, 7, 1988, 111-126.
220. S. Cambanis and M. Maejima, Two classes of self-similar stable processes with stationary increments, Jan. 88.
221. H.P. Huckle, G. Kallianpur and R.L. Karandikar, Smoothness properties of the conditional expectation in finitely additive white noise filtering, Jan. 88. *J. Multivariate Anal.*, to appear.

ON THE RATE OF CONVERGENCE IN STRASSEN'S FUNCTIONAL LAW
OF THE ITERATED LOGARITHM

by

Joop Mijnheer
Department of Mathematics
University of Leiden
The Netherlands

and

Center for Stochastic Processes
Department of Statistics
University of North Carolina
Chapel Hill, NC 27599-3260 USA

Accession For	
NTIS GRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input checked="" type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By _____	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-1	

Summary

An improvement of the rate of convergence in the functional law of the iterated logarithm (F.L.I.L) is given.



AMS Mathematics subject classification: 60J65, 60F17.

Key words and phrases: rate of convergence, functional limit theorem.

Research partially supported by the Air Force Office of Scientific Research
Contract No. F49620 85C 0144.

1. Introduction.

Let $\{W(t): 0 \leq t < \infty\}$ be a Brownian motion or a Wiener process on (Ω, \mathcal{F}, P) . We define the functions $\{f_T: T \geq 20\}$

$$f_T: [0, 1] \times \Omega \rightarrow \mathbb{R}$$

by

$$f_T(t, \omega) = W(Tt, \omega) (2T \log \log T)^{-1/2}.$$

Let K be the subset of absolutely continuous functions $x \in C[0, 1]$ such that

$$x(0) = 0$$

and

$$\int_0^1 \{\dot{x}(t)\}^2 dt \leq 1.$$

The functional law of the iterated logarithm (F.L.I.L.) states

Theorem. (Strassen 1964) W.p.1. $\{f_n; n \in \mathbb{N}\}$ is relatively compact with limit set K .

Two results about the rate of convergence are known.

$$P(d_\infty(f_n, K) \geq (\log \log n)^{-\alpha} \text{ i.o.}) = \begin{cases} 1 & \text{according as } \alpha \geq 1, \\ 0 & \alpha < 1/2. \end{cases}$$

(Bolthausen 1978) and the following sharpening of the foregoing result.

$$P(d_\infty(f_T, K) \geq (\log \log T)^{-\alpha} \text{ i.o.}) = \begin{cases} 1 & \alpha > 2/3, \\ 0 & \text{according as } \alpha < 2/3. \end{cases}$$

(Grill 1987). The distance d_∞ is the usual distance in $C[0, 1]$ and will be

defined in section 2.

The main result of this paper will be an improvement of this last result.

Theorem. Let $\{f_n\}$ and K be defined as above. Then

$$P(d_\omega(f_n, K) \geq \psi_\delta(n) \text{ i.o.}) = \begin{cases} 1 & \delta < 0, \\ 0 & \delta > 0, \end{cases} \text{ according as}$$

where

$$\psi_\delta(n) = \frac{1}{6} (1+\delta)(\log_3 n)(\log_2 n)^{-2/3}.$$

A simple proof of Strassen's F.L.I.L. is given in Chover (1967). A detailed proof can be found in the book of Freedman (1971). An extensive discussion of the different formulations of functional laws of the iterated logarithm is given in Taqqu and Czado (1985). In section 5 we compare the approach in this paper with several other approaches.

2. Some results of Gaussian processes.

The set K in Strassen's F.L.I.L. is the unit ball of the Hilbert space

$$H = \{f: [0,1] \rightarrow \mathbb{R}, f(t) = \int_0^t \dot{f}(s)ds, \int_0^1 \{\dot{f}(s)\}^2 ds < \infty\}$$

with inner product

$$(f, g) = \int_0^1 \dot{f}(s)\dot{g}(s)ds, \quad f, g \in H.$$

The sequence $\{\varphi_n: n=0,1,\dots\}$, where

$$\varphi_n(t) = \frac{2\sqrt{2}}{\pi(2n+1)} \sin((2n+1)\pi t/2) \quad 0 \leq t \leq 1,$$

is a complete orthonormal system in H . Let $\{X_n: n=0,1,\dots\}$ be a sequence of i.i.d. $N(0,1)$ distributed random variables. The Karhunen-Loève expansion of the Brownian motion $\{W(t): 0 \leq t \leq 1\}$ states

$$(2.1) \quad W(t) = \sum_{n=0}^{\infty} X_n \varphi_n(t).$$

For more details see Loève (1963) or Jain and Marcus (1978). In theorem 3.3 of the last paper they prove that the expansion (2.1) converges uniformly a.s. The space H is also used by Kuelbs and LePage (1973) in order to prove functional laws.

In our theorem we use two norms. The sup norm in $C[0,1]$

$$\|f\|_{\infty} = \sup_{0 \leq t \leq 1} |f(t)|$$

and the norm in H

$$\|f\|_H = \left\{ \int_0^1 \{\dot{f}(s)\}^2 ds \right\}^{1/2}.$$

We have the following relation between these two norms

$$\|f\|_{\infty} \leq \|f\|_H.$$

The corresponding metrics are denoted by d_{∞} and d_H . For each natural number m we define the (Gaussian) processes $\{U_m(t): 0 \leq t \leq 1\}$ and $\{V_m(t): 0 \leq t \leq 1\}$ by

$$V_m(t) = \sum_{n=0}^{m-1} X_n \varphi_n(t)$$

and

$$U_m(t) + V_m(t) = W(t),$$

i.e.

$$U_m(t) = \sum_{n=m}^{\infty} X_n \varphi_n(t).$$

Then we have

$$EU_m(t) = EV_m(t) = 0,$$

$$\sigma^2(U_m(t)) = \sum_{n=m}^{\infty} \varphi_n^2(t) \quad \text{and} \quad \sigma^2(V_m(t)) = \sum_{n=0}^{m-1} \varphi_n^2(t).$$

In the case $t=1$ we have for $m \rightarrow \infty$

$$(2.2a) \quad \sigma^2(V_m(1)) = 2\pi^{-2} \sum_{n=0}^{m-1} 4(2n+1)^{-2} = 1 - \frac{2}{\pi^2 m} + o(m^{-2})$$

and

$$(2.2b) \quad \sigma^2(U_m(1)) = \frac{2}{\pi^2 m} + o(m^{-2}).$$

We also have

$$\|V_m\|_H^2 = 2 \sum_{n=0}^{m-1} X_n^2 \int_0^1 \cos^2\{(2n+1)\pi t/2\} dt = \sum_{n=0}^{m-1} X_n^2.$$

Thus $\|V_m\|_H^2$ has a chi-square distribution with m degrees of freedom.

We note that U_m and V_m are independent Gaussian processes. There exists a rich literature on the maximum of Gaussian processes. See for example Berman (1985), Talagrand (1988), Piterbarg and Prisjaznjuk (1978) and the references in those papers. We shall not use the results out of one of these papers but prove the following lemma. Because the processes U_m and V_m have such a nice structure we give new proofs. (Of course making use of the ideas of the other papers.) Note that the processes have no independent increments. It is easy to see that the variance takes its maximum value for $t=1$. The processes will reach their maximum near $t=1$, as we can conclude from the following lemma. In this lemma we shall compare the tail of distribution of the maximum with the tail of the distribution of the process at $t=1$.

Lemma 2.1. Let the Gaussian processes U_m and V_m be defined as above. Let $\epsilon > 0$. Then we have for $m, u \rightarrow \infty$ and $m = o(u)$

$$(2.3.a) \quad P\left(\max_{0 \leq t \leq 1} V_m(t) > u\right) \leq (1 + o(1))P(V_m(1) > u(1-\epsilon))$$

and

$$(2.3.b) \quad P\left(\max_{0 \leq t \leq 1} U_m(t) > u\right) \leq (1 + o(1))P(U_m(1) > u(1-\epsilon)).$$

The distribution of the maximum of a Brownian motion is given by

$$(2.4) \quad P\left(\sup_{0 \leq t \leq 1} W(t) > u\right) = 2P(W(1) > u)$$

See Freedman (1971) corollary 29. Thus, for $u \rightarrow \infty$, we have

$$P\left(\sup_{0 \leq t \leq 1} W(t) > u\right) \sim P(W(1) > u - u^{-1} \log 2)$$

by application of the expansion

$$(2.5) \quad P(W(1) > u) \sim (2\pi)^{-1/2} u^{-1} e^{-u^2/2} \quad \text{for } u \rightarrow \infty.$$

See Freedman (1971) lemma (4.a).

In (2.3.a) we have the trivial lower bound

$$(2.6) \quad P(V_m(1) > u) \leq P\left(\max_{0 \leq t \leq 1} V_m(t) > u\right) \quad \text{for all } u.$$

Similarly for U_m .

Proof of lemma 2.1.

Part a. The mean value theorem implies

$$V_m(t+h) = V_m(t) + \sqrt{2} h \sum_{k=0}^{m-1} X_k \cos\left\{\frac{(2k+1)\pi}{2}(t+h_t)\right\}$$

when mh is small. Divide the interval $[0,1]$ in Δ^{-1} (integer) intervals of length Δ . Then we have

$$(2.7) \quad \begin{aligned} P\left(\max_{0 \leq t \leq 1} V_m(t) > u\right) &\leq \sum_{j=1}^{\Delta^{-1}-1} P\left(\max_{j\Delta \leq t \leq (j+1)\Delta} V_m(t) > u\right) \\ &= \sum_{j=0}^{\Delta^{-1}-1} P(V_m(j\Delta) + \max_{j\Delta \leq t \leq (j+1)\Delta} (V_m(t) - V_m(j\Delta)) > u). \end{aligned}$$

We also have

$$\max_{j\Delta \leq t \leq (j+1)\Delta} |V_m(t) - V_m(j\Delta)| \leq \Delta \sqrt{2} \sum_{k=0}^{m-1} |X_k|.$$

Note that this (random) bound is independent of t and j . For Δ_m^2 small we can apply the central limit theorem in order to obtain

$$(2.8) \quad P\left(\max_{j\Delta \leq t \leq (j+1)\Delta} V_m(t) > u\right) \leq P(V_m(j\Delta) > u - \epsilon u) + r_j$$

where the error r_j is asymptotically small with respect to $P(V_m(1) > u - \epsilon u)$ for $u \rightarrow \infty$. Using (2.7), $\sigma^2(V_m(t))$ is maximal for $t=1$. (2.8) and (2.5) we obtain

$$\begin{aligned} P\left(\max_{0 \leq t \leq 1} V_m(t) > u\right) &\leq \Delta^{-1} P(V_m(1) > u - \epsilon u) + \sum r_j \\ &\leq P(V_m(1) > u(1 - 2\epsilon)). \end{aligned}$$

Part b. $U_m(t)$ is an infinite series. We write

$$U_m(t) = \sum_{k=m}^{m^2-1} \frac{2\sqrt{2}}{\pi(2k+1)} X_k \sin(2k+1) \frac{\pi}{2} t + U_{m^2}(t).$$

We have seen that $\sigma^2(U_{m^2}(t)) \leq cm^{-2}$ uniformly in t . Thus we obtain (uniformly in t)

$$P(|U_{m^2}(t)| > u\delta) = o(P(U_m(1) > u(1-\epsilon)))$$

for $m, u \rightarrow \infty$.

The further proof is similar to that of part a. □

We apply the following asymptotic expansion for the right tail of the chi-square distribution with increasing degrees of freedom.

Lemma 2.2. For $m \rightarrow \infty$ and $m = o(x)$ for $x \rightarrow \infty$ we have

$$P(\chi_m^2 > x) = \{1 + o(1)\} \{\pi m\}^{-1/2} e^{-1/2x} e^{(-1/2m-1)\log(xe/m)}.$$

Proof of Lemma 2.2.

The assertion follows easily after some calculus and the application of Stirling's formula. □

The following assertions are well-known for gamma distributions but we shall only apply them for chi-square distributions.

Lemma 2.3.

Let X (resp. Y) be χ_n^2 (resp. χ_m^2) distributed. X and Y are independent. Then

- a) $X + Y$ has a χ_{n+m}^2 distribution
- b) $X + Y$ and $X/(X + Y)$ are independent
- c) $X/(X + Y)$ has a $B(\frac{n}{2}, \frac{m}{2})$ distribution.

Proof.

See Rohatgi (1976) Section 5.3 Theorem 4 resp. Th. 6 and Th. 15. □

Define the projection Π_m of H onto the finite-dimensional subspace with base $\{\varphi_0, \dots, \varphi_{m-1}\}$ by

$$\Pi_m \left(\sum_{n=0}^{\infty} a_n \varphi_n(t) \right) = \sum_{n=0}^{m-1} a_n \varphi_n(t).$$

Thus we have

$$V_m = \Pi_m W.$$

We use the notation Lx resp. L_2x for $\log x$ resp. $\log \log x$.

3. Lower bound.

Define the sequence $n_k = \exp(k\varphi(k))$ where φ is slowly varying at infinity and $\lim_{x \rightarrow \infty} \varphi(x) = \infty$. Then we have, for $k \rightarrow \infty$, $n_k/n_{k+1} \sim \exp(-\varphi(k)) \rightarrow 0$ and

$$P((2n_{k+1} L_{2n_{k+1}})^{-1/2} \|W(n_k \cdot)\|_\infty > \epsilon \psi_0(n_k)) = \\ \leq 4P(|U| > \epsilon \psi_0(n_k) (2 L_{2n_{k+1}} (n_{k+1}/n_k))^{1/2})$$

using (2.4) and the scaling property of a Brownian motion

$$= O(e^{-\epsilon^2 \psi_0^2(n_k) e^{\varphi(k)} L_{2n_k}} (\psi_0(n_k)^{-1} (L_{2n_k})^{-1/2} e^{-1/2 \varphi(k)}))$$

by (2.5). We choose $\varphi(k)$ such that the summation of these probabilities converges. For example take $\varphi(k) = 4/3 L_2 k$.

Now we define for $k=1,2,\dots$ the sequence of functions f_k^*

$$f_k^* : [0,1] \times \Omega \rightarrow \mathbb{R}$$

by

$$f_k^*(t, \omega) = (2n_{k+1} L_{2n_{k+1}})^{-1/2} \{W((n_{k+1} - n_k)t + n_k, \omega) - W(n_k, \omega)\}.$$

We easily see that $f_k^*(t)$ has the same distribution as $(2n_{k+1} L_{2n_{k+1}})^{-1/2} (1 - n_k/n_{k+1})^{1/2} W(n_{k+1} t)$. We can write, for each k ,

$$f_k^*(t) = \sum_{j=0}^{\infty} X_{k,j}^* \varphi_j(t) \cdot \{2L_{2n_{k+1}}\}^{-1/2} \{1 - n_k/n_{k+1}\}^{1/2}$$

where $X_{k,j}^*$, $j=0,1,\dots$, are i.i.d. $N(0,1)$ distributed random variables.

Remark that the random variables $X_{k,j}^*$ depend on k . Take $m_k = (L_{2n_k})^{1/3}$. Then

we have

$$(3.1) \quad P(\| \prod_{m_k} f_k^* \|_H > 1 + \psi_\delta(n_k)) \\ = P(\chi_{m_k}^2 > (2L_{2n_{k+1}}) (1 + \psi_\delta(n_k))^2 (1 - n_k/n_{k+1})^{-1})$$

which can be estimated using lemma 2.2. For $\delta < 0$ we have

$$\sum_k P(\| \Pi_{m_k} f_k^* \|_H > 1 + \psi_\delta(n_k)) = \infty.$$

Using the property that the increments of a Wiener process are independent, the Borel-Cantelli lemma implies that

$$P(\| \Pi_{m_k} f_k^* \|_H > 1 + \psi_\delta(n_k) \text{ i.o.}) = 1.$$

Therefore, w.p. 1 we have

$$\Pi_{m_k} f_k^* \notin K \text{ i.o.}$$

For $\Pi_{m_k} f_k^* \notin K$ the projection onto K is given by $\| \Pi_{m_k} f_k^* \|_H^{-1} \Pi_{m_k} f_k^*$. Thus w.p. 1 we have

$$d_H(\Pi_{m_k} f_k^*, K) = d_H(\Pi_{m_k} f_k^*, \| \Pi_{m_k} f_k^* \|_H^{-1} \Pi_{m_k} f_k^*) = \| \Pi_{m_k} f_k^* \|_H^{-1} > \psi_\delta(n_k) \text{ i.o.}$$

Now we shall show that w.p. 1 we have, for $0 < \delta_1 < \delta$, $d_\infty(f_k^*, K) > (1-\delta_1)\psi_0(n_k)$ i.o. Similarly as above we can show that for $\delta > 0$

$$\sum_k P(\| \Pi_{m_k} f_k^* \|_H > 1 + (1+\delta)\psi_0(n_k)) < \infty.$$

Then the Borel-Cantelli lemma implies

$$P(\| \Pi_{m_k} f_k^* \|_H > 1 + (1+\delta)\psi_0(n_k) \text{ i.o.}) = 0.$$

Define the events A_k , $k = 1, 2, \dots$ by

$$(3.2) \quad 1 + (1-\delta)\psi_0(n_k) \leq \| \Pi_{m_k} f_k^* \|_H \leq 1 + (1+\delta)\psi_0(n_k)$$

or

$$(2L_2 n_k)(1 + (1-\delta)\psi_0(n_k))^2 (1 - n_k^{-1} n_{k+1}^{-1})^{-2} \leq \sum_{j=0}^{m_k-1} (X_{k,j}^*)^2 \leq (2L_2 n_k) \cdot$$

$$\cdot (1 + (1+\delta)\psi_0(n_k))^2 (1 - n_k n_{k+1}^{-1})^{-2}.$$

Using lemma 2.1. part a we have that, for estimating the right tail probabilities of $\| \Pi_{m_k} f_k^* \|_\infty$, we may use the r.v.

$$\Pi_{m_k} f_k^*(1) = \left\{ \sum_{j=0}^{m_k-1} X_{k,j}^* \varphi_j(1) \right\} \{2L_{2n_{k+1}}\}^{-1/2} (1 - n_k n_{k+1}^{-1})^{1/2}.$$

The Cauchy-Schwarz inequality implies

$$\begin{aligned} (\Pi_{m_k} f_k^*(1))^2 &\leq (1 - n_k n_{k+1}^{-1}) \left\{ \sum_{j=0}^{m_k-1} (X_{k,j}^*)^2 \right\} \{2L_{2n_{k+1}}\}^{-1} \left\{ \sum_{j=0}^{m_k-1} \varphi_j^2(1) \right\} \\ &\leq \| \Pi_{m_k} f_k^* \|_H^2 \cdot \sigma^2(V_m(1)), \end{aligned}$$

where $\sigma^2(V_m(1))$ is given in (2.2.a). The r.v. $\Pi_{m_k} f_k^*(1)$ has a normal distribution with $E \Pi_{m_k} f_k^*(1) = 0$ and

$$\sigma^2(\Pi_{m_k} f_k^*(1)) = \sigma^2(V_m(1)) (2L_{2n_{k+1}})^{-1} (1 - n_k n_{k+1}^{-1}).$$

The vector $X^* = (X_{k,0}^*, \dots, X_{k, m_k-1}^*)$ has a m_k -dimensional normal distributor.

There exists an orthogonal transformation P such that the first row vector of

PX^* becomes $\{\sigma(V_m(1))\}^{-1} \left\{ \sum_{j=0}^{m_k-1} X_{k,j}^* \varphi_j(1) \right\}$. It follows from lemma 2.3. part c

that

$$(3.3) \quad \{\sigma^2(V_m(1))\}^{-1} \{1 - n_k n_{k+1}^{-1}\}^{-1} \{\Pi_{m_k} f_k^*(1)\}^2 \| \Pi_{m_k} f_k^* \|_H^{-2}$$

has a $B(\frac{1}{2}, \frac{1}{2}m_k - \frac{1}{2})$ distribution and by lemma 2.3. part b is the r.v. given in

$$(3.3) \text{ independent of } \| \Pi_{m_k} f_k^* \|_H^2.$$

Consider

$$\begin{aligned} P(\|\Pi_{m_k} f_k^*\|_\infty > 1 - \epsilon \wedge A_k) &= P(\|\Pi_{m_k} f_k^*\|_\infty > 1 - \epsilon | A_k) P(A_k) = \\ &\geq P(B(\frac{1}{2}, \frac{1}{2}m_k - \frac{1}{2}) > 1 - \epsilon_1) P(A_k) \end{aligned}$$

for some positive ϵ_1 . With the estimate

$$P(B(\frac{1}{2}, \frac{1}{2}m_k - \frac{1}{2}) > 1 - \epsilon_1) \sim cm_k^{-\frac{1}{2}} e^{-\frac{1}{2}m_k \log \epsilon_1^{-1}}$$

for $k \rightarrow \infty$ we obtain

$$\sum_k P(\|\Pi_{m_k} f_k^*\|_\infty > 1 - \epsilon \wedge A_k) = \infty.$$

The Borel-Cantelli lemma implies that w.p.1 we have i.o.

$$\|\Pi_{m_k} f_k^*\|_\infty > 1 - \epsilon \wedge \|\Pi_{m_k} f_k^*\|_H \geq 1 + (1-\delta)\psi_0(n_k).$$

Or, w.p. 1 we have i.o.

$$\|\Pi_{m_k} f_k^*\|_\infty > 1 - \epsilon \wedge d_H(\Pi_{m_k} f_k^*, K) = \|\Pi_{m_k} f_k^*\|_H - 1 > (1-\delta)\psi_0(n_k).$$

Next we want to conclude that w.p.1 we have

$$d_\infty(\Pi_{m_k} f_k^*, K) > (1-\delta_1)\psi_0(n_k) \quad \text{i.o.}$$

One may apply results from the theory of linear spaces. See, for example, Banach (1932) chapter XI § 4 or Köthe (1960) § 26.4. We indicate a simple proof using the structure of H and $\Pi_m H$.

It is well-known that H is isometric isomorphic with ℓ_2 . Let

$$e = (e_0, \dots, e_{m-1}) \in \mathbb{R}^m \text{ with } e_j = \varphi_j(1) \text{ and } \Pi_m f = \sum_{j=0}^{m-1} f_j \varphi_j. \text{ Then}$$

$$\Pi_m f(1) = \sum f_j e_j = (f, e)_m, \text{ where } (\cdot, \cdot)_m \text{ is the inner product in } \mathbb{R}^m. \text{ Lemma 2.1}$$

implies that we may consider $(f, e)_m$ instead of $\|\Pi_m f\|_\infty$.

W.p.1 we have $(\Pi_{m_k} f_k^*, e)_{m_k} > 1 - \epsilon$ and $\|\Pi_{m_k} f_k^*\|_H > 1 + (1 - \delta)\psi_0(n_k)$. We write

$\Pi_{m_k} f_k^* = ae + bd$ where $d \in \mathbb{R}^{m_k}$ perpendicular to e (i.e. $(e, d)_{m_k} = 0$). Thus

$(\Pi_{m_k} f_k^*, e)_{m_k} = a > 1 - \epsilon$. Then we have

$$\begin{aligned} d_\infty(\Pi_{m_k} f_k^*, K) &\geq ((1 - \|\Pi_{m_k} f_k^*\|_H^{-1}) (\Pi_{m_k} f_k^*, e)_{m_k}) \\ &= (1 - \|\Pi_{m_k} f_k^*\|_H^{-1}) a (e, e)_{m_k} > (1 - \delta_1)\psi_0(n_k). \end{aligned}$$

To complete the proof we consider

$$\begin{aligned} &P(\|\Pi_{m_k} f_k^*\|_H > 1 + (1 - \delta)\psi_0(n_k) \wedge \|f_k^* - \Pi_{m_k} f_k^*\|_\infty > (\psi_0(n_k)/m_k)^{1/2}) \\ &= P(\|\Pi_{m_k} f_k^*\|_H > 1 + (1 - \delta)\psi_0(n_k) P(\|f_k^* - \Pi_{m_k} f_k^*\|_\infty > (\psi_0(n_k)/m_k)^{1/2}) \end{aligned}$$

because of the independence of $\Pi_{m_k} f_k^*$ and $f_k^* - \Pi_{m_k} f_k^*$. Applying (3.1), lemma 2.2

and (2.3.b) we obtain

$$\sum_k P(\|\Pi_{m_k} f_k^*\|_H > 1 + (1 - \delta)\psi_0(n_k) \wedge \|f_k^* - \Pi_{m_k} f_k^*\|_\infty > (\psi_0(n_k)/m_k)^{1/2}) < \infty$$

Thus w.p.1 we have i.o.

$$\|\Pi_{m_k} f_k^*\|_H > 1 + (1 - \delta)\psi_0(n_k)$$

and

$$\|f_k^* - \Pi_{m_k} f_k^*\|_\infty \leq (\psi_0(n_k)/m_k)^{1/2}.$$

One easily sees that for k sufficiently large

$$(\psi_0(n_k)/m_k)^{1/2} < \frac{1}{3} \delta \psi_0(n_k).$$

4. Upper bound.

In this section we use the subsequences $n_k = \exp(k/\varphi(k))$ where $\varphi(k) = (\log k)^{4/3}$ and $m_k = (L_2 n_k)^{1/3}$. We write

$$P(d_\infty(f_{n_k}, K) > \psi_\delta(n_k)) =$$

$$P(d_\infty(f_{n_k}, K) > \psi_\delta(n_k) \wedge \prod_{m_k} f_k \notin K) +$$

$$P(d_\infty(f_{n_k}, K) > \psi_\delta(n_k) \wedge \prod_{m_k} f_{n_k} \in K \cap \bar{K}^{1-(L_2 n_k)^{-1/6}}) +$$

$$P(d_\infty(f_{n_k}, K) > \psi_\delta(n_k) \wedge \prod_{m_k} f_{n_k} \in K^{1-(L_2 n_k)^{-1/6}}) =$$

$$P_{1,k} + P_{2,k} + P_{3,k}.$$

The event in $P_{3,k}$ implies $f_{n_k} \in K$ for n_k sufficiently large. This follows from the following result.

Lemma 4.1. Let n_k and m_k be defined as above. Then

$$\sup_{m_k < m \leq m_{k+1}} \sup_{0 \leq t \leq 1} |U_m(t)| (2L_2 n_k)^{-1/2} \leq (L_2 n_k)^{-1/6} \text{ a.s.}$$

for k sufficiently large.

Proof. Using lemma 2.1.b we have

$$\begin{aligned} P\left(\sup_{0 \leq t \leq 1} U_{m_k}(t) > \frac{1}{2} \sqrt{2} (L_2 n_k)^{1/3}\right) &\leq (1+o(1))P(U > \frac{1}{2}\pi(1-\epsilon)(L_2 n_k)^{1/2}) \\ &= o((\varphi(k)/k)^{\pi^2(1-\epsilon)^2/8} (Lk)^{-1/2}) \text{ by (2.5).} \end{aligned}$$

The Borel-Cantelli lemma implies: w.p.1 for k sufficiently large

$$\sup_{0 \leq t \leq 1} (2L_2 n_k)^{-1/2} |U_{m_k}(t)| \leq \frac{1}{2} (L_2 n_k)^{-1/6}.$$

For $m_k < m \leq m_{k+1}$ we have $U_m(t) = \sum_{j=m}^{m_{k+1}-1} X_j \varphi_j(t) + U_{m_{k+1}}(t)$. Consider

$$P\left(\sup_{m_k < m \leq m_{k+1}-1} \sup_{0 \leq t \leq 1} (2L_2 n_k)^{-1/2} \left| \sum_{j=m}^{m_{k+1}-1} X_j \varphi_j(t) \right| > \frac{1}{2} (L_2 n_k)^{-1/6}\right)$$

$$\leq \sum_{m=m_k}^{m_{k+1}} P\left(\sup_{0 \leq t \leq 1} \left| \sum_{j=m}^{m_{k+1}-1} X_j \varphi_j(t) \right| > \frac{1}{2} \sqrt{2} (L_2 n_k)^{1/3}\right)$$

$$\leq c(m_{k+1} - m_k) P(|U| > \frac{1}{4} \pi \sqrt{3} (1-\epsilon) k^{1/2} L_k).$$

Application of (2.5) and the Borel-Cantelli lemma gives the desired result. \square

Lemma 4.2. Let n_k be defined as above. Then, w.p.1 and for k sufficiently large, we have

$$\max_{n_k < n \leq n_{k+1}} \sup_{0 \leq t \leq 1} |f_n(t) - f_{n_k}(t)| < \epsilon \psi_0(n_k).$$

Proof. From the definition of f_n we have

$$(4.1) \quad \max_{n_k < n \leq n_{k+1}} \sup_{0 \leq t \leq 1} |f_n(t) - f_{n_k}(t)| =$$

$$\max_{n_k < n \leq n_{k+1}} \sup_{0 \leq t \leq 1} \left| \frac{W(nt)}{(2nL_2n)^{1/2}} - \frac{W(n_k t)}{(2n_k L_2 n_k)^{1/2}} \right| \leq$$

$$\max_{n_k < n \leq n_{k+1}} \sup_{0 \leq t \leq 1} \left| \frac{W(nt) - W(n_k t)}{(2nL_2n)^{1/2}} \right| +$$

$$\max_{n_k < n \leq n_{k+1}} \left| \left[\frac{(2n_k L_2 n_k)^{1/2}}{2nL_2n} \right] - 1 \right| \sup_{0 \leq t \leq 1} \frac{|W(n_k t)|}{(2n_k L_2 n_k)^{1/2}}$$

From the definition of n_k it follows that $(n_{k+1} - n_k)n_{k+1}^{-1} \sim \{\varphi(k)\}^{-1}$. Using the L.I.L. we have that the last term on the right hand side of (4.1) is less than $2\{\varphi(k)\}^{-1}$. Next we consider

$$\begin{aligned} & P\left(\max_{n_k < n \leq n_{k+1}} \sup_{0 \leq t \leq 1} (2nL_2n)^{-1/2} |W(nt) - W(n_k t)| > \epsilon \psi_0(n_k)\right) \\ & \leq P\left(\sup_{0 \leq u \leq 1-h} \sup_{0 \leq v \leq h} |W(u+v) - W(v)| > \epsilon_1 \psi_0(n_k) (2L_2 n_k)^{1/2}\right) \end{aligned}$$

using the scaling property of the Brownian motion where $h \sim (\log k)^{-4/3}$. Now we apply the estimate given in lemma 1.1.1 of Csörgö and Révész (1981) and the Borel-Cantelli lemma in order to obtain the desired result. \square

The assertion in the last lemma gives us that we have only to show that f_{n_k} is close to K .

Lemma 4.3. Take $\delta > 0$. Let n_k and m_k be defined as above. Then

$$\| \prod_{m_k} f_{n_k} \|_H < 1 + \psi_\delta(n_k) \quad \text{a.s.}$$

for k sufficiently large.

Proof. It follows from the definition of $\|\cdot\|_H$ that we have

$$P\left(\| \prod_{m_k} f_{n_k} \|_H > 1 + \psi_\delta(n_k)\right) = P\left(\chi_{m_k}^2 > (2L_2 n_k)(1 + \psi_\delta(n_k))^2\right)$$

Applying lemma 2.2 we have that the summation of the probabilities converges.

The Borel-Cantelli lemma gives the result. \square

Now we can complete the proof for the upper bound. It follows from lemma 4.3 that in the events described in $P_{1,k}$ and $P_{2,k}$ we have, for $\delta > 0$,

$$1 - (L_2 n_k)^{-1/6} \leq \| \Pi_{m_k} f_{n_k} \|_H \leq 1 + (1+\delta)\psi_0(n_k) \quad \text{a.s.}$$

We have

$$\begin{aligned} P_{1,k} &= P(d_\infty(f_{n_k}, K) > (1+\delta)\psi_0(n_k) \wedge \Pi_{m_k} f_{n_k} \notin K) \\ &\leq P(d_\infty(f_{n_k}, K) > (1+\delta)\psi_0(n_k) \wedge 1 \leq \| \Pi_{m_k} f_{n_k} \|_H < 1 + (1 + \frac{1}{2}\delta)\psi_0(n_k)) + r_{n_k} \end{aligned}$$

where the error r_{n_k} is given in the proof of lemma 4.3.

As we have seen in section 2 we can write

$$f_{n_k}(t) = \Pi_{m_k} f_{m_k}(t) + (2L_2 n_k)^{-1/2} U_{m_k}(t)$$

where $\Pi_{m_k} f_{m_k}(t)$ and $U_{m_k}(t)$ are independent. In lemma 4.1 we showed that $(2L_2 n_k)^{-1/2} U_{m_k}(t)$ is small.

$$\begin{aligned} P(d_\infty(f_{n_k}, K) > (1+\delta)\psi_0(n_k) \wedge 1 \leq \| \Pi_{m_k} f_{n_k} \|_H < 1 + (1+\frac{1}{2}\delta)\psi_0(n_k)) \\ \leq P(\| \Pi_{m_k} f_{n_k} \|_H \geq 1) P((2L_2 n_k)^{-1/2} |U_{m_k}(1)| > \frac{1}{2} \delta \psi_0(n_k)) \\ = O(k^{-1} (Lk)^{7/6} e^{(L_2 k)^{1/3} L_3 k/3} \cdot e^{-\delta^2 \pi^2 (L_2 k)^2 / 8}) \end{aligned}$$

by lemma 2.2 and estimate (2.5). Since $\sum_k P_{1,k} < \infty$ the Borel Cantelli lemma gives the desired result.

For the event considered in $P_{2,k}$ we define the stopping time M by

$$\{M=m\} = \{ \| \Pi_m f_{n_k} \|_H \leq 1 + \psi_0(n_k) < \| \Pi_{m+1} f_{n_k} \|_H \}$$

Then we have

$$P(M=m) = P(\chi_m^2 \leq 2L_2 n_k (1+\psi_0(n_k))^2 \leq \chi_m^2 + U^2)$$

where χ_m^2 and U^2 are independent and chi-square distributed with respect to m

and 1 degrees of freedom.

If $\prod_{m_k} f_{n_k} \in K \cap \bar{K}^{1-(L_2 n_k)^{-1/6}}$ we have $M = m > m_k$. The Borel-Cantelli

lemma gives us an upper bound for M .

$$\begin{aligned} P_k^* &= P(1 - (L_2 n_k)^{-1/6} \leq \| \prod_{m_k} f_{n_k} \|_H \leq \| \prod_{m_k^{5/2}} f_{n_k} \|_H \leq 1 + \psi_0(n_k)) \\ &= P(2L_2 n_k (1 - (L_2 n_k)^{-1/6})^2 \leq \chi_{m_k}^2 \leq \chi_{m_k^{5/2}}^2 \leq (2L_2 n_k)(1 + \psi_0(n_k))^2) \\ &\leq P(\chi_{m_k^{5/2}}^2 \leq 2L_2 n_k (1 - (L_2 n_k)^{-1/6})^2) P(B(\frac{1}{2} m_k, \frac{1}{2} m_k^{5/2} - \frac{1}{2} m_k) \geq (1 - (L_2 n_k)^{-1/6})^2) \end{aligned}$$

by conditioning on $\chi_{m_k^{5/2}}^2$ and lemma 2.3 parts b and c

$$\begin{aligned} &= O((Lk)^{5/6} k^{-1} e^{2(L_2 n_k)^{5/6}} e^{-(L_2 n_k)^{2/3}} \frac{1}{2} m_k^{5/2} L(2eL_2 n_k / m_k^{5/2})) \\ &\cdot O(m_k^{-1/2} e^{3m_k L m_k / 4} m_k^{-5/2} e^{-\frac{1}{2}(m_k^{5/2} - m_k)L((L_2 n_k)^{1/6} / 2)}) \end{aligned}$$

by lemma 2.2 and computation of the probabilities of a beta distributed r.v.

It follows from the upperbound as derived above that

$$\sum P_k^* < \infty.$$

The Borel-Cantelli lemma gives us that from now on we only have to consider those values for M that are less than $m_k^{5/2}$.

When $M = m$ we have $\prod_{m} f_{n_k} \in K$. Finally we shall prove that in this case we

have

$$d_\infty(\prod_{m} f_{n_k}, K) < \epsilon \psi_0(n_k)$$

and also

$$d_\infty(\prod_{m} f_{n_k}, f_{n_k}) < \epsilon \psi_0(n_k).$$

Then we have proved the assertion for the upper bound. Consider

$$P_k^{***} = \sum_{m=m_k}^{5/2} P(\prod_{m_k} f_{n_k} \in K \cap \bar{K} \quad 1-(L_2 n_k)^{-1/6} \wedge M=m \wedge \\ |X_{m+1} \varphi_{m+1}(t)| > \epsilon \psi_0(n_k)) \\ \leq \sum_{m=m_k}^{5/2} P(\chi_{m+1}^2 \geq 2L_2 n_k (1+\psi_0(n_k))^2) \cdot P(B(1/2, 1/2m) \geq \epsilon_1 m \psi_0(n_k))$$

by conditioning on χ_{m+1}^2 , lemma 2.3 parts b and c and we also use that $\chi_{m+1}^2 \in [2L_2 n_k (1+\psi_0(n_k))^2, 2(1+\epsilon)L_2 n_k]$. Using the estimates for χ^2 and beta distribution we obtain $\sum_k P_k^{***} < \infty$.

Finally we consider

$$P_k^{****} = \sum_{m=m_k}^{5/2} P(\prod_{m_k} f_{n_k} \in K \cap \bar{K} \quad 1-(L_2 n_k)^{-1/6} \wedge M=m \wedge \\ \|U_{m+1}(\cdot)\|_{\infty} (2L_2 n_k)^{-1/2} \geq \epsilon \psi_0(n_k)) \\ \leq \sum_{m=m_k}^{5/2} P(\chi_{m+1}^2 \geq 2L_2 n_k (1+\psi_0(n_k))^2) P(\sup_{0 \leq t \leq 1} |U_{m+1}(t)| > (2L_2 n_k)^{1/2} \epsilon \psi_0(n_k))$$

by the independence of the processes U_{m+1} and V_{m+1} . Using the lemmas 2.1 and 2.2 we obtain

$$\sum_k P_k^{****} < \infty.$$

5. Discussion and remarks.

All proofs of Strassen's theorem contain the following assertions.

- i) There exists some sequence $\{n_k\}$ such that w.p.1 $d_{\infty}(f_n, f_{n_k}) < \epsilon$ for all

$n \in (n_k, n_{k+1})$ and k sufficiently large.

ii) Let $\Pi_m f$ be the piecewise linear approximation of f . Then, for fixed m , w.p.1 $d_\infty(\Pi_m f_{n_k}, f_{n_k}) < \epsilon$ for m, k sufficiently large.

iii) $\int_0^1 \left(\frac{d}{dt} \Pi_m f_{n_k}(t) \right)^2 dt$ has the same distribution as $(2L_2 n_k)^{-1} \chi_m^2$ and is w.p.1 less than $(1+\epsilon)^2$ for k sufficiently large. Thus the last assertion implies $\| \Pi_m f_{n_k} \|_H \leq 1 + \epsilon$ w.p.1 for k large.

The projection Π_m as defined in section 2 is a different approximation than the one above. Above we have that $\Pi_m f$ and $f - \Pi_m f$ are dependent. In the approximation used in sections 3 and 4 of this paper we have that

$\Pi_m f_n = (2L_2 n)^{-1/2} V_m$ and $f_n - \Pi_m f_n = (2L_2 n)^{-1/2} U_m$ are independent.

As far as I know is m fixed in all proofs of the F.L.I.L.

Remark 1. In order to prove an integral test for the rate of convergence in the F.L.I.L. one needs asymptotic expansions in lemma 2.1. The lower class result becomes more difficult to prove.

Remark 2. For the Brownian bridge we have the following expansion

$$B(t) = \sum_{k=1}^{\infty} (k\pi)^{-1} X_k \sqrt{2} \sin(k\pi t) \quad 0 \leq t \leq 1$$

where X_1, X_2, \dots are i.i.d. $N(0,1)$. See Shorack and Wellner (1986) chapter 1 exercise 15. By the same method as given in this paper one can obtain the rate of convergence in Finkelstein's F.L.I.L. See Finkelstein (1971) or Shorack and Wellner (1986) chapter 13 section 3 theorem 1.

References

Banach, S.: Théorie des opérations linéaires. Z. Subwencji. Funduszu Kultury Narodowej, Warszawa (1932)

- Berman, S.M.: The maximum of a Gaussian process with non-constant variance. *Ann. Inst. Henri Poincaré.* 21, 383-391 (1985)
- Bolthausen, E.: On the speed of convergence in Strassen's law of the iterated logarithm. *Ann. Probab.* 6, 668-672 (1978)
- Chover, J.: On Strassen's version of the log log law. *Z. Wahrscheinlichkeitstheorie und Verw. Geb.* 8, 83-90 (1967)
- Csáki, E.: A Relation Between Chung's and Strassen's Laws of the Iterated Logarithm. *Z. Wahrscheinlichkeitstheorie und Verw. Geb.* 54, 287-301 (1980)
- Csörgö, M. and P. Révész: Strong approximations in probability and statistics. Academic Press, New York (1981)
- Feller, W.: An introduction to probability theory and its applications. Vol. II. Second edition. Wiley & Sons, New York (1971)
- Finkelstein, H.: The law of the iterated logarithm for empirical distributions. *Ann. Math. Statist.* 42, 607-615 (1971)
- Freedman, D.: Brownian motion and diffusion. Holden-Day, San Francisco (1971)
- Grill, K.: On the Rate of Convergence in Strassen's Law of the Iterated Logarithm. *Prob. Th. Rel. Fields* 74, 583-589 (1987)
- Jain, N.C. and M.B. Marcus: Continuity of sub Gaussian processes. *Advances in Probability and related topics. Vol. 4. Probability in Banach spaces*, Ed. J. Kuelbs (1978)
- Köthe, G.: Topologische lineare Räume I. Springer-Verlag, Berlin (1960)
- Kuelbs, J. and R. LePage: The law of the iterated logarithm for Brownian motion in a Banach space. *Trans. Amer. Math. Soc.* 185, 253-264 (1973)
- Loève, M.: Probability Theory 3d ed., Van Nostrand, Princeton, N.J. (1963)
- Piterbarg, V.I. and V.P. Prisjaznjuk.: Asymptotics of the probability of large excursions for a non-stationary Gaussian process. *Theor. Prob. Math. Statist.* 19, 131-144 (1978)
- Rohatgi, V.K.: An Introduction to Probability Theory and Mathematical Statistics. Wiley & Sons, New York (1976)
- Shorack, G.R. and J.A. Wellner.: Empirical processes with applications to statistics. Wiley & Sons, New York (1986)
- Strassen, V.: An invariance principle for the law of the iterated logarithm. *Z. Wahrscheinlichkeitstheor. Ver. Geb.* 3, 211-216 (1964)

222. I. Mitom, Weak solution of the Langevin equation on a generalized functional space, Feb. 88.
223. L. de Haan, S.I. Resnick, H. Rootzén and C. de Vries, Extremal behaviour of solutions to a stochastic difference equation with applications to arch-processes, Feb. 88.
224. O. Kallenberg and J. Szulga, Multiple integration with respect to Poisson and Lévy processes, Feb. 88. *Prob. Theor. Zet. Fields*, 1989, to appear.
225. D.A. Dawson and L.G. Corostiza, Generalized solutions of a class of nuclear space valued stochastic evolution equations, Feb. 88.
226. G. Samorodnitsky and J. Szulga, An asymptotic evaluation of the tail of a multiple symmetric α -stable integral, Feb. 88. *Ann. Probability*, to appear.
227. J.J. Hunter, The computation of stationary distributions of Markov chains through perturbations, Mar. 88.
228. H.C. Ho and T.C. Sun, Limiting distribution of nonlinear vector functions of stationary Gaussian processes, Mar. 88.
229. R. Brigoia, On functional estimates for ill-posed linear problems, Apr. 88.
230. M.R. Leadbetter and S. Mandagopalan, On exceedance point processes for stationary sequences under mild oscillation restrictions, Apr. 88. *Oberwolfach Conf. on Extremal Value Theory*, J. Misler and R. Weiss, eds., Springer, to appear.
231. S. Cambanis, J. P. Nolan and J. Rosinski, On the oscillation of infinitely divisible processes, Apr. 88.
232. G. Hardy, G. Kallianpur and S. Ramasubramanian, A nuclear space-valued stochastic differential equation driven by Poisson random measures, Apr. 88.
233. D.J. Daley, T. Rojcki, Light traffic approximations in queues (II), May 88.
234. G. Kallianpur, I. Mitom, R.L. Wolpert, Diffusion equations in duals of nuclear spaces, July 88.
235. S. Cambanis, Admissible translates of stable processes: A survey and some new models, July 88.
236. E. Platen, On a wide range exclusion process in random medium with local jump intensity, Aug. 88.
237. R.L. Smith, A counterexample concerning the extremal index, Aug. 88.
238. G. Kallianpur and I. Mitom, A Langevin-type stochastic differential equation on a space of generalized functionals, Aug. 88.
239. C. Houdré, Harmonizability, V -boundedness, $(2,P)$ -boundedness of stochastic processes, Aug. 88.
240. G. Kallianpur, Some remarks on Hu and Meyer's paper and infinite dimensional calculus on finitely additive canonical Hilbert space, Sept. 88.
241. L. de Haan, A Brownian bridge connected with extreme values, Sept. 88.
242. O. Kallenberg, Exchangeable random measures in the plane, Sept. 88.
243. E. Masry and S. Cambanis, Trapezoidal Monte Carlo integration, Sept. 88.
244. L. Pitt, On a problem of H.P. McKean, Sept. 88.
245. C. Houdré, On the prediction theory of multivariate $(2,p)$ -bounded processes, Sept. 88.
246. C. Houdré, Fourier integrals for stochastic processes and dilation of vector measures, Sept. 88.
247. J. Mijneer, On the rate of convergence in Strassen's functional law of the iterated logarithm, Sept. 88.