

NAVAL POSTGRADUATE SCHOOL

Monterey, California

AD-A215 139



A NOTE ON SMALE'S GLOBAL NEWTON METHOD

Allen Goldstein

August 1989

Approved for public release; distribution unlimited
Prepared for:

Naval Postgraduate School
Monterey, CA 93943

DTIC
SERIALS
SECTION
OCT 1989
D

NAVAL POSTGRADUATE SCHOOL
Department of Mathematics

Rear Admiral R. W. West JR.
Superintendent


Harrison Shull
Provost

This report was prepared in conjunction with research conducted for the Naval Postgraduate School and funded by the Naval Postgraduate School. Reproduction of all or part of this report is authorized.

Prepared by:


ALLEN A. GOLDSTEIN
Adjunct Professor of Mathematics

Reviewed by:


HAROLD M. FREDRICKSEN
Chairman
Department of Mathematics

Released by:


KNEALE T. MARSHALL
Dean of Information
and Policy Sciences

SECURITY CLASSIFICATION OF THIS PAGE

REPORT DOCUMENTATION PAGE				Form Approved OMB No 0704-0188	
1a REPORT SECURITY CLASSIFICATION UNCLASSIFIED		1b RESTRICTIVE MARKINGS			
2a SECURITY CLASSIFICATION AUTHORITY		3 DISTRIBUTION/AVAILABILITY OF REPORT Approved for public release; distribution unlimited			
2b DECLASSIFICATION/DOWNGRADING SCHEDULE					
4 PERFORMING ORGANIZATION REPORT NUMBER(S) NPS-53-89-016		5 MONITORING ORGANIZATION REPORT NUMBER(S) NPS-53-89-016			
6a NAME OF PERFORMING ORGANIZATION Naval Postgraduate School	6b OFFICE SYMBOL (If applicable) 53	7a NAME OF MONITORING ORGANIZATION Naval Postgraduate School			
6c ADDRESS (City, State, and ZIP Code) Monterey, CA 93943		7b ADDRESS (City, State, and ZIP Code) Monterey, CA 93943			
8a NAME OF FUNDING/SPONSORING ORGANIZATION Naval Postgraduate School	8b OFFICE SYMBOL (If applicable) 53	9 PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER O&MN Direct Funding			
8c ADDRESS (City, State, and ZIP Code) Monterey, CA 93943		10 SOURCE OF FUNDING NUMBERS			
		PROGRAM ELEMENT NO	PROJECT NO	TASK NO	WORK UNIT ACCESSION NO
11 TITLE (Include Security Classification) A NOTE ON SMALE'S GLOBAL NEWTON METHOD					
12 PERSONAL AUTHOR(S) Allen Goldstein					
13a TYPE OF REPORT	13b TIME COVERED FROM 5/89 TO 8/89	14 DATE OF REPORT (Year, Month, Day) August 18, 1989		15 PAGE COUNT 5	
16 SUPPLEMENTARY NOTATION					
17 COSATI CODES			18 SUBJECT TERMS (Continue on reverse if necessary and identify by block number) Global Newton methods, unconstrained optimization; computational complexity		
FIELD	GROUP	SUB GROUP			
19 ABSTRACT (Continue on reverse if necessary and identify by block number) We present an implementation of Smale's Global Newton method in a simple setting. The iteration count for the algorithm is sensitive only to the quantity ϵ defined below.					
20 DISTRIBUTION/AVAILABILITY OF ABSTRACT <input checked="" type="checkbox"/> UNCLASSIFIED UNLIMITED <input type="checkbox"/> SAME AS RPT <input type="checkbox"/> DTC USERS		21 ABSTRACT SECURITY CLASSIFICATION UNCLASSIFIED			
22a NAME OF RESPONSIBLE INDIVIDUAL Allen Goldstein		22b TELEPHONE (Include Area Code) (408) 646-2664		22c OFFICE SYMBOL 53G0	

A note on Smale's Global Newton Method

A. A. Goldstein*

We have previously considered Smale's [1] global Newton method in conjunction with the Kantorovich inequalities in a simple setting. Our object was to determine a class of problems for which the method could be expected to be efficient. We now repeat this project using Smale's [2] estimates at one point instead of the Kantorovich estimate.

The power of Smale's estimate stems not only from the fact that the information is concentrated at one point. A further advantage is that, in contrast with the Kantorovich inequalities, no estimate of the norm of the inverse of the derivative operator by itself is needed. Newton step lengths can be small even if the derivative operator is nearly singular.

The algorithm below requires only local information for its implementation. To predict its behaviour, however, certain global constants are needed. These constants can be estimated on a thin cylinder containing a segment joining the origin and the initial value of the vector valued function for which we are seeking the root. The iteration count for the algorithm is sensitive only to the parameter $\beta\gamma$ defined below.

Theorem 1 (Smale 86) Assume F is an analytic map between real Banach spaces X and Y . That is, the Fréchet derivatives $F^{(k)}(x)$ exist for all $x \in X$ and $k=1,2,3,\dots$. Given $x_0 \in X$, assume that the inverse of $F'(x)$, which we denote by $F'_{-1}(x)$, exists. Set

$$\beta(x_0) = \|F'_{-1}(x_0)F(x_0)\| \quad \text{and}$$
$$\gamma(x_0) = \sup \left\{ \left\| \frac{1}{k!} F'_{-1}(x_0) F^{(k)}(x_0) \right\|^{\frac{1}{k-1}} : k \geq 2 \right\}$$

If

$$\beta(x_0)\gamma(x_0) < .130707$$

then x_0 is an approximate root of F . That is the Newton sequence

$$x_{k+1} = x_k - F'_{-1}(x_k)F(x_k)$$

is well defined and $\{x_k\}$ converges to say ξ , a root of F at the rate:

$$\|x_{k+1} - x_k\| \leq 2\left(\frac{1}{2}\right)^{2^k} \beta(x_0)$$

* supported by grants NIH RR01243-05 AND NPS LMC-M4E1

Moreover,

$$\|x_k - \xi\| \leq \frac{7}{4} \left(\frac{1}{2}\right)^{2^{k-1}} \beta(x_0)$$

The next result is not given by Smale but it is readily calculated using his ingredients. In what follows we shall often abbreviate $\beta(x_k)$ (and similar expressions) by β_k .

Remark 2

$$\|F'(x_k)\| < 1.85 \|F'(x_0)\| \quad k = 1, 2, 3, \dots$$

Proof. In [2], Proposition 2, we find the formula

$$\alpha_{k+1} \leq \alpha_k / (2\alpha_k^2 - 4\alpha_k + 1),$$

where $\alpha_k = \beta_k \gamma_k \leq 1/8$. Then $\alpha_{k+1} < (2\alpha_k)^2$. In Smale's proof of Lemma 2 we find $\|I - F'_{-1}(x_k)F'(x_{k+1})\| \leq (1 - \alpha_k)^{-2} - 1$. Whence $\|I - F'_{-1}(x_k)F'(x_{k+1})\| \leq 1 + (11/4)\alpha_k$. Let $H_k = F'_{-1}(x_k)F'(x_{k+1})$ then $F'(x_{k+1}) = F'(x_k)H_k$, and

$$\|F'(x_{k+1})\| < \|F'(x_0)\| \prod_{i=0}^k (1 + (11/4)(2\alpha_0)^{2^i})$$

We can improve this result by using sharper bounds for $i=0$ and for $i=1$, namely, .306 and .138, respectively. Then

$$\log \left(\prod_{i=2}^{\infty} (1 + (11/4)(2\alpha_0)^{2^i}) \right) < (11/4) \int_1^{\infty} (1/4)^{2^x} dx < .172$$

Finally we have the estimate

$$\prod_{i=0}^{\infty} \|H_k\| < 1.85.$$

Assume the hypotheses of Theorem 1. Let

$$T = \{x \in X : \|F(x)\| \leq \|F(x_0)\|\}.$$

We assume that $\gamma(x)$, $F(x)$ and $\beta(x)$ are defined and can be calculated throughout T . Assume the existence of numbers β , γ , K and σ that bound $\beta(x)$, $\gamma(x)$, $\|F'(x)\|$ and

$$\sigma(x) = \|F'_{-1}(x) \frac{F(x)}{\|F(x)\|}\|$$

on T . Let $\lceil x \rceil$ denote the smallest integer $\geq x$.

Algorithm 3 Given $x_i \in T$, we define x_{i+1} as follows. Set $t_i = 1 - (8\gamma_i\beta_i)^{-1}$, where $\gamma_i = \gamma(x_i)$ and β_i is defined similarly. Set $\xi_0^i = x_i$ and run the Newton sequence for G starting at ξ_0^i . Let k be the smallest integer satisfying

$$G(\xi_k^i) \leq \|F(x_i)\|/40\gamma_i\beta_i$$

Set $x_{i+1} = \xi_k^i$.

The algorithm can be run and terminated with a posteriori data, without any knowledge of the values of the constants β , γ , σ , and K . Moreover to verify a) and b) below we do not use these constants. However the global behaviour of the algorithm as described in c) and d) is given in terms of these constants. Let $K_i = 1.85 \|F(x_i)\|$.

Claim 4

(a) x_{i+1} can be found in $S = \lceil \log_2(\log_2(80 K_i \gamma_i \beta_i^2 / \|F(x_i)\|)) \rceil$ steps.

(b) $\|F(x_{i+1})\| \leq (1 - (1/10 \gamma_i \beta_i)) \|F(x_i)\|$

(c) Let $S = \log_2(\log_2(80 K \gamma \beta \sigma))$. Given $\epsilon > 0$, if $k \geq \lceil 10 \gamma \beta S \log_2(1/\epsilon) \rceil$ then $(\|F(x_k)\| / \|F(x_0)\|) \leq \epsilon$.

(d) If $k \geq \lceil 10 \gamma \beta S \log(8/\sigma) \rceil$ then x_k is an approximate root of F .

Proof. We show first that x_{i+1} can be chosen as claimed. Let $G^i(x) = t_i F(x_i)$. Then x_i is an approximate root for G^i . Hence if $\xi_0 = x_i$ and $\xi_{k+1} = \xi_k - F'(\xi_k)G(\xi_k)$ then ξ_k converges to ξ^i , a root of G^i . We now prove (b)

We have that $G^i(x_i) - G^i(\xi^i) = G^i(x_i) = (F(x_i)/8\beta_i\gamma_i) = F(x_i) - F(\xi_i)$. Also, $F(x_i) - F(x_{i+1}) = F(x_i) - F(\xi^i) + G^i(\xi^i) - G^i(x_{i+1}) = (F(x_i)/8\beta_i\gamma_i) + G^i(\xi^i) - G^i(x_{i+1})$. Thus (b) is true.

We now prove (a) by counting the steps needed to ensure that

$$\|G^i(\xi^i)\| \leq \|F(x_i)\|/40\gamma_i\beta_i = tmp.$$

Since $G^i(\xi_i) = F'(\xi_i)(\xi_{i+1} - \xi_i)$ we may choose k to ensure that $2K^i(\frac{1}{2})^{2^k}\beta(x_i) \leq tmp$

To prove (c) replace β_i and γ_i by β and γ , respectively. If k satisfies $k \log_2(1 - (1/10 \gamma \beta)) \leq \epsilon$ then $(\|F(x_k)\| / \|F(x_0)\|) \leq \epsilon$.

To go from k to $k+1$ requires no more than S steps.

We now turn to d). If $\|F(x)\| \leq \frac{1}{8\gamma\sigma}$ then x is an approximate root for F . Thus we replace $1/\epsilon$ in c) by $8\sigma\gamma$

Remarks 5

The method is sensitive only to the number $\gamma\beta$!

Note that from the above proof $F(\xi^i) = t_i F(x_i)$. Since x_{i+1} lies near ξ^i the global constants need be estimated only on a slender tube surrounding the segment joining the origin and $F(x_0)$.

Bibliography

- [1] Steve Smale Algorithms for solving Equations, Proceedings of the International Congress of Mathematicians, Vol1, pp172,195. AMS.1986.
- [2] Steve Smale Newton's method estimates from data at one point. The Merging of Disciplines... Springer-Verlag 1986 185-196.

Accession For	
NTIS GRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By _____	
Distribution/	
Availability Codes	
Dist	Avail. and/or Special
A-1	

INITIAL DISTRIBUTION LIST

DIRECTOR (2)
DEFENSE TECH. INFORMATION CENTER
CAMERON STATION
ALEXANDRIA, VA 22314

DIRECTOR OF RESEARCH ADMINISTRATION
CODE 012
NAVAL POSTGRADUATE SCHOOL
MONTEREY, CA 93943

OFFICE OF NAVAL RESEARCH
CODE 422AT
ARLINGTON, VA 22217

PROFESSOR ALLEN GOLDSTEIN (12)
CODE 53GO
DEPARTMENT OF MATHEMATICS
NAVAL POSTGRADUATE SCHOOL
MONTEREY, CA 93943

LIBRARY (2)
NAVAL POSTGRADUATE SCHOOL
MONTEREY, CA 93943

DEPARTMENT OF MATHEMATICS
CODE 53
NAVAL POSTGRADUATE SCHOOL
MONTEREY, CA 93943

CENTER FOR NAVAL ANALYSIS
4401 FORD AVENUE
ALEXANDRIA, VA 22302-0268