

SECURITY CLASSIFICATION OF THIS PAGE

<p>AD-A216 970</p> <p>Unclassified</p>		DOCUMENTATION PAGE		Form Approved OMB No 0704-0188	
		1b RESTRICTIVE MARKINGS			
4. PERFORMING ORGANIZATION REPORT NUMBER(S) N00014-79-C-0647		6b OFFICE SYMBOL (If applicable) JLE		3 DISTRIBUTION / AVAILABILITY OF REPORT Approved for public release; distribution unlimited.	
6a NAME OF PERFORMING ORGANIZATION Colorado State University		7a. NAME OF MONITORING ORGANIZATION		5 MONITORING ORGANIZATION REPORT NUMBER(S)	
6c. ADDRESS (City, State, and ZIP Code) Department of Chemistry Fort Collins, CO 80523		7b. ADDRESS (City, State, and ZIP Code)			
8a. NAME OF FUNDING / SPONSORING ORGANIZATION Office of Naval Research		8b. OFFICE SYMBOL (If applicable)		9 PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER N00014-79-C-0647	
8c. ADDRESS (City, State, and ZIP Code) 800 North Quincy Street Arlington, VA 22217-5000		10 SOURCE OF FUNDING NUMBERS			
		PROGRAM ELEMENT NO	PROJECT NO	TASK NO	WORK UNIT ACCESSION NO
11 TITLE (Include Security Classification) "Observation and Geometry Assignment of the Conformation of Benzyl Alcohol in the Gas Phase"					
12 PERSONAL AUTHOR(S) Jeffrey I. Seeman, Henry V. Secor, Hoong-Sun Im and E. R. Bernstein					
13a TYPE OF REPORT Technical Report		13b TIME COVERED FROM _____ TO _____		14. DATE OF REPORT (Year, Month, Day) December 15, 1989	
15 PAGE COUNT					
16 SUPPLEMENTARY NOTATION					
17 COSATI CODES			18 SUBJECT TERMS (Continue on reverse if necessary and identify by block number)		
FIELD	GROUP	SUB-GROUP	benzyl alcohol, planar conformation, gauche conformation, perpendicular conformation, sterically unencumbered derivatives, gas phase, supersonic jet mass spectroscopy,		
			2-phenethyl alcohol, electron cloud, structure, internal hydrogen bonding.		
19 ABSTRACT (Continue on reverse if necessary and identify by block number) SEE ATTACHED ABSTRACT					
20 DISTRIBUTION / AVAILABILITY OF ABSTRACT <input checked="" type="checkbox"/> UNCLASSIFIED/UNLIMITED <input type="checkbox"/> SAME AS RPT <input type="checkbox"/> DTIC USERS			21 ABSTRACT SECURITY CLASSIFICATION Unclassified		
22a NAME OF RESPONSIBLE INDIVIDUAL Elliot R. Bernstein			22b. TELEPHONE (Include Area Code) (303) 491-6347		22c OFFICE SYMBOL

DTIC
ELECTE
S **D**
JAN 25 1990
B

90 01 24 0 22

OFFICE OF NAVAL RESEARCH

Contract N00014-79-C-0647

TECHNICAL REPORT #65

"Observation and Geometry Assignment of the Conformation of
Benzyl Alcohol in the Gas Phase"

by

Jeffrey I. Seeman, Henry V. Secor,
Hoong-Sun Im and E. R. Bernstein

To be submitted for publication in Chemistry Communications

Department of Chemistry
Colorado State University
Fort Collins, Colorado 80523

December 15, 1989

Reproduction in whole or in part is permitted for
any purpose of the United States Government.

This document has been approved for public release
and sale; its distribution is unlimited

Observation and Geometry Assignment of the Conformation of Benzyl Alcohol in
the Gas Phase

Jeffrey I. Seeman,^{*a} Henry. V. Secor,^a Hoong-Sun Im,^b and E. R. Bernstein^{*b}

^aPhilip Morris Research Center, P.O. Box 26583, Richmond, Virginia 23261,
U.S.A.

^bDepartment of Chemistry, Condensed Matter Sciences Laboratory, Colorado
State University, Fort Collins, Colorado 80523, U.S.A.

Supersonic molecular jet laser spectroscopy is used to establish
the perpendicular conformation of benzyl alcohol and a number of sterically
unencumbered derivatives in the gas phase.

Benzyl alcohol (1) and its derivatives have vast synthetic utility, are
frequently found in natural products, and play a central role in numerous
mechanistic investigations. In spite of this chemical importance, the
conformational preference of the $-\text{CH}_2^{\text{H}}\text{OH}$ group relative to the aromatic ring
has not been unambiguously established: various experimental and theoretical
studies have failed to lead to a unified and consistent picture; Planar^{1,2}

<input checked="" type="checkbox"/>
<input type="checkbox"/>
<input type="checkbox"/>

Codes

Dist


Avail and/or
Special

A-1

1



(2) and gauche^{3,4} (4) conformations as well as a freely rotating model⁴ have been proposed for benzyl alcohols (see Scheme 1).




$$\tau = \tau(C_{ortho}-C_{ipso}-C_{\alpha}-C_{\beta})$$

The geometry of the minimum energy conformations of a variety of aromatic ring substituents has recently been established using molecular jet laser spectroscopy.⁵⁻⁷ The supersonic expansion process results in molecules at near 0 K: intramolecular conformational interconversions are thereby stopped, and electronic transitions become very sharp and well resolved.⁵⁻⁷ Since each stable conformation of a molecule has a specific potential energy minimum in S_0 and S_1 , the energy of the electronic origin (0_0^0 transition) for the $S_1 \leftarrow S_0$ transition for each stable conformer is generally unique. Hence, each stable conformation is associated with its own spectroscopic 0_0^0 transition; the converse is also true. Mass resolved excitation spectra (time of flight mass spectra - TOFMS) can thus be readily obtained for each conformer of the system under study. Structural assignments can then be made by an 'origin counting' procedure. Herein, we report the conformation of benzyl alcohol in the gas phase.


The TOFMS of (1) and benzyl alcohol- d_2 (5) each exhibit four distinct features, as shown in Figures 1(A) and (B). The isotope shifts for the second, third, and fourth transitions of (5) relative to those of (1) are substantial, ranging from 3% to 8% for these features. These data

*
see † page


indicate that only a single 0_0^0 transition is found for benzyl alcohol. (Similar origin assignments have been made using deuterated anisoles⁵ and styrenes⁷.) Hence, only a single stable conformation [(2) or (3) or (4)] exists for benzyl alcohol in the gas phase. 'Structural logic' involving symmetry and potential energy considerations permits a distinction amongst these three conformations to be made. The strategy involves examination of the TOFMS of suitably substituted benzyl alcohols, in particularly 4-ethylbenzyl alcohol (6), 3-fluorobenzyl alcohol (7) and 3-methylbenzyl alcohol (8).



(6)



(7)




(8)


The torsional angle $\tau(\text{C}_{\text{ortho}}-\text{C}_{\text{ipso}}-\text{C}_{\alpha}-\text{C}_{\beta})$ is ca. 90° for a sterically unhindered aromatic ethyl group. The TOFMS of (6), shown in Figure 1(C), exhibits a doubling of all the transitions for benzyl alcohol, indicating that the oxygen atom of these benzyl alcohols must be out of the plane of the aromatic ring, i.e., conformation (2) is excluded from consideration as outlined in Scheme 1. Conformations (3) and (4) can be distinguished by examination of the TOFMS of (7), which is shown in Figure 1(D). The spectra of (1) and (7) are nearly identical, indicating that the C-O bond must lie in the plane perpendicular to the phenyl ring plane and passing through the $\text{C}_{\text{ipso}}-\text{C}_4$ axis. Hence, only conformer (3) is found for these benzyl alcohols. This conclusion is confirmed by the observation of a single

origin for 3-methylbenzyl alcohol (8): a closely spaced doublet with components at 36,944.7 and 36,945.8 cm^{-1} is observed due to the torsional motion of the methyl group.⁶


Meta and para substituents on the benzyl alcohol ring system have apparently little or no essential influence on the $-\text{CH}_2\text{OH}$ conformation. In contrast, ortho fluoro substitution does alter the benzyl alcohol geometrical and electronic properties. The TOFMS of 2-fluorobenzyl alcohol (9) [Figure 2(A)] is considerably different than the TOFMS of benzyl alcohol and the 3- and 4-substituted benzyl alcohols shown in Figure 1. The progression shown in Figure 2(A) is not due to rotation of the hydroxyl group about the $\text{C}_{\text{ipso}}-\text{C}_{\alpha}$ bond, since the TOFMS of the deuterated analogue (10) is identical to that of (9). Importantly, the TOFMS of the $-\text{OD}$ analogue (11) shown in Figure 2(B) indicates significant isotope effects, strongly suggesting that the transitions depicted in Figure 2 are due to oscillatory motion of H(D) in the $\text{O}-\text{H}\cdots\text{F}$ ($\text{O}-\text{D}\cdots\text{F}$) hydrogen bond. Unlike the suggestion that (9) exists in three conformations based on microwave spectroscopy,⁸ our results are consistent with the existence of only a single stable ground state conformation. Intramolecular $\text{CH}_2-\text{OH}\cdots\text{F}$ hydrogen bonding in (9)-(11) differs significantly in the S_0 and S_1 electronic states. The observed spectroscopic transitions in the TOFMS of (9)-(11) reflect a displacement in the stable minimum energy conformations of these molecules in S_0 compared to S_1 .



(9)




(10)



(11)


We draw two major conclusions from this work: sterically unencumbered benzyl alcohols exist in the perpendicular conformation (3) in the gas phase; and the $-\text{CH}_2\text{OH}$ conformation in 2-fluorobenzyl alcohol is altered, relative to (1) by intramolecular hydrogen bonding, which causes different equilibrium positions for the $-\text{OH}$ proton in S_1 as compared to S_0 . A full paper on this work will deal with internal motion of additional heteroatom-substituted benzyl alcohols.

Plane of aromatic




2

planar



3

perpendicular



4

gauche

No. of conformations

Compound	Predicted			Observed
	(2)	(3)	(4)	
Benzyl alcohol (1)	1	1	1	1
4-Ethylbenzyl alcohol (5)	1	2	2	2
3-Fluorobenzyl alcohol (6)	2	1	2	1
3-Methylbenzyl alcohol (8)	2	1	2	1

Scheme 1

Footnote

* Observation of a single origin transition in this work implies either (a) that only a single stable ground state conformation exists or (b) that one conformation is significantly more stable than the others, assuming there is no difference between the conformations' rate of cooling in the expansion gas.

References



- 1 T. Schaefer, W. Danchura, W. Niemczura, and W. J. E. Parr, Can. J. Chem., 1978, 56, 1721.
- 2 M. Ito and M. Hirota, Bull. Chem. Soc., Jpn., 1981, 54, 2093.
- 3 M. Traetteberg, H. Østensen, and R. Seip, Acta Chem. Scand., 1980, 34A, 449.
- 4 R. J. Abraham and J. M. Bakke, Tetrahedron, 1978, 2947.
- 5 P. J. Breen, E. R. Bernstein, H. V. Secor, and J. I. Seeman, J. Am. Chem. Soc., 1989, 111, 1958.
- 6 J. I. Seeman, H. V. Secor, P. J. Breen, V. H. Grassian, and E. R. Bernstein, J. Am. Chem. Soc., 1989, 111, 3140.
- 7 V. H. Grassian, E. R. Bernstein, H. V. Secor, and J. I. Seeman, J. Phys. Chem., 1989, 93, 3470.
- 8 K.-V. Hansen and T. Pedersen, J. Mol. Structure, 1983, 97, 311.

Figure Captions


Figure 1. TOFMS of the 0_0^0 region of (A) benzyl alcohol (1), (B) benzyl alcohol- d_2 (5), 4-ethylbenzyl alcohol (6), and 3-fluorobenzyl alcohol (7). Comparison of the four transitions in (A) and (B) indicates that benzyl alcohol exists in a single conformation, since the transition energy of the second, third, and fourth transitions are all shifted slightly to the red due to the known isotope effect on vibrational motion. These four transitions are each doubled in (C) due to the presence of two conformations for (6). The near identical spectrum of (7) in (D) compared to (A) and (B) indicates that 3-fluorobenzyl alcohol exists in a single conformation.

Figure 2. TOFMS of the 0_0^0 region of (A) 2-fluorobenzyl alcohol (9) and (B) 2-fluorobenzyl alcohol-OD (11). The TOFMS of the 0_0^0 region of (10) is essentially identical to that found for (9). The dissimilarity between these spectra and the spectra shown in Figure 1 indicates that the geometry of the ground and excited states of 2-fluorobenzyl alcohol are different from each other. The isotope shifts seen in the comparison of (A) and (B) indicate that the observed spectrum is due to an oscillatory motion of the hydrogen atom in the H...F intramolecular hydrogen bond.



A




B



C




D




0

100



ENERGY (cm^{-1})



A



B



TECHNICAL REPORT DISTRIBUTION LIST, GENERAL

	<u>No.</u> <u>Copies</u>		<u>No.</u> <u>Copies</u>
Office of Naval Research Chemistry Division, Code 1113 800 North Quincy Street Arlington, VA 22217-5000	3	Dr. Ronald L. Atkins Chemistry Division (Code 385) Naval Weapons Center China Lake, CA 93555-6001	1
Commanding Officer Naval Weapons Support Center Attn: Dr. Bernard E. Douba Crane, IN 47522-5050	1	Chief of Naval Research Special Assistant for Marine Corps Matters Code OOMC 800 North Quincy Street Arlington, VA 22217-5000	1
Dr. Richard W. Drisko Naval Civil Engineering Laboratory Code L52 Port Hueneme, California 93043	1	Dr. Bernadette Eichinger Naval Ship Systems Engineering Station Code 053 Philadelphia Naval Base Philadelphia, PA 19112	1
Defense Technical Information Center Building 5, Cameron Station Alexandria, Virginia 22314	2 <u>high</u> <u>quality</u>	Dr. Sachio Yamamoto Naval Ocean Systems Center Code 52 San Diego, CA 92152-5000	1
David Taylor Research Center Dr. Eugene C. Fischer Annapolis, MD 21402-5067	1	David Taylor Research Center Dr. Harold H. Singerman Annapolis, MD 21402-5067 ATTN: Code 283	1
Dr. James S. Murday Chemistry Division, Code 6100 Naval Research Laboratory Washington, D.C. 20375-5000	1		

TECHNICAL REPORT DISTRIBUTION LIST, GENERAL

	<u>No. Copies</u>		<u>No. Copies</u>
Office of Naval Research Chemistry Division, Code 1113 800 North Quincy Street Arlington, VA 22217-5000	3	Dr. Ronald L. Atkins Chemistry Division (Code 385) Naval Weapons Center China Lake, CA 93555-6001	1
Commanding Officer Naval Weapons Support Center Attn: Dr. Bernard E. Douda Crane, IN 47522-5050	1	Chief of Naval Research Special Assistant for Marine Corps Matters Code OOMC 800 North Quincy Street Arlington, VA 22217-5000	1
Dr. Richard W. Drisko Naval Civil Engineering Laboratory Code L52 Port Hueneme, California 93043	1	Dr. Bernadette Eichinger Naval Ship Systems Engineering Station Code 053 Philadelphia Naval Base Philadelphia, PA 19112	1
Defense Technical Information Center Building 5, Cameron Station Alexandria, Virginia 22314	2 <u>high quality</u>		
David Taylor Research Center Dr. Eugene C. Fischer Annapolis, MD 21402-5067	1	Dr. Sachio Yamamoto Naval Ocean Systems Center Code 52 San Diego, CA 92152-5000	1
Dr. James S. Murday Chemistry Division, Code 6100 Naval Research Laboratory Washington, D.C. 20375-5000	1	David Taylor Research Center Dr. Harold H. Singerman Annapolis, MD 21402-5067 ATTN: Code 283	1