

2

FILE

USAARL Report No. 90-1

AD-A219 282



Six-Month Evaluation of Extended Wear Soft Contact Lenses Among Armored Troops: Part I: Clinical Findings (Reprint)

By

William G. Bachman
Bruce C. Leibrecht
John K. Crosley
Dudley R. Price
Patrick M. Leas
Gerald A. Bentley

Sensory Research Division

December 1989

DTIC ELECTE. MAR 15 1990 S E D

Approved for public release; distribution unlimited.

90 03 15 055

United States Army Aeromedical Research Laboratory
Fort Rucker, Alabama 36362-5292

Notice

Qualified requesters

Qualified requesters may obtain copies from the Defense Technical Information Center (DTIC), Cameron Station, Alexandria, Virginia 22314. Orders will be expedited if placed through the librarian or other person designated to request documents from DTIC.

Change of address

Organizations receiving reports from the U. S. Army Aeromedical Research Laboratory on automatic mailing lists should confirm correct address when corresponding about laboratory reports.

Disposition

Destroy this report when it is no longer needed. Do not return it to the originator.

Disclaimer

The views, opinions, and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy, or decision, unless so designated by other official documentation. Citation of trade names in this report does not constitute an official Department of the Army endorsement or approval of the use of such commercial items.

Human use

Human subjects participated in these studies after giving their free and informed voluntary consent. Investigators adhered to AR 70-25 and USAMRDC Reg 70-25 on Use of Volunteers in Research.

Reviewed:




BRUCE C. LEIBRECHT, Ph.D.
LTC, MS
Director, Sensory Research
Division



J.D. LAMOTHE, Ph.D.
COL, MS
Chairman, Scientific
Review Committee

Released for publication:



DAVID H. KARNEY
Colonel, MC
Commanding

REPORT DOCUMENTATION PAGE

1a REPORT SECURITY CLASSIFICATION		1b RESTRICTIVE MARKINGS	
2a SECURITY CLASSIFICATION AUTHORITY		3 DISTRIBUTION / AVAILABILITY OF REPORT Approved for public release; distribution unlimited	
2b DECLASSIFICATION / DOWNGRADING SCHEDULE			
4 PERFORMING ORGANIZATION REPORT NUMBER(S) USAARL Report No. 90-1		5 MONITORING ORGANIZATION REPORT NUMBER(S)	
6a NAME OF PERFORMING ORGANIZATION Sensory Research Division U.S. Army Aeromedical Rsch Lab	6b OFFICE SYMBOL <i>(If applicable)</i> SGRD-UAS-VS	7a NAME OF MONITORING ORGANIZATION U.S. Army Medical Research and Development Command	
6c ADDRESS (City, State, and ZIP Code) P.O. Box 577 Fort Rucker, AL 36362-5292		7b ADDRESS (City, State, and ZIP Code) Fort Detrick Frederick, MD 21701-5012	
8a NAME OF FUNDING / SPONSORING ORGANIZATION	8b OFFICE SYMBOL <i>(If applicable)</i>	9 PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER	
8c ADDRESS (City, State, and ZIP Code)		10 SOURCE OF FUNDING NUMBERS	
		PROGRAM ELEMENT NO. 62787A	PROJECT NO. 3E162787 A879
		TASK NO.	WORK UNIT ACCESSION NO. 168
11 TITLE (Include Security Classification) Six-Month Evaluation of Extended-Wear Soft Contact Lenses Among Armor Troops: Part I, Clinical Findings (U) (Reprint)			
12 PERSONAL AUTHOR(S) William G. Bachman, Bruce C. Leibrecht, John K. Crosley, Dudley R. Price, Patrick M. Leas, and Gerard A. Bentley			
13a TYPE OF REPORT	13b TIME COVERED FROM _____ TO _____	14 DATE OF REPORT (Year, Month, Day) December 1989	15 PAGE COUNT 13
16 SUPPLEMENTARY NOTATION This report is a reprint of an article published in International Contact Lens Clinic, June 1989, 16(6):162-168			
17 COSATI CODES		18 SUBJECT TERMS (Continue on reverse if necessary and identify by block number)	
FIELD	GROUP	(U) Contact lenses, (U) Military environments, (U) Vision correction, (U) Ocular physiology	
06	04		
20	06		
19 ABSTRACT (Continue on reverse if necessary and identify by block number)			
<p>This report addresses the clinical aspects of wearing contact lenses in an operational military environment. Male volunteers in an armored division wore extended-wear soft contact lenses (SCLs) or spectacles for up to 6 months, participating fully in their units' normal activities. Seventy-four percent of those successfully fitted with SCLs wore their lenses for the duration of the study, when administrative losses were factored out. More than one-third of the SCL wearers experienced one or more ocular conditions requiring at least a temporary suspension of lens wear. Corneal edema and corneal staining occurred rarely at clinically significant levels. Higher than expected rates of corneal vascularization were most likely influenced by measurement criteria. Relatively frequent conjunctival injection appeared to be largely due to local environmental factors.</p>			
20 DISTRIBUTION / AVAILABILITY OF ABSTRACT <input checked="" type="checkbox"/> UNCLASSIFIED/UNLIMITED <input type="checkbox"/> SAME AS RPT <input type="checkbox"/> DTIC USERS		21 ABSTRACT SECURITY CLASSIFICATION Unclassified	
22a NAME OF RESPONSIBLE INDIVIDUAL Chief, Scientific Information Center		22b TELEPHONE (Include Area Code) (205) 255-6907	22c OFFICE SYMBOL SGRD-UAX-SI

Accession For	
NTIS GRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-1	20



Six-Month Evaluation of Extended Wear Soft Contact Lenses among Armor Troops: Part I, Clinical Findings

William G. Bachman, OD, Bruce C. Leibrecht, PhD, John K. Crosley, OD, Dudley R. Price, MD, Patrick M. Leas, OD, and Gerard A. Bentley, OD

ABSTRACT

This report addresses the clinical aspects of wearing contact lenses in an operational military environment. Male volunteers in an armored division wore extended wear soft contact lenses (SCL) or spectacles for up to six months, participating fully in their units' normal activities. Seventy-four percent of those successfully fitted with SCL wore their lenses for the duration of the study when administrative losses were factored out. More than one-third of the SCL wearers experienced one or more ocular conditions requiring at least a temporary suspension of lens wear. Corneal edema and corneal staining occurred rarely at clinically significant levels. High rates of corneal vascularization were influenced by reporting criteria. Relatively frequent conjunctival injection appeared to be largely due to environmental factors.

1 The views of the authors do not purport to reflect the position of the Department of the Army or the Department of Defense.
 2 Citation of trade names does not constitute official Department of the Army endorsement or approval of the use of such commercial items.
 3 Human subjects participated in the study after giving their free and informed voluntary consent. The investigators adhered to AR 70-25 and USAMRDC Regulation 70-25 on Use of Volunteers in Research.

INTRODUCTION

A large proportion of U.S. Army personnel wear spectacles to correct for ametropia. Reinke¹ estimates the proportion of spectacle wearers to be greater than 48%. Unfortunately, spectacle wearing soldiers frequently face problems when interfacing with military hardware, since spectacles are minimally compatible or outright incompatible with many military systems. Examples of such systems include protective masks, binoculars, weapons sights, night vision goggles, and helmet mounted displays. To make matters worse, rain, dust, sweat, and condensation clinging to spectacle lenses can compound operational problems for ametropic troops.

Contact lenses, especially extended wear lenses, offer an appealing alternative for solving the compatibility and environmental problems faced by spectacle wearing soldiers. A number of investigators have studied contact lens wear among military personnel. As early as 1952, McGee² and Enoch² evaluated clinical, environmental, and performance aspects among 10 enlisted soldiers tested on several types of contact lenses available at the time. In the years since, military aviators have received the most frequent attention; several researchers³⁻⁹ used mainly clinical measures with sample sizes ranging from 1 to 55, while one investigator¹⁰ used a retrospective questionnaire among 7 British Army pilots. Some studies have focused on ground troops,¹¹⁻¹⁶ and reports addressing contact lens wear among

civilians in extreme environments¹⁷⁻²⁰ offer information relevant to military applications. However, the literature furnishes little data from large sample studies combining clinical and survey methodologies with military personnel.

This study was initiated to assess the safety and utility of soft contact lenses (SCL) when worn by troops in an armored division while performing their normal military duties. In the armor environment, the sighting devices found in tanks and other fighting vehicles provide excellent examples of the interface problems confronting the spectacle wearing soldier. The problems are compounded when the sights must be used in moving vehicles. The specific study objectives were: (1) estimate success rates in wearing selected extended wear contact lenses; (2) determine the impact of extended wear on ocular physiology; (3) evaluate the acceptability of wear and care aspects; and (4) assess the impact on military job performance. This report addresses findings relevant to the first two objectives; a follow-on article (to appear in the July 1989 issue) will present subjective patient responses pertaining to the latter two objectives. Bachman et al.²¹ have provided a full account of the study's methods and results.

MATERIALS AND METHODS

Subjects A total of 311 volunteers participated in this study; 215 were fitted with extended wear SCL, while 96 served as spectacle-wearing controls. Ranging in age from 18-43, all were male soldiers stationed at Ford Hood, Texas. Most of the subjects were crew members of tanks, combat vehicles (tracked personnel and weapons carriers), or air defense artillery weapons (antiaircraft missiles or guns).

The 311 subjects resulted from prescreening a large number of health records of ametropic troops. Prescreening was followed by thorough clinical evaluation to identify conditions that would medically contraindicate participation as a subject. These conditions included, but were not limited to, acute and/or subacute inflammations of the anterior segment of the eye; any disease that affected the cornea, conjunctiva, or sclera; corneal hypoesthesia; low tear breakup time or insufficient lacrimation; a requirement to take certain medications, such as diuretics and decongestants, which might adversely affect tear production; a history of moderate to severe allergy; any systemic disease that might affect the eye or be aggravated by wearing contact lenses; and refractive errors that could not be compensated adequately by available contact lens powers.

Contact Lens Materials At the time this study was initiated, soft contact lenses worn for extended periods of time offered the greatest potential to solve soldier-system interface problems. Three different types of extended wear SCL were used: (1) 71% water content (Permalens XL; CooperVision); (2) 55% water content (Hydrocurve II; Barnes-Hind); and (3) 38.5% water content (CSI T; Sola-Syntex). This mix provided reasonably broad fitting capabilities. The Permalens XL and CSI T lenses were available to correct myopia in 0.50

diopter steps from -0.50 to -8.00 diopters. The Hydrocurve II lens was available in the same myopic corrections, as well as in 0.50 diopter steps to correct hyperopia from +1.00 to +5.00 diopters. Lenses were available in sufficient quantities so that they could be dispensed to the subjects directly from stock. Cleaning solutions, cases, storage materials, and fitting procedures recommended by the respective manufacturer were used.

The SCL subjects were instructed to wear their lenses continuously for a period of 7 days plus or minus 1 day. On the 7th day of continuous wear, the lenses were to be removed 2 hours prior to bedtime, cleaned, and stored in the cases until the following morning. This conservative approach to wearing time minimized physiological risks and interference with the individual's performance of duties. Six months of wearing time were targeted for each subject. To minimize problems related to lens deposits, subjects wore the same lenses no longer than four months, at the end of which the old lenses were replaced with new ones.

Clinical Procedures Following the initial examination, each SCL wearer was fitted with lenses to provide a comfortable, stable acuity of at least 20/25 binocularly. Any volunteer who could not achieve adequate comfort, acuity, and lens stability with any of the three available lenses was eliminated. All subjects began immediately with extended lens wear. Followup visits at 24 hours, 7 days, and every 30 days thereafter were scheduled routinely.

Spectacle-wearing subjects received an initial examination similar to the initial exam for SCL participants. Each participant's refractive prescription was verified and a new pair of standard issue spectacles provided, if necessary. Two followup exams were scheduled—one 30 days after the first, and another at the end of the study.

RESULTS AND DISCUSSION

The SCL wearers were fit across a period of 14 weeks. The study spanned the months of May through December, with individual participants varying in their starting and ending dates. Consequently, each participant encountered a broad range of climatic conditions. Temperatures ranged from 102°F to 31°F, with conditions generally dry and dusty. Rainfall during the first five months of the test period averaged 1.57 inches per month, while the average during October through December was 4.67 inches per month. Relative humidity generally ranged between 35% and 70%.

The SCL and spectacle groups were quite comparable in terms of age distribution: in each group, 60% of the participants were age 25 and below. The median age of the spectacle wearers was 23.5 years (range, 18-41), while it was 24.0 years (range, 18-43) for the SCL wearers.

Visual Status The range of uncorrected visual acuity of the two groups was the same, 20/20 to 20/450. However, the mean for the SCL wearers was 20/180, while that of the spectacle wearers was 20/100. This

difference is related directly to spherical refractive errors, which for the SCL wearers ranged from +4.75 diopters to -7.50 diopters, with the mean being -2.18 diopters. For the spectacle wearers, the range was from +7.75 diopters to -6.00 diopters and the mean was -0.51 diopter.

The cylindrical refractive error data for the SCL wearers reflected the imposed limits on correctable cylindrical error allowed for this group. The range was rather narrow, from plano to a high of -1.75 diopters. The spectacle wearers, on the other hand, exhibited a wide range extending to -5.00 diopters. For the SCL wearers the mean cylinder was -0.39 diopters, while the mean for the spectacle wearers was -1.45 diopters.

A summary of the corrected binocular acuities for selected exams is contained in Table 1. Acuities recorded for the SCL wearers' initial exam were obtained as part of the refraction; the acuities for the remaining exams were recorded through the habitually worn contact lenses. Both the initial and the final exam acuities for the spectacle wearers were obtained as part of a complete eye exam.

During their initial exam, 99% of the SCL wearers were correctable to 20/20 or better. Among the spectacle wearers at the initial exam, 95% were corrected to 20/20 or better. At the 7-day exam, the acuities of the SCL subjects were somewhat reduced. This reduction may have been related to the inability of the soft lenses to fully compensate for allowable astigmatism, initial adjustment to SCL wear, and/or a possible need to change lens parameters.

The improvement in SCL wearers' acuity noted at the 90-day visit could have been due partially to the attrition of subjects who were having problems with acuity. It was also likely related to progressive adaptation to SCL wear.

A comparison of the final corrected acuities of both groups shows the proportion achieving 20/20 or better was 95% for SCL subjects and 97% for spectacle wearers.

Ocular Physiology Summary data from selected biomicroscopy examinations (initial, 90-day, and final) appear in Table 2. The classification codes are those

recommended by the Food and Drug Administration for clinical investigations.

Vascularization. Table 2 shows the percentages of eyes exhibiting vascularization. It is readily apparent that vessel ingrowth increased over the course of the study. The reason for the high rates of vascularization found is the stringent criterion to report any measurable amount of vascularization. Zucarro, Thayer, and Poland²² in a 5-year study of SCL wearers reported vascularization in only 3% of all followup examinations, but recorded only occurrences of at least 1.5 mm vessel extension inside the limbus. Nilsson and Persson²³ reported no vascularization at all in a 2-year study of extended wear contact lens patients. They defined vascularization as growth greater than 1.25 mm. It appears then that extensions of 1-1.5 mm into the cornea have not been considered significant.

Injection. Table 2 summarizes the percentages of eyes exhibiting injection over the course of the study. The incidence of injection is much higher than found in the studies of Zucarro, Thayer, and Poland,²² Nilsson and Persson,²³ and Rengstorff et al.²⁴ The SCL wearing soldiers in this study seemed predisposed to injection, as did their spectacle wearing counterparts. This may have been related to the environment in which they worked and their constant exposure to local irritants, such as dust, wind, smoke and fumes.

Staining. As can be seen in Table 2, the percentage of eyes exhibiting staining of any kind was very low.

Edema. As Table 2 shows, the percentage of eyes exhibiting moderate degrees of microedema or gross edema was very small.

Other Complications. This classification includes observations not discussed above. Table 2 shows the only unusual occurrence was the high incidence of follicular hypertrophy. This is attributed to the endemic occurrence of mild vernal conjunctivitis at the test installation during the study period. The "other" classification in this category includes such observations as papillae, pingueculae, blepharitis, and coated lenses.

Suspension of Lens Wear In accordance with accepted clinical practice and the terms of the approved research protocol, SCL wear was suspended tempo-

TABLE 1
Corrected Acuity (Binocular) at Selected Examinations

Visual Acuity	SCL Wearers				Spectacle Wearers	
	Initial (n = 215)	7-day (n = 176)	90-day (n = 109)	Final (n = 84)	Initial (n = 96)	Final (n = 70)
20/20 or better	99%	90%	95%	95%	95%	97%
20/25	1%	8%	5%	3%	4%	2%
20/30	0	1%	0	2%	1%	1%
20/40 or poorer	0	1%	0	0	0	0

TABLE 2
Percentage of Eyes Exhibiting Biomicroscopy Classifications (SCL Wearers)

Classification	Initial (n = 430)	90-day (n = 236)	Final (n = 240)*
VASCULARIZATION			
Ingrowth, 1 quadrant	16%	28%	24%
Ingrowth, >1 quadrant	4%	20%	33%
Continuing growth, <2 mm	—	3%	3%
Continuing growth, >2 mm	—	<1%	—
INJECTION			
Mild congestion	28%	51%	31%
Severe congestion	6%	12%	20%
Hyperemia	—	2%	4%
STAINING			
Minimal stippling	1%	5%	3%
Superficial punctate	—	3%	<1%
Epithelial abrasions	—	<1%	—
EDEMA			
Slight, localized	2%	4%	13%
Slight, generalized	—	<1%	2%
Moderate, localized	—	—	1%
Moderate, generalized	—	—	2%
Vertical striae	—	<1%	—
OTHER COMPLICATIONS			
Increased sebaceous secretion	3%	—	<1%
Follicular hypertrophy	10%	5%	10%
Other (see text)	9%	<1%	5%

*Includes some cases of suspension that were deferred for disposition until the final exam.

rarily when ocular complications developed. At least one period of suspended wear occurred for 69 SCL wearers (42% of the average census) during the course of the study. The various conditions resulting in suspended SCL wear are presented in Table 3. Some individuals were suspended more than once (none more than three times), resulting in 87 cases. The most common cause of suspension was inflammation of some segment of the anterior portion of the eye or ocular adnexa, accounting for 41 percent of the total number of suspensions. Abrasions, staining, and epithelial de-

fects of the cornea collectively accounted for 29% of the total number of suspensions.

Attrition Ocular conditions that posed unacceptable jeopardy to the SCL wearers occurred occasionally, necessitating removal of subjects from the study. In addition, a number of SCL wearers were discontinued before the end of the study due to administrative circumstances or self-withdrawal. The cases of attrition are listed in Table 4 according to the nature of the cause. A total of 64 SCL subjects failed to complete the study for administrative or personal reasons.

TABLE 3
Medically-Related SCL Wear Suspensions

Cause	Number of Suspensions
Conjunctivitis	23
Corneal abrasion	18
Corneal staining	9
Overwear syndrome	6
Giant papillary conjunctivitis	6
Corneal edema	4
Iritis	3
Neovascularization	3
Keratitis	3
Corneal ulcer	1
Other*	11

*Includes dermatitis (eyelids), phlyctenule, ocular hypertension, eye trauma, use of medication, sensitivity to solutions, and decreased visual acuity.

The number of SCL participants failing to complete the study for medically-related reasons was 40. This figure was compared to the census at the start of the study, excluding those subjects who eventually withdrew for nonmedical reasons, to yield a six-month attrition rate of 26%. Three conditions—discomfort, dissatisfaction with acuity, and giant papillary conjunctivitis—accounted for 68% of the cases of medically-related attrition.

Suspension and attrition both reflect the occurrence of ocular complications. Accordingly, a comprehensive picture of ocular complications can be obtained by combining the data for suspension and attrition. A total of 79 SCL subjects developed at least one ocular complication. This translates into a proportion equal to 47% when the average census across the entire study is used for computation. In other words, on a six-month equivalent basis, 47% of those wearing SCL developed one or more ocular condition(s) requiring at least a short suspension of SCL wear.

The rate of occurrence of ocular complications declined as the study progressed. The trend for both suspension-precipitating complications and total complications can be seen in Table 5, where rates are based on average monthly census figures. A similar declining trend has been noted elsewhere in the literature²⁰ and may be related to progressive attrition of complication-prone subjects as cumulative SCL wearing time increases.

TABLE 4
Attrition among SCL Wearers

Cause	Number of Attritions
A. Nonmedical	
Missed appointments	18
Discharge from service	21
Reassignment	13
Lack of interest	6
Lenses not available	3
Lost or damaged lenses:	2
Extended travel	1
	<hr/> 64
B. Medically related	
Discomfort	14
Dissatisfaction with acuity	5
Discomfort and dissatisfaction with acuity	2
Giant papillary conjunctivitis	6
Neovascularization (>2 mm)	6
Decreased visual acuity (>7 days duration)	2
Blepharitis	1
Corneal staining	1
Corneal stromal infiltrates	1
Tight lens syndrome	1
Insertion problems	1
	<hr/> 40

Among the spectacle wearers, 16 subjects were discontinued for administrative and personal reasons. No ocular complications or cases of medically-related attrition occurred among the spectacle wearers. It should be noted the spectacle wearing participants were not examined as often as the SCL wearers.

SCL Wear Success Rates Those SCL participants at the end of the study who had not been removed for medical reasons were defined as medically successful SCL wearers. However, those SCL participants, who were discontinued for administrative or personal reasons, can be labelled neither medically unsuccessful nor successful. Consequently, nonmedical attritions should be factored out when computing success rates for this study. Of the 215 SCL subjects who started the study, a total of 151 remained in the study to a definitive disposition. Of these, 111 (74%) successfully completed the study. In other words, when nonmedical attritions were factored out, three out of every four SCL wearers reached the end of the study without being discontinued for medically-related reasons. This six-month rate was somewhat artificially constrained by the limited types and parameters of SCL used. On the other hand, the rate may have been elevated by the well-motivated participants.

TABLE 5
Monthly Rates of Ocular Complications among SCL Wearers

Day	Average Census	SCL Wear Suspensions		Total Ocular Complications	
		Number	Rate	Number	Rate
1-30	208	27	13.0%	37	17.8%
31-60	194	15	7.7%	23	11.9%
61-90	180.5	17	9.4%	21	11.6%
91-120	162.5	11	6.8%	17	10.5%
121-150	141.5	8	5.7%	11	7.8%
151-180	125.5	7	5.6%	13*	10.4%*
>180	—	2	—	5	—

*Includes several cases of suspension which were deferred for final disposition until the end of the study

CONCLUSIONS

Because of methodological limitations, the results of this study should be generalized with caution. The major findings obtained in the armor environment support the following conclusions.

1. When SCL wearers discontinued for administrative reasons were factored out, 74% of those fitted successfully completed the study.
2. More than one-third of the SCL wearers experienced one or more ocular conditions requiring at least a temporary suspension of SCL wear.
3. Both corneal edema and corneal staining occurred rarely at clinically significant levels.
4. Corneal vascularization occurred frequently. This was influenced by stringent classification criteria.
5. Conjunctival injection was common, apparently due largely to environmental factors.

Acknowledgments

Special thanks go to the 2d Armored Division, Fort Hood, Texas for providing volunteer participants and administrative support. The U.S. Army Training and Doctrine Command's Combined Arms Test Activity, Fort Hood, Texas, provided extensive support in planning and conducting the entire study, with Mr. Terry Glover serving as project officer. Dr. Stephen Risk and Dr. Theodore Felton provided invaluable clinical optometric support. We thank Ms. Carolyn Johnson and Ms. Jimmie Henderson for reproof typing, and Staff Sergeant Nonion Fallana and Sergeant Rosalinda Ibanez for technical support.

References

1. Reinke AR. *Contact Lenses for Military Environment Requirements: Survey and Study of Feasibility (ACN 17167)*. Fort Sam Houston, Texas: U.S. Army Combat Developments Command, 1970.
2. McGraw JL, Enoch JM. *Contact Lenses: An Evaluation Study*. Fort Knox, Kentucky: U.S. Army Medical Research Laboratory USAMRI, Report No. 99-1952.

3. Crosley JK, Braun EG, Bailey RW. *Soft (Hydrophilic) Contact Lenses in U.S. Army Aviation: An Investigative Study of the Bausch and Lomb Solfens(TM)*. Fort Rucker, Alabama: USAARL Report No. 74-10, 1974.
4. Nilsson K, Rengstorff RH. Continuous wearing of Duragel (R) contact lenses by Swedish air force pilots. *Am J Optom Physiol Opt* 1979;56(6):356-8.
5. Polishuk A, Raz D. Soft hydrophilic contact lenses in civil and military aviation. *Aviat Space Environ Med* 1975;46(9):1188-90.
6. Brennan DH, Girvin JK. The flight acceptability of soft contact lenses—an environmental trial. *Aviat Space Environ Med* 1985;56(1):43-8.
7. Tredici TJ, Flynn WJ. The use of contact lenses by USAF aviators. *Aviat Space Environ Med* 1987;58(5):438-43.
8. Flynn WJ, Block MG, Tredici TJ, Provines WF. Effect of positive acceleration on soft contact lens wear. *Aviat Space Environ Med* 1987;58(6):581-7.
9. Flynn WJ, Miller RE, Tredici TJ, Block MG. Soft contact lens wear at altitude: effects of hypoxia. *Aviat Space Environ Med* 1989;59(1):44-8.
10. Braithwaite MG. The use of contact lenses by Army air crew. *J R Army Med Corps* 1983;129:43-5.
11. van Norren D. Contact lenses in the military service. *Am J Optom Physiol Opt* 1984;61(7):441-7.
12. Rouwen AJ, Pinckers AJ, van Pad Bosch AA, Punt H, Doesburg WH, Lemmens WA. Visual acuity, spectacle blur and slit-lamp biomicroscopy on asymptomatic contact-lens-wearing recruits. *Graefes Arch Clin Exp Ophthalmol* 1983;221(2):73-7.
13. Rouwen AJ, Rosenbrand RM. High water content soft lenses used for flexible daily/extended wear for military personnel. *Int Eye Care* 1986;2(8):435-9.
14. Rengstorff RH. Contact lenses and basic training in the U.S. Army. *Milit Med* 1965;130(4):419-21.
15. Rouwen AJ, Punt H, Pinckers AJ, Doesburg WH, Lemmens WA. Contact lens wear during military field maneuvers. *Contactologia* 1986;8D:136-41.
16. Rippel W. Contact lenses and paratroopers. *Klin Mbl Augenheilkunde* 1979;174:284-6.
17. Kok-van-Aalphen C, van der Linden J, Visser R, Bol A. Protection of the police against tear gas with soft lenses. *Milit Med* 1985;150:451-4.
18. Hart LG. Wearing contact lenses in space shuttle operations. *Aviat Space Environ Med* 1985;56(12):1224-5.

19. Socks JF. *Use of Contact Lenses for Cold Weather Activities Results of a Survey* Groton, Connecticut: Naval Submarine Medical Research Laboratory NSMRL Report No 969, 1981
20. Coe JE, Douglas RB. The effect of contact lenses on ocular responses to sulphur dioxide. *J Soc Occup Med* 1982;32(2) 92-4
21. Bachman WG, Leibrecht BC, Crosley JK, Price DR, Bentley GA, Leas PM. *An Operational Evaluation of Extended-Wear Soft Contact Lenses in an Armored Division* Fort Rucker, Alabama. U.S. Army Aeromedical Research Laboratory USAARL Report No 87-12, 1987
22. Zucarro V, Thayer T, Poland J. Hydrocurve II extended-wear a five-year study. *Int Eyecare* 1985;1(5) 379-85
23. Nilsson E, Persson G. Low complication rate in extended-wear of contact lenses. *Acta Ophthalmol* 1986;64 88-92.
24. Rengstorff RH, Nilsson KT, Sylvanuer AE. 2000 extended wear cases: a retrospective survey of contact lens complications. *J Br CL Assoc* 1987;10(1) 13-15
25. Koetting R. A retrospective study of patients fitted with extended-wear lenses. *Contacto* 1983; 27 7-8

Clinical Implications

This observational study focused on the success rate and ocular physiological response associated with the extended wearing of soft contact lenses (SCL) by armor troops over the course of six months. Reported results indicate that a significant number of people (74%) completing the study were able to tolerate the extended wear of the study's SCL and that a substantial number of SCL wearers (better than 40%) did develop periodic problems that required at least temporary suspension of SCL wear.

The medically related reasons for suspension of SCL wear were generally of a serious nature, reminding us that the application of the extended wearing of the SCL does offer significant ocular risks along with potential benefits. The impact of lens care system and patient compliance on the study results were not reported. Care system and compliance issues prompt the thought that disposable SCL for extended wear (a modality not available at the time that this study was conducted) may improve the success rate for a patient population such as those in armor troops.

I must agree with the authors that the results of the study should be generalized with caution due to limitations in various areas of the study methodology.

Chris Snyder, OD
School of Optometry
The University of Alabama, Birmingham
Birmingham, AL 35294

About the Authors

William G. Bachman received the Doctor of Optometry degree from Southern College of Optometry and the Master of Science in Physiological Optics from the University of Alabama, Birmingham. He recently retired after serving 20 years in the Army as both a clinician and a researcher, particularly in the area of contact lenses. He is currently an Assistant Professor at the School of Optometry, University of Missouri, St. Louis.



Bruce C. Leibrecht holds a PhD in Experimental Psychology from Michigan State University. He has served as a Research Psychologist with the U.S. Army Medical Research and Development Command since 1969. He is currently Director of the Sensory Research Division of the U.S. Army Aeromedical Research Laboratory, a position he has held since 1981. Having been active in several areas of biomedical research, he is the author of a number of published scientific articles.



Initial distribution

Commander
U.S. Army Natick Research
and Development Center
ATTN: Documents Librarian
Natick, MA 01760

Naval Submarine Medical
Research Laboratory
Medical Library, Naval Sub Base
Box 900
Groton, CT 06340

Commander/Director
U.S. Army Combat Surveillance
& Target Acquisition Lab
ATTN: DELCS-D
Fort Monmouth, NJ 07703-5304

Commander
10th Medical Laboratory
ATTN: Audiologist
APO New York 09180

Commander
Naval Air Development Center
Biophysics Lab
ATTN: G. Kydd
Code 60B1
Warminster, PA 18974

Naval Air Development Center
Technical Information Division
Technical Support Detachment
Warminster, PA 18974

Commanding Officer
Naval Medical Research
and Development Command
National Naval Medical Center
Bethesda, MD 20014

Under Secretary of Defense
for Research and Engineering
ATTN: Military Assistant
for Medical and Life Sciences
Washington, DC 20301

Commander
U.S. Army Research Institute
of Environmental Medicine
Natick, MA 01760

U.S. Army Avionics Research
and Development Activity
ATTN: SAVAA-P-TP
Fort Monmouth, NJ 07703-5401

U.S. Army Research and Development
Support Activity
Fort Monmouth, NJ 07703

Chief, Benet Weapons Laboratory
LCWSL, USA ARRADCOM
ATTN: DRDAR-LCB-TL
Watervliet Arsenal, NY 12189

Commander
Man-Machine Integration System
Code 602
Naval Air Development Center
Warminster, PA 18974

Commander
Naval Air Development Center
ATTN: Code 6021 (Mr. Brindle)
Warminster, PA 18974

Commanding Officer
Harry G. Armstrong Aerospace
Medical Research Laboratory
Wright-Patterson
Air Force Base, OH 45433

Director
Army Audiology and Speech Center
Walter Reed Army Medical Center
Washington, DC 20307-5001

Director
Walter Reed Army Institute
of Research
Washington, DC 20307-5100

HQ DA (DASG-PSP-0)
5109 Leesburg Pike
Falls Church, VA 22041-3258

Naval Research
Laboratory Library
Code 1433
Washington, DC 20375

Harry Diamond Laboratories
ATTN: Technical Infor-
mation Branch
2800 Powder Mill Road
Adelphi, MD 20783-1197

U.S. Army Materiel Systems
Analysis Agency
ATTN: Reports Processing
Aberdeen Proving Ground
MD 21005-5017

U.S. Army Ordnance Center
and School Library
Building 3071
Aberdeen Proving Ground,
MD 21005-5201

U.S. Army Environmental Hygiene
Agency
Building E2100
Aberdeen Proving Ground,
MD 21010

Technical Library
Chemical Research
and Development Center
Aberdeen Proving Ground,
MD 21010-5423

Commander
U.S. Army Institute
of Dental Research
Walter Reed Army Medical Center
Washington, DC 20307-5300

Naval Air Systems Command
Technical Air Library 950D
Rm 278, Jefferson Plaza II
Department of the Navy
Washington, DC 20361

Naval Research Laboratory Library
Shock and Vibration Infor-
mation Center, Code 5804
Washington, DC 20375

Director
U.S. Army Human Engineer-
ing Laboratory
ATTN: Technical Library
Aberdeen Proving Ground,
MD 21005-5001

Commander
U.S. Army Test
and Evaluation Command
ATTN: AMSTE-AD-H
Aberdeen Proving Ground,
MD 21005-5055

Director
U.S. Army Ballistic
Research Laboratory
ATTN: DRXBR-OD-ST Tech Reports
Aberdeen Proving Ground,
MD 21005-5066

Commander
U.S. Army Medical Research
Institute of Chemical Defense
ATTN: SGRD-UV-AO
Aberdeen Proving Ground,
MD 21010-5425

Commander
U.S. Army Medical Research
and Development Command
ATTN: SGRD-RMS (Ms. Madigan)
Fort Detrick, Frederick,
MD 21701

Commander
U.S. Army Medical Research
Institute of Infectious Diseases
Fort Detrick, Frederick,
MD 21701

Director, Biological
Sciences Division
Office of Naval Research
600 North Quincy Street
Arlington, VA 22217

Commander
U.S. Army Materiel Command
ATTN: AMCDE-XS
5001 Eisenhower Avenue
Alexandria, VA 22333

Commandant
U.S. Army Aviation
Logistics School
ATTN: ATSQ-TDN
Fort Eustis, VA 23604

U.S. Army Training
and Doctrine Command
ATTN: ATCD-ZX
Fort Monroe, VA 23651

Structures Laboratory Library
USARTL-AVSCOM
NASA Langley Research Center
Mail Stop 266
Hampton, VA 23665

Naval Aerospace Medical
Institute Library
Bldg 1953, Code 102
Pensacola, FL 32508

Command Surgeon
U.S. Central Command
MacDill Air Force Base
FL 33608

Air University Library
(AUL/LSE)
Maxwell AFB, AL 36112

Commander
U.S. Army Biomedical Research
and Development Laboratory
ATTN: SGRD-UBZ-I
Fort Detrick, Frederick,
MD 21701

Defense Technical
Information Center
Cameron Station
Alexandria, VA 22313

U.S. Army Foreign Science
and Technology Center
ATTN: MTZ
220 7th Street, NE
Charlottesville, VA 22901-5396

Director,
Applied Technology Laboratory
USARTL-AVSCOM
ATTN: Library, Building 401
Fort Eustis, VA 23604

U.S. Army Training
and Doctrine Command
ATTN: Surgeon
Fort Monroe, VA 23651-5000

Aviation Medicine Clinic
TMC #22, SAAF
Fort Bragg, NC 28305

U.S. Air Force Armament
Development and Test Center
Eglin Air Force Base, FL 32542

U.S. Army Missile Command
Redstone Scientific
Information Center
ATTN: Documents Section
Redstone Arsenal, AL 35898-5241

U.S. Army Research and Technology
Laboratories (AVSCOM)
Propulsion Laboratory MS 302-2
NASA Lewis Research Center
Cleveland, OH 44135

AFAMRL/HEX
Wright-Patterson AFB, OH 45433

University of Michigan
NASA Center of Excellence
in Man-Systems Research
ATTN: R. G. Snyder, Director
Ann Arbor, MI 48109

John A. Dellinger,
Southwest Research Institute
P. O. Box 28510
San Antonio, TX 78284

Product Manager
Aviation Life Support Equipment
ATTN: AMCPM-ALSE
4300 Goodfellow Blvd.
St. Louis, MO 63120-1798

Commander
U.S. Army Aviation
Systems Command
ATTN: AMSAV-ED
4300 Goodfellow Blvd
St. Louis, MO 63120

Commanding Officer
Naval Biodynamics Laboratory
P.O. Box 24907
New Orleans, LA 70189

U.S. Army Field Artillery School
ATTN: Library
Snow Hall, Room 14
Fort Sill, OK 73503

Commander
U.S. Army Health Services Command
ATTN: HSOP-SO
Fort Sam Houston, TX 78234-6000

U.S. Air Force Institute
of Technology (AFIT/LDEE)
Building 640, Area B
Wright-Patterson AFB, OH 45433

Henry L. Taylor
Director, Institute of Aviation
University of Illinois-
Willard Airport
Savoy, IL 61874

COL Craig L. Urbauer, Chief
Office of Army Surgeon General
National Guard Bureau
Washington, DC 50310-2500

Commander
U.S. Army Aviation
Systems Command
ATTN: SGRD-UAX-AL (MAJ Lacy)
4300 Goodfellow Blvd., Bldg 105
St. Louis, MO 63120

U.S. Army Aviation Systems Command
Library and Information
Center Branch
ATTN: AMSAV-DIL
4300 Goodfellow Blvd
St. Louis, MO 63120

Federal Aviation Administration
Civil Aeromedical Institute
CAMI Library AAC 64D1
P.O. Box 25082
Oklahoma City, OK 73125

Commander
U.S. Army Academy
of Health Sciences
ATTN: Library
Fort Sam Houston, TX 78234

Commander
U.S. Army Institute
of Surgical Research
ATTN: SGRD-USM (Jan Duke)
Fort Sam Houston, TX 78234-6200

Director of Professional Services
AFMSC/GSP
Brooks Air Force Base, TX 78235

U.S. Army Dugway Proving Ground
Technical Library
Bldg 5330
Dugway, UT 84022

U.S. Army Yuma Proving Ground
Technical Library
Yuma, AZ 85364

AFFTC Technical Library
6520 TESTG/ENXL
Edwards Air Force Base,
CAL 93523-5000

Commander
Code 3431
Naval Weapons Center
China Lake, CA 93555

Aeromechanics Laboratory
U.S. Army Research
and Technical Labs
Ames Research Center,
M/S 215-1
Moffett Field, CA 94035

Sixth U.S. Army
ATTN: SMA
Presidio of San Francisco,
CA 94129

Commander
U.S. Army Aeromedical Center
Fort Rucker, AL 36362

U.S. Air Force School
of Aerospace Medicine
Strughold Aeromedical Library
Documents Section, USAFSAM/TSK-4
Brooks Air Force Base, TX 78235

Dr. Diane Damos
Department of Human Factors
ISSM, USC
Los Angeles, CA 90089-0021

U.S. Army White Sands
Missile Range
Technical Library Division
White Sands Missile Range,
NM 88002

U.S. Army Aviation Engineering
Flight Activity
ATTN: SAVTE-M (Tech Lib)
Stop 217
Edwards Air Force Base,
CA 93523-5000

Ms. Sandra G. Hart
Ames Research Center
MS 239-5
Moffett Field, CA 94035

Commander
Letterman Army Institute
of Research
ATTN: Medical Research Library
Presidio of San Francisco,
CA 94129

Mr. Frank J. Stagnaro, ME
Rush Franklin Publishing
300 Orchard City Drive
Campbell, CA 95008

Commander
U.S. Army Medical Materiel
Development Activity
Fort Detrick, Frederick,
MD 21701-5009

Commander, U.S. Army
Aviation Center
Directorate
of Combat Developments
Bldg 507
Fort Rucker, AL 36362

Chief
Army Research Institute
Field Unit
Fort Rucker, AL 36362

Commander
U.S. Army Safety Center
Fort Rucker, AL 36362

U.S. Army Aircraft Development
Test Activity
ATTN: STEBG-MP-QA
Cairns AAF
Fort Rucker, AL 36362

Commander
U.S. Army Medical Research
and Development Command
ATTN: SGRD-PLC (COL Sedge)
Fort Detrick, Frederick
MD 21701

MAJ John Wilson
TRADOC Aviation LO
Embassy of the United States
APO New York 09777

Netherlands Army Liaison Office
Building 602
Fort Rucker, AL 36362

British Army Liaison Office
Building 602
Fort Rucker, AL 36362

Italian Army Liaison Office
Building 602
Fort Rucker, AL 36362

Australian Army Liaison Office
Building 602
Fort Rucker, AL 36362

Directorate
of Training Development
Bldg 502
Fort Rucker, AL 36362

Chief
Human Engineering Laboratory
Field Unit
Fort Rucker, AL 36362

Commander
U.S. Army Aviation Center
and Fort Rucker
ATTN: ATZQ-T-ATL
Fort Rucker, AL 36362

President
U.S. Army Aviation Board
Cairns AAF
Fort Rucker, AL 36362

Dr. William E. McLean
Human Engineering Laboratory
ATTN: SLCHE-BR
Aberdeen Proving Ground,
MD 21005-5001

Canadian Army Liaison Office
Building 602
Fort Rucker, AL 36362

German Army Liaison Office
Building 602
Fort Rucker, AL 36362

French Army Liaison Office
Building 602
Fort Rucker, AL 36362

Brazilian Army Liaison Office
Building 602
Fort Rucker, AL 36362

Commandant
Royal Air Force Institute
of Aviation Medicine
Farnborough Hants UK GU14 6SZ