

REPORT DOCUMENTATION PAGE

Form Approved  
OMB No. 0704-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.

1. AGENCY USE ONLY (Leave blank) 2. REPORT DATE 3. REPORT TYPE AND DATES COVERED  
Journal Article

4. TITLE AND SUBTITLE  
Ab Initio and Semi-empirical Calculations on the Oxyhalides of Silicon

5. FUNDING NUMBERS  
--

AUTHOR(S)  
James J.P. Stewart and Mark S. Gordon

PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES)  
Frank J. Seiler Research Laboratory  
USAF Academy  
Colorado 80840-6528

8. PERFORMING ORGANIZATION REPORT NUMBER  
FJSRL-JR-90-0004

SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)  
Air Force Office of Scientific Research  
Boiling AFB DC 20332

10. SPONSORING/MONITORING AGENCY REPORT NUMBER

6. SUPPLEMENTARY NOTES

REF ID: A111111  
SEP 17 1990

12a. DISTRIBUTION/AVAILABILITY STATEMENT  
DISTRIBUTION UNLIMITED

12b. DISTRIBUTION CODE

13. ABSTRACT (Maximum 200 words)  
In 1982 the heats of formation of the silicon oxyhalides were determined by experiment to be -267.9 Kcal/mole (SiOF<sub>2</sub>), -167.8 (SiOCl<sub>2</sub>), -137.4 (SiOBr<sub>2</sub>), and -99.4 (SiOI<sub>2</sub>). MNDO-PM3 semiempirical calculations indicate that the heats of formation are nearer to -229.4, -121.2, -94.1, and -50.2 Kcal/mole respectively. Ab-initio calculations using a 6-31G(d) basis set confirm the semiempirical results for the fluoride and chloride systems. *Keywords:*

**BEST AVAILABLE COPY**

14. SUBJECT TERMS  
PM3; thermochemical data; silicon oxyhalides; SiOF<sub>2</sub>; SiOCl<sub>2</sub>; SiOBr<sub>2</sub>; SiOI<sub>2</sub>  
*Keywords: (IS)*

15. NUMBER OF PAGES  
4

16. PRICE CODE

17. SECURITY CLASSIFICATION OF REPORT  
UNCLASSIFIED

18. SECURITY CLASSIFICATION OF THIS PAGE  
UNCLASSIFIED

19. SECURITY CLASSIFICATION OF ABSTRACT  
UNCLASSIFIED

20. LIMITATION OF ABSTRACT  
NONE

AD-A226 590

DTIC FILE COPY

# AB INITIO AND SEMI-EMPIRICAL CALCULATIONS ON THE OXYHALIDES OF SILICON

JAMES J. P. STEWART

*The Frank J. Seiler Research Laboratory, U.S. Air Force Academy,  
Colorado Springs, CO 80840*

and

MARK S. GORDON

*Department of Chemistry, North Dakota State University,  
Fargo, North Dakota 58105*

*(Received January 20, 1989; in final form April 30, 1989)*


In 1982 the heats of formation of the silicon oxyhalides were determined by experiment to be  $-267.9$  Kcal/mole ( $\text{SiOF}_2$ ),  $-167.8$  ( $\text{SiOCl}_2$ ),  $-137.4$  ( $\text{SiOBr}_2$ ), and  $-99.4$  ( $\text{SiOI}_2$ ). MNDO-PM3 semiempirical calculations indicate that the heats of formation are nearer to  $-229.4$ ,  $-121.2$ ,  $-94.1$ , and  $-50.2$  Kcal/mole, respectively. *Ab-initio* calculations using a 6-31G(d) basis set confirm the semiempirical results for the fluoride and chloride systems.

*Key words:* PM3; thermochemical data; silicon oxyhalides;  $\text{SiOF}_2$ ;  $\text{SiOCl}_2$ ;  $\text{SiOBr}_2$ ;  $\text{SiOI}_2$ .

## INTRODUCTION

While surveying the applicability of a new semi-empirical model,<sup>1</sup> MNDO-PM3, to silicon systems, various anomalies in thermodynamic data became evident. In their investigations on halogen-filled incandescent lamps, Dittmer and Niemann had been frustrated by the lack of reliable thermochemical data, and had determined the experimental heats of formation for a large number of main-group compounds including the silicon oxyhalides.<sup>2</sup> Their recommended heats of formation for several silicon compounds, based on their thermochemical measurements and critically revised literature data, are given in Table I, along with the results of PM3 calculations. In order to allow comparison with other work, we include their cited literature data. References put in parentheses relate to data which differs from their recommended values by more than 2.4 Kcal/mole.

From these experimental results, the predicted heats for the metathetic reactions of the type



are  $-78.3$  (F),  $-91.8$  (Cl),  $-94.1$  (Br), and  $-95.5$  (I) Kcal/mole. From PM3 calculations, the corresponding heats of reaction are  $+24.0$  (F),  $+3.3$  (Cl),  $+8.6$  (Br), and  $+2.7$  (I) Kcal/mole. The differences between calculated and experimental heats of reaction were surprising and indicate that either the

TABLE I  
Reported recommended experimental heats of formation

Compound	H <sub>f</sub> (298) Kcal/mole	Ref.*	Calc(PM3) Kcal/mole
SiO <sub>2</sub>	-72.9	a, b	-88.9
SiF <sub>4</sub>	-384.6	c, d, e	-390.6
SiCl <sub>4</sub>	-170.9	(a, f, g, h, i)	-156.4
SiBr <sub>4</sub>	-107.8	(j, d, h, k, l)	-107.9
SiI <sub>4</sub>	-30.4	c, g, h, m, j	-14.2
SiOF <sub>2</sub>	-267.9	a	-229.4
SiOCl <sub>2</sub>	-167.8		-121.2
SiOBr <sub>2</sub>	-137.4		-94.1
SiOI <sub>2</sub>	-99.4		-50.2

\* Taken from Reference 1.

<sup>a</sup>JANAF Thermochemical Tables, 2nd ed., Dow Chemicals Midland, Michigan, (1971); <sup>b</sup>K. F. Zmbov, L. L. Ames and J. L. Margrave, *High Temp. Sci.* **5**, 235 (1973); <sup>c</sup>JANAF Thermochemical Tables, 1978 Supplement, *J. Phys. Chem. Ref. Data* **7**, 793 (1978); <sup>d</sup>J. D. McDonald, C. H. Williams, J. C. Thompson, and J. L. Margrave, *Adv. in Chem. Ser.* **72**, 261 (1968); <sup>e</sup>A. F. Vorob'ev V. P. Kolesov and S. M. Skuratov, *Zh. Neorgan. Khim.* **5**, 1902 (1960); <sup>f</sup>I. Barin and O. Knacke, "Thermochemical Properties of Inorganic Substances," Springer-Verlag, Berlin 1973; <sup>g</sup>H. Schafer, H. Bruderreck and B. Morcher, *Z. Anorg. Allg. Chem.*, **352**, 122, (1967); <sup>h</sup>H. Schafer and H. Heine, *Z. Anorg. Allg. Chem.*, **332**, 25 (1964); <sup>i</sup>E. Wolf, *Z. Anorg. Allg. Chem.*, **313**, 228 (1961); <sup>j</sup>I. Barin, O. Knacke and O. Kubaschewski, "Thermochemical Properties of Inorganic Substances," Supplement, Springer-Verlag, Berlin 1977; <sup>k</sup>E. Wolf and C. Herbst, *Z. Anorg. Allg. Chem.*, **347**, 113 (1966); <sup>l</sup>E. Wolf, W. Stahn and M. Schonherr, *Z. Anorg. Allg. Chem.*, **319**, 168 (1962); <sup>m</sup>G. A. Kokovin and T. Thieme, *Z. Phys. Chemie (Leipzig)* **259**, 1129 (1978).

experimental or computational results are severely in error. Dittmer and Niemann commented on the fact that their enthalpies of formation were more negative than those assumed by JANAF, but noted that their values were supported by tungsten transport measurements in an SiF<sub>4</sub>/O<sub>2</sub> mixture. There was nothing in the computational method to suggest that errors of this magnitude were likely. The average error in heats of formation for 763 compounds was 8.6 Kcal/mole and the standard deviation 12.4 Kcal/mole.<sup>1</sup> For 24 normal-valent silicon-halogen compounds, the average error (Reference 1) at the PM3 level was 6.5 Kcal/mol, while for three normal-valent compounds involving an Si-O bond the average error was 7.6 Kcal/mol. Interestingly, of the three Si-O systems studied, SiO<sub>2</sub> had the largest error in the heat of formation, being too stable by 16 Kcal/mol. From this we conclude that there is nothing to indicate that errors in the calculated results on the order of 80-100 Kcal/mol are likely.

Nonetheless, since the theoretical results were based on parametric fitting of semi-empirical functions, we decided to carry out considerably more reliable 6-31G(d)//6-31G(d) calculations using the Gaussian-86(3) program on the fluoride and chloride systems. The geometries for these systems optimized at the 6-31G(d) and PM3 level are shown in Table II. With the exception of SiBr<sub>4</sub>, in which the calculated Si-Br bond is far too short, the agreement between the PM3

TABLE II  
Calculated and experimental geometries\* for silicon species

Molecule (Point Gp)	Geometric quantity	PM3	6-31G(d)	Exp <sup>a</sup>
SiX <sub>4</sub> (Td)	Si—F	1.580	1.557	1.552
	Si—Cl	2.039	2.029	2.017
	Si—Br	1.795		2.150
	Si—I	2.435		2.430
SiO <sub>2</sub> (D <sub>∞h</sub> )	Si—O	1.495	1.482	
SiOX <sub>2</sub> (C <sub>2v</sub> )	Si—F	1.570	1.556	
	Si—O	1.508	1.482	
	F—Si—O	126.1	127.6	
	Si—Cl	2.008	2.008	
	Si—O	1.497	1.487	
	Cl—Si—O	125.7	125.5	
	Si—Br	1.853		
	Si—O	1.492		
	Br—Si—O	130.3		
	Si—I	2.445		
	Si—O	1.482		
I—Si—O	126.6			

\* Distances in Angstroms, angles in degrees.

<sup>a</sup> M. W. Chase, C. A. Davies, J. R. Downey, D. R. Frurip, R. A. McDonald, and A. N. Syverud, JANAF Thermochemical Tables, Third Edition, *J. Phys. Chem. Ref. Data*, **14**, Suppl. 1 (1985).

and the experimental or 6-31G(d) geometries is quite satisfactory. The heat of reaction for the chloride system was 3.3 Kcal/mole at the SCF level; 5.1 Kcal/mole at the MP2 level, and after adding in the zero point vibrational energy, 5.6 Kcal/mole. The corresponding values for the fluoride system are +19.9, +22.8, and +23.2 kcal/mole, respectively. All these calculations refer to the system at 0 K; however, the results for the system at 298 K would be similar. The very close agreement between the semi-empirical and *ab-initio* calculations is remarkable, and lends credence to the semi-empirical results for the higher halides.

## DISCUSSION

Although Dittmer and Nieman assume an uncertainty in their reported results of 7.2 Kcal/mole, the actual error is likely to be considerably higher. Their SiO<sub>2</sub>, SiF<sub>4</sub> and SiI<sub>4</sub> results are in agreement with other published values, while the SiCl<sub>4</sub> value recommended is only 12.5 kcal more negative than the JANAF value,<sup>4</sup> and the SiBr<sub>4</sub> value is 8.5 more negative than the NBS value.<sup>5</sup> On the other hand, there appear to be no previously reported values for the heats of formation of the oxyhalides other than the JANAF value of -231.0 for SiOF<sub>2</sub>. As a result, we conclude that it is likely that the reported values for the oxyhalides are substantially too negative. Assuming the JANAF values for SiO<sub>2</sub> (-73 Kcal/mole) and SiCl<sub>4</sub>, and the *ab initio* value for the heat of reaction, then

the heat of formation of  $\text{SiOCl}_2$  is predicted to be  $-110.1$  Kcal/mole. As the heats of most metathetic reactions are in the order of a few Kcal/mole, and high-level *ab initio* calculations indicate that this is the case for one of the reactions in question, we conclude that the experimental values for the oxyhalides of silicon reported by Dittmer and Nieman are in error by approximately 50 Kcal/mole.

Experimental measurements of very unstable species are fraught with difficulty. At present there does not exist any simple method for checking the accuracy of these methods. On the other hand, semiempirical methods are now becoming very easy to use, and are of known, although at present not very high, accuracy. For computational methods, an unstable or exotic species is no more difficult to calculate than a stable or well-behaved species. For this reason, in the future, experimentalists are likely to use the available rapid computational methods to survey the systems they plan to investigate. Additional caution would then be suggested if there was a large difference between the calculated and experimental results. If the accuracy of the calculated result was questioned, then more rigorous techniques, such as high level *ab initio* methods, as used here, could be used to confirm or refute the semiempirical results.

## REFERENCES

1. J. J. P. Stewart, *J. Comp. Chem.* **10**, 221 (1989).
2. G. Dittmer and U. Niemann, *Philips J. Res.* **37**, 1 (1982).
3. Gaussian 86, M. J. Frisch, J. S. Binkley, H. B. Schelegel, K. Raghavachari, C. F. Melius, R. L. Martin, J. J. P. Stewart, F. W. Bobrowicz, C. M. Rohlfing, L. R. Kahn, D. J. Defrees, R. Seeger, R. A. Whiteside, D. J. Fox, E. M. Fleuder, and J. A. Pople, Carnegie-Mellon Quantum Chemistry Publishing Unit, Pittsburgh PA, 1984.
4. M. W. Chase, C. A. Davies, J. R. Downey, D. R. Frurip, R. A. McDonald, and A. N. Syverud, JANAF Thermochemical Tables, Third Edition, *J. Phys. Chem. Ref. Data*, **14**, Suppl. 1 (1985).
5. D. D. Wagman, W. H. Evans, V. B. Parker, R. H. Schumm, II, Halow, S. M. Bailey, K. L. Churney, and R. L. Nuttall, *J. Phys. Chem. Ref. Data*, Suppl. 11, **2**, (1982).



Accession For	
NTIS	<input checked="" type="checkbox"/>
DTIC	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By _____	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
<b>A-1 20</b>	