

1

USAARL Report No. 91-12

AD-A233 252



**Contact Lens Anterior Surface pH
(Reprint)**

By

Morris R. Lattimore

Sensory Research Division

February 1991

SEARCHED
SERIALIZED
APR 01 1991
S G D

Approved for public release; distribution unlimited.

91 3 12 091

**United States Army Aeromedical Research Laboratory
Fort Rucker, Alabama 36362-5292**

Notice

Qualified requesters

Qualified requesters may obtain copies from the Defense Technical Information Center (DTIC), Cameron Station, Alexandria, Virginia 22314. Orders will be expedited if placed through the librarian or other person designated to request documents from DTIC.

Change of address

Organizations receiving reports from the U.S. Army Aeromedical Research Laboratory on automatic mailing lists should confirm correct address when corresponding about laboratory reports.

Disposition

Destroy this report when it is no longer needed. Do not return to the originator.

Disclaimer

The views, opinions, and/or findings contained in this report are those of the authors and should not be construed as an official Department of the Army position, policy, or decision, unless so designated by other official documentation. Citation of trade names in this report does not constitute an official Department of the Army endorsement or approval of the use of such commercial items.

Human use

Human subjects participated in these studies after giving their free and informed voluntary consent. Investigators adhered to AR 70-25 and USAMRDC Reg 70-25 on Use of Volunteers in Research.

Reviewed:


THOMAS L. FREZELL
LTC, MS
Director, Sensory Research
Division


ROGER W. WILEY, O.D., Ph.D.
Chairman, Scientific
Review Committee

Released for publication:


DAVID H. KARNEY
Colonel, MC, SFS
Commanding

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE

REPORT DOCUMENTATION PAGE				Form Approved OMB No. 0704-0188	
1a. REPORT SECURITY CLASSIFICATION Unclassified		1b. RESTRICTIVE MARKINGS			
2a. SECURITY CLASSIFICATION AUTHORITY		3. DISTRIBUTION/AVAILABILITY OF REPORT Approved for public release; distribution unlimited			
2b. DECLASSIFICATION/DOWNGRADING SCHEDULE					
4. PERFORMING ORGANIZATION REPORT NUMBER(S) USAARL Report No. 91-12		5. MONITORING ORGANIZATION REPORT NUMBER(S)			
6a. NAME OF PERFORMING ORGANIZATION U.S. Army Aeromedical Research Laboratory		6b. OFFICE SYMBOL (If applicable) SGRD-UAS-VS	7a. NAME OF MONITORING ORGANIZATION U.S. Army Medical Research and Development Command		
6c. ADDRESS (City, State, and ZIP Code) P.O. Box 577 Fort Rucker, AL 36362-5292		7b. ADDRESS (City, State, and ZIP Code) Fort Detrick Frederick, MD 21702-5012			
8a. NAME OF FUNDING/SPONSORING ORGANIZATION		8b. OFFICE SYMBOL (If applicable)	9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER		
8c. ADDRESS (City, State, and ZIP Code)		10. SOURCE OF FUNDING NUMBERS			
		PROGRAM ELEMENT NO. 62787A	PROJECT NO. 3M162787A879	TASK NO. BG	WORK UNIT ACCESSION NO. 168
11. TITLE (Include Security Classification) Contact Lens Anterior Surface pH (U)					
12. PERSONAL AUTHOR(S) Lattimore, Morris R.					
13a. TYPE OF REPORT Final		13b. TIME COVERED FROM _____ TO _____	14. DATE OF REPORT (Year, Month, Day) 1991 February		15. PAGE COUNT 4
16. SUPPLEMENTARY NOTATION This report is a reprint of a publication in International Contact Lens Clinic, Vol. 17, September/October 1990, p 228-231.					
17. COSATI CODES			18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)		
FIELD	GROUP	SUB-GROUP			
06	04				
06	05		Hydrogel contact lenses, tear film pH, extended wear		
19. ABSTRACT (Continue on reverse if necessary and identify by block number) Recent reports of CO ₂ accumulation under hydrogel lenses, paired with the detection of a decrease in stromal pH following contact lens wear, have highlighted the potential for tear pH assessment as a clinical tool. The in situ anterior hydrogel lens surface pH was measured with a flat-surfaced, self-referenced pH electrode in order to indirectly evaluate fluid exchange between the precorneal tear film and hydrogel lenses. Volunteer human subjects were fitted with moderate water content (58%), disposable extended wear hydrogel lenses. Measurements were recorded from the lens in its packaged state (pH 6.99), from the lens in situ 5 minutes after initial lens application (pH 7.17), 24 hours later (pH 7.34), and at the end of 7 days continuous contact lens wear (pH 7.43). Possible cornea-tear film-hydrogel lens interactions could explain certain hydrogel lens-associated contrast sensitivity deficits and transient endothelial changes.					
20. DISTRIBUTION/AVAILABILITY OF ABSTRACT <input checked="" type="checkbox"/> UNCLASSIFIED/UNLIMITED <input type="checkbox"/> SAME AS RPT. <input type="checkbox"/> DTIC USERS			21. ABSTRACT SECURITY CLASSIFICATION Unclassified		
22a. NAME OF RESPONSIBLE INDIVIDUAL Chief, Scientific Information Center			22b. TELEPHONE (Include Area Code) (205) 255-6907	22c. OFFICE SYMBOL SGRD-UAX-SI	

Contact Lens Anterior Surface pH

Morris R. Lattimore, Jr., OD, PhD

Recent reports of CO₂ accumulation under hydrogel lenses, paired with the detection of a decrease in stromal pH following contact lens wear, have highlighted the potential for tear pH assessment as a clinical tool. The *in situ* anterior hydrogel lens surface pH was measured with a flat-surfaced, self-referenced pH electrode in order to indirectly evaluate fluid exchange between the precorneal tear film and hydrogel lenses. Volunteer human subjects were fitted with moderate water content (58%), disposable extended wear hydrogel lenses. Measurements were recorded from the lens in its packaged state (pH 6.99), from the lens *in situ* 5 minutes after initial lens application (pH 7.17), 24 hours later (pH 7.34), and at the end of 7 days continuous contact lens wear (pH 7.43). Possible cornea-tear film-hydrogel lens interactions could explain certain hydrogel lens-associated contrast sensitivity deficits and transient endothelial changes.

Keywords: Hydrogel contact lenses; tear film pH; extended wear

The views of the author do not purport to reflect the position of the Department of the Army or the Department of Defense.

Citation of trade names does not constitute official Department of the Army endorsement or approval of the use of such commercial items.

Human subjects participated in the study after giving their free and informed voluntary consent. The investigator adhered to AR 70-25 and USAMRDC Regulation 70-25 on Use of Volunteers in Research.

Address reprint requests to Dr. Lattimore at the U.S. Army Aeromedical Research Laboratory, Visual Sciences Branch, P.O. Box 577, Fort Rucker, AL 36362-5292.

Accepted for publication August 1990.

Introduction

The anterior corneal surface is associated closely with an overlying canopy of moisture known as the precorneal tear film. Traditionally, clinicians have been concerned with how certain characteristics of the tears can influence corneal integrity; tear film formation problems¹ and tear osmolarity issues² represent two examples of purported tear film influence on the cornea. However, the tear film can be susceptible to influence by the cornea, as evidenced by the presence of both glycolytic and tricarboxylic acid cycle enzymes within the tear layer. The source of these enzymes has been shown not to be the lacrimal gland, but, rather, the underlying corneal tissue.³ Therefore, tear chemistry is affected directly by the cornea. Consequently, clinicians should be reminded that although anatomically distinct the cornea and its tear film are functionally interactive.

Attempts at quantifying the normal tear pH value have yielded varying results. Although one cause of variation appears to be due to instrumentation differences, the primary cause of the variation appears to be the location or source of the tear sample. In the past, the tear film has been approached as a unitary entity independent of whether or not a sample or pH reading was obtained from the fornix, cul-de-sac, inferior meniscus, or limbus. Based on this variety of pH results, shown in *Table 1*,⁴⁻¹⁰ it can be concluded that tear pH is location-specific. Discussions stemming from this investigation are limited to the precorneal tear film.

Efforts at documenting the pH of the precorneal tear film (i.e., that canopy of mucin, aqueous, and oil directly anterior to the cornea) have resulted in a mean value range of 7.45 (Ref. 9) to 7.83 (Ref. 10). Since measurements of precorneal tear film pH under the extended open-eye condition (i.e., nonblinking state) have been shown to match

Table 1. Recent Tear pH Studies

Author(s)	Year	Location	Instrument	N (subjects)	Mean \pm error
Norn ⁴	1988	Inferior fornix	Microglass electrode	41	6.93 \pm 0.24
Coles and Jaros ⁵	1984	Lateral fornix	Direct contact microelectrode	133	7.11 \pm 1.50
Fischer and Wiederholt ⁶	1982	Limbus (1 o'clock)	Micro-pH electrode	4	7.60 \pm 0.09
		Limbus (5 o'clock)	Micro-pH electrode	4	7.50 \pm 0.08
Abelson et al. ⁷	1981	Inferior cul-de-sac	Microcombination glass pH probe	44	7.00 \pm 0.20
Andres et al. ⁸	1988	Precorneal	Micro-pH electrode	71	7.51 \pm 0.18
Carney and Hill ⁹	1976	Meniscus	Microelectrode	16	7.45 \pm 0.16
Chen and Maurice ¹⁰	1990	Precorneal	Fluorescent probe	6	7.83 \pm 0.10

that predicted by CO₂ cornea-tear film equilibration calculations,⁶ it is likely the above values are very close to the true precorneal tear film pH.

Initial research indicated hydrogel contact lenses may provide a barrier to carbon dioxide (CO₂) efflux from the cornea, although at the time this was considered to be insignificant in terms of corneal physiology.¹¹ However, recent measurements of tear CO₂ accumulation under hydrogel lenses,¹² paired with the detection of a decrease in both subcontact lens¹⁰ and stromal pH following contact lens wear,¹³ indicates yet another functional link between the precorneal tear film and corneal physiology. Indeed, Holden et al.^{12,14} have tied the issues of subhydrogel lens CO₂ accumulation and tissue pH changes to the endothelial bleb response. Since the issue of anterior segment CO₂ expiration has been associated with one aspect of the precorneal tear film (i.e., the subcontact lens tear film), it is possible other aspects of the precorneal tear film may be influenced as well. The purpose of this study was to evaluate fluid exchange interactions between hydrogel lenses and the precorneal tear film in an attempt to indirectly monitor corneal and subcontact lens pH changes resulting from hydrogel contact lens wear.

Materials and Methods

A self-referenced pH electrode (Orion Research, Model SA 230), designed for pH recording from semisolid materials, was used to assess the in situ anterior contact lens surface pH response to continuous wear of a 58% water, disposable soft contact lens. The recorded pH reading was the peak value of a transient response. Upon initial probe application, the measured pH value was within 0.2 of the final or peak value. However, a gradual drift in the alkaline direction led to stabilization of the reading, presumably due to temperature changes at the probe surface. If the probe was kept in contact with the lens beyond the stabilization period, a gradual shift in the acidic direction was noted. This has been attributed to CO₂ accumulation under the probe (Fatt, personal communication).

Subjects were on a 1-week wearing cycle, after which time the lenses were removed, disposed of, and replaced after at least one night of lens-free sleep. The pH electrode was calibrated with a 7.00 and a 10.00 pH standard solution at 35°C and disinfected by alcohol swab and surface drying between each assessment. Probe calibration was then maintained at 35°C. Measurements were recorded from the contact lens in its storage packet immediately after opening, then 5 minutes after initial lens application onto the volunteer subject's eye, 24 hours after initial lens application, and 7 days after initial lens application. Additional anterior lens surface pH recordings were made during the course of follow-up examinations after 1, 3, and 6 months of contact lens-wearing experience using the weekly wearing paradigm detailed above. Each measurement for any one individual was taken at the same time of day in order to minimize error from individual diurnal variations.⁹ However, pH assessments across individuals occurred at varying times of day, thereby eliminating any group diurnal effect.

Results

Figure 1 provides a graphical data representation. The contact lens in solution is very near a neutral pH of 7.00. Within the first 5 minutes of contact lens wear, the pH reading started to rise into the alkaline region (7.17); a further increase in pH is noted after 24 hours of wear (7.34). Stabilization of pH (7.43) is apparent at day 7 near established norms for the nonlens-wearing precorneal tear film. Subsequent pH measurements after 1, 3, and 6 months of weekly disposable contact lens wear fall between the pH values found on day 1 and day 7 (7.38). Baseline, 5-minute, and 24-hour data are statistically significant by the *t*-test ($p < 0.05$). Subsequent measurements (7 day and 1-, 3-, and 6-month follow-ups) are not statistically different from the 24-hour pH value ($p > 0.20$). However, an analysis of variance (ANOVA) over the 1-week initial period elicits a statistically significant trend for pH shift over the entire initial 7-day time period.




Figure 1. Anterior lens surface pH vs. continuous wearing time.

Discussion

The initial in situ pH reading of 7.17, taken just 5 minutes after lens application, suggests that a fluid exchange between the anterior tear film and the contact lens occurs very quickly. However, pH values obtained on subsequent follow-up evaluations (1, 3, and 6 months postfitting) documented the pH status of lenses that had been worn 2-7 days prior to those pH measurements. Since the average long-term follow-up pH value (7.38) falls between the initial week's pH values for day 1 (7.34) and day 7 (7.43), it would be reasonable to accept the concept of a long-term pattern of fluid exchange reaching equilibrium somewhere within a 7-day range of hydrogel lens wear. It should be noted here that the use of this pH electrode methodology assumes the anterior contact lens surface pH measurement accurately represents both the prelens tear film pH and the pH of the anterior water component of the hydrogel contact lens. However, it is possible these two entities could have slightly different pH values.

The final pH data for day 7 of the initial week of lens wear are not much different from the accepted published norms for the precorneal tear film.^{6,9} The initial data (days 0 and 1) are less alkaline compared to precorneal tear film norms, possibly due to the starting lens pH of 7.00; if the lenses were packaged in a storage solution of a more alkaline nature near 7.45, this pattern of pH adjustment might not be exhibited. In any event, the data do not support the use of this system as a useful indirect monitor of corneal and subcontact lens pH changes related to hydrogel lens wear. However, it may be possible to estimate CO₂ expiration rates by monitoring the anterior lens surface pH over a lengthy continuous probe application period. In future studies, the combined knowledge of CO₂ expiration rates and O₂ uptake rates might provide clinically useful information.

Accepting previous reports of pH decrease/CO₂ trapping or buildup under hydrogel lens,^{10,12,14} it is possible that a pH gradient exists within the matrix of a hydrogel lens (Figure 2). Moreover, this gradient, bordered by different




Figure 2. The graphic represents the proposed pH gradient that could be present within the in situ hydrogel lens matrix as a result of carbon dioxide accumulation. This pH gradient, in turn, would reflect the presence of underlying water content and refractive index gradients as well.

pH environments at each hydrogel lens surface, would preclude a lens from being considered as simply a unitary piece of plastic. It previously has been shown that soft lens hydration is directly influenced by the pH of its solution.¹⁵ Therefore, a lens in close approximation with a cornea, with differing pH solutions at each surface, could have a transitional water content from one surface to the other. Consequently, there would be a varying index of refraction as well. This pH gradient then would create layers of "lenses" between the physical confines of the anterior and posterior lens surfaces. This laminar arrangement of varying water content and refractive indices could be responsible for the optical issues linked to certain contrast sensitivity deficits of hydrogel lens wear.¹⁶⁻¹⁸

Lastly, is the initial, packaged lens pH significant to the physiological integrity of the cornea? It is accepted that the maintenance of corneal thickness and transparency, by way of active ion transport, is pH dependent.¹⁹ In addition, induced relative acidic pH changes at the level of the endothelium have been linked to the transient endothelial bleb response.^{12,14} Finally, a number of possible effects of an acidic shift in the cornea have previously been suggested.²⁰ Therefore, it is not unreasonable to suggest that the application of a moderate- to high-water-content hydrogel lens of a 7.0 pH or lower, weighing approximately 0.013 g, holding roughly 7.5 μl of water (if a 58% water material), would immediately create a stressful environment for the cornea. Within the context of this study (58% water content lens), application of a hydrogel lens to the anterior surface of the cornea effectively doubles the volume of fluid anterior to the cornea, since the typical precorneal tear film is 7-8 μl in volume. If the lens matrix possesses a pH that is relatively acidic compared with the precorneal tear film norm, then a significant metabolic challenge could be presented to the cornea proper. The pH-mediated transient endothelial bleb response could therefore be a reflection of this initial challenge. Upon initial lens application, the pH-induced stress would be at a peak and then begin to decline as the water component slowly equilibrates with the tear film. However, concurrent CO₂ expiration and trapping would elicit a supplementary pH-induced stress. With

the endothelial bleb response being related to pH challenge, it is reasonable to conclude that the application of a hydrogel lens, exhibiting pH characteristics relatively acidic compared with the normal precorneal tear film, is the trigger for hydrogel lens-related transient endothelial changes. If this untested hypothesis is valid, then immediate, transient endothelial changes (i.e., the bleb response) could be bypassed by packaging hydrogel lenses at a slightly alkaline pH.

In summary, simple pH measurement of the anterior lens surface does not appear to provide clinically useful information, although a pH profile monitored over an extended time period may provide information concerning CO₂ expiration. A number of questions can be raised concerning both the susceptibility of visual performance and corneal physiology to external influence by the physical state of the hydrogel material when placed on the cornea.

Acknowledgments

The investigator is deeply indebted to the Army Aviation Center and all the AH-64 "Apache" aviators who participated in this research effort; they are truly "above the best." Special thanks is extended to Ms. Carolyn Johnson, SSG Nonilon Fallaria, and SGT Vincent Reynoso for their essential administrative and technical support.

References

- Grayson M: Diseases of the cornea. St. Louis, C.V. Mosby, 1983, pp 339-343.
- Farris RL, Gilbard JP, Stuckell RN, Mandel ID: Diagnostic tests in keratoconjunctivitis sicca. *CLAO J* 1983;9:23-28.
- Fullard RJ, Carney LG: Diurnal variation in human tear enzymes. *Exp Eye Res* 1984;38:15-26.
- Norn MS: Tear fluid pH in normals, contact lens wearers, and pathological cases. *Acta Ophthalmol* 1988;66:485-489.
- Coles WH, Jaros PA: Dynamics of ocular surface pH. *Br J Ophthalmol* 1984;68:549-552.
- Fischer FH, Wiederholt M: Human precorneal tearfilm pH measured by microelectrodes. *Graefes Arch Ophthalmol* 1982;218:168-170.
- Abelson MB, Udell IJ, Weston JH: Normal human tear pH by direct measurement. *Arch Ophthalmol* 1981;99:301-304.
- Andres S, Garcia ML, Espina MM, Valero J, Valls O: Tear pH, air pollution, and contact lenses. *Am J Optom Physiol Opt* 1988;65:627-631.
- Carney LG, Hill RM: Human tear pH. *Arch Ophthalmol* 1976;94:821-824.
- Chen FS, Maurice DM: The pH in the precorneal tearfilm and under a contact lens measured with a fluorescent probe. *Exp Eye Res* 1990;50:251-259.
- Fatt I, Bieber MT, Pye SD: Steady state distribution of oxygen and carbon dioxide in the in vivo cornea of an eye covered by a gas-permeable contact lens. *Am J Optom Arch Am Acad Optom* 1969;64:3-14.
- Holden BA, Ross R, Jenkins J: Hydrogel contact lenses impede carbon dioxide efflux from the human cornea. *Curr Eye Res* 1987;6:1283-1290.
- Bonano JA, Polse KA: Effect of contact lens wear on stromal pH. *Am J Optom Physiol Opt* 1986;63:58-62.
- Holden BA, Williams L, Zantos SG: The etiology of transient endothelial changes in the human cornea. *Invest Ophthalmol Vis Sci* 1985;26:1354-1359.
- McCarey BE, Wilson LA: pH, osmolarity, and temperature effects on the water content of hydrogel contact lenses. *Contact Intraoc Lens Med J* 1982;8:158-167.
- Woo GCS, Hess RF: Contrast sensitivity function and soft contact lenses. *ICLC* 1979;6:171-176.
- Bernstein IH, Brodrick J: Contrast sensitivity through spectacles and soft contact lenses. *Am J Optom Physiol Opt* 1981;58:309-313.
- Grey CP: Changes in contrast sensitivity during the first hour of soft lens wear. *Am J Optom Physiol Opt* 1986;63:702-707.
- Fischer FH, Weiderholt M: The pH dependency of sodium and chloride transport in the isolated human cornea. *Invest Ophthalmol Vis Sci* 1978;17:810-813.
- Efron N: The greenhouse effect and contact lenses, in *Transactions of the British Contact Lens Association Conference*, 1989, p. 7.

Morris R. Lattimore is a graduate of the University of Wisconsin (BA, 1972), Illinois College of Optometry (OD, 1976), and the University of Houston (PhD, 1987). His 14 years of active military service include 8 years of clinical work (Ft. Leonard Wood, MO; Camp Casey, Korea; and Ft. Ord, CA), plus 6 years of postgraduate education and research experience. Although Dr. Lattimore's current work concerning the feasibility of contact lens wear by Army aviators is primarily clinically oriented, his personal research interest centers on the determination of underlying mechanisms responsible for exhibited clinical conditions.



Initial distribution

**Commander, U.S. Army Natick Research,
Development and Evaluation Center
ATTN: STRNC-MIL (Documents
Librarian)
Natick, MA 01760-5040**

**Naval Submarine Medical
Research Laboratory
Medical Library, Naval Sub Base
Box 900
Groton, CT 06340**

**Commander/Director
U.S. Army Combat Surveillance
and Target Acquisition Lab
ATTN: DELCS-D
Fort Monmouth, NJ 07703-5304**

**Commander
10th Medical Laboratory
ATTN: Audiologist
APO New York 09180**

**Naval Air Development Center
Technical Information Division
Technical Support Detachment
Warminster, PA 18974**

**Commanding Officer, Naval Medical
Research and Development Command
National Naval Medical Center
Bethesda, MD 20814-5044**

**Deputy Director, Defense Research
and Engineering
ATTN: Military Assistant
for Medical and Life Sciences
Washington, DC 20301-3080**

**Commander, U.S. Army Research
Institute of Environmental Medicine
Natick, MA 01760**

**U.S. Army Avionics Research
and Development Activity
ATTN: SAVAA-P-TP
Fort Monmouth, NJ 07703-5401**

**U.S. Army Communications-Electronics
Command
ATTN: AMSEL-RD-ESA-D
Fort Monmouth, NJ 07703**

**Library
Naval Submarine Medical Research Lab
Box 900, Naval Sub Base
Groton, CT 06349-5900**

**Commander
Man-Machine Integration System
Code 602
Naval Air Development Center
Warminster, PA 18974**

**Commander
Naval Air Development Center
ATTN: Code 602-B (Mr. Brindle)
Warminster, PA 18974**

**Commanding Officer
Harry G. Armstrong Aerospace
Medical Research Laboratory
Wright-Patterson
Air Force Base, OH 45433**

**Director
Army Audiology and Speech Center
Walter Reed Army Medical Center
Washington, DC 20307-5001**

**Commander, U.S. Army Institute
of Dental Research
ATTN: Jean A. Setterstrom, Ph. D.
Walter Reed Army Medical Center
Washington, DC 20307-5300**

Naval Air Systems Command
Technical Air Library 950D
Room 278, Jefferson Plaza II
Department of the Navy
Washington, DC 20361

Naval Research Laboratory Library
Shock and Vibration
Information Center, Code 5804
Washington, DC 20375

Director, U.S. Army Human
Engineering Laboratory
ATTN: Technical Library
Aberdeen Proving Ground, MD 21005

Commander, U.S. Army Test
and Evaluation Command
ATTN: AMSTE-AD-H
Aberdeen Proving Ground, MD 21005

Director
U.S. Army Ballistic
Research Laboratory
ATTN: DRXBR-OD-ST Tech Reports
Aberdeen Proving Ground, MD 21005

Commander
U.S. Army Medical Research
Institute of Chemical Defense
ATTN: SGRD-UV-AO
Aberdeen Proving Ground,
MD 21010-5425

Commander, U.S. Army Medical
Research and Development Command
ATTN: SGRD-RMS (Ms. Madigan)
Fort Detrick, Frederick, MD 21702-5012

Director
Walter Reed Army Institute of Research
Washington, DC 20307-5100

HQ DA (DASG-PSP-O)
5109 Leesburg Pike
Falls Church, VA 22041-3258

Naval Research Laboratory
Library Code 1433
Washington, DC 20375

Harry Diamond Laboratories
ATTN: Technical Information Branch
2800 Powder Mill Road
Adelphi, MD 20783-1197

U.S. Army Materiel Systems
Analysis Agency
ATTN: AMXSY-PA (Reports Processing)
Aberdeen Proving Ground
MD 21005-5071

U.S. Army Ordnance Center
and School Library
Simpson Hall, Building 3071
Aberdeen Proving Ground, MD 21005

U.S. Army Environmental
Hygiene Agency
Building E2100
Aberdeen Proving Ground, MD 21010

Technical Library Chemical Research
and Development Center
Aberdeen Proving Ground, MD
21010--5423

Commander
U.S. Army Medical Research
Institute of Infectious Disease
SGRD-UIZ-C
Fort Detrick, Frederick, MD 21702

Director, Biological
Sciences Division
Office of Naval Research
600 North Quincy Street
Arlington, VA 22217

Commander
U.S. Army Materiel Command
ATTN: AMCDE-XS
5001 Eisenhower Avenue
Alexandria, VA 22333

Commandant
U.S. Army Aviation
Logistics School ATTN: ATSQ-TDN
Fort Eustis, VA 23604

Headquarters (ATMD)
U.S. Army Training
and Doctrine Command
Fort Monroe, VA 23651

Structures Laboratory Library
USARTL-AVSCOM
NASA Langley Research Center
Mail Stop 266
Hampton, VA 23665

Naval Aerospace Medical
Institute Library
Building 1953, Code 03L
Pensacola, FL 32508-5600

Command Surgeon
HQ USCENTCOM (CCSG)
U.S. Central Command
MacDill Air Force Base FL 33608

Air University Library
(AUL/LSE)
Maxwell Air Force Base, AL 36112

U.S. Air Force Institute
of Technology (AFIT/LDEE)
Building 640, Area B
Wright-Patterson
Air Force Base, OH 45433

Henry L. Taylor
Director, Institute of Aviation
University of Illinois-Willard Airport
Savoy, IL 61874

Chief, Nation Guard Bureau
ATTN: NGB-AR (COL Urbauer)
Room 410, Park Center 4
4501 Ford Avenue
Alexandria, VA 22302-1451

Commander
U.S. Army Aviation Systems Command
ATTN: SGRD-UAX-AL (MAJ Gillette)
4300 Goodfellow Blvd., Building 105
St. Louis, MO 63120

U.S. Army Aviation Systems Command
Library and Information Center Branch
ATTN: AMSAV-DIL
4300 Goodfellow Boulevard
St. Louis, MO 63120

Federal Aviation Administration
Civil Aeromedical Institute
Library AAM-400A
P.O. Box 25082
Oklahoma City, OK 73125

Commander
U.S. Army Academy
of Health Sciences
ATTN: Library
Fort Sam Houston, TX 78234

Commander
U.S. Army Institute of Surgical Research
ATTN: SGRD-USM (Jan Duke)
Fort Sam Houston, TX 78234-6200

AAMRL/HEX
Wright-Patterson
Air Force Base, OH 45433

University of Michigan
NASA Center of Excellence in Man-
Systems Research
ATTN: R. G. Snyder, Director
Ann Arbor, MI 48109

John A. Dellinger,
Southwest Research Institute
P. O. Box 28510
San Antonio, TX 78284

Product Manager
Aviation Life Support Equipment
ATTN: AMCPM-ALSE
4300 Goodfellow Boulevard
St. Louis, MO 63120-1798

Commander
U.S. Army Aviation
Systems Command
ATTN: AMSAV-ED
4300 Goodfellow Boulevard
St. Louis, MO 63120

Commanding Officer
Naval Biodynamics Laboratory
P.O. Box 24907
New Orleans, LA 70189-0407

Assistant Commandant
U.S. Army Field Artillery School
ATTN: Morris Swott Technical Library
Fort Sill, OK 73503-0312

Commander
U.S. Army Health Services Command
ATTN: HSOP-SO
Fort Sam Houston, TX 78234-6000

Director of Professional Services
HQ USAF/SGDT
Bolling Air Force Base, DC 20332-6188

U.S. Army Dugway Proving Ground
Technical Library, Building 5330
Dugway, UT 84022

U.S. Army Yuma Proving Ground
Technical Library
Yuma, AZ 85364

AFFTC Technical Library
6510 TW/TSTL
Edwards Air Force Base,
CA 93523-5000

Commander
Code 3431
Naval Weapons Center
China Lake, CA 93555

Aeromechanics Laboratory
U.S. Army Research and Technical Labs
Ames Research Center, M/S 215-1
Moffett Field, CA 94035

Sixth U.S. Army
ATTN: SMA
Presidio of San Francisco, CA 94129

Commander
U.S. Army Aeromedical Center
Fort Rucker, AL 36362

U.S. Air Force School
of Aerospace Medicine
Strughold Aeromedical Library Technical
Reports Section (TSKD)
Brooks Air Force Base, TX 78235-5301

Dr. Diane Damos
Department of Human Factors
ISSM, USC
Los Angeles, CA 90089-0021

U.S. Army White Sands
Missile Range
ATTN: STEWS-IM-ST
White Sands Missile Range, NM 88002

U.S. Army Aviation Engineering
Flight Activity
ATTN: SAVTE-M (Tech Lib) Stop 217
Edwards Air Force Base, CA 93523-5000

Ms. Sandra G. Hart
Ames Research Center
MS 262-3
Moffett Field, CA 94035

Commander, Letterman Army Institute
of Research
ATTN: Medical Research Library
Presidio of San Francisco, CA 94129

Mr. Frank J. Stagnaro, ME
Rush Franklin Publishing
300 Orchard City Drive
Campbell, CA 95008

Commander
U.S. Army Medical Materiel
Development Activity
Fort Detrick, Frederick, MD 21702-5009

Commander
U.S. Army Aviation Center
Directorate of Combat Developments
Building 507
Fort Rucker, AL 36362

U. S. Army Research Institute
Aviation R&D Activity
ATTN: PERI-IR
Fort Rucker, AL 36362

Commander
U.S. Army Safety Center
Fort Rucker, AL 36362

U.S. Army Aircraft Development
Test Activity
ATTN: STEBG-MP-P
Cairns Army Air Field
Fort Rucker, AL 36362

Commander U.S. Army Medical Research
and Development Command
ATTN: SGRD-PLC (COL Sedge)
Fort Detrick, Frederick, MD 21702

MAJ John Wilson
TRADOC Aviation LO
Embassy of the United States
APO New York 09777

Netherlands Army Liaison Office
Building 602
Fort Rucker, AL 36362

British Army Liaison Office
Building 602
Fort Rucker, AL 36362

Italian Army Liaison Office
Building 602
Fort Rucker, AL 36362

Directorate of Training Development
Building 502
Fort Rucker, AL 36362

Chief
USAHEL/USAAVNC Field Office
P. O. Box 716
Fort Rucker, AL 36362-5349

Commander U.S. Army Aviation Center
and Fort Rucker
ATTN: ATZQ-CG
Fort Rucker, AL 36362

Commander/President
TEXCOM Aviation Board
Cairns Army Air Field
Fort Rucker, AL 36362

Dr. William E. McLean
Human Engineering Laboratory
ATTN: SLCHE-BR
Aberdeen Proving Ground,
MD 21005-5001

Canadian Army Liaison Office
Building 602
Fort Rucker, AL 36362

German Army Liaison Office
Building 602
Fort Rucker, AL 36362

