

1

USAARL Report No. 91-18

AD-A239 992



The Hazard of Exposure to Impulse Noise as a Function of Frequency

Volume I

By



James H. Patterson, Jr.
Melvin Carrier, Jr.
Kevin Bordwell
Ilia M. Lomba Gautier

Sensory Research Division

and

Roger P. Hamernik
William A. Ahroon
George Turrentine
C. E. Hargett, Jr.

Auditory Research Laboratory
State University of New York
at Plattsburgh



June 1991

91-08939



Approved for public release; distribution unlimited.

United States Army Aeromedical Research Laboratory
Fort Rucker, Alabama 36362-5292

Notice

Qualified requesters

Qualified requesters may obtain copies from the Defense Technical Information Center (DTIC), Cameron Station, Alexandria, Virginia 22314. Orders will be expedited if placed through the librarian or other person designated to request documents from DTIC.

Change of address

Organizations receiving reports from the U.S. Army aeromedical Research Laboratory on automatic mailing lists should confirm correct address when corresponding about laboratory reports.

Disposition

Destroy this report when it is no longer needed. Do not return it to the originator.

Disclaimer

The views, opinions, and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy, or decision, unless so designated by other official documentation. Citation of trade names in this report does not constitute an official Department of the Army endorsement or approval of the use of such commercial items.

Animal use

In conducting the research described in this report, the investigators adhered to the Guide for care and use of laboratory animals, as promulgated by the Committee on Care and Use of Laboratory Animals of the Institute of Laboratory Resources Commission on Life Sciences, National Academy of Sciences-National Research Council.

Reviewed:



THOMAS L. FREZELL
LTC, MS
Director, Sensory Research Division

Released for publication:



ROGER W. WILBY
Chairman, Scientific
Review Committee



DAVID H. KARNEY
Colonel, MC, SFS
Commanding

REPORT DOCUMENTATION PAGE				Form Approved OMB No. 0704-0188		
1a. REPORT SECURITY CLASSIFICATION UNCLASSIFIED			1b. RESTRICTIVE MARKINGS			
2a. SECURITY CLASSIFICATION AUTHORITY			3. DISTRIBUTION/AVAILABILITY OF REPORT Approved for public release; distribution unlimited			
2b. DECLASSIFICATION/DOWNGRADING SCHEDULE						
4. PERFORMING ORGANIZATION REPORT NUMBER(S) USAARL Report No. 91-18, Volume I			5. MONITORING ORGANIZATION REPORT NUMBER(S) U.S. Army Medical Research and Development Command			
6a. NAME OF PERFORMING ORGANIZATION Sensory Research Division U.S. Army Aeromedical Rsch Lab		6b. OFFICE SYMBOL (If applicable) SGRD-UAS-AS	7a. NAME OF MONITORING ORGANIZATION			
6c. ADDRESS (City, State, and ZIP Code) P.O. Box 577 Fort Rucker, AL 36362-5292			7b. ADDRESS (City, State, and ZIP Code) Fort Detrick Frederick, MD 21701-5012			
8a. NAME OF FUNDING/SPONSORING ORGANIZATION		8b. OFFICE SYMBOL (If applicable)	9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER			
8c. ADDRESS (City, State, and ZIP Code)			10. SOURCE OF FUNDING NUMBERS			
			PROGRAM ELEMENT NO. 62777A 0601102A	PROJECT NO. 3E162777A87B 3M161102BS15	TASK NO. 	WORK UNIT ACCESSION NO. 157 282
11. TITLE (Include Security Classification) The hazard of exposure to impulse noise as a function of frequency						
12. PERSONAL AUTHOR(S) James H. Patterson, Jr., Melvin Carrier, Jr., Kevin Bordwell, Ilia M. Lomba Gautier, Roger P. Hamernik, William A. Ahroon, George Turrentine, and C. E. Hargett, Jr.						
13a. TYPE OF REPORT Final		13b. TIME COVERED FROM _____ TO _____		14. DATE OF REPORT (Year, Month, Day) 1991 June	15. PAGE COUNT 69	
16. SUPPLEMENTARY NOTATION						
17. COSATI CODES			18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)			
FIELD	GROUP	SUB-GROUP	Impulse noise, hearing, chinchilla, audiometry, and histology			
20	01					
19. ABSTRACT (Continue on reverse if necessary and identify by block number) The energy spectrum of a noise is known to be an important variable in determining the effects of a traumatic exposure. However, existing criteria for exposure to impulse noise do not consider the frequency spectrum of an impulse as a variable in the evaluation of the hazards to the auditory system. This report presents the results of a study that was designed to determine the relative potential that impulsive energy concentrated at different frequencies has in causing auditory system trauma. One hundred and eighteen (118) chinchilla, divided into 20 groups with 5 to 7 animals per group, were used in these experiments. Pre- and post-exposure hearing thresholds were measured at 10 test frequencies between 0.125 and 8 kHz on each animal using avoidance conditioning procedures. Quantitative histology (cochleograms) was used to determine the extent and pattern of the sensory cell damage. The noise exposure stimuli consisted of six different computer-generated narrow band tone bursts having center frequencies located at 0.260, 0.775, 1.025, 1.350, 2.450, and 3.550 kHz. Each narrow band exposure stimulus was presented at two to four different intensities. An analysis of the audiometric and histological data allowed a frequency weighting function to be derived.						
20. DISTRIBUTION/AVAILABILITY OF ABSTRACT <input checked="" type="checkbox"/> UNCLASSIFIED UNLIMITED <input type="checkbox"/> SAME AS RPT <input type="checkbox"/> DTIC USERS			21. ABSTRACT SECURITY CLASSIFICATION			
22a. NAME OF RESPONSIBLE INDIVIDUAL Chief, Scientific Information Center			22b. TELEPHONE (Include Area Code) (205) 255-6907	22c. OFFICE SYMBOL SGRD-UAX-SI		

9. (Cont'd)

The weighting function clearly demonstrates that equivalent amounts of impulsive energy concentrated at different frequencies is not equally hazardous to auditory function. Comparison of this weighting function with the A-weighting function showed that A-weighting overestimates the hazard from low frequency impulses. Volume 1 of the report describes the study and discusses the results. Volume 2 contains individual threshold shifts and sensory cell loss data. It is available upon request from the SIC USAARL.

Table of contents

Introduction. 11

Methods and procedures. 12

Results and discussion. 14

Histological results. 20

Conclusions 21

References. 67

Appendix A. 69

List of tables

1. Definition of experimental groups. 13

2. Summary of mean preexposure thresholds (dB)
for all animals (N = 118) compared
to published norms. 14

3. Preexposure threshold means (dB) and standard
deviations for all groups. 15

List of figures

1. The narrow band impulse pressure-time waveform
(inset) and the relative frequency spectrum
of the impulse for the 0.260 kHz CF, 146 dB
peak SPL stimulus. 16

2. The narrow band impulse pressure-time waveform
(inset) and the relative frequency spectrum
of the impulse for the 0.775 kHz CF,
144 dB peak SPL stimulus. 16

3. The narrow band impulse pressure-time waveform
(inset) and the relative frequency spectrum
of the impulse for the 1.025 kHz CF,
139 dB peak SPL stimulus. 17

4. The narrow band impulse pressure-time waveform
(inset) and the relative frequency spectrum
of the impulse for the 1.350 kHz CF,
139 dB peak SPL stimulus. 17

5.	The narrow band impulse pressure-time waveform wave (inset) and the relative frequency spectrum of the impulse for the 2.450 kHz CF, 144 dB peak SPL stimulus.	18
6.	The narrow band impulse pressure-time waveform (inset) and the relative frequency spectrum of the impulse for the 3.550 kHz CF, 139 dB peak SPL stimulus.	18
7.	The mean preexposure audiogram for all 118 chinchillas (symbols) and for published norms (dashed line). The error bars represent one standard deviation above and below the plotted point.	19
8.	The mean threshold recovery curves for the groups exposed to the 0.260 kHz CF impulse at the indicated peak SPL for the 0.125 kHz audiometric test frequency.	23
9.	The mean threshold recovery curves for the groups exposed to the 0.775 kHz CF impulse at the indicated peak SPL for the 0.125 kHz audiometric test frequency.	23
10.	The mean threshold recovery curves for the groups exposed to the 1.025 kHz CF impulse at the indicated peak SPL for the 0.125 kHz audiometric test frequency.	24
11.	The mean threshold recovery curves for the groups exposed to the 1.350 kHz CF impulse at the indicated peak SPL for the 0.125 kHz audiometric test frequency.	24
12.	The mean threshold recovery curves for the groups exposed to the 2.450 kHz CF impulse at the indicated peak SPL for the 0.125 kHz audiometric test frequency.	25
13.	The mean threshold recovery curves for the groups exposed to the 3.550 kHz CF impulse at the indicated peak SPL for the 0.125 kHz audiometric test frequency.	25
14.	The mean threshold recovery curves for the groups exposed to the 0.260 kHz CF impulse at the indicated peak SPL for the 0.250 kHz audiometric test frequency.	26

15.	The mean threshold recovery curves for the groups exposed to the 0.775 kHz CF impulse at the indicated peak SPL for the 0.250 kHz audiometric test frequency	26
16.	The mean threshold recovery curves for the groups exposed to the 1.025 kHz CF impulse at the indicated peak SPL for the 0.250 kHz audiometric test frequency.	27
17.	The mean threshold recovery curves for the groups exposed to the 1.350 kHz CF impulse at the indicated peak SPL for the 0.250 kHz audiometric test frequency.	27
18.	The mean threshold recovery curves for the groups exposed to the 2.450 kHz CF impulse at the indicated peak SPL for the 0.250 kHz audiometric test frequency.	28
19.	The mean threshold recovery curves for the groups exposed to the 3.550 kHz CF impulse at the indicated peak SPL for the 0.250 kHz audiometric test frequency.	28
20.	The mean threshold recovery curves for the groups exposed to the 0.260 kHz CF impulse at the indicated peak SPL for the 0.500 kHz audiometric test frequency.	29
21.	The mean threshold recovery curves for the groups exposed to the 0.775 kHz CF impulse at the indicated peak SPL for the 0.500 kHz audiometric test frequency.	29
22.	The mean threshold recovery curves for the groups exposed to the 1.025 kHz CF impulse at the indicated peak SPL for the 0.500 kHz audiometric test frequency.	30
23.	The mean threshold recovery curves for the groups exposed to the 1.350 kHz CF impulse at the indicated peak SPL for the 0.500 kHz audiometric test frequency.	30
24.	The mean threshold recovery curves for the groups exposed to the 2.450 kHz CF impulse at the indicated peak SPL for the 0.500 kHz audiometric test frequency.	31

25.	The mean threshold recovery curves for the groups exposed to the 3.550 kHz CF impulse at the indicated peak SPL for the 0.500 kHz audiometric test frequency.	31
26.	The mean threshold recovery curves for the groups exposed to the 0.260 kHz CF impulse at the indicated peak SPL for the 1.000 kHz audiometric test frequency.	32
27.	The mean threshold recovery curves for the groups exposed to the 0.775 kHz CF impulse at the indicated peak SPL for the 1.000 kHz audiometric test frequency.	32
28.	The mean threshold recovery curves for the groups exposed to the 1.025 kHz CF impulse at the indicated peak SPL for the 1.000 kHz audiometric test frequency.	33
29.	The mean threshold recovery curves for the groups exposed to the 1.350 kHz CF impulse at the indicated peak SPL for the 1.000 kHz audiometric test frequency.	33
30.	The mean threshold recovery curves for the groups exposed to the 2.450 kHz CF impulse at the indicated peak SPL for the 1.000 kHz audiometric test frequency.	34
31.	The mean threshold recovery curves for the groups exposed to the 3.550 kHz CF impulse at the indicated peak SPL for the 1.000 kHz audiometric test frequency.	34
32.	The mean threshold recovery curves for the groups exposed to the 0.260 kHz CF impulse at the indicated peak SPL for the 1.400 kHz audiometric test frequency.	35
33.	The mean threshold recovery curves for the groups exposed to the 0.775 kHz CF impulse at the indicated peak SPL for the 1.400 kHz audiometric test frequency.	35
34.	The mean threshold recovery curves for the groups exposed to the 1.025 kHz CF impulse at the indicated peak SPL for the 1.400 kHz audiometric test frequency.	36

35.	The mean threshold recovery curves for the groups exposed to the 1.350 kHz CF impulse at the indicated peak SPL for the 1.400 kHz audiometric test frequency.	36
36.	The mean threshold recovery curves for the groups exposed to the 2.450 kHz CF impulse at the indicated peak SPL for the 1.400 kHz audiometric test frequency.	37
37.	The mean threshold recovery curves for the groups exposed to the 3.550 kHz CF impulse at the indicated peak SPL for the 1.400 kHz audiometric test frequency.	37
38.	The mean threshold recovery curves for the groups exposed to the 0.260 kHz CF impulse at the indicated peak SPL for the 2.000 kHz audiometric test frequency.	38
39.	The mean threshold recovery curves for the groups exposed to the 0.775 kHz CF impulse at the indicated peak SPL for the 2.000 kHz audiometric test frequency.	38
40.	The mean threshold recovery curves for the groups exposed to the 1.025 kHz CF impulse at the indicated peak SPL for the 2.000 kHz audiometric test frequency.	39
41.	The mean threshold recovery curves for the groups exposed to the 1.350 kHz CF impulse at the indicated peak SPL for the 2.000 kHz audiometric test frequency.	39
42.	The mean threshold recovery curves for the groups exposed to the 2.450 kHz CF impulse at the indicated peak SPL for the 2.000 kHz audiometric test frequency.	40
43.	The mean threshold recovery curves for the groups exposed to the 3.550 kHz CF impulse at the indicated peak SPL for the 2.000 kHz audiometric test frequency.	40
44.	The mean threshold recovery curves for the groups exposed to the 0.260 kHz CF impulse at the indicated peak SPL for the 2.800 kHz audiometric test frequency.	41

45.	The mean threshold recovery curves for the groups exposed to the 0.775 kHz CF impulse at the indicated peak SPL for the 2.800 kHz audiometric test frequency.	41
46.	The mean threshold recovery curves for the groups exposed to the 1.025 kHz CF impulse at the indicated peak SPL for the 2.800 kHz audiometric test frequency.	42
47.	The mean threshold recovery curves for the groups exposed to the 1.350 kHz CF impulse at the indicated peak SPL for the 2.800 kHz audiometric test frequency.	42
48.	The mean threshold recovery curves for the groups exposed to the 2.450 kHz CF impulse at the indicated peak SPL for the 2.800 kHz audiometric test frequency.	43
49.	The mean threshold recovery curves for the groups exposed to the 3.550 kHz CF impulse at the indicated peak SPL for the 2.800 kHz audiometric test frequency.	43
50.	The mean threshold recovery curves for the groups exposed to the 0.260 kHz CF impulse at the indicated peak SPL for the 4.000 kHz audiometric test frequency.	44
51.	The mean threshold recovery curves for the groups exposed to the 0.775 kHz CF impulse at the indicated peak SPL for the 4.000 kHz audiometric test frequency.	44
52.	The mean threshold recovery curves for the groups exposed to the 1.025 kHz CF impulse at the indicated peak SPL for the 4.000 kHz audiometric test frequency.	45
53.	The mean threshold recovery curves for the groups exposed to the 1.775 kHz CF impulse at the indicated peak SPL for the 4.000 kHz audiometric test frequency.	45
54.	The mean threshold recovery curves for the groups exposed to the 2.450 kHz CF impulse at the indicated peak SPL for the 4.000 kHz audiometric test frequency.	46

55.	The mean threshold recovery curves for the groups exposed to the 3.550 kHz CF impulse at the indicated peak SPL for the 4.000 kHz audiometric test frequency.	46
56.	The mean threshold recovery curves for the groups exposed to the 0.260 kHz CF impulse at the indicated peak SPL for the 5.700 kHz audiometric test frequency.	47
57.	The mean threshold recovery curves for the groups exposed to the 0.775 kHz CF impulse at the indicated peak SPL for the 5.700 kHz audiometric test frequency.	47
58.	The mean threshold recovery curves for the groups exposed to the 1.025 kHz CF impulse at the indicated peak SPL for the 5.700 kHz audiometric test frequency.	48
59.	The mean threshold recovery curves for the groups exposed to the 1.350 kHz CF impulse at the indicated peak SPL for the 5.700 kHz audiometric test frequency.	48
60.	The mean threshold recovery curves for the groups exposed to the 2.450 kHz CF impulse at the indicated peak SPL for the 5.700 kHz audiometric test frequency.	49
61.	The mean threshold recovery curves for the groups exposed to the 3.550 kHz CF impulse at the indicated peak SPL for the 5.700 kHz audiometric test frequency.	49
62.	The mean threshold recovery curves for the groups exposed to the 0.260 kHz CF impulse at the indicated peak SPL for the 8.000 kHz audiometric test frequency.	50
63.	The mean threshold recovery curves for the groups exposed to the 0.775 kHz CF impulse at the indicated peak SPL for the 8.000 kHz audiometric test frequency.	50
64.	The mean threshold recovery curves for the groups exposed to the 1.025 kHz CF impulse at the indicated peak SPL for the 8.000 kHz audiometric test frequency.	51

65.	The mean threshold recovery curves for the groups exposed to the 1.350 kHz CF impulse at the indicated peak SPL for the 8 000 kHz audiometric test frequency.	51
66.	The mean threshold recovery curves for the groups exposed to the 2.450 kHz CF impulse at the indicated peak SPL for the 8.000 kHz audiometric test frequency.	52
67.	The mean threshold recovery curves for the groups exposed to the 3.550 kHz CF impulse at the indicated peak SPL for the 8.000 kHz audiometric test frequency.	52
68.	The group mean maximum threshold shift for each audiometric test frequency following exposure to the 0.260 kHz CF impulse at the indicated peak SPL.	53
69.	The group mean maximum threshold shift for each audiometric test frequency following exposure to the 0.775 kHz CF impulse at the indicated peak SPL.	53
70.	The group mean maximum threshold shift for each audiometric test frequency following exposure to the 1.025 kHz CF impulse at the indicated peak SPL.	54
71.	The group mean maximum threshold shift for each audiometric test frequency following exposure to the 1.350 kHz CF impulse at the indicated peak SPL.	54
72.	The group mean maximum threshold shift for each audiometric test frequency following exposure to the 2.450 kHz CF impulse at the indicated peak SPL.	55
73.	The group mean maximum threshold shift for each audiometric test frequency following exposure to the 3.550 kHz CF impulse at the indicated peak SPL.	55
74.	The group mean permanent threshold shift (PTS) for each audiometric test frequency following exposure to the 0.260 kHz CF impulse at the indicated peak SPL.	56

75.	The group mean permanent threshold shift (PTS) for each audiometric test frequency following exposure to the 0.775 kHz CF impulse at the indicated peak SPL.	56
76.	The group mean permanent threshold shift (PTS) for each audiometric test frequency following exposure to the 1.025 kHz CF impulse at the indicated peak SPL.	57
77.	The group mean permanent threshold shift (PTS) for each audiometric test frequency following exposure to the 1.350 kHz CF impulse at the indicated peak SPL.	57
78.	The group mean permanent threshold shift (PTS) for each audiometric test frequency following exposure to the 2.450 kHz CF impulse at the indicated peak SPL.	58
79.	The group mean permanent threshold shift (PTS) for each audiometric test frequency following exposure to the 3.550 kHz CF impulse at the indicated peak SPL.	58
80.	The group mean outer hair cell (OHC) loss and inner hair cell (IHC) loss within octave band lengths of the basilar membrane at the indicated frequencies following exposure to the 0.260 kHz CF impulse at the indicated peak SPL.	59
81.	The group mean outer hair cell (OHC) loss and inner hair cell (IHC) loss within octave band lengths of the basilar membrane at the indicated frequencies following exposure to the 0.775 kHz CF impulse at the indicated peak SPL.	59
82.	The group mean outer hair cell (OHC) loss and inner hair cell (IHC) loss within octave band lengths of the basilar membrane at the indicated frequencies following exposure to the 1.025 kHz CF impulse at the indicated peak SPL.	60
83.	The group mean outer hair cell (OHC) loss and inner hair cell (IHC) loss within octave band lengths of the basilar membrane at the indicated frequencies following exposure to the 1.350 kHz CF impulse at the indicated peak SPL.	60

- 84. The group mean outer hair cell (OHC) loss and inner hair cell (IHC) loss within octave band lengths of the basilar membrane at the indicated frequencies following exposure to the 2.450 kHz CF impulse at the indicated peak SPL. 61
- 85. The group mean outer hair cell (OHC) loss and inner hair cell (IHC) loss within octave band lengths of the basilar membrane at the indicated frequencies following exposure to the 3.550 kHz CF impulse at the indicated peak SPL. 61
- 86. The group mean PTS evaluated at 1, 2 and 4 kHz ((PTS)_{1,2,4}) as a function of the total sound exposure level for the twenty groups of animals exposed to the narrow band impulses. 62
- 87. The ((PTS)_{1,2,4}) as a function of the P-weighted total sound exposure level. The regression line yields a correlation coefficient of 0.904. 63
- 88. The empirical P-weighting function (symbols) derived from the narrow band exposure data along with the conventional A-weighting function (solid line). . . . 64
- 89. The group mean total outer (upper) and inner (lower) hair cell loss as a function of the total sound exposure level for the twenty groups of animals exposed to the narrow band impulses. 65
- 90. The group mean total outer (upper) and inner (lower) hair cell loss as a function of the P-weighted total sound exposure level. The regression lines yield correlation coefficients of 0.858 and 0.849 for outer and inner hair cell loss respectively. . . 66

Introduction

It has long been recognized that exposures to steady noise at different frequencies were not equally hazardous. The National Academy of Sciences Committee on Hearing, Bioacoustics and Biomechanics (CHABA) damage risk criteria for steady noise provided weighting functions which represented the relative hazard associated with noise exposures of different spectral content (Kryter et al., 1966). More recently, these weighting curves have been simplified to the A-weighting function which has been incorporated into many sound level meters. To a first approximation, A-weighting of steady noise provides a good estimate of the relative hazard to hearing of the various frequencies which comprise a steady noise exposure (Mills et al., 1983).

In contrast, the early damage risk criteria for impulse noise [e.g., Coles et al., (1968); OSHA Department of Labor, (1974); Smoorenburg, (1982); Pfander et al., (1980)] used no spectral weighting functions. For example, the Coles et al., (1968) suggestions which were incorporated into the CHABA (1968) document were formulated in terms of peak pressure and two measures of duration. The two measures of duration, the A-duration and the B-duration, resolve in practice to a limit based on only B-duration for most real world impulses. Such a criterion totally ignores any differences in the distribution of energy across frequencies. On purely theoretical grounds, this neglect of the spectrum does not seem reasonable. The auditory system transfers a pressure wave in air to fluid-borne stimulation of the auditory receptors with an efficiency which varies as a function of frequency.

Data have been reported by Price et al. (1988) showing that impulses from large bore versus small bore weapons produce a threshold of injury in cats at different peak pressures. Large bore weapons have most of their energy at low frequencies, while small bore weapons have their spectral peaks at a higher frequencies. Those results have been interpreted as indicating that a frequency selective mechanism has a strong influence over the injury resulting from exposure to impulse noise. In an earlier study, Price (1979) used tone pips to show as the energy in an impulse shifted to very high frequencies, with pressure held constant, the injuries resulting from exposure were less severe when the higher frequency tone pips were used. However, this study did not explore pulses with energy at center frequencies below 1.0 kHz.

No systematic study of injury from low frequency impulses has been reported. The objective of this study was to explore the hearing loss and cochlear injury produced by narrow band impulses with energy concentrated in the low to mid frequencies, i.e., from approximately 0.2 to 4.0 kHz.

Methods and procedures

A detailed presentation of all the methods that were used in these experiments can be found in Patterson et al., (1986). Only brief highlights of the procedures are presented below.

a. Subjects:

The subjects were 118 male chinchilla villadera. The animals were made monaural by surgical destruction of the left cochlea under halothane anesthesia, (Miller, 1970) and were allowed to recover for 2 weeks before any further experimentation took place.

b. Procedures:

Behavioral audiometry was performed using a shock avoidance procedure (Burdick, et al., 1978). A baseline audiogram was established as the average of the last five audiograms prior to exposure. The audiometric test frequencies used were: 0.125, 0.250, 0.5, 1.0, 1.4, 2.0, 2.8, 4.0, 5.7, and 8.0 kHz. Postexposure audiograms were obtained at 2 minutes, 1 hour, 3, 6, 24, 48, and 72 hours after exposure and then twice weekly until 30 days postexposure. Three audiograms were obtained during the final postexposure week. At 30 days postexposure, the animals were anesthetized and killed by decapitation. The cochleas were removed and fixed with a 2.5 percent buffered glutaraldehyde. The fixed tissue was evaluated for loss of sensory cells using surface preparation histological techniques.

c. Exposure Stimuli:

The narrow band impulse noise was produced by exciting an Altec model 515B speaker* in a model 815 enclosure with a signal produced by the PDP 11/34 computer system*. The computer synthesized signal was obtained by exciting a digital band pass filter of the 4-pole Learner-type (Gold and Rader, 1969) with a digital impulse. This filter has a bandwidth of approximately 400 Hz independent of center frequency and steep attenuation

* See Appendix A.

outside the pass band. This permitted the synthesis of a set of impulses of equivalent energy at a variety of center frequencies while assuring minimal spread of energy to other frequencies.

The center frequencies were selected to explore the range from 0.200 to 4.000 kHz. The general strategy was to vary the energy levels of impulses at each center frequency to span the range of injury from large permanent hearing loss to little or no hearing loss. At the lowest frequency, equipment limitations prevented the synthesis of impulses intense enough to produce large permanent threshold shifts. Center frequencies, peak pressures, and sound exposure levels for each experimental group are shown in Table 1. Sound exposure levels (SEL) (Young, 1970) were computed from:

$$SEL = 10 \log_{10} \int_{-\infty}^{\infty} (P^2(t)/P_r^2 t_r) dt$$

where $t_r = 1s$ and $P_r = 20 \mu Pa$.

 Table 1

Definition of experimental groups.

Group	N	Center frequency (Hz)	Center frequency (dB)	Peak SPL (dB)	Total SEL
1	6		260	139	132.5
2	6		260	146	139.8
3	5		775	134	124.8
4	6		775	139	129.4
5	6		775	144	134.8
6	6		1025	129	119.8
7	6		1025	134	124.2
8	5		1025	139	129.1
9	6		1025	144	134.6
10	6		1350	129	119.8
11	6		1350	134	124.2
12	6		1350	139	129.0
13	6		2450	129	120.6
14	6		2450	134	124.9
15	7		2450	139	129.6
16	5		2450	144	135.0
17	6		3550	124	113.0
18	6		3550	129	119.9
19	6		3550	134	124.2
20	6		3550	139	129.5
Total	118				

Figures 1 through 6 show the impulse pressure-time histories and relative frequency amplitude spectra for the six center frequencies used. Each subject was exposed to 100 impulses at a fixed intensity and center frequency (Table 1). Five to seven subjects were exposed individually to one combination of intensity and center frequency. The impulses were delivered at a rate of one every 3 seconds. All exposures were delivered to the unprotected right ear at a normal angle of incidence.

Results and discussion

Audiometric results:

The mean preexposure thresholds for all 118 animals are reported in Table 2 along with the data of Miller (1970) and are shown plotted as a function of frequency in Figure 7. There is a good agreement between the two audiograms. The mean preexposure thresholds for each group and at each audiometric test frequency is presented in Table 3. A two-way analysis of variance with repeated measures on one factor (frequency) showed that there were no statistically significant differences in the mean preexposure thresholds among groups.

Table 2

Summary of mean preexposure thresholds (dB) for all animals (N = 118) compared to published norms.

	Test frequency (Hz)										
	125	250	500	1000	1400	2000	2800	4000	5700	8000	
Present study	23.6	7.8	2.4	1.7	1.7	1.4	2.0	2.0	2.7	4.4	\bar{x}
	2.2	3.8	4.4	3.7	4.1	3.9	3.7	3.2	2.8	3.7	s
Miller (1970)	19.9	8.8	5.1	3.0	2.2	2.7	-0.2	1.9	1.9	5.8	\bar{x}
	5.4	3.9	6.1	4.1	6.6	4.7	4.9	7.1	6.7	5.4	s
	36	36	36	36	34	36	35	36	35	36	N

Table 3

Preexposure threshold means (dB) and standard deviations for all groups.

Group	N	Test frequency (Hz)										\bar{X}	s
		125	250	500	1000	1400	2000	2800	4000	5700	8000		
1	6	24.83	10.33	2.00	1.50	1.83	3.17	3.17	2.50	2.50	3.83	\bar{X}	
		1.17	0.82	2.97	2.07	2.48	1.72	1.72	1.64	2.43	1.72	s	
2	6	23.17	5.50	2.17	1.17	1.17	-0.17	1.83	1.50	1.50	3.17	\bar{X}	
		1.72	2.07	3.06	2.79	2.32	1.72	1.17	2.59	1.05	1.72	s	
3	5	23.20	6.40	1.20	3.20	1.00	1.00	0.40	0.60	2.40	6.20	\bar{X}	
		2.17	3.05	2.77	1.30	3.54	1.87	1.82	1.52	0.55	4.32	s	
4	6	25.00	10.67	3.00	0.67	1.67	3.00	2.33	2.33	3.17	5.83	\bar{X}	
		1.67	0.52	2.28	1.37	2.50	1.41	1.37	1.21	0.98	1.94	s	
5	6	22.50	7.33	2.17	1.83	1.17	0.17	1.50	-0.17	3.17	3.67	\bar{X}	
		1.52	2.34	2.71	1.60	2.40	1.33	1.76	0.98	2.71	1.97	s	
6	6	21.17	5.33	1.17	1.83	0.17	0.83	1.17	0.83	1.17	5.50	\bar{X}	
		3.54	1.97	2.64	2.14	1.47	1.72	3.19	1.72	1.17	2.07	s	
7	6	23.17	5.17	2.17	0.83	2.50	-1.17	0.50	1.17	2.17	3.17	\bar{X}	
		0.98	2.04	3.25	1.60	1.76	4.92	2.66	2.40	1.33	1.60	s	
8	5	24.20	9.80	3.00	1.40	2.20	4.60	2.40	3.40	2.00	3.80	\bar{X}	
		2.59	1.79	2.55	2.97	0.84	1.67	2.61	1.67	1.22	1.79	s	
9	6	24.50	10.83	6.00	5.33	4.00	5.17	7.67	4.33	6.17	10.33	\bar{X}	
		1.64	10.19	16.01	14.19	16.32	14.41	13.50	11.13	8.47	10.76	s	
10	6	23.67	8.50	2.67	3.17	2.33	0.83	2.50	2.50	2.00	3.17	\bar{X}	
		2.16	1.76	4.03	2.56	1.97	3.60	0.84	2.51	2.61	2.04	s	
11	6	23.67	6.00	3.33	2.00	3.33	1.00	2.33	2.67	2.67	2.17	\bar{X}	
		1.51	3.52	2.94	1.67	1.37	2.28	1.75	2.58	1.21	2.64	s	
12	6	24.50	10.50	2.17	2.00	0.50	2.83	2.00	2.50	1.50	5.33	\bar{X}	
		1.64	1.76	3.66	1.41	2.26	0.98	0.89	1.76	2.59	2.07	s	
13	6	23.67	8.33	2.17	2.67	0.33	1.00	1.67	1.00	3.00	5.33	\bar{X}	
		2.94	3.61	2.56	2.16	2.34	1.67	0.82	1.41	3.22	2.58	s	
14	6	24.00	6.33	2.33	1.33	1.67	0.67	0.17	4.00	4.67	1.67	\bar{X}	
		0.63	4.18	3.44	3.61	3.27	1.97	2.04	2.28	2.42	2.58	s	
15	7	24.86	7.43	3.00	1.57	3.14	2.71	3.57	2.43	3.57	3.00	\bar{X}	
		2.41	4.16	3.83	1.13	1.95	1.38	0.79	1.90	1.51	1.63	s	
16	5	22.80	8.60	1.40	-0.20	1.60	-0.40	0.40	0.00	1.80	4.40	\bar{X}	
		2.59	3.21	2.88	2.17	1.82	1.52	1.82	2.12	1.30	3.05	s	
17	6	22.67	6.33	2.00	0.00	0.67	1.67	-0.67	1.00	2.00	3.83	\bar{X}	
		2.25	1.86	2.76	2.45	2.25	1.51	3.27	1.09	2.28	2.99	s	
18	6	23.67	8.00	1.33	0.83	1.00	0.33	1.00	0.67	1.67	5.33	\bar{X}	
		3.01	4.00	2.58	1.17	3.46	1.86	1.79	1.51	1.75	2.34	s	
19	6	23.00	7.33	2.00	2.00	2.67	-0.67	2.33	2.33	3.67	3.67	\bar{X}	
		1.26	3.72	2.97	0.89	1.97	2.66	2.73	3.27	3.67	2.25	s	
20	6	22.83	7.83	2.83	1.33	0.83	1.50	2.17	3.17	2.83	4.83	\bar{X}	
		2.79	4.17	1.72	2.42	1.94	1.87	2.14	2.14	2.79	5.11	s	




Figure 1. The narrow band impulse pressure-time waveform (inset) and the relative frequency spectrum of the impulse for the 0.260 kHz CF, 146 dB peak SPL stimulus.




Figure 2. The narrow band impulse pressure-time waveform (inset) and the relative frequency spectrum of the impulse for the 0.775 kHz CF, 144 dB peak SPL stimulus.




Figure 3. The narrow band impulse pressure-time waveform (inset) and the relative frequency spectrum of the impulse for the 1.025 kHz CF, 139 dB peak SPL stimulus.




Figure 4. The narrow band impulse pressure-time waveform (inset) and the relative frequency spectrum of the impulse for the 1.350 kHz CF, 139 dB peak SPL stimulus.




Figure 5. The narrow band impulse pressure-time waveform (inset) and the relative frequency spectrum of the impulse for the 2.450 kHz CF, 144 dB peak SPL stimulus.




Figure 6. The narrow band impulse pressure-time waveform (inset) and the relative frequency spectrum of the impulse for the 3.550 kHz CF, 139 dB peak SPL stimulus.




Figure 7. The mean preexposure audiogram for all 118 chinchillas (symbols) and for published norms (dashed line). The error bars represent one standard deviation above and below the plotted point.

Figures 8 through 67 illustrate the group mean threshold shifts (TS) measured over a period of 30 days postexposure (threshold recovery functions) for each of the 20 groups of animals in this study, and at each of the 10 audiometric test frequencies. In general for each noise stimulus there is an orderly relation between threshold shift and the peak SPL. Many of these recovery curves, especially those from groups exposed to the higher level stimuli, show a pronounced growth of TS or a lengthy delayed recovery, both of which have been shown to be indications of permanent changes (Hamernik et al., 1988). Because of the growth of the TS recovery functions, the first postexposure threshold measurement often is not the most appropriate index of the severity of early postexposure noise trauma. In Figures 68 through 73 the group mean maximum threshold shift (TS_{max}) for each exposure condition is shown plotted as a function of audiometric test frequency. As might be anticipated from the recovery functions, the TS_{max} shows an orderly relation with stimulus intensity for each of the stimulus center frequencies that were used. One surprising feature seen in the TS_{max} data is the lack of a clear effect of stimulus center frequency. For the noise impulses having CFs at and below 2.450 kHz, the TS_{max} audiograms are surprisingly flat between about 0.250 and 8 kHz. Only the 3.550 kHz CF stimulus produced a clear high frequency effect in the TS_{max} audiogram.

The permanent threshold shift (PTS) was computed for each animal by calculating the mean of the last five postexposure audiograms and subtracting from this the preexposure audiogram for that animal. The group mean PTS audiograms for each exposure stimulus center frequency are shown in Figures 74 through 79. The shape and ordering of these audiograms across stimulus intensity and audiometric test frequency is basically the same as that described for the TS_{max} audiograms. At the lowest CF used (0.260 Hz), the maximum exposure that was possible with the speaker used was not of sufficient intensity to produce more than a 10 dB PTS and this was confined to the test frequencies below 0.5 kHz.

Histological results

Surface preparation histology was obtained for each animal in this study, and cochleograms were plotted. Individual animal cochleograms and group cell loss data are presented in graphical and tabular form in the appendix to this report. In order to analyze the sensory cell data, inner and outer hair cell losses (IHC and OHC) were computed over lengths of the basilar membrane corresponding to octaves with center frequencies from 0.250 to 16.8 kHz, and group mean losses were computed for each exposure group. The results of this type of an analysis for each group is shown in Figures 80 through 85. As in the audiometric results, there was a general ordering of sensory cell loss as a function of stimulus intensity. This effect is evident especially in the OHC data. The inner hair cell losses show some deviation from this ordering at a few locations of the cochlea (e.g., Figure 83). A few other similar anomalies in the IHC losses, such as those evident in Figure 84, are not statistically significant.

In order to understand the relation between the audiometric data, histological data, and the stimulus variables of center frequency and intensity, the data were analyzed as follows: For each of the 20 groups of animals that were exposed to the narrow band impulses, a mean permanent threshold shift evaluated at 1, 2, and 4 kHz ($(PTS)_{1,2,4}$) was computed and the groups were compared on the basis of sound exposure level. The results of this data reduction are shown plotted as a function of SEL in Figure 86. The group mean PTS from each set of the two to four groups of animals that comprise an intensity series for a specific CF impulse behaves in an orderly manner with $(PTS)_{1,2,4}$ increasing in an approximately linear fashion with increasing SEL.

The relative susceptibility to a noise-induced PTS (NIPTS) is seen to be a function of the impulse center frequency, with the lower frequency impulses producing relatively little NIPTS even at the higher SELs. A relative frequency weighting function can be derived from the data presented in Figure 86 by shifting each center frequency intensity series data set along

the SEL axis the amount that is necessary to collapse the data into a single PTS versus SEL function using one of the exposures as a "zero" reference. Such a data shifting process was carried out "by eye" to produce a best fit using the 1.350 kHz series of data as the reference point. The amounts shifted were: 0.260 kHz CF impulses, -20 dB; 0.775 kHz CF impulses, -7.2 dB; 1.025 kHz CF impulses, -4 dB; 1.350 kHz CF impulses, 0 dB; 2.450 kHz CF impulses, -4 dB; and 3.550 kHz CF impulses, +4 dB. The realignment of the data that such a shift produces is shown in Figure 87, and the weighting function, thus obtained, is shown plotted (solid line with symbols) in Figure 88 where it is compared to the conventional A-weighting function (solid line). The new empirical weighting function is referred to as P-weighting in the text that follows. A linear regression through the shifted data set shown in Figure 87 showed a correlation coefficient of 0.894 with a slope of 2.6 dB dB PTS/dB SEL P-weighted sound exposure level. The empirical function derived from the narrow band impulse data is seen to differ from the A-weighting function by as much as 10 dB at the low frequencies. Also evident in this figure is the anomalous behavior of the data point produced by the exposures to the 2.450 kHz, CF impulses.

The histological data were reduced in a similar manner. For each of the 20 exposure conditions, the total IHC and OHC loss in each cochlea was determined and a group average total OHC and IHC loss computed. The group mean total OHC and IHC losses are shown plotted in Figure 89. As with the $(PTS)_{1,2,4}$ data shown in Figure 86, there is an orderly and approximately linear increase of cell loss with increasing stimulus intensity for each center frequency impulse. Also evident from this figure is that for a given impulse intensity, the amount of cell loss is related to the center frequency of the narrow band impulse. The effect of applying the P-weighting function derived from the audiometric data to the data in Figure 89 is shown in Figure 90. The slope of the regression line through the shifted data is 264.3 dB PTS/dB SEL and the correlation coefficient is 0.858 for total OHC loss and 12.4 dB PTS/dB SEL and 0.849 for total IHC loss.

Conclusions

Exposure to narrow band impulses whose primary energy is centered at low frequencies is much less hazardous to the auditory system than impulses with energy concentrated in the mid range frequencies. This implies that the estimation of the hazard to the auditory system from exposure to impulse noise requires a weighting function which de-emphasizes the low frequencies. A-weighting, which is commonly used in the assessment of hazard from steady noise, does not de-emphasize the low frequencies enough. Thus, A-weighted SEL will tend to overestimate the hazard from impulses with large amounts of low

frequency energy. The results of this study suggest a function more like the P-weighting function will more accurately assess the hazard from low frequency impulse noise.

There are some caveats for the conclusions drawn above. While a considerable amount of data (20 exposure conditions) were used to derive the P-weighting function, its generality can not be confirmed until it is tested with a wide variety of impulse noise exposures. In addition, it should be noted that this function was derived from chinchilla data. Data from human exposures will be required before any such function can be used for human hazard assessment.




Figure 8. The mean threshold recovery curves for the groups exposed to the 0.260 kHz CF impulse at the indicated peak SPL for the 0.125 kHz audiometric test frequency.




Figure 9. The mean threshold recovery curves for the groups exposed to the 0.775 kHz CF impulse at the indicated peak SPL for the 0.125 kHz audiometric test frequency.




Figure 10. The mean threshold recovery curves for the groups exposed to the 1.025 kHz CF impulse at the indicated peak SPL for the 0.125 kHz audiometric test frequency.




Figure 11. The mean threshold recovery curves for the groups exposed to the 1.350 kHz CF impulse at the indicated peak SPL for the 0.125 kHz audiometric test frequency.




Figure 12. The mean threshold recovery curves for the groups exposed to the 2.450 kHz CF impulse at the indicated peak SPL for the 0.125 kHz audiometric test frequency.




Figure 13. The mean threshold recovery curves for the groups exposed to the 3.550 kHz CF impulse at the indicated peak SPL for the 0.125 kHz audiometric test frequency.




Figure 14. The mean threshold recovery curves for the groups exposed to the 0.260 kHz CF impulse at the indicated peak SPL for the 0.250 kHz audiometric test frequency.




Figure 15. The mean threshold recovery curves for the groups exposed to the 0.775 kHz CF impulse at the indicated peak SPL for the 0.250 kHz audiometric test frequency.




Figure 16. The mean threshold recovery curves for the groups exposed to the 1.025 kHz CF impulse at the indicated peak SPL for the 0.250 kHz audiometric test frequency.




Figure 17. The mean threshold recovery curves for the groups exposed to the 1.350 kHz CF impulse at the indicated peak SPL for the 0.250 kHz audiometric test frequency.




Figure 18. The mean threshold recovery curves for the groups exposed to the 2.450 kHz CF impulse at the indicated peak SPL for the 0.250 kHz audiometric test frequency.




Figure 19. The mean threshold recovery curves for the groups exposed to the 3.550 kHz CF impulse at the indicated peak SPL for the 0.250 kHz audiometric test frequency.




Figure 20. The mean threshold recovery curves for the groups exposed to the 0.260 kHz CF impulse at the indicated peak SPL for the 0.500 kHz audiometric test frequency.




Figure 21. The mean threshold recovery curves for the groups exposed to the 0.775 kHz CF impulse at the indicated peak SPL for the 0.500 kHz audiometric test frequency.




Figure 22. The mean threshold recovery curves for the groups exposed to the 1.025 kHz CF impulse at the indicated peak SPL for the 0.500 kHz audiometric test frequency.




Figure 23. The mean threshold recovery curves for the groups exposed to the 1.350 kHz CF impulse at the indicated peak SPL for the 0.500 kHz audiometric test frequency.




Figure 24. The mean threshold recovery curves for the groups exposed to the 2.450 kHz CF impulse at the indicated peak SPL for the 0.500 kHz audiometric test frequency.




Figure 25. The mean threshold recovery curves for the groups exposed to the 3.550 kHz CF impulse at the indicated peak SPL for the 0.500 kHz audiometric test frequency.




Figure 26. The mean threshold recovery curves for the groups exposed to the 0.260 kHz CF impulse at the indicated peak SPL for the 1.000 kHz audiometric test frequency.




Figure 27. The mean threshold recovery curves for the groups exposed to the 0.775 kHz CF impulse at the indicated peak SPL for the 1.000 kHz audiometric test frequency.




Figure 28. The mean threshold recovery curves for the groups exposed to the 1.025 kHz CF impulse at the indicated peak SPL for the 1.000 kHz audiometric test frequency.




Figure 29. The mean threshold recovery curves for the groups exposed to the 1.350 kHz CF impulse at the indicated peak SPL for the 1.000 kHz audiometric test frequency.




Figure 30. The mean threshold recovery curves for the groups exposed to the 2.450 kHz CF impulse at the indicated peak SPL for the 1.000 kHz audiometric test frequency.




Figure 31. The mean threshold recovery curves for the groups exposed to the 3.550 kHz CF impulse at the indicated peak SPL for the 1.000 kHz audiometric test frequency.




Figure 32. The mean threshold recovery curves for the groups exposed to the 0.260 kHz CF impulse at the indicated peak SPL for the 1.400 kHz audiometric test frequency.




Figure 33. The mean threshold recovery curves for the groups exposed to the 0.775 kHz CF impulse at the indicated peak SPL for the 1.400 kHz audiometric test frequency.




Figure 34. The mean threshold recovery curves for the groups exposed to the 1.025 kHz CF impulse at the indicated peak SPL for the 1.400 kHz audiometric test frequency.




Figure 35. The mean threshold recovery curves for the groups exposed to the 1.350 kHz CF impulse at the indicated peak SPL for the 1.400 kHz audiometric test frequency.




Figure 36. The mean threshold recovery curves for the groups exposed to the 2.450 kHz CF impulse at the indicated peak SPL for the 1.400 kHz audiometric test frequency.




Figure 37. The mean threshold recovery curves for the groups exposed to the 3.550 kHz CF impulse at the indicated peak SPL for the 1.400 kHz audiometric test frequency.




Figure 38. The mean threshold recovery curves for the groups exposed to the 0.260 kHz CF impulse at the indicated peak SPL for the 2.000 kHz audiometric test frequency.




Figure 39. The mean threshold recovery curves for the groups exposed to the 0.775 kHz CF impulse at the indicated peak SPL for the 2.000 kHz audiometric test frequency.




Figure 40. The mean threshold recovery curves for the groups exposed to the 1.025 kHz CF impulse at the indicated peak SPL for the 2.000 kHz audiometric test frequency.




Figure 41. The mean threshold recovery curves for the groups exposed to the 1.350 kHz CF impulse at the indicated peak SPL for the 2.000 kHz audiometric test frequency.




Figure 42. The mean threshold recovery curves for the groups exposed to the 2.450 kHz CF impulse at the indicated peak SPL for the 2.000 kHz audiometric test frequency.




Figure 43. The mean threshold recovery curves for the groups exposed to the 3.550 kHz CF impulse at the indicated peak SPL for the 2.000 kHz audiometric test frequency.




Figure 44. The mean threshold recovery curves for the groups exposed to the 0.260 kHz CF impulse at the indicated peak SPL for the 2.800 kHz audiometric test frequency.




Figure 45. The mean threshold recovery curves for the groups exposed to the 0.775 kHz CF impulse at the indicated peak SPL for the 2.800 kHz audiometric test frequency.




Figure 46. The mean threshold recovery curves for the groups exposed to the 1.025 kHz CF impulse at the indicated peak SPL for the 2.800 kHz audiometric test frequency.




Figure 47. The mean threshold recovery curves for the groups exposed to the 1.350 kHz CF impulse at the indicated peak SPL for the 2.800 kHz audiometric test frequency.




Figure 48. The mean threshold recovery curves for the groups exposed to the 2.450 kHz CF impulse at the indicated peak SPL for the 2.800 kHz audiometric test frequency.




Figure 49. The mean threshold recovery curves for the groups exposed to the 3.550 kHz CF impulse at the indicated peak SPL for the 2.800 kHz audiometric test frequency.




Figure 50. The mean threshold recovery curves for the groups exposed to the 0.260 kHz CF impulse at the indicated peak SPL for the 4.000 kHz audiometric test frequency.




Figure 51. The mean threshold recovery curves for the groups exposed to the 0.775 kHz CF impulse at the indicated peak SPL for the 4.000 kHz audiometric test frequency.




Figure 52. The mean threshold recovery curves for the groups exposed to the 1.025 kHz CF impulse at the indicated peak SPL for the 4.000 kHz audiometric test frequency.




Figure 53. The mean threshold recovery curves for the groups exposed to the 1.775 kHz CF impulse at the indicated peak SPL for the 4.000 kHz audiometric test frequency.




Figure 54. The mean threshold recovery curves for the groups exposed to the 2.450 kHz CF impulse at the indicated peak SPL for the 4.000 kHz audiometric test frequency.




Figure 55. The mean threshold recovery curves for the groups exposed to the 3.550 kHz CF impulse at the indicated peak SPL for the 4.000 kHz audiometric test frequency.




Figure 56. The mean threshold recovery curves for the groups exposed to the 0.260 kHz CF impulse at the indicated peak SPL for the 5.700 kHz audiometric test frequency.




Figure 57. The mean threshold recovery curves for the groups exposed to the 0.775 kHz CF impulse at the indicated peak SPL for the 5.700 kHz audiometric test frequency.




Figure 58. The mean threshold recovery curves for the groups exposed to the 1.025 kHz CF impulse at the indicated peak SPL for the 5.700 kHz audiometric test frequency.




Figure 59. The mean threshold recovery curves for the groups exposed to the 1.350 kHz CF impulse at the indicated peak SPL for the 5.700 kHz audiometric test frequency.




Figure 60. The mean threshold recovery curves for the groups exposed to the 2.450 kHz CF impulse at the indicated peak SPL for the 5.700 kHz audiometric test frequency.




Figure 61. The mean threshold recovery curves for the groups exposed to the 3.550 kHz CF impulse at the indicated peak SPL for the 5.700 kHz audiometric test frequency.




Figure 62. The mean threshold recovery curves for the groups exposed to the 0.260 kHz CF impulse at the indicated peak SPL for the 8.000 kHz audiometric test frequency.




Figure 63. The mean threshold recovery curves for the groups exposed to the 0.775 kHz CF impulse at the indicated peak SPL for the 8.000 kHz audiometric test frequency.




Figure 64. The mean threshold recovery curves for the groups exposed to the 1.025 kHz CF impulse at the indicated peak SPL for the 8.000 kHz audiometric test frequency.




Figure 65. The mean threshold recovery curves for the groups exposed to the 1.350 kHz CF impulse at the indicated peak SPL for the 8.000 kHz audiometric test frequency.




Figure 66. The mean threshold recovery curves for the groups exposed to the 2.450 kHz CF impulse at the indicated peak SPL for the 8.000 kHz audiometric test frequency.




Figure 67. The mean threshold recovery curves for the groups exposed to the 3.550 kHz CF impulse at the indicated peak SPL for the 8.000 kHz audiometric test frequency.




Figure 68. The group mean maximum threshold shift for each audiometric test frequency following exposure to the 0.260 kHz CF impulse at the indicated peak SPL.




Figure 69. The group mean maximum threshold shift for each audiometric test frequency following exposure to the 0.775 kHz CF impulse at the indicated peak SPL.




Figure 70. The group mean maximum threshold shift for each audiometric test frequency following exposure to the 1.025 kHz CF impulse at the indicated peak SPL.




Figure 71. The group mean maximum threshold shift for each audiometric test frequency following exposure to the 1.350 kHz CF impulse at the indicated peak SPL.




Figure 72. The group mean maximum threshold shift for each audiometric test frequency following exposure to the 2.450 kHz CF impulse at the indicated peak SPL.




Figure 73. The group mean maximum threshold shift for each audiometric test frequency following exposure to the 3.550 kHz CF impulse at the indicated peak SPL.




Figure 74. The group mean permanent threshold shift (PTS) for each audiometric test frequency following exposure to the 0.260 kHz CF impulse at the indicated peak SPL.




Figure 75. The group mean permanent threshold shift (PTS) for each audiometric test frequency following exposure to the 0.775 kHz CF impulse at the indicated peak SPL.




Figure 76. The group mean permanent threshold shift (PTS) for each audiometric test frequency following exposure to the 1.025 kHz CF impulse at the indicated peak SPL.




Figure 77. The group mean permanent threshold shift (PTS) for each audiometric test frequency following exposure to the 1.350 kHz CF impulse at the indicated peak SPL.




Figure 78. The group mean permanent threshold shift (PTS) for each audiometric test frequency following exposure to the 2.450 kHz CF impulse at the indicated peak SPL.




Figure 79. The group mean permanent threshold shift (PTS) for each audiometric test frequency following exposure to the 3.550 kHz CF impulse at the indicated peak SPL.




Figure 80. The group mean outer hair cell (OHC) loss and inner hair cell (IHC) loss within octave band lengths of the basilar membrane at the indicated frequencies following exposure to the 0.260 kHz CF impulse at the indicated peak SPL.




Figure 81. The group mean outer hair cell (OHC) loss and inner hair cell (IHC) loss within octave band lengths of the basilar membrane at the indicated frequencies following exposure to the 0.775 kHz CF impulse at the indicated peak SPL.




Figure 82. The group mean outer hair cell (OHC) loss and inner hair cell (IHC) loss within octave band lengths of the basilar membrane at the indicated frequencies following exposure to the 1.025 kHz CF impulse at the indicated peak SPL.




Figure 83. The group mean outer hair cell (OHC) loss and inner hair cell (IHC) loss within octave band lengths of the basilar membrane at the indicated frequencies following exposure to the 1.350 kHz CF impulse at the indicated peak SPL.




Figure 84. The group mean outer hair cell (OHC) loss and inner hair cell (IHC) loss within octave band lengths of the basilar membrane at the indicated frequencies following exposure to the 2.450 kHz CF impulse at the indicated peak SPL.




Figure 85. The group mean outer hair cell (OHC) loss and inner hair cell (IHC) loss within octave band lengths of the basilar membrane at the indicated frequencies following exposure to the 3.550 kHz CF impulse at the indicated peak SPL.




Figure 86. The group mean PTS evaluated at 1, 2 and 4 kHz ($PTS_{1,2,4}$) as a function of the total sound exposure level for the twenty groups of animals exposed to the narrow band impulses.




Figure 87. The $\overline{PTS}_{1,2,4}$ as a function of the P-weighted total sound exposure level. The regression line yields a correlation coefficient of 0.904.




Figure 88. The empirical P-weighting function (symbols) derived from the narrow band exposure data along with the conventional A-weighting function (solid line).




Figure 89. The group mean total outer (upper) and inner (lower) hair cell loss as a function of the total sound exposure level for the twenty groups of animals exposed to the narrow band impulses.




Figure 90. The group mean total outer (upper) and inner (lower) hair cell loss as a function of the P-weighted total sound level. The regression lines yield correlation coefficients of 0.858 and 0.849 for outer and inner hair cell loss respectively.

References

- Burdick, C. K., Patterson, J. H. Jr., Mozo, B. T., and Camp, R. T., Jr. 1978. Threshold shifts in chinchillas exposed to octave bands of noise centered at 63 and 1000 Hz for three days. Journal of the acoustical society of America. 64: 458-466.
- Coles, R. R. A., Garinther, G. R., Hodge, D. C., and Rice, C. G. 1968. Hazardous exposure to impulse noise. Journal of the acoustical society of America. 43: 336-343.
- Gold, B., and Rader, C. M. 1969. Digital processing of signals, McGraw Hill Publishers, New York, NY.
- Hamernik, R. P., Patterson, J. H., Jr. and Salvi, R. J. 1986. The effects of impulse intensity and the number of impulses on hearing and cochlear pathology in the chinchilla. Journal of the acoustical society of America. 81: 1118-1129.
- Hamernik, R. P., Ahroon, W. A., and Patterson, J. H., Jr. 1988. Threshold recovery functions following impulse noise trauma. Journal of the acoustical society of America. 84: 941-950.
- Kryter, K. D., Ward, W. D., Miller, J. D., and Eldredge, D. H. 1966. Hazardous exposure to intermittent and steady-state noise. Journal of the acoustical society of America. 39: 451-464.
- OSHA, Department of Labor. 1974. Occupational noise exposure. Proposed requirements and procedures. Federal register, 39, 207: 155-159.
- Miller, J. D. 1970. Audibility curve of the chinchilla. Journal of the acoustical society of America. 48: 513-523.
- Mills, J. H., Osguthorpe, J. D., Burdick, C. K., Patterson, J. H. Jr., and Mozo, B. 1983. Temporary threshold shifts produced by exposure to low-frequency noises. Journal of the acoustical society of America. 73: 918-923.
- NAS-NRC committee on hearing, bioacoustics and biomechanics (CHABA, 1968). Proposed damage risk criterion for impulse noise (gunfire). Washington, DC.
- Patterson, J. H., Jr., Lomba Gautier, I. M., Curd, D. L., and Hamernik, R. P. 1986. The role of peak pressure in determining the auditory hazard of impulse noise. Fort Rucker, AL: U.S. Army Aeromedical Research Laboratory. USAARL Report No. 86-7.

- Pfander, F., Bongartz, H., Brinkmann, H., and Kietz, H. 1980. Danger of auditory impairment from impulse noise: A comparative study of the CHABA damage-risk criteria and those of the Federal Republic of Germany. Journal of acoustical society of America. 67: 628-633.
- Price, G. R., Kim, H. N., Lim, D. J., and Dunn, D. 1988. Hazard from weapons impulses: histological and electrophysiological evidence. Journal of the acoustical society of America. 85: 1245-1254.
- Price, G. R. 1979. Loss of auditory sensitivity following exposure to spectrally narrow impulses. Journal of the acoustical society of America. 66: 456-465.
- Smooenburg, G. F. 1982. Damage risk criteria for impulse noise. In: New perspectives on noise-induced hearing loss, eds. R. P. Hamernik, D. Henderson, and R. J. Salvi (Raven Press, New York). pp. 471-490.
- Young, R. W. 1970. On the energy transported with a sound pulse. Journal of the acoustical society of America. 47: 441-442.

Appendix A

List of manufacturers

Altec Lansing Corporation
1515 South Manchester Avenue
Anaheim, CA 92803

Digital Equipment Corporation
Maynard, MA 01754

Initial distribution

Commander, U.S. Army Natick Research,
Development and Evaluation Center
ATTN: STRNC-MIL (Documents
Librarian)
Natick, MA 01760-5040

Naval Submarine Medical
Research Laboratory
Medical Library, Naval Sub Base
Box 900
Groton, CT 06340

Commander/Director
U.S. Army Combat Surveillance
and Target Acquisition Lab
ATTN: DELCS-D
Fort Monmouth, NJ 07703-5304

Commander
10th Medical Laboratory
ATTN: Audiologist
APO New York 09180

Naval Air Development Center
Technical Information Division
Technical Support Detachment
Warminster, PA 18974

Commanding Officer, Naval Medical
Research and Development Command
National Naval Medical Center
Bethesda, MD 20814-5044

Deputy Director, Defense Research
and Engineering
ATTN: Military Assistant
for Medical and Life Sciences
Washington, DC 20301-3080

Commander, U.S. Army Research
Institute of Environmental Medicine
Natick, MA 01760

U.S. Army Avionics Research
and Development Activity
ATTN: SAVAA-P-TP
Fort Monmouth, NJ 07703-5401

U.S. Army Communications-Electronics
Command
ATTN: AMSEL-RD-ESA-D
Fort Monmouth, NJ 07703

Library
Naval Submarine Medical Research Lab
Box 900, Naval Sub Base
Groton, CT 06349-5900

Commander
Man-Machine Integration System
Code 602
Naval Air Development Center
Warminster, PA 18974

Commander
Naval Air Development Center
ATTN: Code 602-B (Mr. Brindle)
Warminster, PA 18974

Commanding Officer
Harry G. Armstrong Aerospace
Medical Research Laboratory
Wright-Patterson
Air Force Base, OH 45433

Director
Army Audiology and Speech Center
Walter Reed Army Medical Center
Washington, DC 20307-5001

Commander, U.S. Army Institute
of Dental Research
ATTN: Jean A. Setterstrom, Ph. D.
Walter Reed Army Medical Center
Washington, DC 20307-5300

Naval Air Systems Command
Technical Air Library 950D
Room 278, Jefferson Plaza II
Department of the Navy
Washington, DC 20361

Naval Research Laboratory Library
Shock and Vibration
Information Center, Code 5804
Washington, DC 20375

Director, U.S. Army Human
Engineering Laboratory
ATTN: Technical Library
Aberdeen Proving Ground, MD 21005

Commander, U.S. Army Test
and Evaluation Command
ATTN: AMSTE-AD-H
Aberdeen Proving Ground, MD 21005

Director
U.S. Army Ballistic
Research Laboratory
ATTN: DRXBR-OD-ST Tech Reports
Aberdeen Proving Ground, MD 21005

Commander
U.S. Army Medical Research
Institute of Chemical Defense
ATTN: SGRD-UV-AO
Aberdeen Proving Ground,
MD 21010-5425

Commander, U.S. Army Medical
Research and Development Command
ATTN: SGRD-RMS (Ms. Madigan)
Fort Detrick, Frederick, MD 21702-5012

Director
Walter Reed Army Institute of Research
Washington, DC 20307-5100

HQ DA (DASG-PSP-O)
5109 Leesburg Pike
Falls Church, VA 22041-3258

Naval Research Laboratory
Library Code 1433
Washington, DC 20375

Harry Diamond Laboratories
ATTN: Technical Information Branch
2800 Powder Mill Road
Adelphi, MD 20783-1197

U.S. Army Materiel Systems
Analysis Agency
ATTN: AMXSY-PA (Reports Processing)
Aberdeen Proving Ground
MD 21005-5071

U.S. Army Ordnance Center
and School Library
Simpson Hall, Building 3071
Aberdeen Proving Ground, MD 21005

U.S. Army Environmental
Hygiene Agency
Building E2100
Aberdeen Proving Ground, MD 21010

Technical Library Chemical Research
and Development Center
Aberdeen Proving Ground, MD
21010-5423

Commander
U.S. Army Medical Research
Institute of Infectious Disease
SGRD-UIZ-C
Fort Detrick, Frederick, MD 21702

Director, Biological
Sciences Division
Office of Naval Research
600 North Quincy Street
Arlington, VA 22217

Commander
U.S. Army Materiel Command
ATTN: AMCDE-XS
5001 Eisenhower Avenue
Alexandria, VA 22333

Commandant
U.S. Army Aviation
Logistics School ATTN: ATSQ-TDN
Fort Eustis, VA 23604

Headquarters (ATMD)
U.S. Army Training
and Doctrine Command
Fort Monroe, VA 23651

Structures Laboratory Library
USARTL-AVSCOM
NASA Langley Research Center
Mail Stop 266
Hampton, VA 23665

Naval Aerospace Medical
Institute Library
Building 1953, Code 03L
Pensacola, FL 32508-5600

Command Surgeon
HQ USCENTCOM (CCSG)
U.S. Central Command
MacDill Air Force Base FL 33608

Air University Library
(AUL/LSE)
Maxwell Air Force Base, AL 36112

U.S. Air Force Institute
of Technology (AFIT/LDEE)
Building 640, Area B
Wright-Patterson
Air Force Base, OH 45433

Henry L. Taylor
Director, Institute of Aviation
University of Illinois-Willard Airport
Savoy, IL 61874

Chief, Nation Guard Bureau
ATTN: NGB-ARS (COL Urbauer)
Room 410, Park Center 4
4501 Ford Avenue
Alexandria, VA 22302-1451

Commander
U.S. Army Aviation Systems Command
ATTN: SGRD-UAX-AL (MAJ Gillette)
4300 Goodfellow Blvd., Building 105
St. Louis, MO 63120

U.S. Army Aviation Systems Command
Library and Information Center Branch
ATTN: AMSAV-DIL
4300 Goodfellow Boulevard
St. Louis, MO 63120

Federal Aviation Administration
Civil Aeromedical Institute
Library AAM-400A
P.O. Box 25082
Oklahoma City, OK 73125

Commander
U.S. Army Academy
of Health Sciences
ATTN: Library
Fort Sam Houston, TX 78234

Commander
U.S. Army Institute of Surgical Research
ATTN: SGRD-USM (Jan Duke)
Fort Sam Houston, TX 78234-6200

AAMRL/HEX
Wright-Patterson
Air Force Base, OH 45433

John A. Dellinger,
Southwest Research Institute
P. O. Box 28510
San Antonio, TX 78284

Product Manager
Aviation Life Support Equipment
ATTN: AMCPM-ALSE
4300 Goodfellow Boulevard
St. Louis, MO 63120-1798

Commander
U.S. Army Aviation
Systems Command
ATTN: AMSAV-ED
4300 Goodfellow Boulevard
St. Louis, MO 63120

Commanding Officer
Naval Biodynamics Laboratory
P.O. Box 24907
New Orleans, LA 70189-0407

Assistant Commandant
U.S. Army Field Artillery School
ATTN: Morris Swott Technical Library
Fort Sill, OK 73503-0312

Commander
U.S. Army Health Services Command
ATTN: HSOP-SO
Fort Sam Houston, TX 78234-6000

Director of Professional Services
HQ USAF/SGDT
Bolling Air Force Base, DC 20332-6188

U.S. Army Dugway Proving Ground
Technical Library, Building 5330
Dugway, UT 84022

U.S. Army Yuma Proving Ground
Technical Library
Yuma, AZ 85364

AFFTC Technical Library
6510 TW/TSTL
Edwards Air Force Base,
CA 93523-5000

Commander
Code 3431
Naval Weapons Center
China Lake, CA 93555

Aeromechanics Laboratory
U.S. Army Research and Technical Labs
Ames Research Center, M/S 215-1
Moffett Field, CA 94035

Sixth U.S. Army
ATTN: SMA
Presidio of San Francisco, CA 94129

Commander
U.S. Army Aeromedical Center
Fort Rucker, AL 36362

U.S. Air Force School
of Aerospace Medicine
Strughold Aeromedical Library Technical
Reports Section (TSKD)
Brooks Air Force Base, TX 78235-5301

Dr. Diane Damos
Department of Human Factors
ISSM, USC
Los Angeles, CA 90089-0021

U.S. Army White Sands
Missile Range
ATTN: STEWS-IM-ST
White Sands Missile Range, NM 88002

U.S. Army Aviation Engineering
Flight Activity
ATTN: SAVTE-M (Tech Lib) Stop 217
Edwards Air Force Base, CA 93523-5000

Ms. Sandra G. Hart
Ames Research Center
MS 262-3
Moffett Field, CA 94035

Commander, Letterman Army Institute
of Research
ATTN: Medical Research Library
Presidio of San Francisco, CA 94129

Mr. Frank J. Stagnaro, ME
Rush Franklin Publishing
300 Orchard City Drive
Campbell, CA 95008

Commander
U.S. Army Medical Materiel
Development Activity
Fort Detrick, Frederick, MD 21702-5009

Commander
U.S. Army Aviation Center
Directorate of Combat Developments
Building 507
Fort Rucker, AL 36362

U. S. Army Research Institute
Aviation R&D Activity
ATTN: PERI-IR
Fort Rucker, AL 36362

Commander
U.S. Army Safety Center
Fort Rucker, AL 36362

U.S. Army Aircraft Development
Test Activity
ATTN: STEBG-MP-P
Cairns Army Air Field
Fort Rucker, AL 36362

Commander U.S. Army Medical Research
and Development Command
ATTN: SGRD-PLC (COL Sedge)
Fort Detrick, Frederick, MD 21702

MAJ John Wilson
TRADOC Aviation LO
Embassy of the United States
APO New York 09777

Netherlands Army Liaison Office
Building 602
Fort Rucker, AL 36362

British Army Liaison Office
Building 602
Fort Rucker, AL 36362

Italian Army Liaison Office
Building 602
Fort Rucker, AL 36362

Directorate of Training Development
Building 502
Fort Rucker, AL 36362

Chief
USAHEL/USAAVNC Field Office
P. O. Box 716
Fort Rucker, AL 36362-5349

Commander U.S. Army Aviation Center
and Fort Rucker
ATTN: ATZQ-CG
Fort Rucker, AL 36362

Commander/President
TEXCOM Aviation Board
Cairns Army Air Field
Fort Rucker, AL 36362

MAJ Terry Newman
Canadian Army Liaison Office
Building 602
Fort Rucker, AL 36362

German Army Liaison Office
Building 602
Fort Rucker, AL 36362

LTC Patrick Laparra
French Army Liaison Office
USAAVNC (Building 602)
Fort Rucker, AL 36362-5021

Australian Army Liaison Office
Building 602
Fort Rucker, AL 36362

Dr. Garrison Rapmund
6 Burning Tree Court
Bethesda, MD 20817

Commandant Royal Air Force
Institute of Aviation Medicine
Farnborough Hampshire GU14 6SZ UK

Dr. A. Kornfield, President
Biosearch Company
3016 Revere Road
Drexel Hill, PA 29026

Commander
U.S. Army Biomedical Research
and Development Laboratory
ATTN: SGRD-UBZ-I
Fort Detrick, Frederick, MD 21702

Defense Technical Information Center
Cameron Station
Alexandria, VA 22313

Commander, U.S. Army Foreign Science
and Technology Center
AIFRTA (Davis)
220 7th Street, NE
Charlottesville, VA 22901-5396

Director,
Applied Technology Laboratory
USARTL-AVSCOM
ATTN: Library, Building 401
Fort Eustis, VA 23604

U.S. Air Force Armament
Development and Test Center
Eglin Air Force Base, FL 32542

Aviation Medicine Clinic
TMC #22, SAAF
Fort Bragg, NC 28305

Commander, U.S. Army Missile
Command
Redstone Scientific Information Center
ATTN: AMSMI-RD-CS-R/ILL
Documents Redstone Arsenal, AL 35898

U.S. Army Research and Technology
Laboratories (AVSCOM)
Propulsion Laboratory MS 302-2
NASA Lewis Research Center
Cleveland, OH 44135

Dr. H. Dix Christensen
Bio-Medical Science Building, Room 753
Post Office Box 26901
Oklahoma City, OK 73190

Col. Otto Schramm Filho
c/o Brazilian Army Commission
Office-CEBW
4632 Wisconsin Avenue NW
Washington, DC 20016

Dr. Christine Schlichting
Behavioral Sciences Department
Box 900, NAVUBASE NLON
Groton, CT 06349-5900

COL Eugene S. Channing, O.D.
Brooke Army Medical Center
ATTN: HSHE-EAH-O
Fort Sam Houston, TX 78234-6200

U.S. Army Training
and Doctrine Command
ATTN: Surgeon
Fort Monroe, VA 23651-5000