

2

DTIC  
S ELECTE D  
OCT 24 1991  
C

AD-A241 880



CHEMICAL  
RESEARCH,  
DEVELOPMENT &  
ENGINEERING  
CENTER

CRDEC-TR-301

FIELD EMISSION SCANNING ELECTRON MICROSCOPY  
OF THE INTERACTION OF CLOSTRIDIUM DIFFICILE  
WITH HUMAN EMBRYONIC INTESTINAL CELLS

Sheila Wood  
Erica Petersen

RESEARCH DIRECTORATE

August 1991

Approved for public release; distribution is unlimited.



U.S. ARMY  
ARMAMENT  
MUNITIONS  
CHEMICAL COMMAND

Aberdeen Proving Ground, Maryland 21010-5423

91-13916



91 10 23 032

Disclaimer

The findings in this report are not to be construed as an official Department of the Army position unless so designated by other authorizing documents.

# REPORT DOCUMENTATION PAGE

Form Approved  
OMB No. 0704-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.

<b>1. AGENCY USE ONLY (Leave blank)</b>		<b>2. REPORT DATE</b> 1991 August	<b>3. REPORT TYPE AND DATES COVERED</b> Final, 84 Jan - 91 Jan	
<b>4. TITLE AND SUBTITLE</b> Field Emission Scanning Electron Microscopy of the Interaction of <u>Clostridium Difficile</u> with Human Embryonic Intestinal Cells			<b>5. FUNDING NUMBERS</b>  PR-1FJ1-2-R	
<b>6. AUTHOR(S)</b>  Wood, Sheila, and Petersen, Erica				
<b>7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES)</b>  CDR, CRDEC, ATTN: SMCCR-RSB, APG, MD 21010-5423			<b>8. PERFORMING ORGANIZATION REPORT NUMBER</b>  CRDEC-TR-301	
<b>9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)</b>			<b>10. SPONSORING/MONITORING AGENCY REPORT NUMBER</b>	
<b>11. SUPPLEMENTARY NOTES</b>				
<b>12a. DISTRIBUTION/AVAILABILITY STATEMENT</b>  Approved for public release; distribution is unlimited.			<b>12b. DISTRIBUTION CODE</b>	
<b>13. ABSTRACT (Maximum 200 words)</b>  An in vitro assay system, consisting of monolayers of human embryonic intestinal cells (HEI) and <u>Clostridium difficile</u> ( <u>C. difficile</u> ), was used to observe interactions of microorganisms with the cell surface and cytoplasm. Tests were performed using Toxin B positive <u>C. difficile</u> Strain 938, and Toxin B negative Strain 789, harvested from casamino acids agar (CAS), and CAS containing subinhibitory concentrations of clindamycin (CAS-CL). Adherence to the surface and internalization into HEI cells was quantified at 30, 60, and 90 min. Cryopreserved and freeze fractured preparations were made at 60 min and observed for adherence and internalization using field emission scanning electron microscopy. Quantitative microorganism determinations, confirmed by the observed microorganism activity harvested from CAS and CAS-CL, revealed their capability to adhere to and/or enter HEI cells. The Toxin B producer, Strain 938, harvested from CAS-CL entered the cytoplasm and remained intact within it.				
<b>14. SUBJECT TERMS</b> Internalization                      Clindamycin Scanning electron <u>C. difficile</u> Microscopy			<b>15. NUMBER OF PAGES</b> 16	
			<b>16. PRICE CODE</b>	
<b>17. SECURITY CLASSIFICATION OF REPORT</b> UNCLASSIFIED	<b>18. SECURITY CLASSIFICATION OF THIS PAGE</b> UNCLASSIFIED	<b>19. SECURITY CLASSIFICATION OF ABSTRACT</b> UNCLASSIFIED	<b>20. LIMITATION OF ABSTRACT</b>  UL	

Blank

## PREFACE

The work described in this report was authorized under Project No. 1FJ1-2-R, Biotechnology. This work was started in January 1984\* and completed in January 1991.

The use of trade names or manufacturers' names in this report does not constitute an official endorsement of any commercial products. This report may not be cited for purposes of advertisement.

Reproduction of this document in whole or in part is prohibited except with permission of the Commander, U.S. Army Chemical Research, Development and Engineering Center, ATTN: SMCCR-SPS-T, Aberdeen Proving Ground, MD 21010-5423. However, the Defense Technical Information Center and the National Technical Information Service are authorized to reproduce the document for U.S. Government purposes.

This report has been approved for release to the public.

### Acknowledgments

The authors acknowledge Dr. Bill Wergin and the members of the electron microscopy facility at the U.S. Department of Agriculture (Beltsville, MD) for the use of their facilities, and also Deryck J. Mills, Applications Engineer for Oxford Instruments (Oxford, England), for his instructive help in instrument use.

Approved for  
Distribution  
By  
Distribution  
A-1

---

\*This work started when the primary author was a student at the Medical College of Virginia, Richmond, VA.

Blank

## CONTENTS

	Page
1. INTRODUCTION .....	7
2. MATERIALS AND METHODS .....	8
2.1 Organisms Used .....	8
2.2 Organism Preparation for Adherence Assay .....	8
2.3 Tissue Cells Used .....	8
2.4 Assay for Cytotoxicity .....	8
2.5 Adherence Assay .....	9
2.6 Internalization Assay .....	9
2.7 Field Emission Scanning Electron Microscopy, Cryopreservation, and Freeze Fracture .....	9
3. RESULTS .....	10
3.1 Cytotoxicity Assay .....	10
3.2 Adherence Assay .....	10
3.3 Internalization Assay .....	10
3.4 Field Emission Scanning Electron Microscopy .....	10
4. DISCUSSION .....	10
LITERATURE CITED .....	15

Blank

# FIELD EMISSION SCANNING ELECTRON MICROSCOPY OF THE INTERACTION OF CLOSTRIDIUM DIFFICILE WITH HUMAN EMBRYONIC INTESTINAL CELLS

## 1. INTRODUCTION

Clostridium difficile (*C. difficile*), the etiologic agent of pseudomembranous colitis, unpredictably colonizes the bowels of certain hospitalized patients following the administration of antibiotics or during the course of a hospital stay.<sup>1</sup> Enterotoxins produced by this microorganism are capable of damaging the bowel mucosa;<sup>2</sup> but until now, it has not been determined whether [as with microorganisms such as Escherichia coli (*E. coli*)] adhesive factors help this microorganism colonize and proliferate before toxin production.<sup>3-7</sup>

Previous studies using the hamster model clarified some assumptions about proper conditions for colonization. In the hamster, diet, colonization, and competition for colonization sites, were shown to affect the outcome of an experimentally induced infection. More specifically, administration of a high percentage of protein in the diet (or total food deprivation for several days) hampered the animal's ability to resist establishment of C. difficile colitis in the presence or absence of antibiotics.<sup>8</sup> Also, when antibiotics were administered during the course of a high percentage protein diet, experimental lifespans were shortened compared to the administration of antibiotics or increased protein alone. Evidence suggests a synergistic action of diet with antibiotics.<sup>4,8</sup>

Indirectly, a colonization mechanism has been elucidated in the following experiments. The presence of a nonpathogenic strain of C. difficile in the bowel, before challenge with a pathogenic strain, protected the bowel from colonization.<sup>1,2,9,10</sup> This protection was shown not to be immunologic but resulted from exclusion of the pathogenic strain from the mucosa.<sup>11</sup> Although increased survival time of cefoxitin-treated hamsters was observed when prior treatment with a noncytotoxic strain was employed, no protection was seen when cytotoxic and noncytotoxic strains were administered simultaneously. These observations show that attachment sites were blocked by nonpathogenic variants.<sup>8</sup> Colonization appears to be site dependent and may be influenced by the selective production of cell wall components resulting from microbial consumption of available nutrients.<sup>5,12</sup>

This in vitro study notes the effects of a high percentage of protein in the growth media and exposure to clindamycin on the behavior of C. difficile on human embryonic intestinal (HEI) cells. The role of selected nutrients and clindamycin in the predisposition of C. difficile to attach to cell surfaces, enter, and survive within HEI cells is shown. A combination of selected nutrients and the presence of clindamycin, in addition to the microenvironment, may be critical to the development of pseudomembranous colitis. As no valid explanation for relapse has yet been postulated, one explanation that must now be considered is survival within nonphagocytic cells in response to selected microbial nutrients and clindamycin.

## 2. MATERIALS AND METHODS

### 2.1 Organisms Used.

Clostridium difficile strains, isolated from the patient population at the Medical College of Virginia, Virginia Commonwealth University (Richmond, VA), include C. difficile Strain 938 from a symptomatic adult and C. difficile Strain 789 isolated from an asymptomatic child. Assays for the production of Toxin B<sup>7</sup> showed that Strain 938 caused rounding of HEI cells in 24 hr, and Strain 789 did not cause rounding after 48 hr. Rounding was inhibited by C. difficile antitoxin obtained from Dr. Tracey Wilkins, Virginia Polytechnic Institute and State University (Blacksburg, VA).

Identification was based on the appearance of the colony, fluorescence under a 366-nm lamp, gram stain morphology, the absence of lipase or lecithinase, and gas-liquid chromatography profiles. The C. difficile isolates had minimum inhibitory concentrations of 1.0 µg/mL clindamycin when tested by agar dilution susceptibility.<sup>13</sup>

### 2.2 Organism Preparation for Adherence Assay.

Microorganisms were grown on casamino acids agar (CAS) consisting of 1% casamino acids (DIFCO Labs, Detroit MI), 2% agar (DIFCO), and 0.5% sheep blood (Flow Labs, McLean, VA). To expose the organisms to subinhibitory concentrations of clindamycin, 0.05 µg/mL clindamycin was added to the CAS media. Organisms were harvested at 48 hr in Eagle's minimal essential media (MEM) (Flow Labs) containing 1% fetal calf serum (FCS) (GIBCO, Grand Island, NY), pH 5.5, washed once, and resuspended in MEM 1% FCS. Antitoxin at a 1:10 dilution in 0.05 mL MEM was used in an attempt to block adherence. This mixture was incubated at 37 °C in an anaerobe jar with the atmosphere generated by an anaerobe gas pack [Becton Dickinson Microbiology Systems (BDMS), Cockeysville, MD] for 15 min. The preparations were centrifuged at 1500 rpm for 5 min and resuspended in MEM with 1% FCS, pH 5.5.

### 2.3 Tissue Cells Used.

Human embryonic intestinal cells #CCL6 were obtained from the American Type Culture Collection (Rockville, MD). They were maintained in HAM's F12 medium (GIBCO) containing 10% FCS (GIBCO) and penicillin/streptomycin (DIFCO), 50 units and 50 µg/mL, respectively.

### 2.4 Assay for Cytotoxigenicity.

Procedures for the determination of cytotoxigenicity were followed as previously described.<sup>7</sup> Briefly, the cytotoxin assay tested supernatant dilutions from each microorganism for rounding of HEI cells from 4 to 48 hr after inoculation. If rounding was observed, and this rounding was blocked using antitoxin to Toxin B, then the organisms were considered cytotoxigenic.

## 2.5 Adherence Assay.

The adherence assay was performed using C. difficile grown on CAS as previously described. Earlier studies measured adherence of seven different isolates under five different conditions, in duplicate.<sup>7</sup> Briefly, microorganisms were inoculated onto cell monolayers, incubated anaerobically on a rotating platform for 15 min at 37 °C, washed, stained with Giemsa stain, and read. The percentage of cells (number of cells out of 100 with adherent microorganisms) with microorganisms adhering was calculated. Areas with low background adherence were chosen for these counts.

To determine whether adherence depended on microbe viability, viable and nonviable microorganisms were used initially. Viable C. difficile were taken from CAS plates, which had been incubated anaerobically for 48 hr. Nonviable microorganisms were taken from blood agar plates, which had been stored aerobically for 3 days. Subcultures from aerobic plates failed to produce growth when plated on sheep blood agar (SBA) and incubated anaerobically at 37 °C for 48 hr. Gram stains confirmed intact organisms.

## 2.6 Internalization Assay.

Adherence assay procedures were followed with several exceptions: assays were carried out in 24-well Falcon plates (BDMS); confluent monolayers were inoculated with approximately  $5 \times 10^5$  organisms; five wells per test condition were harvested; and incubation was in gas pak pouches (BDMS) generating an anaerobic environment. At 30, 60, and 90 min, monolayers were washed to remove loose microorganisms, and a 1% solution of Triton X100 was placed on the monolayer and incubated at room temperature for 5 min. After 5 min, the microorganisms were harvested, dilutions were made, and microorganisms were subcultured to SBA and incubated anaerobically for 48 hr at which time colony counts were established. To the same 24-well tissue culture plate, in different test wells, clindamycin at 10 µg/mL was added to kill any microorganisms attached to the monolayer surface. After incubation in an anaerobe pouch for 90 min, these test wells were washed to remove dead organisms, and a 1% solution of Triton X100 was added for 5 min. The microorganisms were harvested, diluted, and plated on SBA for subsequent colony counts. To determine the percentage of internalized microorganisms, a ratio of internalized organisms to total cell associated organisms  $\times 100$  was obtained. Microorganisms 789 and 938, harvested from CAS and CAS containing subinhibitory concentrations of clindamycin (CAS-CL), were used.

## 2.7 Field Emission Scanning Electron Microscopy, Cryopreservation, and Freeze Fracture.

The adherence assay was performed as previously described.<sup>7</sup> Preparations were rinsed in buffer and fixed in 3% glutaraldehyde at room temperature after a 60-min incubation at 37 °C. Cryopreservation (Strain 938, CAS) in liquid nitrogen was performed, and etching was used to achieve clarity of the image. Freeze fracture (Strain 938, CAS-CL) was performed by freezing the coverslip containing the cell/microorganism layer in liquid nitrogen, then manually fracturing the coverslip inside a cold chamber.

### 3. RESULTS

#### 3.1 Cytotoxicity Assay.

The in vitro assay for the presence of Toxin B showed rounding of 100% HEI cells at 24 hr by Strain 938. Strain 789, however, caused no rounding of HEI cells after 48 hr.

#### 3.2 Adherence Assay.

Adherence assays previously performed, in duplicate, on seven different isolates gave a 60% average of cells with microorganisms attached. The preparative adherence assay performed prior to this set of experiments showed 62 and 67% attachment for Strains 938 and 789, respectively. In preliminary experiments comparing attachment of viable microorganisms to attachment of nonviable microorganisms, only viable microorganisms attached, which indicated that an active process was at work in the attachment phenomena.

#### 3.3 Internalization Assay.

As seen in the table and Figure 1, the ratio of internal to total cell associated organisms was highest with Strain 938 harvested from CAS-CL at 60 min. Data from the in vitro adherence assay was substantiated by reproducibility of the in vitro results (data not shown) and electron micrograph results.

#### 3.4 Field Emission Scanning Electron Microscopy.

In Figure 2, the cryopreserved preparation of Strain 938 harvested from CAS, showed intimate interaction with HEI cells. In Figure 3, Strain 938 harvested from CAS-CL, shown by freeze fracture, is intact within HEI cells. (Microorganisms are shown with arrows.)

### 4. DISCUSSION

Adults develop C. difficile associated colitis following the administration of antibiotics; whereas, infants harbor C. difficile in the bowel but rarely show signs of disease. In addition, historical data has shown that intravenous and intramuscular administration of antibiotics are associated with the development of pseudomembranous colitis; but, a direct correlation of disease symptoms to dose does not exist.<sup>14</sup> Predisposing environmental conditions in the bowel, which are indirectly related to antibiotic administrations, have been implicated in the development of pseudomembranous colitis. These conditions include disruptions in the protective layer of the bowel, selected antibiotic administration, changes in health or nutritional status of the host, food deprivation, and a designated diet or change in diet.<sup>10</sup> Changes in the organism instead of in the environmental conditions in the bowel have not yet been considered. Clostridium difficile strains differ in their ability to produce cytotoxin, in the presence of cell wall components such as adhesins, in their response to the immediate environment, in the use of selected nutrients, and in their proximity to the bowel mucosa. Organism changes in response to growth conditions cannot be discounted as contributive to the development of pseudomembranes.

TABLE. Numbers of Total Cell Associated *C. difficile* and Internalized *C. difficile*. Percentages were obtained by dividing the number of internalized organisms by the number of total cell associated organisms and multiplying by 100.

Organism/ conditions	Total cell- associated organisms CFU/mL	Internalized organisms CFU/mL	Percentage internal/ total x 100
<u>938 CAS</u>			
30 min	$5.20 \times 10^4$	$1.30 \times 10^4$	25.0
60 min	$5.80 \times 10^4$	$1.74 \times 10^4$	30.0
90 min	$1.55 \times 10^5$	$4.50 \times 10^3$	2.9
<u>938 CAS-CL</u>			
30 min	$4.00 \times 10^4$	$1.20 \times 10^4$	30.0
60 min	$3.50 \times 10^4$	$4.40 \times 10^4$	125.7
90 min	$1.00 \times 10^6$	$4.75 \times 10^3$	0.5
<u>789 CAS</u>			
30 min	$1.40 \times 10^4$	$4.80 \times 10^2$	3.4
60 min	$5.00 \times 10^3$	$1.75 \times 10^3$	35.0
90 min	$2.85 \times 10^3$	$1.30 \times 10^2$	4.5
<u>789 CAS-CL</u>			
30 min	$2.00 \times 10^4$	$8.00 \times 10^2$	4.0
60 min	$7.20 \times 10^3$	$1.78 \times 10^3$	24.7
90 min	$1.75 \times 10^4$	$4.50 \times 10^2$	2.6




Figure 1. Internalization of *C. difficile* by Human Embryonic Intestinal Cells. Ratio of internalized microorganisms versus total cell associated microorganisms at 30, 60, and 90 min. The greatest ratio was observed with *C. difficile* Strain 938 harvested from CAS-CL.




Figure 2. Field Emission Scanning Electron Microscopy of *C. difficile* Strain 938, Harvested from CAS. Intimate association with the cells is apparent. Interaction was stopped by fixation at 60 min, and the sample was cryopreserved prior to observation. Bar = 2.73  $\mu$ .




Figure 3. Field Emission Scanning Electron Micrograph of a Freeze Fractured Preparation of *C. difficile*, Harvested from CAS Containing 0.05  $\mu$ g/mL Clindamycin, Intact Within Human Embryonic Intestinal Cells After 60 min. Organisms, as shown by the arrows, are numerous and appear unharmed. Bar = 6.0  $\mu$ .

Initial results, presented here, suggest a direct effect of available nutrients (amino acids) and clindamycin on the microorganisms. Entrance into and survivability within HEI cells occurred with Toxin B producer, Strain 938, when harvested from media containing amino acids and clindamycin. Ratios of internalized microorganisms to total cell associated microorganisms were determined. Clostridium difficile Strain 938, harvested from CAS-CL were internalized and survived in high numbers at 60 min. When grown on CAS containing no clindamycin, levels of internalization and survival were not as great.

Adhesins have been implicated as an important initiating factor in colonization by enterotoxigenic E. coli.<sup>15</sup> In the case of C. difficile, such an adhesin may play a role in disease production by serving as the initial anchor to the intestinal epithelium. Response of the microorganism to an environmental signal and/or regulatory control and its role in disease production remain to be determined.<sup>15-17</sup>

Blank

## LITERATURE CITED

1. Borriello, S.P., Larson, H.E., Honour, P., and Barclay, F.E., "Antibiotic Associated Diarrhea and Colitis," In Clinical Research Center Report, Medical Research Council, London, England, pp 96-98, 1982.
2. Borriello, S.P., "Clostridium difficile and Gut Disease," In Microbes and Infections of the Gut, Blackwell Scientific Publications, Oxford, England, pp 327-346, 1980.
3. Finley, B.B., Heffron, F., and Falkow, S., "Epithelial Cell Surfaces Induce Salmonella Proteins Required for Bacterial Adherence Invasion," Science Vol. 243, pp 940-941 (1989).
4. Freter, R., "Mechanisms of Association of Bacteria with Mucosal Surfaces," In Adhesion and Microorganism Pathogenicity, Pitman Medical Publishers, London, England, pp 36-55, 1981.
5. Hu, P.C., Cole, R.M., and Huang, Y.S., "Mycoplasma pneumoniae Infection: Role of a Surface Protein in the Attachment Organelle," Science Vol. 216, p 313 (1982).
6. Levine, M.M., "Adhesion of Enterotoxigenic Escherichia coli in Animals and Humans," In Adhesion and Microorganism Pathogenicity, Pitman Medical Publishers, London, England, pp 142-160, 1981.
7. Wood-Helie, S.J., Dalton, H.P., and Shadomy, S., "Hydrophobic and Adherence Properties of Clostridium difficile," Eur. J. Clin. Microbiol Vol. 5, pp 441-445 (1986).
8. Silva, J., Fekety, R., Werk, C., Ebrifgt, J., Cudmore, M., Batts, D., Syrjamaki, C., and Lukens, J., "Inciting and Etiological Agents of Colitis," Rev. Infect. Dis. Vol. 6, pp S214-S220 (1984).
9. Borriello, S.P., and Barclay, F.E., "Clostridium difficile Infection Protection by Prior Colonization with Nonpathogenic Strains," In Models of Anaerobic Infection, Martinus Nijhoff Publishers, Boston, MA, pp 239-240, 1984.
10. Lee, A., "Normal Flora of Animal Intestinal Surfaces," In Adsorption of Microorganisms to Surfaces, Wiley and Sons, New York, NY, pp 145-173, 1980.
11. Wilson, K.H., and Sheagren, J.N., "Antagonism of Toxigenic Clostridium difficile," J. Infect. Dis. Vol. 147, pp 733-736 (1983).
12. Woese, C.R., Maniloff, J., and Zablen, L.B., "Phylogenic Analysis of the Mycoplasmas," Proc Natl Acad Sci USA Vol. 77, p 494 (1980).
13. The National Committee for Clinical Laboratory Standards, Reference Agar Dilution Procedure for Antimicrobial Susceptibility Testing of Anaerobic Bacteria, Vol. 5, pp 25-29, National Institute of Health, Bethesda, MD, 1985.

14. Tedosco, F.J., Barton, R.W, and Alpers, D.H., "Clindamycin Associated Colitis: A Prospective Study," Ann Intern Med Vol. 81, pp 429-433 (1974).

15. Finley, B.B., Heffron, F., and Falkow, S., "Epithelial Cell Surfaces Induce Salmonella Proteins Required for Bacterial Adherence and Invasion," Science Vol. 243, pp 940-941 (1989).

16. Isberg, R.R., Voorhis, D.L., and Falkow, S., "Identification of Invasion: A Protein that Allows Enteric Bacteria to Penetrate Culture Mammalian Cells," Cell Vol. 50, pp 769-778 (1987).

17. Miller, J.F., Mekalanos, J.J., and Falkow, S., "Coordinate Regulation and Sensory Transduction in the Control of Bacterial Virulence," Science Vol. 243, pp 916-921 (1989).