

2

NAVAL POSTGRADUATE SCHOOL

Monterey, California



DTIC
SELECTE
JAN 06 1993
S B D

INVERSE PROBLEMS FOR ORTHOGONAL
MATRICES, TODA FLOWS,
AND SIGNAL PROCESSING
by
L. Faybusovich
G. S. Ammar
W. B. Gragg
Technical Report For Period
July 1992 - September 1992

Approved for public release; distribution unlimited

Prepared for: Naval Postgraduate School
Monterey, CA 93943

422 33 8
93-00347



NAVAL POSTGRADUATE SCHOOL
MONTEREY, CA 93943

Rear Admiral R. W. West, Jr.
Superintendent

Harrison Shull
Provost

This report was prepared in conjunction with research conducted for the Naval Postgraduate School and funded by the Naval Postgraduate School.

Reproduction of all or part of this report is authorized.

This report was prepared by:

L. Faybusovich
L. FAYBUSOVICH
Professor

G. S. Ammar
G. S. AMMAR
Professor

W. B. Gragg
WILLIAM B. GRAGG
Professor

Reviewed by:

Richard Franke
For RICHARD FRANKE
Chairman
Department of Mathematics

Released by:

Paul J. Marto
PAUL J. MARTO
Dean of Research

UNCLASSIFIED


SECURITY CLASSIFICATION OF THIS PAGE

REPORT DOCUMENTATION PAGE

Form Approved
OMB No 0704-0188

1a REPORT SECURITY CLASSIFICATION UNCLASSIFIED			1b RESTRICTIVE MARKINGS			
2a SECURITY CLASSIFICATION AUTHORITY			3 DISTRIBUTION / AVAILABILITY OF REPORT Approved for public release; Distribution unlimited			
2b DECLASSIFICATION / DOWNGRADING SCHEDULE						
4. PERFORMING ORGANIZATION REPORT NUMBER(S) NPS-MA-93-001			5 MONITORING ORGANIZATION REPORT NUMBER(S) NPS-MA-93-001			
6a NAME OF PERFORMING ORGANIZATION Naval Postgraduate School		6b OFFICE SYMBOL (if applicable) MA	7a. NAME OF MONITORING ORGANIZATION Naval Postgraduate School			
6c. ADDRESS (City, State, and ZIP Code) Monterey, CA 93943			7b ADDRESS (City, State, and ZIP Code) Monterey, CA 93943			
8a. NAME OF FUNDING / SPONSORING ORGANIZATION Naval Postgraduate School		8b. OFFICE SYMBOL (if applicable) MA	9 PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER O & M,N Direct Funding			
8c. ADDRESS (City, State, and ZIP Code) Monterey, CA 93943			10 SOURCE OF FUNDING NUMBERS			
			PROGRAM ELEMENT NO	PROJECT NO	TASK NO	WORK UNIT ACCESSION NO
11 TITLE (Include Security Classification) Inverse problems for orthogonal matrices, Toda flows, and signal processing						
12. PERSONAL AUTHOR(S) L. Faybusovich, G. S. Ammar, W. B. Gragg						
13a TYPE OF REPORT Technical		13b TIME COVERED FROM 7/92 TO 9/92		14 DATE OF REPORT (Year, Month, Day) 921013	15 PAGE COUNT 6	
16 SUPPLEMENTARY NOTATION						
17 COSATI CODES			18 SUBJECT TERMS (Continue on reverse if necessary and identify by block number) orthogonal matrices, Toda flows, signal processing			
FIELD	GROUP	SUB-GROUP				
19 ABSTRACT (Continue on reverse if necessary and identify by block number) We consider Toda flows induced on the set of orthogonal upper Hessenberg matrices. The explicit formulas for the evolution of Schur parameters are given.						
20 DISTRIBUTION / AVAILABILITY OF ABSTRACT <input checked="" type="checkbox"/> UNCLASSIFIED/UNLIMITED <input type="checkbox"/> SAME AS RPT <input type="checkbox"/> DTIC USERS			21 ABSTRACT SECURITY CLASSIFICATION UNCLASSIFIED			
22a NAME OF RESPONSIBLE INDIVIDUAL WILLIAM B. GRAGG			22b TELEPHONE (Include Area Code) (408) 646-2194		22c OFFICE SYMBOL MA/Gr	

SECURITY CLASSIFICATION OF THIS PAGE



Inverse problems for orthogonal matrices, Toda flows, and signal processing

L. Faybusovich

Department of Mathematics, University of Notre Dame

G.S. Ammar

Department of Math. Sciences, Northern Illinois University

W.B. Gragg

Department of Mathematics, Naval Postgraduate School

Abstract

We consider Toda flows induced on the set of orthogonal upper Hessenberg matrices. The explicit formulas for the evolution of Schur parameters are given.

1 Introduction

Any symmetric nonnegative definite Toeplitz matrix T_{n+1} of order $n + 1$ can be modeled as an autocorrelation matrix of a stationary signal [12]

$$x_m = \sum_{l=1}^p \alpha_l \cos(m\omega_l + \theta_l) + y_m,$$

where θ_l are arbitrary phase shifts and y_m is a zero mean white noise process whose variance equals the smallest eigenvalue λ_{\min} of T_{n+1} . Assume that the eigenvalue λ_{\min} is simple, and let (η_0, \dots, η_n) be a corresponding eigenvector. Then [12] the polynomial

$$\psi_n(\lambda) = \eta_0 + \dots + \eta_n \lambda^n$$

has n distinct roots $\lambda_1, \dots, \lambda_n$ on the unit circle, and the frequencies of x_m are given by $\{\exp(\pm i\omega_l)\}_{l=1}^p = \{\lambda_j\}_{j=1}^n$, where i denotes the imaginary unit. One can construct [2] an orthogonal Hessenberg matrix O with characteristic polynomial proportional to ψ_n . Moreover,

the amplitudes α_l can be recovered from the first components of the normalized eigenvectors of O . One can then use any of several algorithms designed for unitary and orthogonal Hessenberg eigenproblems [9, 6, 1, 10, 4]

to calculate the frequencies and amplitudes. In the present paper we investigate another aspect of orthogonal Hessenberg matrices. Namely, we consider Toda flows on these matrices (referred to as Schur flows in [3]) and obtain explicit formulas for the evolution of the so-called Schur parameters under the Toda flow. Since Schur parameters determine orthogonal Hessenberg matrices uniquely, we actually obtain an explicit description of the evolution of a given orthogonal Hessenberg matrix under the Toda flow.

2 Inverse problem for orthogonal Hessenberg matrices

Let M be the set of positive Borel measures on C which have the following properties. For any $\mu \in M$ the support of μ , which we denote by Λ_μ , consists of exactly n points which lie on the unit circle U and is such that: i) if $\lambda \in \Lambda_\mu$, then the complex conjugate $\bar{\lambda}$ is also in Λ_μ and $\mu\{\lambda\} = \mu\{\bar{\lambda}\}$; ii) $\mu\{\lambda\} > 0$ for any λ in Λ_μ ; iii) $\mu(C) = 1$; iv) $-1 \notin \Lambda_\mu$. We further introduce a class OH_+ of orthogonal matrices

$O = \| o_{ij} \|$ such that $o_{ij} = 0$ if $i - j > 1$, $o_{i+1,i} > 0$ for all i and $\det O = 1$. Finally, given a vector $\tau = (\tau_0, \dots, \tau_{n-1})^T \in R^n$ introduce a corresponding Toeplitz matrix $T(\tau) = \| t_{ij} \|$, where $t_{ij} = \tau_{|i-j|}$.

Theorem 2.1 Given a positive definite symmetric Toeplitz matrix $T(\tau)$ with $\tau_0 = 1$ there exist exactly one measure $\mu \in M$ and exactly one $O \in OH_+$ such that

$$\int_C \lambda^i d\mu(\lambda) = \tau_i = \langle e_1, O^i e_1 \rangle, \quad (2.1)$$

$i = 0, \dots, n-1$. Here e_1, \dots, e_n is the canonical basis in R^n and \langle, \rangle is the standard scalar product. Conversely, for any $\mu \in M$ and any $O \in OH_+$ the matrices $T(\tau)$, $T(\tau')$ are Positive definite Toeplitz matrices. Here

$$\tau_i = \int_C \lambda^i d\mu(\lambda), \tau'_i = \langle e_1, O^i e_1 \rangle,$$

$i = 0, \dots, n-1$.

Remark 2.2 Theorem 2.1 is more or less known to the experts (see e.g. [8], [11]). We nevertheless give an independent proof to clarify relationships between introduced objects.

Remark 2.3 There is nothing mysterious about the number -1 which we have excluded from the support of each measure in M . This simplifies notations a little bit.

We need the following elementary lemma.

Lemma 2.4 Let v_1, \dots, v_{n-1} be an orthonormal system of vectors in R^n . There exists exactly one orthogonal matrix O such that $Oe_i = v_i, i = 1, \dots, n-1$ and $\det O = 1$.

We can now outline a proof of Theorem 2.1.

Proof: Denote by P_n the vector space of real polynomials of degree less or equal $n-1$. Set

$$\langle \lambda^i, \lambda^j \rangle = \int_C \lambda^{i-j} d\mu(\lambda). \quad (2.2)$$

We prove that (2.2) defines a positive definite scalar product on P_n . Observe that

$$\int \lambda^i \bar{\lambda}^j d\mu = \int \lambda^{i-j} d\mu = \int \bar{\lambda}^i \lambda^j d\mu.$$

Indeed, $\int \lambda^i \bar{\lambda}^j d\mu = \sum_{\lambda \in \Lambda_\mu} \lambda^i \bar{\lambda}^j \mu\{\lambda\} = \sum_{\lambda \in \Lambda_\mu} \lambda^{i-j} \mu\{\lambda\}$, since $\bar{\lambda} = \lambda^{-1}$ for $\lambda \in U$. Further, since $\mu\{\lambda\} = \mu\{\bar{\lambda}\}$ we have

$$\sum_{\lambda \in \Lambda_\mu} \lambda^{i-j} \mu\{\lambda\} = \sum_{\lambda \in \Lambda_\mu} \bar{\lambda}^{i-j} \mu\{\bar{\lambda}\} = \int \bar{\lambda}^i \lambda^j d\mu.$$

Let $q = a_0 + \dots + a_{n-1} \lambda^{n-1} \in P_n$. We have

$$\langle q, q \rangle = \sum_{i,j=0}^{n-1} a_i a_j \int \lambda^i \bar{\lambda}^j d\mu = \int |q|^2 d\mu \geq 0.$$

Further, $\int |q|^2 d\mu = 0$ if and only if $q(\lambda) = 0$ for any $\lambda \in \Lambda_\mu$. Since $\deg q < n = \text{card}(\Lambda_\mu)$, this is possible only if $q = 0$. Consider the polynomial $\xi(\lambda) = \prod_{t \in \Lambda_\mu} (\lambda - t) = b_0 + \dots + b_{n-1} \lambda^{n-1} + \lambda^n$. Since all roots of ξ lie on the unit circle we clearly have $\lambda^n \xi(1/\lambda) = b_0 \xi(\lambda), b_0 = \pm 1$. Further, all coefficients of ξ are real because $\bar{\Lambda}_\mu = \Lambda_\mu$. Consider the linear operator $O : P_n \rightarrow P_n$ defined as follows: $O\lambda^i = \lambda^{i+1}, i = 0, \dots, n-2, O\lambda^{n-1} = -b_0 - b_1 \lambda - \dots - b_{n-1} \lambda^{n-1}$. We now prove that O is orthogonal relative to the scalar product \langle, \rangle . We should prove that

$$\langle O\lambda^i, \lambda^j \rangle = \langle \lambda^i, O^{-1}\lambda^j \rangle$$

for any $i, j = 0, \dots, n-1$. The only non-trivial case is $i = n-1, j = 0$. We have $\langle O\lambda^{n-1}, 1 \rangle = -b_0 - b_1 \tau_1 - \dots - b_{n-1} \tau_{n-1}$, where $\tau_i = \int_C \lambda^i d\mu(\lambda)$. Let $O^{-1}1 = c_0 + \dots + c_{n-1} \lambda^{n-1}$. Then $\langle \lambda^{n-1}, O^{-1}1 \rangle = c_0 \tau_{n-1} + c_1 \tau_{n-2} + \dots + c_{n-1}$. Thus, it is sufficient to prove that $c_i = -b_{n-1-i}, i = 0, \dots, n-1$. We clearly have $1 = c_0 O1 + \dots + c_{n-1} O\lambda^{n-1} = c_0 \lambda + \dots + c_{n-2} \lambda^{n-1} + c_{n-1} (-b_0 - b_1 \lambda - \dots - b_{n-1} \lambda^{n-1})$ or $1 = -c_{n-1} b_0, c_0 - c_{n-1} b_1 = 0, c_1 - c_{n-1} b_2 = 0, \dots, c_{n-2} - c_{n-1} b_{n-1} = 0$. This yields $b_1 = -c_0/b_0, b_2 = -c_1/b_0, \dots, b_{n-1} = -c_{n-2}/b_0$. We now use the relation $\lambda^n \xi(1/\lambda) = b_0 \xi(\lambda)$. It follows that $b_{n-i} = b_0 b_i, i = 0, \dots, n$. Thus $b_{n-i}/b_0 = -c_{i-1}/b_0, i = 1, \dots, n$. These are exactly the required conditions. Thus we have constructed an orthogonal operator O such that $\int_C \lambda^i d\mu = \langle 1, O^i 1 \rangle, i = 0, \dots, n-1$. Observe that the characteristic

polynomial of O coincides with ξ . Thus the spectrum of O is Λ_μ . In particular, $\det O = 1$ (here we use the assumption that $-1 \notin \Lambda_\mu$). Let $p_0 = 1, \dots, p_{n-1}$ be an orthonormal basis in P_n obtained by the orthonormalization of the basis $1, \lambda, \dots, \lambda^{n-1}$. It is clear that the matrix \bar{O} of the operator O is upper Hessenberg in this basis. Moreover, the entries $\bar{o}_{i+1,i}$ are all nonzero (otherwise, $\text{span}(p_0, \dots, p_{i-1}) = \text{span}(1, \dots, \lambda^{i-1})$ is an invariant subspace of O which is not true). Without loss of generality one can suppose that $\bar{o}_{i+1,i} > 0$ for all i . Otherwise one can take $\text{diag}(\epsilon_1, \dots, \epsilon_n) \bar{O} \text{diag}(\epsilon_1, \dots, \epsilon_n)$.

Suppose we are given a positive definite Toeplitz matrix $T(\tau)$ and an orthogonal matrix $O \in OH_+$ such that $\tau_i = \langle e_1, O^i e_1 \rangle, i = 0, \dots, n-1$. Then

$$T(\tau) = V^T V, \quad (2.3)$$

where V is the upper triangular matrix $[e_1, Oe_1, \dots, O^{n-1}e_1]$ with positive entries on the main diagonal. But (2.3) is the Cholesky decomposition of $T(\tau)$. Hence it is uniquely defined by $T(\tau)$. In other words, the vectors $Oe_1, \dots, O^{n-1}e_1$ are uniquely defined by $T(\tau)$. Since these vectors form a basis, the vectors Oe_1, \dots, Oe_{n-1} are uniquely defined by our Toeplitz matrix. Thus by Lemma 2.4 the matrix O is uniquely defined by $T(\tau)$. Given a positive definite Toeplitz matrix $T(\tau)$ we can endow P_n with a scalar product \langle, \rangle and the shift operator defined on $\text{span}(1, \lambda, \dots, \lambda^{n-2})$ as we did before. Then using Lemma 2.4 we can extend this operator to the orthogonal operator O , defined on P_n such that $\det O = 1$. Then the matrix of O in the basis obtained by orthonormalization of the basis $1, \lambda, \dots, \lambda^{n-1}$ belongs to OH_+ and $\tau_i = \langle e_1, O^i e_1 \rangle, i = 0, \dots, n-1$. Consider now the rational function

$$f(z) = \langle 1, (zI - O)^{-1} 1 \rangle.$$

As is easily seen

$$f(z) = \sum_{i=1}^n \frac{r_i}{z - \lambda_i},$$

where all $r_i > 0$. We then can define the measure $\mu \in M$ by the conditions $\mu\{\lambda_i\} = r_i$ and equal to zero otherwise. We immediately see that equations (2.1) are satisfied. It remains to prove that the measure μ is defined uniquely by conditions (2.1). Let $\mu_k \in M, k = 1, 2$ be such that

$$\int_C \lambda^i d\mu_1 = \int_C \lambda^i d\mu_2,$$

$i = 0, \dots, n-1$. Then we can construct $O_k, k = 1, 2$ such that conditions (2.1) are satisfied. But then $O_1 = O_2$. In particular, $\Lambda_{\mu_1} = \Lambda_{\mu_2}$, i.e., $\mu_1 = \mu_2$ because we have for $\mu\{\lambda\}$ the following system of Vandermonde equations:

$$\sum_{\lambda \in \Lambda_\mu} \lambda^i \mu\{\lambda\} = \tau_i,$$

$i = 0, \dots, n-1$. ■

Let $T(\tau)$ be a positive definite $n \times n$ Toeplitz matrix and \langle, \rangle be the corresponding scalar product on P_n . Let

$$p_i(\lambda) = \delta_i \lambda^i + \dots, \delta_i > 0, i = 0, \dots, n-1,$$

be the basis obtained by the orthonormalization procedure from the basis $1, \lambda, \dots, \lambda^{n-1}$. Since p_i is orthogonal to $\text{span}(1, \lambda, \dots, \lambda^{i-1})$, we have: $\lambda p_i(\lambda)$ is orthogonal to $\text{span}(\lambda, \dots, \lambda^i)$. Further, $r = \lambda p_i(\lambda)/\delta_i - p_{i+1}/\delta_{i+1} \in P_{i+1}$. Let $\varphi_i \in P_{i+1}$ be such that $\langle q, \varphi_i \rangle = q(0)$ for any $q \in P_{i+1}$. Since p_i is orthogonal to P_i and both r and φ_i are orthogonal to λP_i , we obtain

$$\lambda p_i(\lambda)/\delta_i = p_{i+1}(\lambda)/\delta_{i+1} + \gamma_i \varphi_i, \quad (2.4)$$

for some real $\gamma_i, i = 0, \dots, n-2$. An easy calculation shows that $\varphi_i = \delta_i \lambda^i p_i(1/\lambda)$. Hence

$$1 = \delta_i^2 / \delta_{i+1}^2 + \gamma_i^2 \delta_i^4. \quad (2.5)$$

In other words, if we know $\gamma_0, \dots, \gamma_{n-2}$, we can find $\delta_1, \dots, \delta_{n-1}$. Then using (2.4), one can determine p_1, \dots, p_{n-1} and consequently using

again (2.4) the corresponding upper Hessenberg orthogonal matrix O . We have by (2.4) $\langle \lambda p_i(\lambda), p_i(\lambda) \rangle = \gamma_i \delta_i p_i(0)$, $i = 0, \dots, n-2$. Evaluating (2.4) at 0, we obtain $p_{i+1}(0) = -\gamma_i \delta_i^2 \delta_{i+1}$, $i = 0, \dots, n-2$. Thus $\sigma_{i+1, i+1} = \langle \lambda p_i(\lambda), p_i(\lambda) \rangle = -\gamma_i \gamma_{i-1} \delta_{i-1}^2 \delta_i^2$, $i = 1, \dots, n-2$. Further, $\sigma_{1,1} = -\gamma_0 p_0(0) = -\gamma_0$. Let us set

$$\sigma_i = \sigma_{i+1, i} = \delta_{i-1} / \delta_i, \quad \nu_i = \gamma_{i-1} \delta_{i-1}^2,$$

$i = 1, \dots, n-1$. We obviously have

$$\sigma_i^2 + \nu_i^2 = 1, \quad \sigma_{i, i} = -\nu_{i-1} \nu_i,$$

$i = 1, \dots, n-1$, $\nu_0 = 1$. Further, $\sigma_{n, n} = \pm \sqrt{1 - \sigma_{n-1}^2}$. The sign is defined by the condition $\det O = 1$. The quantities ν_i, σ_i are called Schur parameters and auxiliary Schur parameters, respectively. As we saw above the Schur parameters ν_i , $i = 1, \dots, n-1$, determine O uniquely.

On the other hand, if we know the entries $\sigma_{i+1, i} = \delta_i / \delta_{i+1}$, $i = 1, \dots, n-1$ of the matrix O we can determine γ_i by (2.5) up to a sign. In other words, the entries $\sigma_{i+1, i}$ (auxiliary Schur parameters) determine O almost uniquely.

3 Explicit formulas for the evolution of auxiliary Schur parameters under the Toda flow

Let $O(t) = \| \sigma_{ij}(t) \|$ be the solution to the Toda flow

$$\dot{O} = [O, \pi O],$$

such that $O(0)$ is upper Hessenberg orthogonal and irreducible. Here $\pi O = O_- - O_-^T$ and O_- is strictly lower triangular part of O . Then $O(t)$ possesses the same properties and $O(t)$ converges when $t \rightarrow \infty$ to a block diagonal matrix. Each two by two block corresponds to a pair of complex conjugate eigenvalues. The blocks are arranged in the decreasing order of real parts of eigenvalues [7, 5]. From the previous discussion we know that $O(t)$ is almost

uniquely defined by its auxiliary Schur parameters $\sigma_i(t) = \sigma_{i+1, i}(t)$. We now describe explicitly how these parameters evolve under the Toda flow.

Theorem 3.1

$$\sigma_i(t) = \frac{\sqrt{\Delta_{i+1}(t) \Delta_{i-1}(t)}}{\Delta_i(t)} \sigma_i(0),$$

$i = 1, \dots, n-1$, $\Delta_0 = 1$. Here $\Delta_i(t)$ is the i -th principal minor of the matrix $\Gamma(t) = \exp((O(0) + O(0)^T)t)$.

Proof:

We know [7] that $O(t) = R(t)Q(0)R(t)^{-1}$, where $\exp(O(0)t) = Q(t)R(t)$, $Q(t)$ is orthogonal, and $R(t)$ is an upper triangular matrix with positive entries on the main diagonal. We then clearly have

$$\sigma_i(t) = \frac{r_{i+1, i+1}(t)}{r_{i, i}(t)} \sigma_i(0), \quad (3.1)$$

$i = 1, \dots, n-1$. Here $R(t) = \| r_{ij}(t) \|$. The operator $\wedge^i R(t)$ naturally acts on the i -th exterior power $\wedge^i R^n$ by the following rule: $\wedge^i R(t)(v_1 \wedge \dots \wedge v_i) = R(t)v_1 \wedge \dots \wedge R(t)v_i$ for any $v_1, \dots, v_i \in R^n$. We have, further, the following relations:

$$\begin{aligned} r_{11}^2(t) &= \langle R(t)e_1, R(t)e_1 \rangle = \\ &= \langle \exp(O(0)t)e_1, \exp(O(0)t)e_1 \rangle = \\ &= \langle e_1, \Gamma(t)e_1 \rangle = \Delta_1(t). \end{aligned}$$

And more generally

$$\begin{aligned} r_{11}^2(t) \dots r_{ii}^2(t) &= \\ &= \langle e_1 \wedge \dots \wedge e_i, \wedge^i \Gamma(t)(e_1 \wedge \dots \wedge e_i) \rangle = \Delta_i(t), \end{aligned} \quad (3.2)$$

$i = 1, \dots, n$. By (3.2) we easily obtain

$$\frac{r_{i+1, i+1}(t)}{r_{i, i}(t)} = \frac{\sqrt{\Delta_{i+1}(t) \Delta_{i-1}(t)}}{\Delta_i(t)}.$$

The result now follows by (3.1). ■

We have the following differential equations for σ_i :

$$\dot{\sigma}_i = \sigma_i(o_{i+1,i+1} - o_{i,i}),$$

$i = 1, \dots, n-1$. Recalling that $o_{i,i} = -\nu_{i-1}\nu_i$, $i = 1, \dots, n$, $\nu_n = \pm 1$, and $\nu_i^2 + \sigma_i^2 = 1$, we obtain $\dot{\nu}_i\nu_i + \dot{\sigma}_i\sigma_i = 0$ or

$$\dot{\sigma}_i = \sigma_i\nu_i(\nu_{i-1} - \nu_{i+1}),$$

$$\dot{\nu}_i = -\sigma_i^2(\nu_{i-1} - \nu_{i+1}),$$

$i = 1, \dots, n-1$. It is interesting to find how moments $\tau_i(t) = \langle e_1, O(t)^i e_1 \rangle$ evolve under the Toda flow. Consider the family of rational functions

$$f_i(z) = \langle e_1, [zE - O(t)]^{-1} e_1 \rangle.$$

We clearly have

$$f_i(z) = \sum_{i=0}^{\infty} \frac{\tau_i(t)}{z^{i+1}}.$$

On the other hand,

$$\begin{aligned} f_i(z) &= \langle Q(t)e_1, [zE - O(0)]^{-1} Q(t)e_1 \rangle = \\ &= \frac{\langle \exp(O(0)t)e_1, [zE - O(0)]^{-1} \exp(O(0)t)e_1 \rangle}{\langle \exp(O(0)t)e_1, \exp(O(0)t)e_1 \rangle} = \\ &= \sum_{i=0}^{\infty} \frac{h_i(t)}{z^{i+1} h_0(t)}. \end{aligned}$$

Here

$$h_i(t) = \langle \exp(O(0)t)e_1, O(0)^i \exp(O(0)t)e_1 \rangle.$$

Thus $\tau_i(t) = \tau_{-i}(t) = h_i(t)/h_0(t)$, $i \geq 0$. We clearly have

$$\dot{h}_i(t) = h_{i+1}(t) + h_{i-1}(t).$$

Thus,

$$\dot{\tau}_i = \tau_{i+1} + \tau_{i-1} - 2\tau_i\tau_1,$$

$i = 0, 1, \dots$.

References

- [1] G.S. Ammar, W.B. Gragg and L. Reichel, *On the Eigenproblem for Orthogonal Matrices*, Proceedings of the 25th Conference on Decision and Control, IEEE, New York, 1063-1966 (1986).
- [2] G.S. Ammar, W.B. Gragg and L. Reichel, *Determination of Pisarenko frequency estimates as eigenvalues of an orthogonal matrix*, in Advanced Algorithms and Architectures for Signal Processing II (F.T. Luk, ed.), Proc. SPIE vol. 826, 143-145 (1987).
- [3] G.S. Ammar and W.B. Gragg, *Schur Flows*, preprint.
- [4] A. Bunse-Gerstner and L. Elsner, *Schur parameter pencils for the solution of the unitary eigenproblem*, Lin. Alg. Appl., vol. 154-156, 741-778 (1991).
- [5] M.Chu, *On the global convergence of the Toda lattice for real normal matrices and its application to the eigenvalue problem*, SIAM J. Math. Anal., vol. 15, 98-104 (1984).
- [6] P.G. Eberlein and C.P. Huang, *Global convergence of the QR-algorithm for unitary matrices with some results for normal matrices*, SIAM J. Numer. Anal., vol. 12, 97-104 (1975).
- [7] L.E.Faybusovich, *QR-algorithm and generalized Toda flows*, Ukrainian. Mat. J., vol. 45, 944-952 (1989).
- [8] W.B. Gragg, *Positive definite Toeplitz matrices, the Arnoldi process for isometric operators, and Gaussian quadrature on the unit circle (in Russian)*, in Numerical Methods in Linear Algebra (E.S. Nikolaev, ed.), Moscow Univ. Press, 1-8 (1986).

- [9] W.B.Gragg, *The QR-algorithm for unitary Hessenberg matrices*, J. Comput. Appl. Math., vol. 16, 1-8 (1986).
- [10] W.B. Gragg and L. Reichel, *A divide and conquer method for unitary and orthogonal eigenproblems*, Numer. Math., vol. 57, 695-718 (1990).
- [11] H.J. Landau, *Maximum entropy and the moment problem*, Bull. Amer. Math. Soc. vol. 16, 47-77 (1987).
- [12] V.F. Pisarenko, *The retrieval of harmonics from a covariance function*, Geophys. J. R. Astr. Soc., vol. 33, 347-366 (1973).

DISTRIBUTION LIST

Director (2)
 Defense Tech Information Center
 Cameron Station
 Alexandria, VA 22314

Research Office (1)
 Code 81
 Naval Postgraduate School
 Monterey, CA 93943

Library (2)
 Code 52
 Naval Postgraduate School
 Monterey, CA 93943

Professor Richard Franke (1)
 Department of Mathematics
 Naval Postgraduate School
 Monterey, CA 93943

Dr. Neil L. Gerr (1)
 Mathematical Sciences Division
 Office of Naval Research
 800 North Quincy Street
 Arlington, VA 22217-5000

Dr. Richard Lau (1)
 Mathematical Sciences Division
 Office of Naval Research
 800 North Quincy Street
 Arlington, VA 22217-5000

Harper Whitehouse (Code 743) (1)
 NCCOSC RDT&E Division
 271 Catalina Blvd.
 San Diego, CA 92152-5000

Keith Bromley (Code 7601) (1) DTIC QUALITY INSPECTED 5
 NCCOSC RDT&E Division
 271 Catalina Blvd.
 San Diego, CA 92152-5000

John Rockway (Code 804) (1)
 NCCOSC RDT&E Division
 271 Catalina Blvd.
 San Diego, CA 92152-5000

Professor William B. Gragg (15)
 Department of Mathematics
 Naval Postgraduate School
 Monterey, CA 93943

Accession For	
NTIS GRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By _____	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-1	