


Results of the Field Study of the  
Surficial Geology and Paleontologic Resources  
of the Pinon Canyon Maneuver Site,  
Las Animas County, Colorado



19990304 027

by  
Dr. Emmett Evanoff  
University of Colorado Museum  
Boulder, Colorado

Prepared for and funded by  
Department of the Army, Fort Carson Command,  
The Directorate of Environmental Compliance and Management,  
Fort Carson, Colorado



Report submitted to  
Midwest Archeological Center  
National Park Service  
Lincoln, Nebraska



June 1998

**DISTRIBUTION STATEMENT A**

**Approved for public release;  
Distribution Unlimited**

**RESULTS OF THE FIELD STUDY OF THE  
SURFICIAL GEOLOGY AND PALEONTOLOGIC RESOURCES OF  
THE  
PINON CANYON MANEUVER SITE, LAS ANIMAS COUNTY,  
COLORADO**

By

**Dr. Emmett Evanoff**  
Principal Investigator

With Help From:

**Mr. Benjamin Burger**  
**Mr. Evan Hall**  
**Mr. William Hughes**  
**Dr. Alan Lester**  
**Ms Melissa Lester**  
**Mr. Paul Murphey**  
**Ms Sandra Swift**  
**Ms Kathryn Townsend**

University of Colorado Museum  
Boulder, Colorado

Prepared for and funded by  
Department of the Army  
Fort Carson Command  
Directorate of Environmental Compliance and Management  
Fort Carson, Colorado

Report Submitted to  
Midwest Archeological Center  
National Park Service  
Lincoln, Nebraska

June 1998

REPORT DOCUMENTATION PAGE			Form Approved OMB No. 0704-0188	
<small>Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.</small>				
1. AGENCY USE ONLY (Leave blank)	2. REPORT DATE June 1998	3. REPORT TYPE AND DATES COVERED Final Report		
4. TITLE AND SUBTITLE Results of the Field Study of the Surficial Geology and Paleontologic Resources of the Pinon Canyon Maneuver Site, Las Animas County, Colorado			5. FUNDING NUMBERS	
6. AUTHOR(S) Emmett Evanoff				
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) University of Colorado Museum Campus Box 315 Boulder, CO 80309-0315			8. PERFORMING ORGANIZATION REPORT NUMBER  N/A	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) Department of the Army, Fort Carson Command Directorate of Environmental Compliance and Management Fort Carson, CO 80913-5000			10. SPONSORING/MONITORING AGENCY REPORT NUMBER  N/A	
11. SUPPLEMENTARY NOTES Prepared in cooperation with Midwest Archaeology Center, National Park Service, Federal Building, Room 474, 100 Centennial Mall North, Lincoln, Nebraska 685-3873				
12a. DISTRIBUTION AVAILABILITY STATEMENT  Availability Unlimited			12b. DISTRIBUTION CODE	
13. ABSTRACT (Maximum 200 words) The Pinon Canyon Maneuver Site contains a large number of fossils and fossil localities, ranging from dinosaur and plant beds to shell beds that were derived in an ancient sea. The lower sequences of sedimentary rocks that are exposed in the canyons along the Purgatoire River were deposited in wind, river, lake and shoreline environments. The upper sequence was deposited in a shallow seaway, the Western Interior Sea. The fossils of these marine rocks include clams, snails, and ammonoids that lived in the Gulf Coast region and in other continents. Pinon Canyon is one of the few places in the Western Interior Sea that these species of geographically widespread animals lived. The fossils of the lower canyons include fossil logs that accumulated as log jams the bottom of deep valleys. Nowhere else in the western United States are logs of this age known. Dinosaur bones, dinosaur stomach stones (gastroliths), and plant fossils occur in the Morrison Formation. The upper Dakota Group at the rims of the canyons contain abundant plant fossils, including some of the earliest fossils of flowering plants in the region.				
14. SUBJECT TERMS Paleontologic resources; dinosaur fossils, fossil logs, plant fossils, marine invertebrate fossils.			15. NUMBER OF PAGES 35	
			16. PRICE CODE	
17. SECURITY CLASSIFICATION OF REPORT Unclassified	18. SECURITY CLASSIFICATION OF THIS PAGE Unclassified	19. SECURITY CLASSIFICATION OF ABSTRACT Unclassified	20. LIMITATION OF ABSTRACT Unlimited	

## PREFACE

The paleontologic survey reported in this manuscript is an important part of the Fort Carson Cultural Resources Management Program whose goal is to maintain the largest possible area for military training while protecting significant cultural and environmental resources. The current study is a demonstration project funded by a grant awarded by the Legacy Resource Management Program. Legacy was established by Congress in 1991 to provide the Department of Defense (DoD) with an opportunity to enhance the management of resources on lands under DoD jurisdiction.

The Directorate of Environmental Compliance and Management (DECAM) is tasked with maintaining Fort Carson's compliance with federal, state and local environmental laws and mandates. The DECAM holistic management philosophy considers that all resources are interrelated such that decisions affecting one resource will impact other resources. The decisions we make today will affect the condition of Department of Army lands and resources for future training, research and recreation. Mission requirements, training resources, wildlife, range, soil, hydrology, air, and recreation considerations all influence management decisions. Integrating compliance and resource protection concerns into a comprehensive planning process reduces the time and effort expended on the compliance process, minimizes conflicts between resource protection and use, allows flexibility in project design, minimizes costs, and maximizes resource protection.

Federal laws protect the resources on the Pinon Canyon Maneuver Site and Fort Carson; theft and vandalism are federal crimes. Protective measures ensure that Army activity does not inadvertently impact significant cultural and paleontologic sites. Fort Carson does not give out site location information nor are sites developed for public visitation. Similar resources are located in the Picketwire Canyonlands where public visits can be arranged through the U.S. Forest Service, Comanche National Grasslands in La Junta, Colorado.

Fort Carson endeavors to make results of the resource investigations available to the public and scientific communities. Technical reports on cultural resources are on file at the Fort Carson Curation Facility and Colorado State Historic Preservation Office and are available through the National Technical Information Service, Springfield, VA. Selected reports have been distributed to public libraries in Colorado. Three video programs produced by Fort Carson are periodically shown on Public Broadcasting Stations. Fort Carson continues to demonstrate that military training and resource protection are mutually compatible goals.

Stephen A. Chomko  
Cultural Resources Manager  
Directorate of Environmental Compliance  
and Management  
Fort Carson, Colorado

April 1998

## Acknowledgments

The field crew for this project included Benjamin Burger, Melissa Burke, Evan Hall, William Hughes, Alan Lester, Paul Murphey, Sandra Swift, and Kathryn Townsend. The competence and enthusiasm of the field crew was of great help and is greatly appreciated. Unpublished data on the thicknesses of the stratigraphic units in the canyons was provided by Dr. Fred Peterson of the U.S. Geological Survey. Mr. Paul Murphey drafted the final copies of the geologic maps on a map base of 1:24,000 scale.

Anne Vawser of the National Park Service has been the principal government contact for this project. Government contacts while in the field included Stephen Chomko from Fort Carson, and Dan Crouch and Melissa Connor from the National Park Service. I extend my appreciation to these personnel and others at the PCMS and the Midwest Archeological Center, Lincoln, Nebraska, for allowing this study to be made.

Funding for this project was provided by the Department of Defense Legacy Resource Management Program (Contract Number 1443PX6115970067).

Cover illustration is of the skull of *Diplodocus* from Ostrom and McIntosh (1966).

## Technical Abstract

The sedimentary rock sequence in the canyons of the the Pinon Canyon Maneuver Site is composed of nine stratigraphic units, two of which have high paleontologic significance. These two important units are the Bell Ranch and Morrison formations. The Bell Ranch Formation is a series of multicolored breccias, sandstones and siltstones that were deposited in a series of valleys cut into the underlying dune sands of the Entrada Formation. The Bell Ranch contains local accumulations of logs near the base of the deepest paleovalleys and these logs are unique for the Middle Jurassic record of the western United States. The name Bell Ranch for these rocks is an extension of the nomenclature from northeast New Mexico. The Morrison Formation contains a three part sequence including a lower gypsum and gray mudrock interval, a middle limestone and claystone interval, and an upper multicolored mudstone and sandstone interval. The lower interval is unfossiliferous, but the middle interval contains plants, invertebrates (ostracodes) and includes the dinosaur tracks of the adjacent Picketwire Track Site. The upper interval contains scattered accumulations of dinosaur bones, mostly of sauropods. Two localities, both in Welsh Canyon, have large accumulations of bones. Fossils occur in the Cretaceous Dakota Group that caps the rims of the canyons, but none of the fossils are unique for the region. The most fossiliferous unit in the Dakota is the Pajarito Formation, a sequence of alternating sandstones and carbonaceous shales with locally abundant plant fossils. One locality of dinosaur tracks occurs in the upper transition units of the uppermost Dakota. The names Pajarito and Mesa Rica Sandstone are extended to the rocks of the upper Dakota from equivalent units in New Mexico. The oldest two units in the canyons, the upper Dockum Group and the Entrada Sandstone, are unfossiliferous. The Entrada Sandstone is very thick in the canyons, contrary to what has been reported for the area.

Of the 12 fossil localities documented in this study, only four are considered to be of high paleontologic significance. These include the two petrified wood localities (96 Pal 3 and Jb OVM 1) in the Bell Ranch, and the two dinosaur bone localities (Jm OVM 1 and Jm OVM 4) in the Morrison of Welsh Canyon. These localities and the other dinosaur bone localities should be periodically monitored for new fossil materials and to determine the impact of human activities. Any plans for future construction in the area should consider any of the documented fossil localities of this study and of Kauffman (1986). Potential research projects derived from this study include 1) a study of the distribution and origin of the Bell Ranch Formation and its flora, 2) a detailed study of the sedimentology and stratigraphy of the Morrison Formation, 3) a taxonomic and paleoecologic study of the fossil plants of the Pajarito flora, and 4) additional field mapping of the Cretaceous marine rocks.

## Popular Abstract

The Pinon Canyon Maneuver Site contains a large number of fossils and fossil localities, ranging from dinosaur and plant beds to shell beds that were derived in an ancient sea. The lower sequence of sedimentary rocks that are exposed in the canyons along the Purgatoire River were deposited in wind, river, lake, and shoreline environments. The upper sequence was deposited in a shallow seaway, the Western Interior Sea. The fossils of these marine rocks include clams, snails, and ammonoids (extinct squid-like animals with hard external shells) that lived in the Gulf Coast region and in other continents such as Africa and Europe. Pinon Canyon is one of the few places in the Western Interior Seaway that these species of geographically widespread animals lived. The fossils of the lower canyons include fossil logs that accumulated as log jams at the base of deep valleys. Nowhere else in the western United States are logs of this age known. The most important unit for dinosaurs in the area is the Morrison Formation, for it contains dinosaur bones and dinosaur stomach stones (gastroliths). The Picketwire Track Site just to the east of the Pinon Canyon Maneuver Site is within the Morrison Formation. Plant fossils also occur in the Morrison, but plant fossils are more abundant in the uppermost rocks that support the rims of the canyons (the upper Dakota Group). These plants include some of the earliest fossils of flowering plants known from the region. At least one dinosaur trackway occurs in the rocks just above those containing the plants, near the top of the Dakota Group.

## Table Of Contents

Preface	ii
Acknowledgments	iii
Technical Abstract	iv
Popular Abstract	v
Methods	1
Stratigraphy	3
Paleontologic Resources	14
Stratigraphic Units with High Paleontologic Significance	16
Stratigraphic Units with Moderate Paleontologic Significance	20
Stratigraphic Units with Little or No Paleontologic Significance	22
Management Recommendations	23
Suggestions For Future Research	23
Comments On The Present Status Of Geologic Mapping	24
References Cited	26

## List of Tables

Table 1. Lithostratigraphic units in the Pinon Canyon Maneuver Site	5
Table 2. Fossils known from the stratigraphic units exposed at the Pinon Canyon Maneuver Site	15
Table 3. Fossil plants from the Pajarito Formation at Pinon Canyon Maneuver Site and adjacent areas	21

## List of Figures

Figure 1. Map of the eastern half of the Pinon Canyon Maneuver Site	2
Figure 2. Stratigraphic column of the sedimentary rock units exposed in the Pinon Canyon Maneuver Site	4
Figure 3. A partial history of stratigraphic nomenclature of the sedimentary rocks in the Pinon Canyon Maneuver Site	8
Figure 4. Stratigraphic column of the sedimentary rock units exposed in the canyons of the eastern part of Pinon Canyon Maneuver Site	9
Figure 5. View of the upper Dockum Group, Entrada Sandstone, and Bell Ranch Formation in the lower part of Bravo Canyon	10

Figure 6. Large massive cliffs of the Entrada Sandstone at the mouth of Red Rock Canyon _____	10
Figure 7. View of the Bell Ranch conglomerates overlying bleached sandstones of the Entrada Sandstone in Welsh Canyon _____	11
Figure 8. Alternating limestone and claystone beds of the middle Morrison Formation near the mouth of Taylor Arroyo _____	12
Figure 9. Variegated mudstones and a large ribbon sandstone in the upper Morrison Formation in Minnie Canyon _____	12
Figure 10. A history of the stratigraphic nomenclature of the Dakota Group in southeast Colorado and northeast New Mexico _____	13
Figure 11. The Dakota Group exposed in the north cliffs at the mouth of Taylor Arroyo _____	13
Figure 12. Fossil logs from the Lost Korgel Forest locality _____	16
Figure 13. Gastroliths from the mudstones of the upper Morrison Formation _____	17
Figure 14. Large dinosaur bone fragments from the upper Morrison at Jm OVM 5 _____	18
Figure 15. A cross-section view of a large fragment of a dinosaur bone from 96 Pal 1 _____	18
Figure 16. Dinosaur fossils from Jm OVM 4 _____	19
Figure 17. A diagram of the skull of <i>Diplodocus</i> _____	19
Figure 18. Overlapping shells of the inoceramid clam <i>Mytiloides</i> from the Bridge Creek Limestone Member of the Niobrara Formation _____	20
Figure 19. A fragment of fossil wood from the upper transition units of the Dakota Formation _____	21
Figure 20. The shells of freshwater clams and snails that occur in the alluvium of Taylor Arroyo _____	22
Figure 21. Explanation of symbols for the geologic maps and the stratigraphic columns of the Pinon Canyon Maneuver Site _____	25

# RESULTS OF THE FIELD STUDY OF THE SURFICIAL GEOLOGY AND PALEONTOLOGIC RESOURCES OF THE PINON CANYON MANEUVER SITE, LAS ANIMAS COUNTY, COLORADO

Emmett Evanoff

The Pinon Canyon Maneuver Site (PCMS) in Las Animas County, Colorado (Figure 1) contains a 400 m sequence of sedimentary rocks, some of which contain abundant fossils. These fossils include a wide variety of vertebrate, invertebrate, plant, and trace fossils. The purpose of this report is to: 1) document the known occurrences of fossils in the lower portion of these rocks as derived from field studies and 2) to amend the surficial geology and paleontologic resources maps of the PCMS (Evanoff 1997). This work represents the results of a field evaluation of the geology and fossil resources of a part of the area not examined extensively by a previous paleontological survey (Kauffman 1986). Specifically, the rocks examined were those exposed in the canyons in the eastern part of the area. This work will help land managers of the PCMS to manage the paleontologic resources.


Previous geologic maps of part or all of the PCMS include those of Stose (1912), Duce (1924), Gould (1934), Bass et al. (1947), Maher and Collins (1953), Scott (1968), Johnson (1969), Taylor (1974), and Kauffman (1986). The stratigraphy of the rocks have been documented by Lee (1901), Stose (1912), Duce (1924), Keyes (1940), Maher (1946), Oriel and Mudge (1956), Long (1966), Cobban (1968), Cobban and Scott (1972), Taylor (1974, 1980), Kauffman (1986), and Mateer (1987). The paleontology of the PCMS has been documented by Cobban (1971), Cobban and Scott (1972), Taylor (1974), and Kauffman (1986). Geomorphological and geoarcheological investigations of the area have been reported by Schuldenrein (1985), and the geoarcheology of the PCMS has been reported by McFaul and Reider (1990).

## Methods

This project included examining the outcrops of the eastern canyons to determine the paleontologic resources and to check the validity of the previous mapping provided by Evanoff (1997). Rocks of all the major canyons on the eastern part of the PCMS were examined including Taylor Arroyo, Spring Canyon, Lockwood Arroyo, Red Rock Canyon, Welsh Canyon, Bravo Canyon, Bent Canyon, Sugarloaf Canyon, Horse Canyon, Stage Canyon, Iron Canyon, Minnie Canyon, and all the smaller unnamed canyons adjacent to and within these larger canyons (Figure 1). The rocks examined included the oldest rock units in the area, determined to be of the Triassic upper Dockum Group, to the uppermost sandstones of the Cretaceous Dakota Group. The younger marine rocks of the central and western parts of the area were not examined, except to clarify the stratigraphy

---

Emmett Evanoff, University of Colorado Museum, Campus Box 315, Boulder, CO, 80309-0315



**Figure 1. Map of the eastern half of the Pinon Canyon Maneuver Site showing the distribution of canyons and arroyos west of the Purgatoire River examined during this project. UTM Zone 13; Public Land Survey 6<sup>th</sup> Principal Meridian.**

(such as the presence of the Smoky Hill Member of the Niobrara Formation) and relocate problematic localities given by Kauffman (such as Kauffman's Greenhorn locality #22). Kauffman's (1986) paleontological survey was quite comprehensive concerning the marine faunas in the rocks above the Dakota Group, but questions remained concerning the distribution and amount of fossils in the lower, primarily terrestrial, sequence.


The fieldwork that is compiled in this report was performed on 30 May to 12 June and 2-9 July 1997. The field crew divided into groups of two to three workers that hiked the canyons examining the rock outcrops for fossils and making notes on the distribution of rock exposures and fossil localities. Small representative collections from the important localities were made for helping future researchers recognize the fossil resources. Subsequent work in the office has included sorting and labeling the collections, revising the maps using the field data to more accurately represent the geology of the eastern canyons, and to plot the new fossil localities on the maps to add to the previous paleontologic resources data base. The revised mapping includes parts of the following 7½ quadrangle maps: Beaty Canyon (1972, rev. 1984), Doss Canyon North (1972, rev. 1984), Johnson Canyon (1972, rev. 1984), OV Mesa (1993), Packers Gap (1993), Riley Canyon (1993), Rock Crossing (1993), Sheep Canyon (1993), and Stage Canyon (1993).

### Stratigraphy

Nine major Mesozoic sedimentary units, one Tertiary shallow intrusive unit, and five mappable Quaternary deposits occur in the PCMS (Figure 2). These units are described in Table 1. The nomenclature used is a composite of those used by Scott (1968), Johnson (1969), Taylor (1974), and Kauffman (1986) with a few new additions. A history of the nomenclature is shown in Figure 3. The current study has emphasized the geology and paleontology of the eastern canyons of the PCMS and the detailed stratigraphy of these canyons is presented in Figure 4.

The oldest rocks exposed in the PCMS are those of the upper part of the Triassic Dockum Group. These fine-grained, horizontally bedded red sandstones occur in the lower part of Bravo Canyon (Figure 5) in the center of the E½, section 34, T. 28 S., R. 56 W. The Entrada Sandstone is much thicker than what has been previously stated in the literature (88 m thick, relative to the 4 m thick reported in Evanoff 1997) and supports massive cliffs in the lower canyons (Figure 6). Previous workers confused most of the Entrada Sandstone with sandstones of the upper Dockum Group, a problem first recognized by Fred Peterson (written communication, May, 1997). The Entrada is truncated by a paleovalley system that was filled with breccia and sandstones of the Bell Ranch Formation (Figure 7). The Bell Ranch is a Middle Jurassic unit that occurs between the Entrada and Morrison formations in northeast New Mexico (Griggs and Read 1959; Lucas, Hunt and Kues 1987). Previous workers have placed these sandstones and the overlying gypsum beds into the Ralston Creek Formation (see Taylor 1972), but recent regional stratigraphic studies indicates that the Ralston Creek to be the lower beds of the Morrison Formation, and the underlying variegated sandstones and mudstones are separated from the gypsum beds by a major unconformity (Fred Peterson, written communication, May 1997). The name Ralston Creek Formation is not used herein for

## Stratigraphy of the Pinon Canyon Maneuver Site



**Figure 2. Stratigraphic column of the sedimentary rock units exposed in the Pinon Canyon Maneuver Site. Thickness are from Kauffman (1986) and Peterson (written communication, May 1997).**

**TABLE 1. LITHOSTRATIGRAPHIC UNITS IN THE PINON CANYON MANEUVER SITE.**

Qa	<b>Modern alluvium.</b> Holocene. Primarily clayey silt and sand along modern drainages. Can locally contain shells of freshwater clams and snails and land snails. Low to moderate fossil potential.
Qls	<b>Landslide deposits.</b> Upper Quaternary. Poorly sorted deposits in debris slides and earthflows, primarily below Dakota Sandstone and along terrace scarps. Low fossil potential.
Qsd	<b>Sand dunes.</b> Upper Quaternary. Yellowish gray sand occurring in longitudinal dunes on the western slopes of the area. Low fossil potential.
Qog	<b>Older gravels.</b> Lower Quaternary and Pliocene? Gravels on high terrace near the mouth of Taylor Arroyo, containing gneiss, chert, and basalt cobbles. Low fossil potential.
Qu	<b>Upper Quaternary Sediments Undifferentiated.</b> Deposits that include older alluvium, slopewash and colluvium in valley bottoms and on slopes. Includes the "Pitty gravels" of Helgren et al. (1985) on divides above the canyons of the Purgatoire River. Low to moderate fossil potential.
Ti	<b>Intrusive rocks.</b> Oligocene. Dark-gray finely crystalline dikes of olivine basalt that supports ridges in the southwest margin of the area. No fossil potential.
	<b>Niobrara Formation.</b> Upper Cretaceous. Abundant light colored limestones with interbedded shales. Moderate fossil potential; abundant invertebrate fossils and rare vertebrate fossils. 225 m, 737 ft. thick.
Kns	<b>Smoky Hill Member.</b> Calcareous shale and ledge forming limestone with interbedded chalk. Lower portion includes abundant limestone beds; while upper portion is dark calcareous shales. Limited outcrops occur by the western-most boundary of the area.
Knf	<b>Fort Hays Member.</b> Fine grained micritic limestone separated by thin beds of soft calcareous shale. Limestone is olive-gray to white and weathers to buff. Shales are well laminated, olive-gray and weathers to yellowish-gray. 10 m, 33 ft, thick.
Kc	<b>Carlile Shale.</b> Upper Cretaceous. Darker colored sands and shales with lenses of calcarenite. Moderate fossil potential; abundant invertebrate fossils and rare vertebrate fossils. 46 m, 152 ft, thick.
	<b>Juana Lopez Member.</b> Lenses of dark brownish-gray fine-grained well laminated calcarenite containing patches of light gray glauconite. Contains abundant marine fossils. 3 m, 11 ft, thick
	<b>Blue Hill Shale Member.</b> Dark-gray soft to well indurated well laminated shale. Upper half is sandy and contains two prominent layers of septarian limestone concretions. 28 m, 91 ft, thick.
	<b>Fairport Chalky Member.</b> Dark olive-gray to black, well laminated soft calcareous shale. Yellowish-gray limestone near base. 16 m, 51 ft, thick.
Kgh	<b>Greenhorn Formation.</b> Upper Cretaceous. Calcarenite limestone beds with dark gray platy to fissile shale beds. Moderate fossil potential with abundant invertebrate fossils and rare vertebrate fossils. 37 m, 121 ft. thick.
	<b>Bridge Creek Limestone Member.</b> Gray, shaly weathering micritic limestone beds separated by soft calcareous shale and bentonite layers. 13 m, 43 ft, thick.

**Hartland Shale Member.** Dark gray calcareous platy shale and thin layers of calcarenite composed of tests of foraminifera and prisms of *Inoceramus* shells. 18 m, 60 ft, thick.

**Lincoln Limestone Member.** Layers of dark gray, platy, shaly calcarenite composed of tests of foraminifera and prisms of *Inoceramus* shells; separated by dark gray soft calcareous shale. 5.6 m, 18 ft, thick

**Kg Graneros Shale.** Upper Cretaceous. Dark gray fissile, soft to medium-hard, non-calcareous shale and numerous bentonitic beds. Ferruginous siltstone contains fish scale and bones.. Moderate fossil potential with abundant invertebrates and rare vertebrates. 30 m, 98 ft, thick.

**Thatcher Limestone Member.** Bench-forming micritic limestone separating the upper and lower black shales. Diverse mollusk fossils. 0.2 m, 0.5 ft, thick.

**Kd Dakota Group.** Lower Cretaceous. Fine to coarse-grained sandstones and conglomerates with interbedded claystone, shales and siltstones. Upper contact is conformable, the base is unconformable. Trace fossils are abundant and plant fossils (wood, stems, and leaves) are locally numerous at the top. Dinosaur tracks are rare, but are present in the upper-most sandstone beds. Can be divided into a lower and upper interval, with the Lytle and Glencairn formations in the lower part, and the Mesa Rica, Pajarito, and upper transition units comprising the upper part. The upper part has a moderate fossil significance for abundant fossil plants and rare dinosaur tracks. 71 m, 231 ft, thick.

**upper transition units.** Platy to massive brown fine-grained sandstones with abundant trace fossils and symmetric ripplemarks. Locally has a thick basal sandstone filling scours in the underlying Pajarito Formation. Contains rare dinosaur tracks. 7m, 23 ft, thick.

**Pajarito Formation.** Alternating flaggy brown to white sandstones and gray to brown muddy shales. The shales are typically carbonaceous and locally have fossil leaves. The sandstones are rippled, highly burrowed and contain numerous wood impressions. The fossil wood is typically burrowed by shipworms (*Teredo*). 4.5 m, 15 ft thick.

**Mesa Rica Sandstone.** A cliff-forming quartz-rich cross-bedded sandstone. The sandstones are medium to fine grained and are white to yellow brown that typically weather rusty brown. Many highlands, such as the Black Hills, are capped by this unit. Fossils are rare in these sandstones. 6 to 9 m, 19 to 30 ft, thick

**Glencairn Formation.** Black to dark gray, fissile basal shale grading upwards into interbedded tan siltstones and sandstones. Contains rare shells of marine clams, abundant trace fossils, and shale beds containing numerous fish scales. 33 to 36 m, 107 to 117 ft, thick.

**Lytle Formation.** Coarse to fine-grained white sandstones with alternating mudstone layers towards the top. Capped by a thin interval of conglomeratic, brown-weathering resistant sandstones, called the Plainview by several researchers. The Lytle supports large cliffs in the area. Lytle fossils are rare in the area and are limited to scattered casts of logs. 17 to 29 m, 57 to 95 ft, thick.

**Jm Morrison Formation.** Upper Jurassic. Consists primarily of mudrocks (claystones and mudstones) alternating with thick gypsum beds, thin limestones and sandstone sheets and ribbons. Colors range from white, light grays, blues and green at the base to reds, purples and yellows at the top. Fossil potential is high. Formation strata have yielded fossil plants, dinosaur tracks and bones from the area. Total thickness: 92 m, 302 ft, thick. The Morrison can be divided into three parts:

**upper Morrison.** Composed predominantly of multicolored mudstones and claystones, with interbedded gray to brown sheet and ribbon sandstones, and thin widespread brown to gray limestones that weather to a rusty brown. Swelling clays (smectites) are abundant in the mudrocks and act as slip surfaces for numerous landslides that typically cover the upper Morrison outcrops. Fossils include dinosaur bones, typically in conglomeratic sandstones and rarely in the mudrocks, and scattered gastroliths. 51 m, 167 ft, thick.

**middle Morrison.** Alternating white micritic limestones and greenish gray claystones. The limestones have algal laminations (stromatolitic) in the lower part. Fossil ostracodes and

silicified plant fragments occur in the upper limestones. The Picketwire track site is in the upper-most limestones of this interval. 16 m, 52 ft, thick.

**lower Morrison.** Alternating thick gypsum beds and gray mudrocks, mostly claystones and mudstones, with thin sandstone partings. The gypsum beds are easily eroded and this interval is poorly exposed in a flat bench on top of the Bell Ranch. The top of this interval is marked by a widespread interval of bluish gray to red welded chert locally forming thick masses. No fossils are known from this interval. 25 m, 83 ft, thick.

- Jb Bell Ranch Formation.** Middle Jurassic. Lower part is composed of multicolored sandstones and breccias filling deep paleovalleys. Colors in the lower part range from dark red-brown, purple, brown-orange to green-gray. Upper part is composed predominantly of red to purple siltstones and sandstones. Large silicified logs, up to 1.45 m (4.8 ft) in diameter, are locally present in the deepest part of the paleovalleys. Rare vertebrates are reported from the upper beds. The unit's fossil significance is high. Thickness varies greatly from near 0 m thick to 14 m, 44 ft, thick.
- Je Entrada Sandstone.** Middle Jurassic. Massive, salmon-red, to white, well sorted, fine-grained sandstone. Highly bedded in large to very large crossbed sets. Upper part has a well-developed weathering zone. The Entrada is a prominent cliff supporter in the lower canyons. No fossils are known from the unit in the area. 88 m, 288 ft thick.
- TRd Dockum Group.** Triassic. Coarse- to very fine-grained red sandstone with thick to thin horizontal bedding. Boulder conglomerate and chert beds near top. Fossilized plant material, invertebrates, and vertebrates known from other areas, but no fossils are known from the area. 28 m, 98 ft thick.
- 

the rocks in the PCMS. The Morrison Formation includes a lower gypsum sequence, a middle limestone and green claystone sequence (Figure 8), and an upper variegated mudstone and sandstone sequence (Figure 9).


The nomenclature of the Dakota Group is problematic (Figure 10). The Dakota can be divided into two major divisions, a lower part composed of the Lytle and Glencairn formations, and an upper part that includes a basal sandstone, a middle shale, and an upper sandstone interval. In the field the two subdivisions are marked by the base of the upper sandstone cliffs in the canyons (Figure 11). Early workers named these divisions the Purgatoire Formation and the Dakota Sandstone. Subsequent workers (Kauffman 1986; Mateer 1987) have included all of these units in the Dakota Group and named the upper division the Muddy Formation, a name given to correlative rocks in north-central Wyoming. Workers of the Cretaceous rocks of northeast New Mexico have named the individual units in the upper Dakota and some of these names have herein been extended into the study area. These include the Mesa Rica Sandstone for the lower sandstone interval, and the Pajarito Formation for the middle shale sequence. The name Romeroville Sandstone has not been extended into the area, for the upper sandstone interval includes two units that may not be equivalent to the Romeroville. The lower and upper parts of the Dakota Group are distinct in the aerial photographs and have been mapped.

The Cretaceous rocks of marine origin overlying the Dakota have been discussed extensively in Kauffman (1986) and are summarized in Table 1. The units mapped in this study do not include the subdivisions of the Cretaceous rocks as indicated by Kauffman (1986), because the members of the Graneros, Greenhorn, and Carlile formations are not


Age	Scott, 1968, & Taylor, 1974	Kauffman et. al., 1986	Evanoff, 1997	This Report	Map Symbol			
Cretaceous	Niobrara Formation	Niobrara Formation	Niobrara Formation	Niobrara Formation	Kns			
						Smoky Hill Member	Smoky Hill Member	
	Fort Hays Member	Fort Hays Member	Fort Hays Member	Fort Hays Member	Knf			
						Carillie Shale	Carillie Shale	
	Greenhorn Limestone	Greenhorn Limestone	Greenhorn Limestone	Greenhorn Limestone	Greenhorn Limestone	Kgh		
							Juana Lopez Member	
							Blue Hill Shale Member	
							Fairport Member	
	Graneros Shale	Graneros Shale	Graneros Shale	Graneros Shale	Graneros Shale	Kg		
							Bridge Creek Ls Memb.	
Dakota Sandstone	Dakota Sandstone	Dakota Group	Dakota Group	Dakota Group	Kdu			
						Dry Creek Canyon Member	upper sandstone	
Purgatoire Formation	Purgatoire Formation	Purgatoire Formation	Purgatoire Formation	Purgatoire Formation	Kdl			
						Glencalrn Shale Member	Glencalrn Formation	
Jurassic	Morrison Formation	Morrison Formation	Morrison Formation	Morrison Formation	Jm			
						Entrada Sandstone	Entrada Sandstone	
	Ralston Creek Formation	Ralston Creek Formation	Ralston Creek Formation	Ralston Creek Formation	Ralston Creek Formation	Jeb		
							Entrada Sandstone	Entrada Sandstone
	Dockum Group	Dockum Group	Dockum Group	Dockum Group	Dockum Group	Jd		
							Big Basin Formation	Big Basin Formation
	Day Creek Dolomite	Day Creek Dolomite	Day Creek Dolomite	Day Creek Dolomite	Day Creek Dolomite	Jeb		
							Whitehorse Sandstone	Whitehorse Sandstone
	Triassic	Triassic	Triassic	Triassic	Triassic	Jeb		
							Entrada Sandstone	Entrada Sandstone
Permian	Permian	Permian	Permian	Permian	Jeb			
						Entrada Sandstone	Entrada Sandstone	

Figure 3. A partial history of the stratigraphic nomenclature of the sedimentary rocks in the Pinon Canyon Maneuver Site.

## Stratigraphy of the Canyons, Pinon Canyon Maneuver Site



**Figure 4. Stratigraphic column showing the sedimentary rock units exposed in the canyons of the eastern part of the Pinon Canyon Maneuver Site. The thicknesses are from Kauffman (1986) and Peterson (written communication, May 1997).**



**Figure 5.** View of the horizontal bedded red sandstones of the upper Dockum Group (TRd) under the rounded massive cliffs of the Entrada Sandstone (Je). The red sandstones at the skyline are sandstones in the Bell Ranch Formation (Jb). The white bench below the Bell Ranch sandstones is the weathering zone on the top of the Entrada Sandstone. The view is to the southeast in the lower part of Bravo Canyon. Photograph provided by Paul Murphey.




**Figure 6.** Large massive cliffs of cross bedded sandstones of the upper Entrada Sandstone. The view is to the northwest near the mouth of Red Rock Canyon.




**Figure 7. View to the northeast of the Bell Ranch conglomerates overlying bleached sandstones of the Entrada Sandstone. The view is a cross section through a paleovalley, with the contact between the two formations extending from the field assistant's hand to the upper right of the photograph. The location is up the valley from the Lost Korgel Forest, (96 Pal 3). Sandra Swift is the field assistant.**

easily mapped off of the aerial photographs. Kauffman (1986) did not recognize the Smoky Hill Member of the Niobrara Formation in the area but these rocks are exposed near the western edge of the PCMS south of the Cantonment (in sections 5, 6, and 19, T. 30 S., R. 60 W.).

The Quaternary deposits of the area have been greatly generalized on the maps. Large areas are covered by a combination of older alluvium, pediment sediments, and colluvium that are not easily distinguished on the aerial photographs or on the ground. These deposits have been mapped as Quaternary undifferentiated sediments (Qu) on the maps. In addition, old lag gravels, informally named the "Pitty gravels" by Helgren et al. (1985) that occur on the tops of divides in the area have been included in these undifferentiated Quaternary sediments. Schuldenrein (1985) has determined that all of the unconsolidated sediments in the PCMS is probably Holocene (late Quaternary) in age, except for the "Pitty gravels" that may be lag gravels derived from the Miocene Ogallala Formation that once covered the region. However, older gravels also occur on a high terrace near the mouth of Taylor Arroyo. These gravels include clasts of gneiss, chert, and basalt, lithologies that are not found in the modern drainage basin of Taylor Arroyo. Quaternary debris derived from the Dakota Group covers much of the exposures of the fossiliferous upper Morrison in the canyons. In some cases these Morrison exposures are covered by Quaternary landslides that moved on the mudrocks with swelling clays in the upper Morrison. Only the largest of these landslides have been mapped.



**Figure 8.** Alternating limestone and claystone beds of the middle Morrison Formation near the mouth of Taylor Canyon at Jm DCN 1. The view is to the east and the field assistant is William Hughes.



**Figure 9.** View to the southwest of the upper Morrison in Minnie Canyon. Included are variegated mudstones and a large sandstone ribbon (channel deposit) on the right. The cliff on the skyline is in the Dakota Group. This outcrop is in the SW $\frac{1}{4}$ , NE $\frac{1}{4}$ , SE $\frac{1}{4}$ , sec. 6, T. 28 S., R. 55 W



## Paleontologic Resources

The paleontologic significance of a stratigraphic unit in any area depends on three criteria. The first criterion is the presence of rare taxa, the second is the diversity of the plant and animal fossils (biota), and the third is the abundance of fossils in a stratigraphic unit. Of these three criteria, the first one is the most important for fossils may be diverse and abundant in a rock unit but if the taxa are widespread and common in other areas then their significance is much reduced. Fossils abound in many of the stratigraphic units in the PCMS, but only a few of the units contain rare or unique taxa. Thus, only a few units are of high significance, though additional study may find additional taxa in the stratigraphic units that are now thought to be less significant paleontologically. Fossils are also typically clumped in their distribution, especially those that were deposited in terrestrial environments, so individual localities become significant or not, depending on the three criteria given above. Localities also become important if they are the source of new taxa - such localities become type localities for the newly described taxa. Finally, the traditional criteria for ranking includes an emphasis for vertebrate fossils. Vertebrate fossils tend to be much more rare than invertebrate, plant, or trace fossils, and are considered to be more significant by most researchers. However, the most significant fossils described in this report, the Middle Jurassic logs of the Bell Ranch Formation, reflect their unique occurrence in the rock sequence of the western United States.

Invertebrate, plant, and trace fossils are all locally very abundant in the PCMS (Table 2). Vertebrate fossils are less common, but are important constituents of only one stratigraphic unit, the upper Morrison Formation. Units that have produced rare invertebrates or include type localities for fossil invertebrate species include the Thatcher Limestone Member of the Graneros Shale, the limestone units of the Greenhorn Formation, and the Fort Hays Member of the Niobrara Formation. The individual localities for these significant occurrences have been described by Kauffman (1986, p. 119-121). Trace fossils (the tracks, trails, and burrows of organisms) are very abundant in all the marine units (Dakota through the Niobrara formations), but in most cases these are common fossils. Vertebrate tracks are rare in the PCMS, though one locality in the upper transition units of the Dakota (Locality Kd SC 1) contains a poorly preserved track way. The highly significant Picketwire Track Site occurs adjacent to the PCMS in the upper limestone beds of the middle Morrison Formation. However, large flat outcrops exposing large areas of the tops of these limestones do not occur in the Maneuver PCMS. Plant fossils abound in the Pajarito Shale, but this is characteristic of the Pajarito in New Mexico (Kues and Lucas 1987). The fossil logs at the base of the Bell Ranch Formation are the most significant fossils described in this study, for no other fossilized wood is known from this interval in the western United States. The other stratigraphic units in the area have few or no fossils and are of low paleontologic significance. The following discussion will discuss the stratigraphic units of high, moderate, and low paleontologic significance.

Table 2. Fossils Known from the Stratigraphic Units Exposed in and Adjacent to the Pinon Canyon Maneuver Site.

		Quaternary Alluvium	Smoky Hill Member	Fort Hays Member	Carlile Shale	Greenhorn Formation	Graneros Shale	upper transitional units	Pajarito Formation	Mesa Rica Sandstone	Glencalrn Formation	Lytle Formation	upper	middle	lower	Bell Ranch Formation	Entrada Sandstone	upper Dockum Group	
		Niobrara						Dakota Group					Morrison						
Marine	Marine Reptiles					X													
	Marine Fish		X	X		X													
	Shark Teeth		X	X	X	X													
	Ammonites	X	X	X	X	X													
	Oysters	X	X	X	X	X													
	Marine Clams	X	X	X	X	X				X									
	Marine Snails		X	X	X	X													
	Foraminifers	X	X	X	X	X	X			X									
	Trace Fossils	X	X	X	X	X	X	X	X	X	X	X	X	X	X				
	Terrestrial	Wood						X	X	X		X	X	X	X		X		
Leaves								X				X							
Stromatolites														X					
Ostracodes														X					
Dinosaur Tracks							X							X					
Dinosaurs													X						
Terrestrial Reptiles																X			
Freshwater Clams		X													x				
Freshwater Snails		X																	
Paleontologic Significance		high		X	X <sup>1</sup>	X <sup>1</sup>	X <sup>1</sup>							X			X		
	medium	X	X				X	X		X				X					
	low			X <sup>2</sup>	X <sup>2</sup>	X <sup>2</sup>			X		X				X	X	X	X	

<sup>1</sup> limestone members    <sup>2</sup> black shales

## Stratigraphic Units with High Paleontologic Significance


The stratigraphic unit with the highest paleontologic significance in the PCMS is the Middle Jurassic Bell Ranch Formation. The Bell Ranch is a sequence of multicolored breccias, sandstones, and siltstones that filled a paleovalley sequence cut into the underlying Entrada Sandstone. Near the base of the deepest valleys, the Bell Ranch contains large fossil logs (Figure 12) as large as 1.45 m in diameter and 2.2 m long. The logs are silicified with no internal structure preserved, but they represent ancient gymnosperm trees, probably aracarians or conifers. In the locality with the best preserved logs (96 Pal 3), the *in situ* logs are all nearly horizontal and are preferentially oriented either parallel or perpendicular to the valley trend. The logs at this locality represent an ancient log jam at the base of a deep and narrow valley. The logs and wood in the Bell Ranch are limited to the bottoms of the deepest paleovalleys and so far only two wood localities are known. However, no other Middle Jurassic logs are known from the Western United States, so the Bell Ranch logs at the PCMS are very significant paleontologically. Bones of a turtle and an unidentified vertebrate have been reported



Figure 12. Fossil logs from the Lost Korgel Forest (96 Pal 3). A) A narrow, long fossil log. The length of the rule is 0.35 m. B) a large log, 1.45 m wide.

from the Bell Ranch by Kauffman (1986) but no bones were found during the present study.

The Morrison Formation is also significant paleontologically. The Morrison is the only unit in the PCMS that has numerous vertebrate fossils as well as scattered plant localities. The vertebrate fossils include bones of dinosaurs and gastroliths, highly polished chert and quartz pebbles (Figure 13) that are thought to have been polished in the crops of dinosaurs. All of the *in situ* vertebrate remains found in this study were from the mudstones and sandstones of the upper Morrison. The bones are typically dark gray (Figure 14) but can range from white to light red in color and some are filled with silica cement (Figure 15). They typically occur in conglomerates associated with lenticular sandstones. These conglomerates contain large bone fragments, mudstone rip-up clasts, and chert gravel. One bone locality (Jm OVM 6) has bones within mudstones. No complete bones were found in the area but large bones of sauropods are the most typical remains. The localities with the most abundant and well preserved bones include Jm OVM 1 and Jm OVM 4. Jm OVM 4 contains the only dinosaur tooth and jaw material known from the area (Figure 16). The jaw and tooth are of a diplodocid sauropod (Figure 17). Gastroliths can occur with the fossil bones or are widely scattered in the mudrocks of the upper Morrison. The middle Morrison limestones contain ostracodes, fragments of plant stems, and stromatolites (thin laminated limestones formed by bacterial and algal mats). The stromatolites are restricted to the lower limestones of the middle Morrison. The ostracodes, plant stems, and the dinosaur trackways of the Picketwire Track Site are all in the upper limestones of the middle Morrison. No fossils are known from the lower Morrison gypsum sequence.



**Figure 13. Highly polished gray chert and yellow quartz pebbles found in the mudstones of the upper Morrison Formation. These are interpreted to be gastroliths, or dinosaur crop stones.**



Figure 14. Large dinosaur bone fragments from the upper Morrison at Jm OVM 5.




Figure 15. A cross-section view of a large fragment of a dinosaur bone from 96 Pal 1. The dark material on the left side is red chert.



Figure 16. Dinosaur fossils from Jm OVM 4. On the left is a small bone in a slab of conglomerate. The center contains a tooth of a diplodocid sauropod. The fossil on the right is a mold of a portion of a sauropod jaw.




Figure 17. A diagram of a skull of *Diplodocus*. The tooth from Jm OVM 4 is short and is probably from the rear of the tooth row. From Ostrom and Mc Intosh (1966).

The limestones of the marine Cretaceous sequence (the Graneros through Niobrara formations) contain a very abundant (Figure 18) and significant molluscan fauna as documented by Kauffman (1986). The PCMS is very important for understanding Cretaceous invertebrate paleobiogeography and biostratigraphy because many marine mollusk taxa that are found in the area also occur in the Gulf Coast and other continents. These occurrences are well documented in Kauffman (1986), and little is needed to update this information. The location of Kauffman's Greenhorn Locality #22, a very significant locality in the Bridge Creek Limestone Member, was problematic, but has been relocated during this study.



Figure 18. Overlapping shells of the inoceramid clam *Mytiloides* from the Bridge Creek Limestone Member of the Greenhorn Formation at Kauffman's Locality #22.

### Stratigraphic Units with Moderate Paleontologic Significance

The Pajarito Formation and the upper transitional units of the upper Dakota Group have abundant plant and trace fossils. Taylor (1974) found identifiable plant fossils from the Pajarito at three localities. These localities and their associated plant taxa are listed in Table 3. Fossil wood is abundant in both Pajarito and the upper transitional units, and this wood typically contains burrows of the shipworm, *Teredo* (Figure 19). Burrows of invertebrates are common in both of these units, and at one locality (Kd SC 1) in the upper transitional units are the undertracks of a bipedal dinosaur. These tracks are represented by alternating staggered small depressions on the top of a thick sandstone slab near the head of Red Rock Canyon. Their indistinct form is a result of the dinosaur walking on a sedimentary surface above the sand bed which now shows the tracks. As the dinosaur walked, it deformed the sediment layers within about a half of a meter of the

Table 3. Fossil Plants from the Pajarito Formation at Pinon Canyon Maneuver Site and Adjacent Areas, from Taylor (1974).

---

*Auracarites* sp. (gymnosperm tree)  
*Sterculia* cf. *S. coriacea* Knowlton (angiosperm)  
*Aralia* sp. (angiosperm)  
*Cyatheites* sp. (tree fern)  
*Onoclea fecunda* (Lesquereux) Knowlton (?) (fern)  
*Sequoia* cf. *S. obovata* Knowlton (gymnosperm)

---




Figure 19. A fragment of fossil wood from the upper transition units of the Dakota Formation on the ridge between Horse and Sugarloaf canyons. The wood has almost been destroyed by the burrows of the shipworm *Teredo*, now represented by the overlapping worm-shaped features.

surface including the sand bed that now preserves the tracks. Therefore, these tracks are called undertracks, and are recognized primarily by their spacing. Plants and traces (including dinosaur tracks) are known from the Pajarito Formation and equivalents of the upper transitional units in New Mexico (Kues and Lucas, 1987). Therefore, the paleontologic significance of these units is considered moderate. Few fossils are known from the lower Dakota formations, but the casts of logs are known from the Lytle Formation and shales containing fish scales are known from the Glencairn Formation (Kauffman 1986). The paleontologic significance of the lower Dakota rocks in the PCMS is lower than that of the fossiliferous upper Dakota rocks, but the the Glencairn Formation can be considered to have moderate paleontologic significance. Finally, the Smoky Hill Member of the Niobrara Formation exposed in the southwest part of the PCMS contains abundant shells of inoceramid bivalves. These shells are common in the

Smoky Hill of other regions and, until additional studies of the Smoky Hill fauna determines otherwise, the unit is considered to have moderate paleontologic significance.

The Quaternary stream alluvium can contain shells of freshwater mollusks. These are most abundant near the mouth of Taylor Arroyo, and include two species of freshwater clams and one species of freshwater snail (Figure 20). These mollusk indicate the presence of past perennial waters. The Quaternary alluvium in the lower part of Taylor Arroyo has a moderate paleontologic significance.




Figure 20. The shells of a) *Sphaerium striatinum*, the striated fingernail clam; b) *Pisidium compressum*, a freshwater seed clam; and c) *Physella virgata*, a freshwater snail that occur in the Quaternary alluvium at the mouth of Taylor Arroyo. The diagrams are from Burch (1975, 1980), and the small bars are divided into millimeters.

#### Stratigraphic Units with Little or No Paleontologic Significance

Few fossils have been found in the black shales of the Graneros, Greenhorn, and Carlile formations, and these have been primarily ammonites associated with carbonate nodules (Kauffman 1986). The Mesa Rica Sandstone of the upper Dakota Group contains few fossils, including only trace fossils and scattered wood fragments (Kauffman 1986). The Lytle Formation contains trace fossils and rare casts of logs (Kauffman 1986). These units have low paleontologic significance.

No fossils are known from the Entrada Sandstone and the upper Dockum rocks in the PCMS. The Quaternary older gravels, eolian sand deposits, and the undifferentiated Quaternary deposits are not known to have fossils. Finally, the shallow intrusive dike of the Hogback was implaced below the ground surface and has no fossils. Fossils can occur in the landslide deposits but these represent a jumbled assemblage derived from the upper Morrison and Dakota Group. All of these rock units have essentially no paleontologic significance.

## Management Recommendations

Of the 12 localities newly documented in this study (not including Kauffman's Locality 22), only four localities are considered to be significant. These localities include the Lost Korgel Forest (96 Pal 3), the Red Rock Wood Locality (Jb OVM 1), Sandy Swift's Bone Bed (Jm OVM 4), and Bill Hughes' Sauropod Locality (Jm OVM 1). The two fossil wood localities in the Bell Ranch Formation are significant because they are the only known localities of Middle Jurassic wood in the western United States. The two dinosaur localities have the greatest accumulations of bones of any Morrison localities in the area. The other eight localities have enough significance to be documented, but are not as important paleontologically.

Several recommendations concerning the management of these fossil localities are as follows. All future construction plans should include a consideration of the possible disturbance to any of the 12 localities, and such disturbance should be minimized. The four significant localities should be periodically examined for any new significant fossil materials that may be exposed and monitored to determine the effects of human disturbance. The other eight localities, especially the dinosaur localities, should be examined only after major human activities have occurred in their vicinity. Future researchers in the area should also be vigilant for new fossil localities and any new localities should be adequately documented. If any significant new fossil materials should be collected, then this material should be prepared and curated into a long-term research collection. No materials should be collected if they cannot be entered into a managed collection.

## Suggestions for Future Research

Four major projects involving future research on the paleontology and geology in the PCMS have become apparent from the current study. The first possible project is to determine the origin and distribution of the Bell Ranch Formation. This would involve determining the relations between the Bell Ranch Formation and the underlying Entrada Sandstone, determine the taxonomic affinities of the fossil wood, and attempt to find additional wood and vertebrate localities. This proposed study would include all the outcrops in Red Rock, Welsh, and Bravo canyons and the southern tributaries to Bent Canyon. This project is very important because of the unique Middle Jurassic fossils of the Bell Ranch.

A detailed study of the stratigraphy and origin of the Morrison Formation in the PCMS would provide an important stratigraphic and sedimentologic framework for the various fossil localities. An especially interesting subproject would be to study the changes in the lacustrine environments of the limestone beds of the middle Morrison. This would also include an attempt to identify the fossil plants in the upper limestones.

A study of the plant fossils of the Pajarito Formation may provide important information concerning the diversity, paleoecology, and evolutionary development of the early angiosperms. A paleobotanist could recollect the localities listed by Taylor (1974) and would have numerous opportunities to find additional localities. A detailed sedimentologic and stratigraphic study of the Pajarito Formation and the upper

transitional units would also provide important information concerning the transition from terrestrial to marine conditions.

The final potential study is to map the various members of the Cretaceous marine rocks above the Dakota Group. This would help clarify the locations of the most fossiliferous outcrops, and provide data on the distribution of structures in the PCMS. Detailed mapping is also needed of the Hogback dike, and the older "Pitty gravels." The end result of this work would be a detailed field-checked geologic map of the PCMS.

### **Comments on the Present Status of Geologic Mapping**

The geologic maps supplied by this study include information derived from the fieldwork. This information includes the distribution of fossil localities not included in the preliminary mapping, and the distribution of stratigraphic units better reflecting the actual stratigraphy of the area. The contacts were plotted on the maps using the aerial photos listed in Evanoff (1997). For the most part, the Entrada and Bell Ranch formations were not separated, for the details of this irregular contact cannot be easily seen on the aerial photographs and would involve more field time than this study allowed. Likewise the formations of the Dakota Group were not separated because the contacts are typically covered by debris. The base of the cliff of the Mesa Rica Sandstone was used to separate the lower and upper Dakota units. The new mapping also shows less Quaternary deposits. The long fingers of Quaternary alluvium in the bases of most of the small canyons shown in the older maps have been eliminated in the new maps, for alluvium is a minor constituent of these valley bottoms. Much of the Dakota and upper Morrison is covered by debris which is not shown on these new maps, except for the distribution of large landslides that were identified in the field or are very obvious in the aerial photographs. The edge of mapping mostly coincides with the contacts of the upper Dakota with the undifferentiated Quaternary deposits as mapped in the previous project (Evanoff, 1997). Figure 21 is an explanation of the current maps and the symbols used in the stratigraphic columns (Figures 2 and 4).

## Geologic Maps Explanation

Quaternary	Qa	Qa - Alluvium		Fault with mark on downdropped block												
	Qsd	Qsd - Sand dune														
	Qls	Qls - Landslide deposits		Contact												
	Qu	Qu - Quaternary deposits undifferentiated														
	Qog	Qog - Older gravels		Strike and dip symbol												
Tl	Tl - Tertiary intrusive dike															
Cretaceous	Kns	Kns - Niobrara Formation, Smoky Hill Member														
	Knf	Knf - Niobrara Formation, Fort Hays Member														
	Kc	Kc - Carlile Shale		Fossil locality												
	Kgh	Kgh - Greenhorn Formation				Dinosaur bones										
	Kg	Kg - Graneros Shale						bone scrap								
	Kdu	Kd - upper Dakota Group								gastroliths						
	Kdl	Kdl - lower Dakota Group										tracks				
	Jm	Jm - Morrison Formation												wood		
Jb		Jb - Bell Ranch Formation														plants
Jbe		Jbe - Entrada Sandstone & Bell Ranch Formation														
Triassic	Td	Td - upper Dockum Group														

## Stratigraphic Column Explanation

	gypsum beds		dinosaur bones		wood & logs																
	chert masses						small bones														
	calcarenite								dinosaur tracks												
	limestone										ostracodes										
	shale												trace fossils								
	claystone														leaves						
	mudstone																stems				
	horizontal bedded sandstone																		paleocurrent directions		
	cross-bedded sandstone																				unconformity
	conglomerate and breccia																				

**Figure 21. Explanation of symbols for the geologic maps and the stratigraphic columns of the Pinon Canyon Maneuver Site.**

## References Cited

- Bass, N. W., C. E. Straub, and H. O. Woodbury  
1947 *Structure Contour Map of the Surface Rocks of the Model Anticline, Las Animas County, Colorado*. U.S. Geological Survey Oil and Gas Investigations, Preliminary Map 68.
- Burch, J. B.  
1975 Freshwater Sphaeriacean Clams (Mollusca: Pelecypoda) of North America. *Biota of Freshwater Ecosystems*, U.S. Environmental Protection Agency, Identification Manual No. 3.  
1980. North American Freshwater Snails, Species List, Ranges, and Illustrations. *Transactions of the POETS Society* 3:81-215.
- Cobban, W. A.  
1968 Pierre Shale and Older Cretaceous Rocks in Southeastern Colorado. *Mountain Geologist* 5:25-27.  
1971 *New and Little Known Ammonites from the Upper Cretaceous (Cenomanian and Turonian) of the Western Interior of the United States*. U.S. Geological Survey Professional Paper 699.
- Cobban, W. A., and Scott, G. R.  
1972 *Stratigraphy and Ammonite Fauna of the Graneros Shale and Greenhorn Limestone near Pueblo, Colorado*. U.S. Geological Survey Professional Paper 645.
- Duce, J. T.  
1924 *Geology of Parts of Las Animas, Otero and Bent Counties, Colorado*. U.S. Geological Survey Bulletin, 27:73-102.
- Evanoff, E.  
1997 *Preliminary Mapping and Report on the Surficial Geology and Paleontologic Resources of the Pinon Canyon Maneuver Site, Las Animas County, Colorado*. Boulder, University of Colorado Museum, unpublished report submitted to the Midwest Archeological Center, National Park Center, Lincoln, Nebraska.
- Finlay, G. L.  
1916 *Description of the Colorado Springs Quadrangle, Colorado*. U.S. Geological Survey Atlas, Folio 203.
- Gould, C. N.  
1934 *Report on Black Hills area, Las Animas County, Colorado*. U.S. Geological Survey, Unpublished Geologic Map. Available at Arthur Lakes Library, Colorado School Of Mines, Colorado.
- Griggs, R. L. and C. B. Read  
1959 Revisions in Stratigraphic Nomenclature in Tucumcari-Sabinoso Area, Northeastern New Mexico. *American Association of Petroleum Geologists Bulletin* 43:2003-2007.

- Helgren, D. M., A. Pitty., and J. Schuldenrein  
1985 Geologic and Geomorphic Setting, in Schuldenrein, J. ed., *Geomorphological and Geoarcheological Investigations at the U.S. Army Fort Carson-Pinon Canyon Maneuver Site, Las Animas County, Colorado*: National Park Service, Rocky Mountain Region, Final Report 2655:47-77.
- Johnson, R. B.  
1969 *Geologic Map of the Trinidad Quadrangle, South-Central Colorado*. U.S. Geological Survey Miscellaneous Investigations Map I-558.
- Kauffman, E. G.  
1986 *Geological and Paleontological Site Analysis of the Pinon Canyon Maneuver Site, Las Animas County, Colorado*. Fort Carson, Colorado, U.S. Army unpublished report.
- Keyes, C. R.  
1940 Early Cretacic Purgatoire Formation of Southeastern Colorado: *Pan American Geologist* 74:301-303.
- Kues, B. S., and S. G. Lucas  
1987 Cretaceous Stratigraphy and Paleontology in the Dry Cimmaron Valley, New Mexico, Colorado, and Oklahoma. In *Northeastern New Mexico*, edited by S. G. Lucas and A. P. Hunt, pp. 167-198. New Mexico Geological Society 38th Annual Field Conference.
- Lee, W. T.  
1901 The Morrison Formation of Southeastern Colorado. *Journal of Geology* 9:343-352.
- Long, C. S.  
1966. *Basal Cretaceous Strata, Southeastern Colorado*. Unpublished Ph.D. dissertation, Department of Earth Sciences, University of Colorado, Boulder.
- Lucas, S. G., A. P. Hunt, and B. S. Kues  
1987 Stratigraphic Nomenclature and Correlation Chart for Northeastern New Mexico. In *Northeastern New Mexico*, edited by S. G. Lucas and A. P. Hunt, pp. 351-354. New Mexico Geological Society 38th Annual Field Conference.
- Maher, J. C.  
1946 Correlation of Paleozoic Rocks across the Las Animas Arch in Baca, Las Animas, and Otero counties, Colorado: *American Association of Petroleum Geologists Bulletin* 30:1756-1763.
- Maher, J. C., and J. B. Collins  
1953 *Permian and Pennsylvanian Rocks of Southeastern Colorado and Adjacent Areas*. United States Geological Survey, Oil and Gas Investigation Map OM-135.

Mateer, N. J.

- 1987 The Dakota Group of Northeastern New Mexico and Southern Colorado. In *Northeastern New Mexico*, edited by S. G. Lucas and A. P. Hunt, pp. 223-236. New Mexico Geological Society 38th Annual Field Conference.

McFaul, M., and Reider, R. G.

- 1990 Physical Environment. In *An introduction to the Archeology of Pinon Canyon, Southeastern Colorado, Volume 1: Background and Methods* edited by W. Andresfsky, Jr., pp. II-1 to II-12. Larson and Tibesar Associates, Inc., Laramie, Wyoming. Submitted to the National Park Service, Rocky Mountain Regional Office, Denver, Colorado, Contract no. CX 1200-7-B054.

Oriel, S. S., and Mudge, M. R.

- 1956 Problems of Lower Mesozoic Stratigraphy in Southeastern Colorado. In *Guide Book to the Geology of the Raton Basin, Colorado*, edited by C. J. Mc Ginnis, pp. 19-24. Rocky Mountain Association of Geologists Guidebook for 1956.

Ostrom, J. H., and McIntosh, J. S.

- 1966 *Marsh's Dinosaurs: The Collections from Como Bluff*. Yale University Press, New Haven, Connecticut.

Schuldenrein, J. (editor)

- 1985 *Geomorphological and Geoarcheological Investigations at the U.S. Army Fort Carson-Pinon Canyon Maneuver Site, Las Animas County, Colorado*. National Park Service, Rocky Mountain Region, Final Report 2655.

Scott, G. R.

- 1968 *Geologic and Structure Contour Map of the La Junta Quadrangle, Colorado and Kansas*. U.S. Geological Survey Miscellaneous Investigations Map I-560.

Stose, G. W.

- 1912 *Description of the Apishapa Quadrangle, Colorado*. U.S. Geological Survey Geologic Atlas Folio 186.

Taylor, A. M.

- 1974 Stratigraphy and Sedimentary Environment of the Lower Cretaceous in Portion of Las Animas, Otero, and Bent Counties, Colorado. Unpublished Ph.D. dissertation, Department of Geology, Colorado School of Mines, Golden.
- 1980 *Depositional Environments of the Dakota Sandstone in Southeastern Colorado*. Society of Economic Paleontologists and Mineralogists, Denver, CO, Guidebook for Field Trip No.3.

Waage, K. W.

- 1953 *Refractory Clay Deposits of South-Central Colorado*. U.S. Geological Survey Bulletin 993.