

REPORT DOCUMENTATION PAGE

Form Approved
OMB No. 0704-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.

1. AGENCY USE ONLY (Leave blank)		2. REPORT DATE 2/28/99	3. REPORT TYPE AND DATES COVERED Final Report, 7/1/95-6/30/98	
4. TITLE AND SUBTITLE Structural Aspects of Protein-Metal Recognition and Discrimination			5. FUNDING NUMBERS N00014-95-1-0957	
6. AUTHOR(S) David W. Christianson				
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Department of Chemistry University of Pennsylvania Philadelphia, PA 19104-6323			8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) Office of Naval Research 800 N. Quincy St. Arlington, VA 22217-5000			10. SPONSORING/MONITORING AGENCY REPORT NUMBER	
11. SUPPLEMENTARY NOTES				
12a. DISTRIBUTION/AVAILABILITY STATEMENT Distribution Unlimited			12b. DISTRIBUTION CODE	
13. ABSTRACT (Maximum 200 words) This project utilized the zinc enzyme human carbonic anhydrase II (CAII) as a paradigm for dissecting and understanding the structural basis of protein-transition metal recognition and discrimination. We have successfully redesigned the zinc binding site of CAII so that protein-metal affinity can be modulated by design, by factors of 10 from micromolar to femtomolar affinity. We have explored the binding of zinc and copper to engineered CAII's using high resolution X-ray crystallographic methods, and we have developed a zinc-sensing scheme using fluorescence anisotropy and a designed sulfonamide inhibitor.				
14. SUBJECT TERMS protein crystallography, zinc enzyme, biosensor			15. NUMBER OF PAGES 5	
			16. PRICE CODE	
17. SECURITY CLASSIFICATION OF REPORT Unclassified	18. SECURITY CLASSIFICATION OF THIS PAGE Unclassified	19. SECURITY CLASSIFICATION OF ABSTRACT Unclassified	20. LIMITATION OF ABSTRACT UL	

FINAL REPORT

GRANT #: N00014-95-1-0957

PRINCIPAL INVESTIGATOR: David W. Christianson

INSTITUTION: University of Pennsylvania

GRANT TITLE: Structural Aspects of Protein-Metal Recognition

AWARD PERIOD: 1 July 1995 - 30 June 1998

OBJECTIVE: To develop and employ a structure-based design rationale for the engineering of the zinc binding site of carbonic anhydrase II (CAII); this work is leading to the development and deployment of a metal ion biosensor. This AASERT award complements the research effort funded by the main ONR grant, "Structural Studies of Metalloprotein Design".

APPROACH: Site-specific mutagenesis experiments in the laboratory of collaborator Carol Fierke (Duke University) are guided by molecular modeling experiments with the goal of predicting structure-stability relationships in engineered metal site variants of CAII. To provide proof-of-concept in the design-analysis-redesign cycle, variants are subsequently subject to high resolution X-ray structure determination.

ACCOMPLISHMENTS: The structures of 30 CAII variants relevant to the design of a CAII-based biosensor have been determined. We have determined the most detailed structure-affinity relationships known for any zinc metalloenzyme. Our work has focused on (a) direct metal ligands, (b) indirect metal ligands (i.e., those that accept hydrogen bonds from direct metal ligands), and (c) hydrophobic residues that surround the protein-zinc binding site. Highlights of this work include the structure-based engineering of the tightest protein-zinc binding site: the T199C variants exhibits femtomolar zinc affinity. Additionally, we are now capable of engineering protein-metal affinity by factors of 10 from the micromolar range to the femtomolar range.

Substitution of direct metal ligands H94, H96, and H119 by negatively-charged cysteine, aspartate, and glutamate residues typically maintains the tetrahedral zinc coordination polyhedron, including the zinc-bound solvent molecule. However, these variants usually exhibit compromised affinity and catalytic activity, and the pKa value for zinc-bound solvent is dramatically increased. This is due to the perturbation of the electrostatic environment of the metal ion. Introduction of a new negative charge from the protein compromises the ability of the metal ion to stabilize bound hydroxide ion, which results in an elevated

19990309031

pKa value. Accordingly, introduction of neutral asparagine and glutamine ligands in place of H119 results in CAII variants that retain substantial catalytic activity and exhibit near-normal pKa values for zinc-bound solvent.

Substitution of indirect zinc ligands results in more subtle effects on structure-affinity relationships. The carboxamide side chain of Q92 accepts a hydrogen bond from zinc ligand H94. The structure of the Q92E variants reveals an essentially isosteric replacement of the carboxamide side chain by the carboxylate side chain. However, the pKa of zinc-bound solvent is slightly elevated, due to the perturbation of the electrostatic environment of the metal ion. The introduction of a new negative charge even one residue removed from the metal ion affects the biological role of the metal ion in stabilizing its bound hydroxide ion. In an unusually dramatic example, the isosteric E117Q substitution results in severely-compromised activity (E117 accepts a hydrogen bond from zinc ligand H119). Unexpectedly, the structure of the E117Q variant suggests that this substitution stabilizes H119 as the negatively-charged histidinate anion, reflected by the corresponding increase in the pKa value for zinc-bound solvent.

Recent studies have focused on multiple-site CAII variants with amino acid substitutions in the beta-strand containing metal ligands H94 and H96. Structural comparison of the metal-bound and metal-free forms of selected double- and triple-site variants reveal conformational changes upon metal binding that are responsible for altered and more favorable metal binding kinetics. These CAII variants in particular are well-suited for application in a real-time CAII-based zinc biosensor, and work to this end is continuing in the laboratories of Carol Fierke (Duke University) and Richard Thompson (University of Maryland).

SIGNIFICANCE: Our structure-based redesign experiments with CAII yield valuable information regarding structure-stability relationships of zinc binding sites in proteins. Our collaborators Carol Fierke (Duke University) and Richard Thompson (University of Maryland School of Medicine) are now exploiting this information to optimize the molecular properties of the CAII-based biosensor.

PATENT INFORMATION: A patent application has been filed jointly with Richard Thompson at the University of Maryland on the use of CAII as a metal ion biosensor using a fluorescence anisotropy sensing scheme.

AWARD INFORMATION: Pfizer Award in Enzyme Chemistry from the American Chemical Society, 1999.

PUBLICATIONS:

Ippolito, J.A.; Christianson, D.W. "Structural Consequences of Redesigning a Protein-Zinc Binding Site". **Biochemistry** 1994, 33, 15241-15249.

Ippolito, J.A.; Baird, T.T.; McGee, S.A.; Christianson, D.W.; Fierke, C.A. "Structure-Assisted Redesign of a Protein-Zinc Binding Site with Femtomolar Affinity". **Proc. Natl. Acad. Sci. USA** 1995, 92, 5017-5021.

Lesburg, C.A.; Christianson, D.W. "X-ray Crystallographic Studies of Engineered Hydrogen Bond Networks in a Protein-Zinc Binding Site". **J. Am. Chem. Soc.** 1995, 117, 6838-6844.

Ippolito, J.A.; Nair, S.K.; Alexander, R.S.; Kiefer, L.L.; Fierke, C.A.; Christianson, D.W. "Structure of His-94-Asp Carbonic Anhydrase II in a New Crystalline Form Reveals a Partially-Occupied Zinc Binding Site". **Protein Eng.** 1995, 8, 975-980.

Nair, S.K.; Elbaum, D.; Christianson, D.W. "Unexpected Binding Mode of the Sulfonamide Fluorophore 5-Dimethylamino-1-Naphthalene Sulfonamide to Human Carbonic Anhydrase II: Implications for the Development of a Zinc Biosensor". **J. Biol. Chem.** 1996, 271, 1003-1007.

Huang, C.-C.; Lesburg, C.A.; Kiefer, L.L.; Fierke, C.A.; Christianson, D.W. "Reversal of the Hydrogen Bond to Zinc Ligand Histidine-119 Dramatically Diminishes Catalysis and Enhances Metal Equilibration Kinetics in Carbonic Anhydrase II". **Biochemistry** 1996, 35, 3439-3446.

Elbaum, D.E.; Nair, S.K.; Patchan, M.W.; Thompson, R.B.; Christianson, D.W. "Structure-Based Design of a Sulfonamide Probe for Fluorescence Anisotropy Detection of Zinc with a Carbonic Anhydrase-Based Biosensor". **J. Am. Chem. Soc.** 1996, 35, 8381-8387.

Christianson, D.W.; Fierke, C.A. "Carbonic Anhydrase: Evolution of the Zinc Binding Site by Nature and by Design". **Acc. Chem. Res.** 1996, 29, 331-339.

Thompson, R.B.; Ge, Z.; Patchan, M.W.; Fierke, C.A.; McCall, K.A.; Elbaum, D.; Christianson, D.W. "Determination of Multiple Analytes Using a Fiber Optic Biosensor Based on Fluorescence Energy Transfer". **Prog. Biomed. Optics** (Proceedings of the SPIE Conference on Ultrasensitive Biochemical Diagnostics, Cohn, G.E.; Soper, S.A.; Chen, C.H.W., Eds.) 1996, 2680, 47-56.

Lesburg, C.A.; Huang, C.-C.; Christianson, D.W.; Fierke, C.A.
"Histidine-Carboxamide Ligand Substitutions in the Zinc
Binding Site of Carbonic Anhydrase II Alter Metal
Coordination Geometry but Retain Catalytic Activity".
Biochemistry 1997, 36, 15780-15791.

Hunt, J.A.; Lesburg, C.A.; Christianson, D.W.; Thompson,
R.B.; Fierke, C.A. "Active Site Engineering of Carbonic
Anhydrase and Its Application to Biosensors". In **The
Carbonic Anhydrases: New Horizons**, Chegwiddden, W.R., Carter,
N.D., & Edwards, Y.H., Eds. In press.