

# Towards sub-microsecond rigid Earth nutation series in the Hamiltonian theory

J. Souchay and M. Folgueira<sup>1</sup>

*Observatoire de Paris, 61 Avenue de l'Observatoire. 75014 Paris, France*

**Abstract.** The nonrigid Earth nutation series adopted by the IAU (International Astronomical Union) are based on the works of Kinoshita (1977) and Wahr (1979). In the first one, the rigid Earth nutation series were calculated by the application of the Hamiltonian canonical equations to the rotation of the rigid and elliptical Earth. In the second one, the transfer function for the nutations of an elliptical, elastic and oceanless Earth with fluid core and a solid inner core was obtained. The nonrigid Earth nutation coefficients were derived from the convolution between Wahr's transfer function and Kinoshita's rigid Earth nutation series.

The improvement in the accuracy of the techniques as a Very Long Baseline (VLBI), Lunar Laser Ranging (LLR) and Global Positioning System (GPS) has led in this decade to the extension of Kinoshita's theory and more precise determination of Wahr's transfer function. In the present paper and starting from Kinoshita's work (1977), we present the different steps carried out, during this last decade, to obtain the sub-microsecond rigid Earth nutation series REN 2000 from the Hamiltonian study of the rotation of a rigid Earth (Souchay *et al.*, 1999).

## 1. Introduction

In 1980, the IAU adopted a new series of nutation for a nonrigid Earth model based on Kinoshita's series on the Hamiltonian study of the rotation of a rigid Earth. The nonrigid Earth nutation coefficients were derived from Kinoshita's series (Kinoshita, 1977) and Wahr's transfer function (Wahr, 1979). 0.1 *mas* (milliarcsecond) was the level of accuracy considered for these coefficients of nutation. The effects considered at this level of precision were influence of zonal harmonics:  $J_2$ ,  $J_3$  and  $J_4$ , and the influence of the triaxiality of the Earth.

Kinoshita's work was the first paper to present a rigorous study of the rotation of a rigid body, starting from Hamiltonian theory. This work has been the basis of a large number of papers, and it was the improvement in the accuracy of VLBI, LLR and GPS which has led to extension of Kinoshita's theory. In the following, the different steps carried out are presented.

---

<sup>1</sup>On leave from Instituto de Astronomia y Geodesia (UCM-CSIC). Facultad de Ciencias Matemáticas. Universidad Complutense de Madrid. Ciudad Universitaria, 28040 Madrid, Spain

## Report Documentation Page

*Form Approved*  
*OMB No. 0704-0188*

Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.

1. REPORT DATE <b>MAR 2000</b>	2. REPORT TYPE <b>N/A</b>	3. DATES COVERED <b>-</b>			
4. TITLE AND SUBTITLE <b>Towards Sub-Microsecond Rigid Earth Nutation Series in the Hamiltonian Theory</b>		5a. CONTRACT NUMBER			
		5b. GRANT NUMBER			
		5c. PROGRAM ELEMENT NUMBER			
6. AUTHOR(S)		5d. PROJECT NUMBER			
		5e. TASK NUMBER			
		5f. WORK UNIT NUMBER			
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) <b>U.S. Naval Observatory 3450 Massachusetts Avenue, N.W. Washington, DC 20392-5420</b>		8. PERFORMING ORGANIZATION REPORT NUMBER			
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)		10. SPONSOR/MONITOR'S ACRONYM(S)			
		11. SPONSOR/MONITOR'S REPORT NUMBER(S)			
12. DISTRIBUTION/AVAILABILITY STATEMENT <b>Approved for public release, distribution unlimited</b>					
13. SUPPLEMENTARY NOTES <b>Towards models and constants for sub-microarcsecond astrometry, Proceedings of IAU Colloquium 180 held at the U.S. Naval Observatory, Washington, DC, USA, 27-30 March 2000</b>					
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT <b>UU</b>	18. NUMBER OF PAGES <b>5</b>	19a. NAME OF RESPONSIBLE PERSON
a. REPORT <b>unclassified</b>	b. ABSTRACT <b>unclassified</b>	c. THIS PAGE <b>unclassified</b>			

## 2. The first second-order theory of the rotation of a rigid Earth

Kinoshita and Souchay (1990) published a new series of nutation for a rigid Earth model, considering a new level of truncation for the coefficients of nutation:  $0.005 \text{ mas}$ . This new level made necessary the study of new effects and the extension of some developments. The principal influences can be briefly summed up as (Kinoshita & Souchay, 1990): influence of the  $J_2$  harmonic and of the second-order parts of the potential of the Earth ( $J_3$ , triaxiality and  $J_4$ ), direct and indirect planetary effects, and extension of the theory to the second order: coupling effect between the rotational motion of the Earth and the orbital motion of the Moon.

## 3. New developments in rigid Earth nutation theory

The updated value of the general precession in longitude led to a change of the value of the dynamical ellipticity of the Earth as well as the scaling factors for the nutation coefficients relative to the potential of the Moon and the Sun. For this reason, Souchay and Kinoshita (1996, 1997) recalculated the coefficients of nutation - due to the lunisolar influence on the geopotential ( $J_2$ ,  $J_3$ ,  $C_{22}$ ,  $S_{22}$  and  $J_4$ ) and those coming from the direct torque exerted by the planets - influenced by these changes. Moreover, they presented new contributions to the nutation not contained in (Kinoshita & Souchay, 1990) including the influence of the periodic oscillations of the ecliptic, and the planetary tilt-effect. The agreement of the new coefficients of Souchay & Kinoshita (1996, 1997) with those of Hartmann & Soffel (1994) and Williams (1994, 1995) was remarkable.

## 4. The effects on nutation of the non-zonal harmonics of third and fourth degree

Taking into account the new accuracy of the precession-nutation observations (a few microseconds), Folgueira *et al.* (1998a,b) calculated the coefficients of nutation for a rigid Earth model due to  $C_{3m}$ ,  $S_{3m}$ ,  $C_{4m}$  and  $S_{4m}$  ( $m \neq 0$ ) harmonics of the geopotential, which were not considered in previous studies of nutation involving Hamiltonian theory, with a level of truncation of  $0.1 \mu\text{as}$  (microarcseconds). These non-zonal harmonics, together with  $C_{22}$  and  $S_{22}$ , give birth to short period nutations, the periods being related to the order  $m$  of the corresponding harmonic. That is to say:

- The diurnal terms come from  $C_{31}$ ,  $S_{31}$ ,  $C_{41}$  and  $S_{41}$ .
- The semidiurnal terms are due to the harmonics with  $m = 2$ .
- The terdiurnal terms have their origin in  $C_{33}$ ,  $S_{33}$ ,  $C_{43}$  and  $S_{43}$  harmonics.

The principal diurnal and subdiurnal terms for the nutation in longitude and obliquity are shown in Tables 1 and 2.

Table 1. Principal terms for quasi-diurnal nutations in longitude and obliquity for the figure axis. The unit is  $\mu as$

$\Phi$	Argument					Period	Longitude ( $\Delta\psi$ )		Obliquity ( $\Delta\varepsilon$ )	
	$l_M$	$l_S$	$F$	$D$	$\Omega$		sin	cos	sin	cos
1	0	0	1	0	1	0.96215	-38.2313	-4.6980	-1.8567	15.1063
1	0	0	-1	0	-1	1.03505	-35.4036	-4.3513	-1.5868	12.9109
1	-1	0	1	0	1	0.99696	24.1443	2.9675	1.1636	-9.4678
1	1	0	-1	0	-1	0.99758	-19.9400	-2.4507	-0.9716	7.9055
1	1	0	1	-2	1	0.99216	-7.0629	-0.8681	-0.3452	2.8089

Table 2. Principal terms for sub-diurnal nutations in longitude and obliquity for the figure axis. The unit is  $\mu as$

$\Phi$	Argument					Period	Longitude ( $\Delta\psi$ )		Obliquity ( $\Delta\varepsilon$ )	
	$l_M$	$l_S$	$F$	$D$	$\Omega$		sin	cos	sin	cos
2	0	0	0	0	0	0.49863	31.2522	-17.9397	-7.1157	-12.3961
2	0	0	-2	0	-2	0.51753	-25.5271	14.6533	5.6775	9.8906
2	0	0	-2	2	-2	0.50000	-10.4455	5.9960	2.3682	4.1255
2	-1	0	-2	0	-2	0.52743	-5.0979	2.9264	1.1818	2.0588
2	0	0	-2	0	-1	0.51756	-4.7105	2.7040	1.0816	1.8842

## 5. The sub-microarcsecond rigid Earth nutation series REN 2000

Souhay *et al.* (1999) presented the new tables REN 2000 of the nutation of a rigid Earth, starting from Hamiltonian theory, with sub-microarcsecond level of truncation. All the previous studied influences, together with the second-order effects due to crossed-nutations and spin-orbit coupling, are included in this paper.

## 6. The diurnal and sub-diurnal nutations of REN 2000

Recently other rigid Earth nutation series, calculated using different analytical methods, have come forth. These are SMART97 and RDAN97 series (Bretagnon *et al.*, 1997, 1998; Roosbeek & Dehant, 1998). Folgueira & Souhay (1999) and Folgueira *et al.* (1999) have compared the nutations given by REN 2000 series with the corresponding ones in SMART97 and RDAN97. Table 3 gives the comparison for the short period nutations in the time domain between these three theories.

Moreover, they have computed the corresponding values for a nonrigid Earth of REN 2000 prograde nutations, using a new transfer function developed by Mathews (1999) and showed that some coefficients changed at the level of a few  $\mu as$ . Finally, they have also explained how short period nutations can perturb the analysis of the polar motion (Bizouard *et al.*, 1999a;b).

Table 3. Comparison in time domain between SMART97, RDAN97 and REN 2000 short period nutations. The unit is  $\mu as$

	$\Delta\psi \sin \varepsilon_0$	$\Delta\varepsilon$
SMART97 – REN 2000	1.74	3.52
RDAN97 – REN 2000	2.40	6.80
RDAN97 – SMART97	1.40	6.80

## 7. Conclusions

In the present paper, we have analyzed the different steps carried out, in this last decade, to reach at the final sub-microarcsecond tables of nutation for a rigid Earth model REN 2000 (Souchay *et al.*, 1999). As summary, we can conclude that:

1. The tables REN 2000 of nutation for a rigid Earth model, starting from the Hamiltonian theory, catch all the coefficients at a *sub-microarcsecond* level.
2. There is a good agreement between REN 2000 and other recent works about rigid Earth nutations (Bretagnon *et al.*, 1997, 1998; Roosbeek & Dehant, 1998).
3. The short period nutations are the last most important contribution to reach at the 0.1  $\mu as$  level in REN 2000 tables.
4. The validity of the analytical series REN 2000 has been checked by a numerical integration of the nutation developed by Souchay & Kinoshita (1991) and Souchay (1998) and by the study of the residuals between the results given by these two different methods.

**Acknowledgments.** This investigation was supported by a Postdoctoral Aid for Complutense University of Madrid (Spain) to carry out a research work at the Observatory of Paris (France).

## References

- Bizouard, Ch., Folgueira, M. and Souchay, J., 1999a, *Proc. Journées 1999. Observatoire de Paris* (submitted)
- Bizouard, Ch., Folgueira, M. and Souchay, J., 1999b, *Proc. IAU Colloquium 178*, 613.
- Bretagnon, P., Rocher, P. and Simon, J.-L., 1997, *Astron. Astrophys.*, **319**, 305.
- Bretagnon, P., Francou, G., Rocher, P. and Simon, J.-L., 1998, *Astron. Astrophys.*, **329**, 329.

- Folgueira, M., Souchay, J. and Kinoshita, H., 1998a, *Celest. Mech.*, **69**(4), 373.
- Folgueira, M., Souchay, J. and Kinoshita, H., 1998b, *Celest. Mech.*, **70**(3), 147.
- Folgueira, M. and Souchay, J., 1999, *Proc. Journées 1998. Observatoire de Paris*, 178.
- Folgueira, M., Bizouard, Ch. and Souchay, J., 1999, *Celest. Mech.* (submitted).
- Hartmann, T. and Soffel, M., 1994, *Astron. J.*, **108**, 1115.
- Kinoshita, H., 1977, *Celest. Mech.*, **15**, 277.
- Kinoshita, H. and Souchay, J., 1990, *Celest. Mech.*, **48**, 187.
- Roosbeek, F. and Dehant, V., 1998, *Celest. Mech.*, **70**(4), 215.
- Souchay, J. and Kinoshita, H., 1991, *Celest. Mech.*, **52**, 45.
- Souchay, J. and Kinoshita, H., 1996, *Astron. Astrophys.*, **312**, 1017.
- Souchay, J. and Kinoshita, H., 1997, *Astron. Astrophys.*, **318**, 639.
- Souchay, J., 1998, *Astron. J.*, **116**, 503.
- Souchay, J., Loysel, B., Kinoshita, H. and Folgueira, M., 1999, *Astron. Astrophys. Suppl. Ser.*, **135**, 111.
- Souchay, J., 1999, *Proc. Journées 1998. Observatoire de Paris*, 109.
- Williams, J.G., 1994, *Astron. J.*, **108**, 711.
- Williams, J.G., 1995, *Astron. J.*, **110**, 1420.