

73rd MORSS CD Cover Page

UNCLASSIFIED DISCLOSURE FORM CD Presentation

712CD

For office use only 41205

21-23 June 2005, at US Military Academy, West Point, NY

Please complete this form 712CD as your cover page to your electronic briefing submission to the MORSS CD. Do not fax to the MORS office.

Author Request (To be completed by applicant) - The following author(s) request authority to disclose the following presentation in the MORSS Final Report, for inclusion on the MORSS CD and/or posting on the MORS web site.

Name of Principal Author and all other author(s):

Cheryl Ann Blain
T. Christopher Massey
Brett D. Estrade

Principal Author's Organization and address:

Phone: (228) 688-5450

Oceanography Division (Code 7322)
Naval Research Laboratory
Stennis Space Center, MS 39529

Fax: (228) 688-4759

Email: blain@nrlssc.navy.mil

Original title on 712 A/B: High Resolution Characterization of Riverine and Coastal Currents

Revised title: N/A

Presented in (input and Bold one): (WG___, CG___, Special Session ___, Poster, Demo, or Tutorial): **WG10**

This presentation is believed to be:

UNCLASSIFIED AND APPROVED FOR PUBLIC RELEASE




Report Documentation Page

Form Approved
OMB No. 0704-0188

Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.

1. REPORT DATE 21 JUN 2005	2. REPORT TYPE N/A	3. DATES COVERED -	
4. TITLE AND SUBTITLE High Resolution Characterization of Riverine and Coastal Currents		5a. CONTRACT NUMBER	
		5b. GRANT NUMBER	
		5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)		5d. PROJECT NUMBER	
		5e. TASK NUMBER	
		5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Oceanography Division Naval Research Laboratory Stennis Space Center, MS 39529		8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)		10. SPONSOR/MONITOR'S ACRONYM(S)	
		11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release, distribution unlimited			
13. SUPPLEMENTARY NOTES See also ADM201946, Military Operations Research Society Symposium (73rd) Held in West Point, NY on 21-23 June 2005., The original document contains color images.			
14. ABSTRACT			
15. SUBJECT TERMS			
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified	UU
			18. NUMBER OF PAGES 34
			19a. NAME OF RESPONSIBLE PERSON



High Resolution Characterization of Riverine and Coastal Currents

*Dr. Cheryl Ann Blain
Dr. T. Christopher Massey
Mr. Brett D. Estrade*


*Oceanography Division
Naval Research Laboratory
Stennis Space Center, MS*

*21-23 June, 2005
73rd MORS Symposium, West Point, NY*

What can we do?

Predictive Capability for Coastal Circulation

- High resolution (meters) currents and water levels in littoral environments that include bays, inter-tidal marshes and rivers




ADCIRC Model

- 3D dynamics
- Forcing from tides, wind, waves, and rivers
- Shoreline inundation/recession
- Utilizes unstructured grids (based on finite elements)
- MPI parallelization




Advantage: Mesh Flexibility




- Realistic coastline morphology
- Fine-scale resolution of inter-tidal zones
- Large domains with remote open boundaries

ADCIRC: SE Louisiana Coastal Waters




Tidal forcing along ocean boundary from global tide model




MS River discharge ~ 2.3 times
Atchafalaya River discharge


ADCIRC: SE Louisiana Coastal Waters



ADCIRC: SE Louisiana Coastal Waters



Modeled Currents – March 17-24, 2004








Development of Embayment and River Models

Bay St. Louis Model

Biloxi Bay Model



Mobile Bay Model

Pearl River Model




Mississippi Bight Model

+ L. Ponchartrain




Coupled Wave-Tide Circulation in Bay St. Louis

Pathways of Numerical Drifters over 23 Hours



Tidal Forcing Only




Wave-Tide Interaction

Cobb and Blain, MTS, 2002
Blain and Cobb, NRL Report,
Cobb and Blain, NRL Report,

Sensitivity of Bay-Shelf Exchange to Forcing and Model Dimensionality


3-D CIRCULATION

After 15 days



2-D CIRCULATION

Real-Time Forecast Capability



 Under Development

Goals

- Rapid Re-Location
- Timely forecasts
- Relevant products

Applications


- AUVFEST
Oct-Nov, 2001
Mississippi Sound

- *Prestige* Oil Spill
Nov-Dec, 2002
Spanish coast

- *Operation Iraqi Freedom*
Mar-Apr, 2003
KAA Persian Gulf


Criteria for Success

- Accurate and flexible dynamical model
- Ability to meet operational time constraints
- Automated, rapid relocation
- Generation of meaningful operational products
- Quantification of forecast skill






A Tidal Model in the N. Persian Gulf

Operation Iraqi Freedom, March-April 2003






No. nodes 247,767
No. elements 480,001
Resolution 2 m – 1 km



Validation of the Forecast Elevations

March 26-27, 2003

0 0.125



RMS Error 25 cm

Mean Absolute Error 20 cm


Phase Lag 15-30 min

Time series correlation 0.98




ADCIRC Model Support for OIF, 2003


Resolution
2 m – 400 m




Max 12-hr Current Thresholds



Water Levels




2D Currents (Mag/Dir)




Fine-Scale Circulation Features

Depth-averaged tidal currents north of Um Qsar




Variability on the scale of hours

Fine-Scale Circulation Features



Importance of Including Inundation



**No Shoreline
Inundation/Drying**


Spatial and temporal resolution sufficient to capture shoreline inundation/drying but circulation highly sensitive to bathymetric data.

Why Merge Models and Ocean Color Data?



Remotely-Sensed
Ocean Color Data

Modeled Coastal
Dynamics



Relate Optical and Physical
Properties of the Coastal Ocean

Complementary Aspects of Merging Models and Ocean Color Data

	<u>Imagery</u>	<u>Coastal Models</u>
Temporal Resolution	1-2 times daily	hourly or less
Spatial Resolution	250 m – 1 km	10 m -100 m
Dimensionality	surface information only	3D representation

Models can fill spatial and temporal gaps in ocean color data
Ocean color data provide observations of the real ocean


Together they can unravel the role of various bio-geo-physical processes in the coastal ocean



Diver Visibility

Derived using
beam attenuation
coefficient from
the MODIS
satellite


- AQUA and Terra
250 m channels
- AM: 7- 9 GMT
- PM: 22-24 GMT




March 5 – May 3, 2003



Low Visibility




FLOOD TIDE




Higher velocities result in more mixing and thus increased turbidity



Increased Visibility




SLACK TIDE




Low velocities allow particle settling and reduce turbidity



Low Visibility



EBB TIDE




The magnitudes of the ebb tide are not as strong as for flood tide



Transport Pathways


Passive Numerical Drogues



Where are we headed?

Improvements to the Predictive Capability for Coastal Circulation and Transport

- High resolution (meters) currents and water levels in littoral environments that include bays, inter-tidal marshes and rivers








Near-Term Developments

- Transport (T, S, tracers)
- Nesting to regional models
- Connection to bio-optical properties




Finite Element Model Development

Salinity Fronts in MS River Delta



Discontinuous Galerkin Methods



Storm Surge, Tides




Tidal Bore




Future: Adaptive Mesh Refinement

A Slotted Cylinder




**Error estimates
guide the
refinement
process.**

Uniform Mesh
1089 Nodes



Adaptive Mesh
~800 to 1200 Nodes



To achieve the same resolution as the adaptive mesh, a uniform mesh would require ~16,000 nodes.

Future: Connecting the Watershed, River and Coastal Ocean



Overlap between the watershed model and a coastal circulation model

Watershed

Circulation

Overland flow

River Discharge
Sediment Load

Water in marshes



Inundation/Drying

Vegetation

Inundation/Drying
Sediment Load

Future: Limited Domain Applications

Coupling global/regional scale and coastal models



- Coastal application in northern Gulf of Mexico; determine how to best extract boundary forcing data from regional/global models


Future: Predicting Bio-Optical Properties

Riverine Applications

IMAGERY



MODEL



Pearl River, MS

UAV



Future: Autonomous Chemical (Mine) Hunting

