

Award Number: MIPR 1LCCDD1149

TITLE: Enhancing the Immunogenicity of a Dengue-2 DNA Vaccine with Adjuvants and Anti-FCyRI Antibodies

PRINCIPAL INVESTIGATOR: Kevin R. Porter, M.D.  
Shuenn-Jue Wu, Ph.D.  
Kanakatte Raviprakash, Ph.D.

CONTRACTING ORGANIZATION: Naval Medical Research Center  
Silver Spring, MD 20910

REPORT DATE: October 2005

TYPE OF REPORT: Final

PREPARED FOR: U.S. Army Medical Research and Materiel Command  
Fort Detrick, Maryland 21702-5012

DISTRIBUTION STATEMENT: Approved for Public Release;  
Distribution Unlimited

The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision unless so designated by other documentation.

# REPORT DOCUMENTATION PAGE

*Form Approved*  
*OMB No. 0704-0188*

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number. **PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.**

<b>1. REPORT DATE (DD-MM-YYYY)</b> 01-10-2005		<b>2. REPORT TYPE</b> Final		<b>3. DATES COVERED (From - To)</b> 1 MAR 2005 - 28 FEB 2006	
<b>4. TITLE AND SUBTITLE</b> Enhancing the Immunogenicity of a Dengue-2 DNA Vaccinewith Adjuvants and Anti-FCyRI Antibodies				<b>5a. CONTRACT NUMBER</b>	
				<b>5b. GRANT NUMBER</b> MIPR 1LCCDD1149	
				<b>5c. PROGRAM ELEMENT NUMBER</b>	
<b>6. AUTHOR(S)</b> Kevin R. Porter, M.D. Shuenn-Jue Wu, Ph.D. Kanakatte Raviprakash, Ph. D.  E-mail: porterk@nmrc.navy.mil				<b>5d. PROJECT NUMBER</b>	
				<b>5e. TASK NUMBER</b>	
				<b>5f. WORK UNIT NUMBER</b>	
<b>7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES)</b>  Naval Medical Research Center Silver Spring, MD 20910				<b>8. PERFORMING ORGANIZATION REPORT NUMBER</b>	
<b>9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES)</b> U.S. Army Medical Research and Materiel Command Fort Detrick, Maryland 21702-5012				<b>10. SPONSOR/MONITOR'S ACRONYM(S)</b>	
				<b>11. SPONSOR/MONITOR'S REPORT NUMBER(S)</b>	
<b>12. DISTRIBUTION / AVAILABILITY STATEMENT</b> Approved for Public Release; Distribution Unlimited					
<b>13. SUPPLEMENTARY NOTES</b> Original contains color plates: All DTIC reproductions will be in black and white.					
<b>14. ABSTRACT</b> The objective is to use adjuvants to enhance the immunogenicity of a dengue (DEN) DNA vaccine. The adjuvants to be evaluated in this project included aluminum phosphate, tetanus toxoid and anti-Fc $\alpha$ R monoclonal antibodies chemically linked to the DNA vaccine. Multiple attempts to link the anti-Fc $\alpha$ R monoclonal antibody to the DNA vaccine resulted in a poor efficiency in the linkage reaction using thiol linkage chemistry. An alternative approach using hydrazine and aldehyde linkage methods were used but failed to improve DNA/antibody conjugation efficiency. Experiments were initiated to evaluate the ability of tetanus toxoid (TT) to enhance the immunogenicity of the DEN DNA vaccine. The antibody responses to the vaccine given IM were enhanced slightly by combining it with TT. However, the ELISA and neutralizing antibody levels were inferior to the levels obtained with ID administration of the vaccine alone. When using low doses of the vaccine given by Biojector, TT appeared to have an inhibitory effect. Aluminum Phosphate was demonstrated to slightly enhance anti-dengue neutralizing antibody responses, only when the DNA vaccine was given ID at suboptimal doses. There was essentially no effect when the vaccine was administered IM.					
<b>15. SUBJECT TERMS</b> dengue, DNA vaccine, monoclonal antibodies, Fc receptor, tetanus toxoid, aluminum phosphate					
<b>16. SECURITY CLASSIFICATION OF:</b>			<b>17. LIMITATION OF ABSTRACT</b>	<b>18. NUMBER OF PAGES</b>	<b>19a. NAME OF RESPONSIBLE PERSON</b> USAMRMC
<b>a. REPORT</b> U	<b>b. ABSTRACT</b> U	<b>c. THIS PAGE</b> U			<b>19b. TELEPHONE NUMBER (include area code)</b>
			UU	17	

## Table of Contents

Cover.....	1
SF 298.....	2
Introduction.....	4
Body.....	5
Key Research Accomplishments.....	16
Reportable Outcomes.....	17
Conclusions.....	17
References.....	17

## INTRODUCTION

Department of Defense (DoD) researchers have used the DNA vaccine approach to develop experimental vaccines against DEN-1 and DEN-2. These vaccines contain the envelope (E) and premembrane (prM) genes of the viruses. The E gene was chosen for inclusion in the DNA vaccines because monoclonal antibody mapping and recombinant protein immunization studies with flaviviruses have shown that the E gene contains the major epitopes responsible for eliciting neutralizing antibodies. The prM gene was also included in the vaccine because the expressed protein is thought to prevent low pH-induced irreversible conformational changes in the E protein as it is processed through the acidic compartments of the Golgi complex prior to secretion. The DEN-2 and DEN-1 DNA vaccines have been tested in mice and shown to elicit neutralizing antibody responses.

Two primate DNA immunization/challenge studies were completed. One utilized Rhesus monkeys, the other *Aotus* monkeys. Both studies were very similar in design and outcome. One-third of the animals were injected with control DNA (plasmid that does not express prM/E) and two-thirds of the animals were immunized with the DNA vaccine that expresses the prM/E genes of DEN-1 virus. All of the animals that received the DEN-1 DNA vaccine responded by producing neutralizing antibodies. The antibody levels peaked one month after the last DNA boost and then steadily declined during the following five months at which time the animals were challenged with DEN-1 virus. Upon challenge with live dengue virus, the DEN-1 DNA vaccine-immunized monkeys showed a significant reduction in the mean days of viremia compared to naïve controls, but the protection was not complete. Although successful, these studies do indicate the need to improve the efficacy of DEN DNA vaccines.

Immune responses to DNA vaccines have been enhanced by co-administering them with immune-modulating compounds including cytokines, chemokines, co-stimulatory molecules and cationic lipids. Although these approaches have resulted in improved potency, none of these are approved for use in humans with prophylactic vaccines. Aluminum-based adjuvants have been used successfully in humans to enhance the immune response to protein vaccines and are approved for use in humans. Studies have demonstrated that intramuscular co-administration of DNA vaccines with aluminum phosphate enhanced the antibody response 10- to 100-fold. This response occurred with DNA vaccines for hepatitis B and influenza A. The use of calcium phosphate, aluminum hydroxide and aluminum hydrophosphate as adjuvants had an inhibitory effect on the antibody response to the DNA vaccines. This inhibitory effect was the result of the DNA binding to these adjuvants, making less available for uptake by antigen presenting cells (APCs). The enhancement of vaccine potency was the result of increased recruitment of APCs.

Highly immunogenic carriers such as tetanus toxoid, diphtheria toxoid and cholera toxin have been used successfully to enhance the immunogenicity of polysaccharide vaccines. These proteins work by involving T-cells in the processing of the carrier protein (T-cell help) and by generating both T- and B-cell memory. Little is known about the effect of these carrier proteins on the immune response to DNA vaccines.

Another possible way to enhance the uptake, expression and processing of vaccines is to link them with antibodies that more efficiently direct them to APCs. Gosselin et al. cross-linked anti-

Fc $\gamma$ RI monoclonal antibodies with tetanus toxoid and compared the immunogenicity of this compound with that of tetanus toxoid alone. The results showed that by linking the immunogen with the monoclonal antibody, antigen presentation was increased 100-fold. The mechanism of the increased antibody response was thought to involve the more efficient uptake and processing of the tetanus toxoid by APCs. This approach has yet to be attempted using DNA vaccines as the immunogen. Because only a small portion of DNA vaccines is actually taken up and expressed by APCs, it stands to reason that increasing the efficiency of uptake would result in enhanced cellular and humoral immune responses to nucleic acid vaccines.

This proposal seeks to enhance the immune response to a dengue (DEN) prM/E DNA vaccine by utilizing aluminum phosphate, tetanus toxoid and anti-Fc $\gamma$ R monoclonal antibodies as adjuvants in various formulations. Phase I of the project will be conducted in mice to demonstrate the feasibility of enhancing anti-DEN immune responses with these novel approaches. Phase II will be conducted using non-human primates to evaluate the protective efficacy of DNA vaccine formulations that are shown to be immunogenic in mice.

## **BODY**

### **Year 1**

#### **Statement of Work**


1. DEN DNA vaccine is chemically linked with Fc $\gamma$ R antibody and evaluated for expression *in vitro*.
2. Co-immunize mice with low and high doses of DEN DNA vaccine together with aluminum phosphate, tetanus toxoid and DNA vaccine linked with anti-Fc $\gamma$ R antibody.
3. Analyze antibody response by indirect ELISA and plaque reduction neutralization tests. Perform ELISPOT assays to detect T-cells producing interferon gamma (IFN $\gamma$ ) in response to DEN antigen stimulation to analyze dengue-specific cellular immune responses.

#### **Progress and Accomplishments**

1. DEN DNA vaccine is chemically linked with Fc $\gamma$ R antibody and evaluated for expression *in vitro*.

The first task was to link the dengue DNA vaccine to anti-Fc $\gamma$ R monoclonal antibodies. Prior to performing the chemical linking procedure, we first sought to confirm that the monoclonal antibodies selected for linkage to the DNA vaccine recognize the Fc $\gamma$ R of mammalian immune cells. The first set of anti-Fc $\gamma$ R antibodies, consisted of anti-Fc $\gamma$ R<sub>II</sub> (CD32) and anti-Fc $\gamma$ R<sub>III</sub> (CD16) (Accurate Chemical, Westbury, NY). Indirect immunofluorescence assays (IFA) were performed to evaluate the recognition of Fc $\gamma$  receptors on K562 cells (human monocyte cell line) and P388D1 cells (mouse monocyte cell line). Experimental results indicated poor recognition of mouse and human Fc $\gamma$  receptors by these monoclonal antibodies, as indicated by low level immunofluorescence. A second set of monoclonal antibodies was obtained (Pharmigen, Inc. San Diego CA) and consisted of anti-Fc $\gamma$ R<sub>I</sub> and anti-Fc $\gamma$ R<sub>II</sub>. IFA demonstrated recognition of anti-Fc $\gamma$ R<sub>II</sub> receptors but not Fc $\gamma$ R<sub>I</sub> (CD64) on K562 cells. The P388D1 mouse cells showed no reactivity with either antibody. This pattern of immunoreactivity was confirmed by flow cytometry analysis (Figure 1A&B).

Figure 1



Further studies to find P388D1 immunoreactive anti Fc $\gamma$ R monoclonal antibodies were conducted. Use of another commercial product, with combined anti- Fc $\gamma$ RII and anti- Fc $\gamma$ RIII immunoreactivity, revealed positive results against P388D1 cells by IFA.

Experiments to link the Fc $\gamma$ RII/III monoclonal antibody to DEN-1 DNA vaccine were then conducted. The DEN-1 DNA vaccine was used instead of the DEN-2 vaccine as originally planned because more extensive non-human primate efficacy data were available for the DEN-1 vaccine. Much of these DEN-1 DNA vaccine data became available after the original proposal was submitted. Historical comparisons with the DEN-1 DNA vaccine data would allow for a

more rigorous determination of whether the modifications improve the immunogenicity and protective efficacy of the DNA vaccine.

To conjugate the anti-FcγRII/RIII monoclonal antibody to the DNA vaccine, the DNA was linearized by Xho I restriction enzyme digestion to produce "sticky" 5' and 3' ends. To first confirm that the linearized DNA is capable of expressing the DEN-1 prM and E genes, 293 cells were transfected with the linearized DNA and analyzed by IFA using the anti-DEN monoclonal antibody 4G2. 293 cells are routinely used to evaluate expression of our experimental DNA vaccine constructs. These cells were used initially so that a head-to-head comparison can be made between the levels of expression obtained with the circular versus the linearized DNA vaccine plasmid construct. The results showed that the linearized DNA efficiently expressed the DEN-1 genes and that the level of expression was equivalent to that seen with the closed circular plasmid construct. Since our plan was to use the P388D1 cells to evaluate in vitro uptake and expression of the DNA vaccine/anti-FcγR antibody fusion product, we next wanted to confirm that the P388D1 cells were first capable of efficiently expressing the prM and E genes of the linearized DEN-1 plasmid before proceeding with the conjugation. Despite repeated attempts, we were unable to efficiently transfect the P388D1 cells and thus could not evaluate the expression of the linear DNA in this cell line. Another mouse macrophage/monocyte cell line (Raw 264.7 cells) was then evaluated. Raw 264.7 cells are transformed by Abelson murine leukemia virus. Following transfection with the DEN-1 vaccine plasmid, the Raw 264.7 cells expressed the prM and E proteins at a level similar to that of the 293 cells. Repeat binding studies with the anti-FcγRII/RIII monoclonal antibody showed efficient binding to the Raw 264.7 cells by IFA.

The first attempt to combine the DNA to the antibody was via thiol linkage to the amine groups on the antibody. The first step was to add a thiol group to the linearized DNA vaccine. To accomplish this, T4 DNA polymerase was used to incorporate dASTP (2'-deoxyadenosine 5'-O-(1-thiotriphosphate) into the ends of the linearized DNA. The thiol-DNA was then incubated with the heterobifunctional cross-linker, GMBS (2 mM) for 1 hour at room temperature. The excess GMBS was removed by centrifugation in microcentrifuge tubes at 10,000 rpm for 10 min. The DNA-GMBS was rinsed several times with PBS. Lastly, the DNA-GMBS was incubated with the antibody anti-FcγRII/RIII monoclonal antibody for 2 hours at room temperature. Figure 2 illustrates the fusion reaction using GMBS. The DNA-GMBS-antibody complex (DAPLEX) was applied to a G-10 column equilibrated in PBS. Seven fractions were collected, with fractions 2 and 3 showing increased absorbance at the wavelength of 260. Protein G was then used to absorb the DAPLEX. Fractions 2 and 3 were pooled and 0.25 μg and 0.5 μg of this material was added to 10 μl (20μg) of protein G. After incubation at 4°C for two hours, the protein G was pelleted and the supernatant removed and set aside for analysis. The protein G pellet was resuspended in 10 μl of protein loading buffer containing SDS, and analyzed along with supernatant by agarose gel electrophoresis with ethidium bromide staining. The results revealed that the supernatant contained mostly DNA and that very little DNA was associated with the pelleted fraction (Figure 3). This suggested that the conjugation efficiency between the DNA and monoclonal antibody was extremely poor.

Figure 2





Figure 3



The reason for the poor efficiency was not clear. One possibility is that the reaction to add the thiol group to the DNA vaccine failed due to the fact that we were attempting to label a large linear DNA molecule.


An alternative approach was taken to add a thiol group to the linearized DNA vaccine and involved the use of a commercially available end-labeling kit. It was thought that the use of a standardized kit would increase the efficiency of the reaction. This kit is marketed by Vector Labs Inc. and is called 5' EndTag Nucleic Acid Labeling System (MB 9001). The kit was reported to work also with DNA and RNA molecules in addition to short oligonucleotide primers. The thiol group was added to the linearized DNA according to the manufacture's instruction. A thiol-reactive label (Biotin maleimide) was then added and the material transferred to nitrocellulose membrane. To confirm successful labeling, the membrane was probed with alkaline phosphatase-streptavidin chemiluminescence kit. Despite multiple attempts, the absence of a positive signal indicated that labeling was unsuccessful.

In consultation with Dr. Andrew Lees in the USUHS Department of Medicine, another approach was pursued. Dr. Lees is an immunologist who has experience in antigen targeting, bioconjugation, vaccinology and polysaccharide chemistry. This new approach involved first conjugating the antibody to an oligonucleotide via an aldehyde linkage. Since oligonucleotides are more conducive to labeling, we anticipated greater success with this approach. The anti-FcγR<sub>II/III</sub> would then be linked to the oligo via the aldehyde group and subsequently ligated to the linearized DNA vaccine. To establish proof-of-principal, we attempted to conjugate the FcR antibody to a linearized plasmid that expresses the green fluorescent protein (gfp). To accomplish this, an aldehyde oligonucleotide was synthesized containing an Nsi I restriction site (aldehyde - 5' TGTAGATCCTAGGTACGATGCA 3') and then conjugated to the FcγR<sub>II/III</sub>

antibody. A complementary non-labeled oligonucleotide was synthesized (5' TCGTACCTAGGATCTACA 3') and annealed to the antibody-oligo conjugate, leaving a Nsi I overhang. The gfp plasmid was double-digested with Nsi I and BLN I, creating an Nsi I site downstream of the BGH gene and a BLN I site upstream of the gfp promoter. Multiple attempts were made to ligate that antibody-oligo to the linear gfp DNA at the Nsi I site, but we were unsuccessful as evidenced by lack of fluorescence in cells exposed to the post-ligation reaction product.


Another alternative approach was taken to construct the DNA vaccine-antibody conjugated vaccine. This approach involved bisaryl hydrazone conjugation chemistry. The DNA vaccine plasmid was first linearized with the restriction enzyme XhoI without disrupting the dengue envelope protein-coding region. Using terminal deoxynucleotide transferase, an aldehyde group was added to the linearized plasmid. A SANH molecule was then added to the anti- FcγRII/RIII antibody and the antibody-SANH product was then conjugated to the aldehyde modified linearized plasmid (Figure 4).

Figure 4



The next step was to determine the efficiency of conjugation. The product of the bisaryl hydrazone conjugation reaction was digested with DNase I to cleave the linearized plasmid off the antibody. This material was electrophoresed on a PAGE gel together with untreated conjugate. As a control, FITC Ab was treated with DNase I. The gel pattern showed that the protein signal from the treated conjugate was marginally stronger than the untreated material (Figure 5). This indicated that conjugation occurred, but at a low level.

Figure 5



Although it appeared that the conjugation efficiency was low, we next wanted to evaluate the uptake and expression of the small amount of antibody-DNA conjugate present in the reaction product. To accomplish this, three Fc-bearing cell lines were used; Raw 264.7 cells, P388D1 cells and J774 cells. Each cell line was incubated for two days at 37°C with antibody-DNA conjugate (0.3 µg and 1.0 µg) and controls that consisted of linearized DNA and saline. IFA was performed on the cells using anti-DEN 2H2 monoclonal antibody (detects expression of prM) and 4G2 (flavivirus-specific monoclonal antibody) to look for intracellular expression of the DNA vaccine. A portion of the cells was also pelleted, lysed and subjected to Western blot analysis in case there was low-level expression of the DNA vaccine not detected by IFA. The results showed no expression by IFA or by Western blot analysis.

Because it was theoretically possible that the DNA-antibody conjugate successfully bound to the cells but was not internalized, the cells were washed and subjected to PCR with primers specific for the DEN-1 envelope gene of the DNA plasmid. The appearance of a positive band in the P388D1 cells suggested that the DNA-antibody did attach to the Fc receptor but was not internalized. This finding was unanticipated and signified that more extensive work was needed for this approach to be successful. Additional funding to further research this vaccine development approach will be sought from DoD and Non-DoD agencies.

2. Co-immunize mice with low and high doses of DEN DNA vaccine with aluminum phosphate, tetanus toxoid and DNA vaccine linked with anti-FcγRI antibody.

Pilot mouse studies were conducted to evaluate the effect of aluminum phosphate (AP) on the immunogenicity of a DEN-1 DNA vaccine. Groups of six mice were immunized with 50 µg or 5 µg of DNA vaccine with or without 50 µg AP. The vaccines were administered both intramuscularly (IM) and intradermally (ID). The mice were primed and boosted on day 30. The results are indicated in item 3 below.

Murine studies were initiated to evaluate the effects of tetanus toxoid (TT) on the immunogenicity of the DEN-1 DNA vaccine. Nine groups of five mice each were given various

doses of DNA vaccine and TT as outlined in Table 1. Mice were primed and boosted on day 30. Serum samples were obtained at one, two and three months after administration of the last boost.

### **Table 1**

A (50µg ID)  
B (50µg + 0.1µg TT IM)  
C (50µg + 1.0µg TT IM)  
D (50µg + 10µg TT IM)  
E (5µg + 0.1µg TT IM)  
F (5µg + 1.0µg TT IM)  
G (5µg + 10µg TT IM)  
H (5µg IM)  
I (50µg IM)

A second mouse study was performed to confirm the results obtained from the first mouse experiment. For groupings and results for the second experiment, see item 3 below.

Murine studies to evaluate the immunogenicity of the DNA-Fc antibody were not initiated due to inefficient conjugation of the anti-FcγR antibody to the DEN-1 DNA.

3. Analyze antibody response by indirect ELISA and plaque reduction neutralization tests. Perform ELISPOT assays to detect T-cells producing interferon gamma (IFNγ) in response to DEN antigen stimulation to analyze dengue-specific cellular immune responses.


Mice immunized with the DEN-1 DNA vaccine with and without AP were evaluated for anti-DEN-1 antibody responses. To accomplish this, sera were screened for the presence of anti-DEN antibody using standard indirect ELISA as described previously [1] except that purified DEN-1 virus particles (West Pac strain) were used as antigen. Serum samples were diluted 1:100 in PBS, added to the plates and incubated at 37°C for one hour. The plates were washed and bound antibody was detected using peroxidase-labeled anti-mouse Ig antibody. At 45 and 57 days post-boost, there was no difference in anti-DEN neutralizing antibody levels between to AP group and the non-AP group when the vaccine was administered ID (Table 2). Aluminum phosphate did appear to slightly enhance the DNA vaccine when given IM. Earlier published studies showed that <12 µg of DEN DNA vaccine failed to elicit an ELISA antibody response [2]. When suboptimal doses of DNA vaccine (5µg) were given with AP, the neutralizing antibody titers were higher when the ID route was used compared to the IM route. Overall, aluminum phosphate had a minimal affect on the immunogenicity of the DEN-1 DNA vaccine as evidenced by the lack of improvement in neutralizing antibody when the vaccine was administered by the optimal route at the highest dose.

**Table 2**

VACCINE	ROUTE	DAY 45	DAY 57
50 µg I012D1 + 50 µg AP	IM	35	40
50 µg I012D1	IM	15	<10
5 µg I012D1 + 50 µg AP	IM	<10	13
50 µg I012D1 + 50 µg AP	ID	160	290
50 µg I012D1	ID	190	270
5 µg I012D1 + 50 µg AP	ID	65	70
Vector + 50 µg AP	IM	<10	<10
Vector + 50 µg AP	ID	<10	<10

A mouse study was initiated to evaluate the ability of TT to enhance antibody responses to the DEN-1 DNA vaccine. Analysis of sera obtained from mice at day 90 (approximately 60 days after the last immunization) revealed all doses of TT appeared to enhance anti-dengue ELISA antibody responses, compared to DNA alone, in all groups of animals immunized with 50 µg of DEN-1 DNA vaccine (Figure 6). At the lower dose of DNA vaccine (5 µg), there appeared to be very little effect.

Plaque reduction neutralization tests (PRNT) were performed to assess neutralizing antibody responses to the DEN-1 DNA vaccine and evaluate if adding TT had any effect. Samples for this analysis were obtained at day 120 (90 days after the last immunization). Sera from the mice in each group (n=6) were pooled and tested at serial two-fold dilutions starting at 1:20 and ending at 1:160. The assay was performed as described previously [3] with the 50% endpoint being calculated using log probit analysis (PRNT50). Anti-DEN-1 neutralizing antibodies were only detected in the groups that received the DEN-1 DNA vaccine ID and the DEN-1 DNA vaccine together with 10 µg of TT IM (reciprocal PRNT50 titers of 43 and 18, respectively). This indicated that there was some enhancement of the neutralizing antibody responses by TT. The responses however were not better than DNA alone when given ID.

**Figure 6**

\* Reciprocal PRNT50 titers.

Sera from the animals were analyzed to determine if immune responses were also generated against TT. To accomplish this, 96-well plates were coated with tetanus C fragment recombinant protein [4] at 50 ng per well, and reacted with mouse sera at a dilution of 1:100. Bound antibody was detected using peroxidase-labeled anti-mouse Ig. Anti-TT immune responses were detected by ELISA in all groups receiving TT (all doses) plus DNA vaccine (data not shown). OD values ranged from 3.5 to 3.8. No quantitative comparisons can be made between the anti-TT and anti-DEN-1 immune responses since both the ELISA assays were used only for qualitative analysis of antibody responses.

To confirm the above results indicating a slightly enhanced IM response when TT was used together with DNA vaccine, a repeat mouse study was conducted. The immunization schedule was identical to that of the first mouse study. The groups are indicated below in Table 3. In this repeat experiment, we also compared ID delivery of the vaccine to Biojector delivery. Other published studies showed that the Biojector enhances immune responses over traditional needle IM injection. In this repeat study, we also asked the question if TT can enhance immune responses to the DNA vaccine when administered ID. In the first mouse evaluation, only one dose (50µg) of the DNA vaccine was tested ID.

Table 3


Vaccine	Dose	TT dose	# Pos/Total	Delivery
1012D1ME	50µg	0	5/5	Biojector
1012D1ME	5µg	0	3/5	Biojector
1012D1ME	5µg	10µg	0/5	Biojector
1012D1ME	5µg	1µg	3/5	Biojector
1012D1ME	5µg	0.1µg	2/5	Biojector
1012D1ME	50µg	0	5/5	Needle ID
1012D1ME	5µg	0	5/5	Needle ID
1012D1ME	5µg	10µg	3/4	Needle ID
1012D1ME	5µg	1µg	4/4	Needle ID
1012D1ME	5µg	0.1µg	5/5	Needle ID
1012D1ME	50µg	10µg	2/4	Needle IM

In the group of animals given 50 µg DNA vaccine and 10 µg TT IM, only 50% of the animals responded by ELISA compared to 100% in the previous experiment. Of those that did respond, the OD values were comparable to the OD values in the previous experiment (data not shown).

In the earlier mouse experiment, 5 µg of DNA vaccine administered IM with and without TT, produce little to no ELISA antibody responses (Figure 4). When administered by Biojector, a greater mean ELISA antibody response was obtained (Figure 5), although not all animals responded (Table 3). The addition of TT appeared to have an inhibitory effect on the vaccine as indicated by a decrease in mean ELISA OD values in the groups of animals that received the DNA vaccine together with TT by Biojector. None of the mice given 5 µg DNA vaccine and 10 µg TT by Biojector responded.

A much better ELISA antibody response was seen when the 5 µg DNA vaccine dose was delivered by the ID route. However, TT appeared to again exhibit an inhibitory effect on the ELISA antibody response when given at 10 µg (Figure 7).

Figure 7




Analysis of neutralizing antibody (NT) responses revealed that TT only enhanced the neutralizing antibody response to 5 µg of DNA vaccine given ID with 1.0 µg and 0.1 µg TT (Table 4). The NT using the 1.0 µg dose of TT was longer lasting, but the levels declined between day 60 to day 220. The greatest NT response was achieved using 50 µg of DNA vaccine given ID.

Table 4

Vaccine	Dose	TT dose	Day 60 NT	Day 220 NT	Delivery
1012D1ME	50µg	0	109	15	Biojector
1012D1ME	5µg	0	<20	<20	Biojector
1012D1ME	5µg	10µg	<20	<20	Biojector
1012D1ME	5µg	1µg	<20	<20	Biojector
1012D1ME	5µg	0.1µg	<20	<20	Biojector
1012D1ME	50µg	0	<b>145</b>	<b>115</b>	Needle ID
1012D1ME	5µg	0	<20	<20	Needle ID
1012D1ME	5µg	10µg	<20	<20	Needle ID
1012D1ME	5µg	1µg	<b>46</b>	<b>18</b>	Needle ID
1012D1ME	5µg	0.1µg	<b>81</b>	<20	Needle ID
1012D1ME	50µg	10µg	<20	<20	Needle IM

At day 220, the mice were boosted with a third dose and two weeks later the spleens from each mouse were harvested and spleen cell suspensions prepared for IFN $\gamma$  ELISPOT analysis. A portion of the cells was stimulated for 36 hours with irradiated A20 cells that express DEN-1 envelope antigen and another portion stimulated with TT. The results were expressed as the number of spots generated per  $2 \times 10^5$  spleen cells. The overall IFN $\gamma$  T cell responses to both the DEN DNA vaccine and the TT were poor for all vaccine combinations, with the DEN-specific responses being slightly higher (Figure 8) compared to the TT-specific responses. The best IFN $\gamma$  T cell response was seen in the group of animals that received 50  $\mu$ g of DNA vaccine alone via Biojector. From these data it appears that TT does not improve T cell responses to the vaccine.

Figure 8



## Year 2

### Summary Statement of Work

1. Immunize mice with DNA vaccine and increasing concentrations of aluminum phosphate and tetanus toxoid (TT).
2. Analyze antibody response by indirect ELISA and plaque reduction neutralization tests. Perform ELISPOT assays to detect T-cells producing IFN $\gamma$  and IL-4 to analyze cellular immune responses.

3. Measure uptake and expression of DEN DNA vaccine formulations in vivo.

### **Progress and Accomplishments**

Earlier studies indicated that little to no adjuvant effect was seen using 50 µg of aluminum phosphate when given together with 50 µg of DNA vaccine, although some benefit was observed if the IM route of administration was used. In light of these findings, further mouse dose titration studies with aluminum phosphate were not felt to be warranted. See year-two progress for TT titration experiments. Studies to examine the uptake and expression of the DNA vaccines in vivo, with and without adjuvants, were not performed due to the lack of success with constructing an immunogenic experimental lot of DNA-antibody complex and due to lack of an adjuvant-induced improvement in the immune responses to the DNA vaccine.

### **Year 3**

#### **Statement of Work**

1. Immunize monkeys with two DEN DNA vaccine formulations that were shown to be the most potent in murine studies. If all three formulations show similar potency, then four additional monkeys will be added so that all three DNA vaccine formulations can be evaluated.
2. Analyze antibody responses to the vaccines using ELISA and plaque reduction neutralization tests. ELISPOT assays to detect T-cells producing IFN $\gamma$  (Th1 response) and IL-4 (Th2 response) will be used to analyze cellular immune responses.
3. At six months post-priming, challenge all monkeys with a low passage live DEN-2 virus to evaluate protective efficacy.

### **Progress and Accomplishments**

The vaccine formulations produced and evaluated under this grant did not display any enhanced immunogenicity in mice over the DNA vaccine used alone. Therefore expensive non-human primate studies were not performed.

### **KEY RESEARCH ACCOMPLISHMENTS**

- Anti-Fc $\gamma$ R antibodies that recognize P388D1 and Raw 264.7 cells were identified.
- P388D1 cells failed to express prM and E proteins from the linear DNA vaccine plasmid but the Raw264.7 cells efficiently expressed the proteins. The Raw 264.7 cells will be used in the in vitro binding studies.
- Aluminum Phosphate was demonstrated to slightly enhance anti-dengue neutralizing antibody responses, only when the DNA vaccine was given ID at suboptimal doses. There was essentially no effect when the vaccine was administered IM.
- The antibody responses to the DEN 1 DNA vaccine given IM were enhanced slightly by combining it with TT. The ELISA and neutralizing antibody levels were inferior to the levels obtained with ID administration of the vaccine alone. The Animals also responded to the TT component as illustrated by the detection of anti-TT antibodies by ELISA.
- Administering TT at the highest dose (10 µg) appeared to have an inhibitory effect on the antibody response to the dengue-1 DNA vaccine when administered ID. A more

pronounced inhibitory effect was observed when the DNA vaccine/TT combination was administered by needless Biojector.

#### REPORTABLE OUTCOMES

Studies conducted in mice to date demonstrated that the addition of AP and TT to a DEN-1 DNA vaccine study results in slight improvement in antibody responses when administered IM. Biojector administration of the vaccine (5 µg) with TT at the lower doses (0.1 µg and 1 µg) appeared to enhance anti-dengue antibody responses compared to IM injection of the DNA alone. When compared to ID administration of the DNA vaccine alone, there has been no overall improvement in the neutralizing antibody responses obtained by combining the DNA vaccine with either TT or aluminum phosphate.

#### CONCLUSIONS

1. TT and AP appear to have some adjuvant effect on the neutralizing antibody responses to a DEN-1 prME DNA vaccine when given IM. These antibody responses however were not superior to the antibody responses elicited when the same DNA vaccine was given alone by the ID route.
2. Anti-FcγRII/III monoclonal antibodies were found that recognize Raw 264.7 mouse monocyte/macrophage cells. The efficiency of linking the monoclonal antibodies to linearized DNA vaccine was poor despite using multiple approaches to produce the dengue 1 DNA vaccine-antibody complex.

#### REFERENCES

1. Raviprakash K, Kochel TJ, Ewing D, et al. Immunogenicity of dengue virus type 1 DNA vaccines expressing truncated and full length envelope protein. *Vaccine* 2000;18:2426-34
2. Porter KR, Kochel TJ, Wu SJ, Raviprakash K, Phillips I and Hayes CG. Protective efficacy of a dengue 2 DNA vaccine in mice and the effect of CpG immuno-stimulatory motifs on antibody responses. *Arch Virol* 1998;143:997-1003
3. Russell P, Nisalak A. Dengue Virus Identification by the Plaque Reduction Neutralization Test. *J. Immunol.* 1967;99:291-296
4. Saikh KU, Sesno J, Brandler P and Ulrich RG. Are DNA-based vaccines useful for protection against secreted bacterial toxins? Tetanus toxin test case. *Vaccine* 1998;16:1029-38