

# REPORT DOCUMENTATION PAGE

*Form Approved*  
OMB No. 0704-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.

1. AGENCY USE ONLY <i>(Leave blank)</i>	2. REPORT DATE August 2006	3. REPORT TYPE AND DATES COVERED Journal Article	
4. TITLE AND SUBTITLE Field studies of exercise and food deprivation.		5. FUNDING NUMBERS	
6. AUTHOR(S) Reed W. Hoyt and Karl E. Friedl		8. PERFORMING ORGANIZATION REPORT NUMBER M06-40	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Biophysics and Biomedical Modeling Division U.S. Army Research Institute of Environmental Medicine Building 42 - Kansas Street Natick, MA 01760		10. SPONSORING / MONITORING AGENCY REPORT NUMBER	
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) U.S. Army Medical Research and Materiel Command Fort Detrick, MD 21702		11. SUPPLEMENTARY NOTES	
12a. DISTRIBUTION / AVAILABILITY STATEMENT Approved for public release; distribution unlimited		12b. DISTRIBUTION CODE	
13. ABSTRACT <i>(Maximum 200 words)</i> Purpose of review The increase in obesity in developed societies drives interest in the interplay of energy intake, metabolic energy expenditure, and body energy stores. A better understanding of energy management in physically active and undernourished humans should help guide strategies to manage obesity safely and effectively. This review focuses on field studies of men and women engaged in prolonged strenuous activities, ranging from ranger training to extreme expeditions. Recent findings Although scientifically unconventional and limited, field studies of exercise and food deprivation have yielded interesting findings: 45% body fat is the normal lower limit to fat reserves in physically active underfed young adult men, and in response to exercise and underfeeding, women used more fat mass and less fat-free mass to meet metabolic fuel requirements. Summary Field studies have shown that fat energy reserves in young adult men can be estimated as percentage body fat minus 5%, and initial body fat mass has a significant positive influence on fat oxidation rates per kilogram of fat-free mass during rapid weight loss associated with underfeeding and exercise. Data logging pedometers, activity monitors, global positioning systems, and wireless body and personal-area networks promise to make it easier to study and care for free-living humans.			
14. SUBJECT TERMS body composition, free-living, negative energy balance, physical work, semi-starvation, soldiers			15. NUMBER OF PAGES 6
17. SECURITY CLASSIFICATION OF REPORT Unclassified			16. PRICE CODE
18. SECURITY CLASSIFICATION OF THIS PAGE Unclassified	19. SECURITY CLASSIFICATION OF ABSTRACT Unclassified	20. LIMITATION OF ABSTRACT Unlimited	

# Field studies of exercise and food deprivation

Reed W. Hoyt and Karl E. Friedl

## Purpose of review

The increase in obesity in developed societies drives interest in the interplay of energy intake, metabolic energy expenditure, and body energy stores. A better understanding of energy management in physically active and undernourished humans should help guide strategies to manage obesity safely and effectively. This review focuses on field studies of men and women engaged in prolonged strenuous activities, ranging from ranger training to extreme expeditions.

## Recent findings

Although scientifically unconventional and limited, field studies of exercise and food deprivation have yielded interesting findings: 4–5% body fat is the normal lower limit to fat reserves in physically active underfed young adult men, and in response to exercise and underfeeding, women used more fat mass and less fat-free mass to meet metabolic fuel requirements.

## Summary

Field studies have shown that fat energy reserves in young adult men can be estimated as percentage body fat minus 5%, and initial body fat mass has a significant positive influence on fat oxidation rates per kilogram of fat-free mass during rapid weight loss associated with underfeeding and exercise. Data logging pedometers, activity monitors, global positioning systems, and wireless body and personal-area networks promise to make it easier to study and care for free-living humans.

## Keywords

body composition, free-living, negative energy balance, physical work, semi-starvation, soldiers

## Introduction

Highly motivated athletes, soldiers, and adventurers routinely participate in activities that demand near-maximal rates of sustained metabolic energy production. For both practical and ethical reasons, field studies are often the only way to study these individuals. We review studies of some of the more extreme events, focusing on field studies of soldiers in training. Additional information on the nutritional factors influencing the physical and mental performance of military personnel under extreme environmental conditions is provided by the National Academy of Sciences [1].

Scientists have studied soldiers in training in the hope of better understanding how exercise and food restriction impact health and physical performance [2]. The classic Minnesota starvation study [3] documented weight loss, body composition and physiological changes over 24 weeks in a group of largely sedentary normal male subjects. The effects of underfeeding have also been investigated in soldiers participating in military training courses in which study durations are shorter, at 3–62 days, and sleep deprivation and sustained physical activity are added stressors. Study populations range from Norwegian cadets participating in a training course that involves a week of food and sleep deprivation and approximately 23 h/day of marching and other activities [4,5,6<sup>\*</sup>], to soldiers participating in 8 weeks of physically demanding US Army Ranger training, in which sleep is restricted to 4 h/day and food is in short supply [7,8]. Others have studied long-range patrols by Swedish ski troops [9], hot weather training by Zimbabwean recruits [10], and a variety of other extremes [11–15] (see Table 1).

## Norwegian ranger training

The dramatic physiological effects of 7 days of strenuous exercise, approximately 1 h/day of sleep and little food on the endocrine status and body composition of Norwegian military academy cadets have been studied by Opstad and colleagues [5,6<sup>\*</sup>,16]. The markedly negative energy balance, and not sleep deprivation, appeared to be the primary factor driving changes in endocrine status in the cadets. Substantial reductions in metabolic regulators, such as insulin, androgens, and thyroid hormones, were accompanied by compensatory increases in growth hormone and adrenergic responses to exercise [5]. Male cadets appeared to lose fat from gluteal adipocytes before abdominal sites, with little change in femoral fat [16]. Male cadets had total energy expenditures (TEEs;  $27 \pm 2$  MJ/day) that exceeded those of the female

Curr Opin Clin Nutr Metab Care 9:685–690. © 2006 Lippincott Williams & Wilkins.

US Army Research Institute of Environmental Medicine, Natick, Massachusetts, USA

Correspondence to Reed W. Hoyt, PhD, Biophysics and Biomedical Modeling Division, US Army Research Institute of Environmental Medicine, Kansas Street, Natick, MA 01760-5007, USA  
Tel: +1 508 233 4802; fax: +1 508 233 5298; e-mail: reed.hoyt@us.army.mil

Disclaimer: The opinions and assertions contained in this paper are the personal views of the authors and do not necessarily represent the official views or policy of the US Department of the Army.

**Current Opinion in Clinical Nutrition and Metabolic Care** 2006, 9:685–690

## Abbreviations

**FFM** fat-free mass  
**IGF-1** insulin-like growth factor 1  
**TEE** total energy expenditure


© 2006 Lippincott Williams & Wilkins  
1363-1950

**Table 1** Examples of notable field studies of humans engaged in prolonged strenuous activities

Event (ref.)	Description	TEE (AEE) (MJ/day; kJ/kg per day)	Food energy intake (MJ/day)	ΔEnergy stores (ΔBW; ΔFFM)	Key findings
Tour de France bicycle race [11]	Elite male cyclists ( $n=4$ ); 22 days of high intensity exercise; well fed and normal sleep	DLW 28.4–38.4 MJ/day (individual)	21.5–27.1 MJ/day free food; self-reported food records (26% fat)	ΔBW ~stable	Established upper limit of voluntarily sustainable energy expenditure (PAL 4.3–5.3) I/B inaccurate because of underreported food intake
Norwegian military training exercise [6*]	Military academy cadets ( $n=10$ men & 6 women), 7 days of little or no food & continuous work (23 h/day)	M 26.6 ± 2.0 W 21.9 ± 2.0 MJ/day M 343 ± 26 W 354 ± 18 kJ/kg per day	0.2–1.9 MJ/day	Men ΔBW –7.5 ± 1.1 kg ΔFM –3.5 ± 0.7 kg ΔFFM –4.0 ± 1.2 kg %BF <sub>final</sub> 12.7 ± 3.3  Women ΔBW –6.0 ± 1.3 kg ΔFM –3.4 ± 0.2 kg ΔFFM –2.6 ± 1.1 kg %BF <sub>final</sub> 22.6 ± 4.8 (DEXA)	Characterized endocrine and immunological responses to energy restriction and sustained effort in healthy young men Sex difference in fuel use: ♀ used more fat and less FFM than ♂ to meet TEE needs BUT similar EE/kg body weight
US Army Ranger course [7]	Male soldiers ( $n=55$ ), 56 days of 20 h/day work with semi starvation & continuous work (20.4 h/day)	DLW 16.7 ± 3.5 (13.1–25.0, depending on terrain) MJ/day	11.7 MJ/day (34% fat)	ΔBW –12.1 ± 3.4 kg ΔFM –4.8 ± 2.4 ΔFFM –7.3 ± 3.2 %BF <sub>final</sub> 5.8 ± 1.8 (DEXA)	Established normal lower limit of body fat in healthy young men with uncomplicated weight loss Demonstrated impact of energy deficit on indices of immune function
G-2 trans-Greenland expedition [12]	Two fit 25-year-old Norwegian soldiers; 86-day trek, 3000 km (~9 h/day)	Flat terrain: 14.6, 16.1 MJ/day Rugged terrain: 28.3, 34.6 MJ/day	25.1 MJ/day High fat diet (60% fat)	ΔBW –1.1, –8.6 kg ΔFM –1.6, –7.0 kg ΔFFM –0.6, –1.4 kg %BF <sub>final</sub> 12.7, 13.5 (DEXA)	Strenuous activity can be sustained with adequate food and sleep Body composition of subjects converged with common tasks & diet
Zimbabwean commando training [10]	Soldiers ( $n=8$ ) 12 days of physically demanding (~8 h/day) strenuous hot weather activity (40°C avg daytime temp)	DLW 23 ± 4.2 I/B 26 ± 2.0	17.0 ± 0.8 MJ/day (35% fat)	ΔBW –3.0 ± 0.1 kg ΔFM –3.0 ± 0.3 ΔFFM –0.3 ± 3.0 (skinfolds)	Documented negative energy balance in hot weather training
Trans-Antarctic expedition [13,14]	Two fit adventurers (48 & 37 years old), 95 days ~2300 km trek across Antarctica	DLW 29.6, 24.1 I/B 29.07.3	21.3 MJ/day (57% fat)	ΔBW –24.6, –21.8 kg ΔFM –16.8, –12.5 kg ΔFFM –7.8, –9.3 kg %BF <sub>final</sub> 1.9, 2.5 (UWW)	Demonstrated high level of voluntarily tolerable rate of lean mass catabolism in healthy men
US Marine Corps 'Crucible' recruit training exercise [15]	Male ( $n=29$ ) and female ( $n=20$ ) recruits, 54 h, high-intensity, moderate food restriction	M 25.7 ± 0.8 W 19.8 ± 0.6 MJ/day M 350 ± 40 W 340 ± 40 kJ/kg per day	M 6.0 ± 2.0 MJ/day W 4.8 ± 1.8 MJ/day (36–37% fat)	Men ΔBW –3.1 ± 0.8 kg %BF <sub>final</sub> 15.7 ± 3.3  Women ΔBW –1.6 ± 0.5 kg %BF <sub>final</sub> 26.3 ± 3.2 (skinfolds)	Men and women performing similar high intensity activities had similar PALs and similar weight-specific EEs

AEE, Average energy expenditure; BF, body fat; BW, body weight; DEXA, dual X-ray energy absorptiometry; DLW, doubly labelled water method; EE, energy expenditure; FFM, fat-free mass; FM, fat mass; I/B, intake/balance method of estimating EE; M, men; PAL, physical activity level = TEE/resting metabolic rate; TEE, total energy expenditure; UWW, underwater weighing or hydrodensitometric method of estimating body composition; W, women.

**Figure 1** Hattori plot illustrating mean ( $\pm$ SD) and individual fat mass and fat-free mass values for women (triangles) and men (circles) at the start (solid symbols) and end (open symbols) of 7 days of exercise and starvation in a Norwegian military training course




Body composition was measured by dual X-ray energy absorptiometry. The women lost less fat-free mass (FFM) for a given change in fat mass (FM) than the men. This figure has not been previously published and is based on data from Hoyt *et al.* [6\*].

cadets ( $22 \pm 2$  MJ/day), but weight-specific TEEs ( $\sim 350$  kJ/kg per day), and physical activity levels (TEE/resting metabolic rate =  $\sim 3.5$ ) were similar [6\*]. A study of Marine recruits participating in a demanding physically demanding 54 h training exercise also found that men and women performing similar high-intensity activities had similar weight-specific TEEs (men  $350 \pm 40$  kJ/kg per day; women  $340 \pm 40$  kJ/kg per day) [15]. Notably, the relative contribution of fat mass to TEE among the Norwegian cadets was nearly 90% in the women compared with 74% in the men, indicating marked sex differences in fat substrate utilization during this demanding training [6\*]. A Hattori plot of the fat mass index compared with the fat-free mass (FFM) index illustrates the differences in body composition and fuel use between the male and female cadets (Fig. 1), with the women clearly losing less FFM for a given change in fat mass than the men. This observation is consistent with a variety of laboratory studies showing that women use more fat than men to meet the energy demands of exercise [17].

### United States Army Ranger training

The 2-month-long US Army Ranger training course provided a prolonged period of exercise and nutritional privation over which to study endocrine and body composition changes [7,8,18]. Over 62 days of ranger training, TEEs averaged 16.7 MJ/day whereas dietary energy intake was only approximately 11.7 MJ/day. These motivated young men voluntarily reached the apparent

**Figure 2** Decrease in body fat distribution among a group of soldiers, in response to 8 weeks of exercise and semi-starvation, converging on an apparent minimum percent body fat



The three distributions reflect the percentage of body fat obtained by dual X-ray energy absorptiometry measurements for the same 50 men at the start of the course and at 6 and 8 weeks. Approximately half of the men had lost all available fat by week 6 and were dependent on body protein stores for the remaining 2 weeks. This figure has not previously been published and is based on data from Friedl *et al.* [7].

normal lower limit of 4–5% body fat (fat mass  $4 \pm 2$  kg; Fig. 2). We had expected these soldiers to have more modest energy deficits and perhaps achieve as much as a 10% body weight loss over the 8-week course. We were astonished to discover an average 16% loss of body weight, with an extreme weight loss of 23% in one particularly lean individual [7]. This rate of weight loss was at least twice that of the Minnesota starvation study [3], in which male volunteers were taken to 24% body weight loss over 24 weeks, and half the rate of weight loss reported for Irish hunger strikers in 1981, in which 10 out of 30 individuals died at an average 62 days (range 48–72 days to death) with an average weight loss of approximately 30% (26–38%) at the time of death [19]. These data suggest that with prolonged negative energy balance, both the rate of weight loss and the total amount of body weight lost are important risk factors.

Although the amount of body weight lost in response to Army Ranger training was substantial, it reflected an uncomplicated energy deficit, with no evidence of other nutritional deficiencies [20]. In addition, grip strength was unchanged, and general strength capacity only decreased modestly [21]. Physiological changes in the Ranger students suggested a shift to increased metabolic conservation, notably subclinical suppression of the thyroid axis [8], low core body temperatures [22], greater susceptibility to hypothermia [23], and apparently increased economy of motion [24]. Except for an increased susceptibility to bacterial infection associated with marked effects on indicators of immune function [24], these young men were surprisingly tolerant of large energy deficits and loss of body mass.

Naturally, periods of exercise and energy deprivation that commonly occur with military field training [25<sup>•</sup>] need to be balanced by rest and refeeding [26]. Ranger students, after 5 weeks of limited physical activity and free food consumption, were able to recover fully from the approximately 250 MJ energy deficit and approximately 12% loss of body mass incurred during the 8 weeks of training. After recovery, FFM and physical performance returned to pretraining levels, whereas final fat mass exceeded initial levels [21].

### Underfeeding constraints on voluntary energy expenditure

Soldiers provided an additional 1.7 MJ/day supplement during the 8-week Ranger course converted approximately half of the additional food energy to body energy stores, and used the remainder to fuel higher energy expenditure [8]. This presumably reflects a lifting of the ceiling on TEE that is imposed by inadequate intakes, as in studies of undernourished Columbian school children playing soccer alongside adequately nourished children [27], when energy intake appeared to limit work performance. Earlier studies of laborers, such as sugar cane cutters and road builders, also support the idea that inadequate energy intake constrains energy expenditure [2].

### Body composition influence on fuel metabolism

During weight loss in Army Ranger students, the proportion of fat-to-lean mass loss was related to initial adiposity ( $R^2 = 0.42$ ;  $n = 105$  men with initial 6–26% body fat), indicating that under conditions of extreme energy restriction, men preserve lean mass more effectively if they have a greater initial body fat availability [18]. Whereas the fatter Ranger students derived the majority of their energy from body fat stores, the contribution of fat mass to fuel needs in the leanest individuals was only approximately 20% [18]. The positive influence of initial fat mass on fat oxidation was similarly evident in the starved and physically active Norwegian cadets, in whom fat oxidation per kilogram of FFM was correlated with initial fat mass ( $R^2 = 0.51$ ) [6<sup>•</sup>]. In the Army Rangers, testosterone had little influence on this pattern of weight loss. Average testosterone levels were generally depressed and only weakly correlated ( $R^2 = 0.10$ ) with the ratio of fat-to-lean loss.

### Other consequences of underfeeding and exercise

In a 3-week study of Swedish ski troops, who had estimated TEEs that reached approximately 26 MJ/day but only consumed approximately 18 MJ/day of prepackaged field rations, a preferential sacrifice of fast-twitch type II muscle fibers was evident [9]. Prolonged exercise alone does not appear to produce this effect [28]. This

reduction in fast-twitch muscle fibers is probably a consequence of a decline in activity in the thyroid axis [29], as has been noted in field studies of prolonged work with inadequate dietary intakes [8,30]. Similar findings are evident in obese patients on a very low calorie diet [31]. Other metabolic changes, such as a progressive increase in total cholesterol levels in the Ranger students and increasing insulin resistance in Norwegian cadets, may be related to other counterregulatory mechanisms directed towards preserving lean mass in the face of large energy deficits [8,32].

In a study of soldiers participating in 5 days of French commando training in the Pyrenees, serum testosterone and insulin concentrations progressively declined with decreasing energy intake (the diets provided 7.5, 13.4, or 17.6 MJ/day) and increasing weight losses; plasma free fatty acids and  $\beta$ -hydroxybutyrate were elevated in the group with the greatest deficit [33]. In the Norwegian cadets, 5 days of continuous work without food produced marked increases in free fatty acids and  $\beta$ -hydroxybutyrate that became even more pronounced in response to bicycle exercise on days 3 and 5 [34]. Circulating insulin-like growth factor 1 (IGF-1) appears to be a key metabolic marker of hypocaloria, or more specifically, protein intake and anabolic status. In the Ranger studies, IGF-1 declined substantially [8], but in a shorter-term laboratory study simulating sustained military operations [35], IGF-1 appeared to reflect a more complicated balance of exercise, macronutrient intakes, and energy balance. The physiological responses to hypocaloria, such as elevated growth hormone, increased insulin resistance, and a shift to increased fat utilization, appear to be enhanced by concurrent physical activity.

### Exercise, diet and reproductive function

Controlled laboratory studies of normal women have forced a reconsideration of dogma that high intensity exercise suppresses reproductive function [36]. These recent studies clearly indicate that luteinizing hormone pulsatility is affected by energy deficits and not by exercise *per se*. This is consistent with observations that menstrual dysfunction is not more prevalent in elite female athletes during periods of high-intensity training [37]. Female army recruits working at a lower exercise intensity, with TEEs averaging 11.7 MJ/day, also do not have an increase in menstrual dysfunction, even among the large subset of these women who had in excess of 35% body fat and lost more than 3 kg of fat weight (as measured by dual X-ray energy absorptiometry) over the 2 months of military training [38]. The observations of Loucks [39] will help ensure that women are not excluded from high physical demand job specialties on the basis of concerns for reproductive health or potential bone loss. More importantly to this review, these findings can be used in designing weight loss routines that have a

minimal impact on reproductive function. The role of macronutrient composition on reproductive function has been examined, and even for women training intensely, carbohydrate intake and not protein consumption appears to be a key driver of the observed changes.

### Extreme treks

We have learned from studies of individuals that adequate nutrition is essential to physiological wellbeing at high levels of energy expenditure. On one hand, Mike Stroud and Ranulph Fiennes did not ingest enough food (21.3 MJ/day) and suffered more than 20 kg weight losses over the course of their harsh 2300 km crossing of the Antarctic continent [13,14], whereas Rune Gjeldnes and Torry Larsen lost little weight and suffered no performance breakdowns during a 3000 km crossing of Greenland while consuming at least 25.1 MJ/day in soy oil and oatmeal [12]. The dramatic weight losses and muscle wasting reported in some trekkers can often be ascribed to poor planning and misadventures that include deficient diets. The success of Tour de France cyclists [11], the Greenland trekkers and others suggests that an adequate diet and appropriate work/rest cycles can minimize any degradation in endurance capacity.

### New technologies for studying free-living humans

Pedometry, heart rate monitoring, accelerometry or actigraphy, and differential global positioning systems are among the various methods used individually to characterize TEE, and more specifically the intensity, duration, and frequency of physical activity. Combining these methods can yield interesting benefits. For example, heart rate and foot-ground contact times measured simultaneously during a brief bout of running can be used to estimate maximal aerobic capacity accurately [40]. This is more practical than using a 2-mile maximal effort run time [41] or laboratory treadmill stress testing, and may be a useful way to follow changes in aerobic fitness over time. Similarly, data from multiple accelerometers, placed on legs, torso, and arms, can be used to estimate metabolic energy expenditure and identify activities and activity patterns [42]. In the future, chronic monitoring of free-living individuals will become more practical as wireless body area networks eliminate wires and enable real-time data analysis and remote assessment of physiological status [43].

### Conclusion

Field studies have made important contributions to the scientific understanding of human energy balance, for example, by identifying the normal lower limit to body fat in physically active underfed young men, and helping to show that men and women have different patterns of body fat and lean mass loss during exercise and under-feeding. Future field studies are likely to be fruitful as

scientists apply new methods of quantifying aerobic fitness and exercise intensity and duration.

### References and recommended reading

Papers of particular interest, published within the annual period of review, have been highlighted as:

- of special interest
- of outstanding interest

Additional references related to this topic can also be found in the Current World Literature section in this issue (p. 770).

- 1 Military nutrition research projects. Food and Nutrition Board of the National Academy of Sciences. Available at: <http://www.iom.edu/CMS/3788/4615.aspx>. Accessed: July 2006.
- 2 Spurr GB. Physical work performance under conditions of prolonged hypocaloria. In: Committee on Military Nutrition Research, Predicting Decrements in Military Performance due to Inadequate Nutrition. Washington, DC: National Academy Press; 1986. pp. 99–135.
- 3 Keys A. The biology of human starvation. Minneapolis: University of Minnesota Press; 1950.
- 4 Opstad PK, Aakvaag A. The effect of a high calory diet on hormonal changes in young men during prolonged physical strain and sleep deprivation. *Eur J Appl Physiol Occup Physiol* 1981; 46:31–39.
- 5 Opstad PK. Medical consequences in young men of prolonged physical stress with sleep and energy deficiency. Report no. NDRE-95/05586. Kjeller, Norway: Norwegian Defence Research Establishment; December 1995. 177 pp.
- 6 Hoyt RW, Opstad PK, Haugen AH, *et al*. Negative energy balance in male and female rangers: effects of 7 days of sustained exercise and food deprivation. *Am J Clin Nutr* 2006; 83:1068–1075.
- A comparison of fuel utilization in underfed fit young women and men working at the limits of endurance.
- 7 Friedl KE, Moore RJ, Martinez-Lopez LE, *et al*. Lower limits of body fat in healthy active men. *J Appl Physiol* 1994; 77:933–940.
- 8 Friedl KE, Moore RJ, Hoyt RW, *et al*. Endocrine markers of semistarvation in healthy lean men in a multistressor environment. *J Appl Physiol* 2000; 88:1820–1830.
- 9 Schantz P, Henriksson J, Jansson E. Adaptation of human skeletal muscle to endurance training of long duration. *Clin Physiol* 1983; 3:141–151.
- 10 Mudambo KS, Scrimgeour CM, Rennie MJ. Adequacy of food rations in soldiers during exercise in hot, day-time conditions assessed by doubly labelled water and energy balance methods. *Eur J Appl Physiol Occup Physiol* 1997; 76:346–351.
- 11 Westertep KR, Saris WH, van Es M, ten Hoor F. Use of the doubly labeled water technique in humans during heavy sustained exercise. *J Appl Physiol* 1986; 61:2162–2167.
- 12 Frykman PN, Harman EA, Opstad PK, *et al*. Effects of a three-month endurance event on physical performance and body composition: the G2 Trans-Greenland Expedition. *Wilderness Environ Med* 2003; 14:240–248.
- 13 Stroud MA, Ritz P, Coward WA, *et al*. Energy expenditure using isotope-labelled water (2H2 18O), exercise performance, skeletal muscle enzyme activities and plasma biochemical parameters in humans during 95 days of endurance exercise with inadequate energy intake. *Eur J Appl Physiol* 1997; 76:243–252.
- 14 Stroud MA, Jackson AA, Waterlow JC. Protein turnover rates of two human subjects during an unassisted crossing of Antarctica. *Br J Nutr* 1996; 76:165–174.
- 15 Castellani JW, Delany JP, O'Brien C, *et al*. Energy expenditure in men and women during 54 h of exercise and caloric deprivation. *Med Sci Sports Exerc* 2006; 38:894–900.
- 16 Rognum TO, Rodahl K, Opstad PK. Regional differences in the lipolytic response of the subcutaneous fat depots to prolonged exercise and severe energy deficiency. *Eur J Appl Physiol* 1982; 49:401–408.
- 17 Tarnopolsky MA. Gender differences in lipid metabolism during exercise and at rest. In: Tarnopolsky M, editor. Gender differences in metabolism: practical and nutritional implications. Nutrition in exercise and sport, volume 14. Boca Raton: CRC Press; 1999. pp. 179–199.
- 18 Friedl KE. Variability of fat and lean tissue loss during physical exertion with energy deficit. In: Kinney JM, Tucker HN, editors. Physiology, stress, and malnutrition: functional correlates, nutritional intervention. New York: Lippincott-Raven Publishers; 1997. pp. 431–450.
- 19 Marriott BM, editor. Review of the results of nutritional intervention, U.S. Army Ranger Training Class 11/92 (Ranger II). National Academy Press: Washington DC. 1993.

- 20 Moore RJ, Friedl KE, Tulley RT, Askew EW. Maintenance of iron status in healthy men during an extended period of stress and physical activity. *Am J Clin Nutr* 1993; 58:923–927.
- 21 Nindl BC, Friedl KE, Frykman PN, *et al.* Physical performance and metabolic recovery among lean, healthy men following a prolonged energy deficit. *Int J Sports Med* 1997; 18:317–324.
- 22 Hoyt RW, Young AJ, Matthew WT, *et al.* Warfighter Physiological Status Monitoring (WPSM): body core temperatures during 96 h of swamp phase Ranger training. Technical Report T97-4 (AD A323018). US Army Research Institute of Environmental Medicine: Natick, MA 01760 USA; 1997.
- 23 Young AJ, Castellani JW, O'Brien C, *et al.* Exertional fatigue, sleep loss, and negative energy balance increase susceptibility to hypothermia. *J Appl Physiol* 1998; 85:1210–1217.
- 24 Friedl KE. Military nutritional immunology. In: Hughes DA, Darlington LG, Bendich A, editors. *Diet and human immune function*. Totowa, New Jersey: Humana Press Inc; 2003. pp. 381–396.
- 25 Tharion WJ, Lieberman HR, Montain SJ, *et al.* Energy requirements of military personnel. *Appetite* 2005; 44:47–65.  
This is a comprehensive review of energy balance of soldiers in the field.
- 26 Barboza PS, Hume ID. Physiology of intermittent feeding: integrating responses of vertebrates to nutritional deficit and excess. *Physiol Biochem Zool* 2006; 79:250–264. [E-pub ahead of print 7 February 2006]
- 27 Spurr GB, Reina JC. Influence of dietary intervention on artificially increased activity in marginally undernourished Columbian boys. *Eur J Clin Nutr* 1988; 42:835–846.
- 28 Helge JW, Lundby C, Christensen DL, *et al.* Skiing across the Greenland icecap: divergent effects on limb muscle adaptations and substrate oxidation. *J Exp Biol* 2003; 206:1075–1083.
- 29 Caizzo VJ, Baker MJ, Baldwin KM. Novel transitions in MHC isoforms: separate and combined effects of thyroid hormone and mechanical unloading. *J Appl Physiol* 1998; 85:2237–2248.
- 30 Opstad PK, Falch D, Oktedalen O, *et al.* The thyroid function in young men during prolonged exercise and the effect of energy and sleep deprivation. *Clin Endocrinol* 1984; 20:657–669.
- 31 Russell DM, Walker PM, Leiter LA, *et al.* Metabolic and structural changes in skeletal muscle during hypocaloric dieting. *Am J Clin Nutr* 1984; 39: 503–513.
- 32 Opstad PK. Alterations in the morning plasma levels of hormones and the endocrine responses to bicycle exercise during prolonged strain. The significance of energy and sleep deprivation. *Acta Endocrinol* 1991; 125: 14–22.
- 33 Guezennec CY, Satabin P, Legrand H, Bigard AX. Physical performance and metabolic changes induced by combined prolonged exercise and different energy intakes in humans. *Eur J Appl Physiol* 1994; 68:525–530.
- 34 Rognum TO, Vaage O, Hostmark A, Opstad PK. Metabolic responses to bicycle exercise after several days of physical work and energy deficiency. *Scand J Clin Lab Invest* 1981; 41:565–571.
- 35 Nindl BC, Castellani JW, Young AJ, *et al.* Differential responses of IGF-1 molecular complexes to military operational field training. *J Appl Physiol* 2003; 95:1083–1089.
- 36 Loucks AB, Thuma JR. Luteinizing hormone pulsatility is disrupted at a threshold of energy availability in regularly menstruating women. *J Clin Endocrinol Metab* 2003; 88:297–311.
- 37 Glass AR, Deuster PA, Kyle SB, *et al.* Amenorrhea in Olympic marathon runners. *Fertil Steril* 1987; 48:740–745.
- 38 Friedl KE, Westphal KA, Marchitelli LJ, *et al.* Evaluation of anthropometric equations to assess body-composition changes in young women. *Am J Clin Nutr* 2001; 73:268–275.
- 39 Loucks AB. Energy balance and body composition in sports and exercise. *J Sports Sci* 2004; 22:1–14.
- 40 Weyand PG, Kelly M, Blackadar T, *et al.* Ambulatory estimates of maximal aerobic power from foot-ground contact times and heart rates in running humans. *J Appl Physiol* 2001; 91:451–458.
- 41 Mello RP, Murphy MM, Vogel JA. Relationship between a two-mile run for time and maximal oxygen uptake. *J Appl Sport Sci Res* 1988; 2: 9–12.
- 42 Zhang K, Pi-Sunyer FX, Boozer CN. Improved energy expenditure estimation for physical activity. *Med Sci Sports Exerc* 2004; 36:883–889.
- 43 Hoyt RW, Friedl KE. Current status of field applications of physiological monitoring for the dismounted soldier. In: Poos M, editor. *Metabolic monitoring technologies for military field, applications*. Washington DC: National Academy of Sciences, National Academy Press; 2004. pp. 247–257. Available at: <http://darwin.nap.edu/books/0309091594/html>.