

Using Hydrodynamic Models to Interpret Remote Sensing Images of the Sea Surface

James T. Kirby and Robert A. Dalrymple

Center for Applied Coastal Research

University of Delaware

Newark, DE 19716

phone: (302) 831-2438 fax: (302) 831-1228 email: kirby@udel.edu

Award #: N00014-97-1-0283

LONG-TERM GOALS

The long-term goals are to develop methodologies for determining beach topography in the nearshore, using various types of remotely-sensed images of the water surface as input. We are particularly interested in regions such as the surfzone, where linear theory provides a poor description of the behavior of individual wave crests.

OBJECTIVES

The objectives of the present project are to:

- (1) Develop a synthetic data set, based on Boussinesq model predictions, representing a number of cases of waves propagating over characteristic nearshore bathymetries.
- (2) Compare model results to available SandyDuck data.
- (3) Study the feasibility of solving the inverse problem for bathymetry from measured surface data, using a wave model to determine wave phase speeds and wave-induced height and velocity fields.

APPROACH

Much of the effort in the first year of the project has been concentrated on enhancing the Boussinesq model to the point where it is capable of providing a robust description of surfzone phenomena. Special attention has been paid to problems associated with describing wave breaking and the shoreline runup. The resulting model has been extensively tested. At present, we have started the process of building up the synthetic data set, which will be used to test simple inversion schemes based on linear theory as well as more involved schemes using the nonlinear model as the means for time-stepping solutions.

The anticipated INSAR data set from the SandyDuck experiment was not available, and we have attempted to utilize, instead, other available data sets based on scanning surface radars. Work on inversion techniques has gone forward based on synthetic data sets.

WORK COMPLETED

Report Documentation Page

Form Approved
OMB No. 0704-0188

Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.

1. REPORT DATE 1998		2. REPORT TYPE		3. DATES COVERED 00-00-1998 to 00-00-1998	
4. TITLE AND SUBTITLE Using Hydrodynamic Models to Interpret Remote Sensing Images of the Sea Surface				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) University of Delaware, Center for Applied Coastal Research, Newark, DE, 19716				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited					
13. SUPPLEMENTARY NOTES See also ADM002252.					
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT Same as Report (SAR)	18. NUMBER OF PAGES 4	19a. NAME OF RESPONSIBLE PERSON
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified			

Various enhancements to the Boussinesq model code have been made. A refinement of the eddy-viscosity algorithm for wave breaking and the development of an accurate beach-slot formulation have improved the model's accuracy in predicting surfzone wave processes. We have also developed an internal wave-source algorithm which generates a wave field which propagates entirely in the incident-wave direction, eliminating the need for much of the extensive sponge layer regions needed at the offshore boundary of many models with symmetric internal sources (Chawla and Kirby, 1998).

A version of the model code in Fortran 90 with HPF extensions has been developed for application on parallel computers, and has been tested on an 8 node PC cluster. Typical speedups relative to a single processor application are typically in the range of 5 - 5.5 times. This model code is ready for testing on massively parallel machines such as the Cray T3.


Various iterative approaches to the solution of the depth inversion problem using the fully nonlinear model equations have been investigated, and a rapid convergence to the correct answer has been obtained for cases where the entire dynamical input (surface and velocity) is specified. We are presently investigating the case where the data set is incomplete.

RESULTS

The refinement of the Boussinesq model's algorithms for shoreline runup and wave breaking have lead to a model which performs well in a number of situations, as judged by comparison to laboratory data (Chen et al 1998a,b; Kennedy et al 1998). The following figure illustrates a comparison between a time-averages velocity field of a rip current measured in the lab (right) and as obtained from the Boussinesq model Funwave (left).

Two inversion methods have been examined to determine bathymetry from surface elevation information (an image or a pair of images of the water surface), and are described in Dalrymple et al (1998). The first method examined the ability of linear dispersion relationship model to determine bathymetry in cases with refraction, diffraction, and wave nonlinearity, and the second method is based on the lagged correlation method (and several images), which is more generally useful for application.

The linear dispersion relationship method is reasonably robust in determining water depth in the presence of nonlinearity and noise, provided that the wave period is accurately known. The second method, based on cross-correlations and a maximum entropy method, is almost insensitive to noise and provides a reasonable method for field applications.



IMPACT/APPLICATIONS

The work on the project to date has led to a more robust version of the Boussinesq model code which is capable of accurately representing wave shoaling, decay and runup as well as wave- induced circulation, in comparison to laboratory data. The model code is available to the research and engineering community.

TRANSITIONS

The work on this project is closely related to an ongoing Army Research Office project (Kirby and Dalrymple, P.I.'s) which is addressing the development of a curvilinear version of the Boussinesq model code for application to complex coastal regions. The model formulation from this project serves as the basis for the ARO project developments, and the resulting ARO code will allow for various optimizations in model applications which are not presently available to users of the existing code,

including the ability to specify a variable cross-shore grid spacing. The resulting curvilinear grid code will also be useful in interpreting remote-sensing imagery of complex coastal areas.

REFERENCES

- Chen, Q., Dalrymple, R. A., Kirby, J. T., Kennedy, A. and Haller, M. C., ``Boussinesq modelling of a rip current system'', submitted to J. Geophys. Res., June 1998a.
- Chen, Q., Kirby, J. T., Dalrymple, R. A., Kennedy, A. B. and Chawla, A., ``Boussinesq modeling of wave transformation, breaking and runup. II: Two-horizontal dimensions'', in preparation for submission to J. Waterway, Port, Coastal and Ocean Engrng., 1998b
- Dalrymple, R. A., Kennedy, A. B., Kirby, J. T. and Chen, Q., ``Determining depth from remotely-sensed images'', Proc. 26th Intl. Conf. Coastal Engrng., Copenhagen, June 1998, in press.
- Dalrymple, R.A., Chen, Q. Kirby, J.T., A. Kennedy, M.C. Haller, Boussinesq Modelling of Rip Currents, AGU Fall, 1998
- Kennedy, A. B., Chen, Q., Kirby, J. T. and Dalrymple, R. A., ``Boussinesq modeling of wave transformation, breaking and runup. I: One horizontal dimension'', in preparation for submission to J. Waterway, Port, Coastal and Ocean Engrng., 1998