

# **Quantitative Chemical Mass Transfer in Coastal Sediments During Early Diagenesis: Effects of Biological Transport, Mineralogy, and Fabric**

Dawn Lavoie  
Code 7431

Naval Research Laboratory  
Stennis Space Center MS 39529

phone: (601) 688-4659 fax: (601) 688-5752 email: [dawn.lavoie@nrlssc.navy.mil](mailto:dawn.lavoie@nrlssc.navy.mil)

Award #: N0001498WX30240

## **LONG-TERM GOAL**

The long-term goal of this study is to develop a better mechanistic and quantitative understanding of the effects of biologically-enhanced transport, mineralogy, sediment fabric, and particle surface chemistry on biogeochemical reactions occurring in coastal sediments. Specifically, we plan to provide quantitative expressions of biogeochemical processes from field and laboratory studies and to develop a numerical model of early diagenesis which explicitly accounts for the complex interactions between the structural, mineralogical and biological components of coastal marine sediments

## **OBJECTIVES**

The short term objective for NRL during FY98 was to identify appropriate sediment fabric and benthic macrofaunal parameters, and to define a fabric format that can be incorporated into the existing reactive transport model at Georgia Institute of Technology (GIT). Our hypothesis is that the fabric of bioturbated sediments will affect the permeability of fluids and exchange functions of various solutes.

## **APPROACH**

The approach is a combination of field collection; laboratory analyses of bulk physical properties; macro/microscopic analyses of sediment fabric utilizing x-radiography. CT scanning and transmission electron microscopy; and permeability modeling using effective medium theory. Specifically, NRL (in conjunction with USM) has established a new field site in St. Louis Bay, Mississippi, a fined-grained estuarine site, for studying bioturbation effects on fluid flow using techniques developed during the FY97 Dry Tortugas effort. We have begun characterizing the mineral-organic matter relationship and micromineralogy of the field samples utilizing the new, high-resolution TEM. This work is closely coordinated with Y. Furukawa, funded under Award # N00014-98-1-0200, who is analyzing pore fluids, bulk geochemistry and vertical distribution of  $^{13}\text{C}$ ; and P. VanCappellan who is incorporating our results into the RT model, STEADYSED.

## **WORK COMPLETED**

- a. Physical properties analyses from the FY97 Dry Tortugas experiment have been completed. Initial image analysis and EMT modeling of permeability have been completed. These results were presented as part of an invited paper at a geotechnical workshop in Japan in March 1998.
- b. Sediments from three tracer experimental sites in St. Louis Bay have been partially characterized. Index properties analyses suggest similar mineralogy, laboratory analyses of permeability have been

# Report Documentation Page

Form Approved  
OMB No. 0704-0188


Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.

1. REPORT DATE <b>1998</b>		2. REPORT TYPE		3. DATES COVERED <b>00-00-1998 to 00-00-1998</b>	
4. TITLE AND SUBTITLE <b>Quantitative Chemical Mass Transfer in Coastal Sediments During Early Diagenesis: Effects of Biological Transport, Mineralogy, and Fabric</b>				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) <b>Naval Research Laboratory, Code 7431, Stennis Space Center, MS, 39529</b>				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT <b>Approved for public release; distribution unlimited</b>					
13. SUPPLEMENTARY NOTES <b>See also ADM002252.</b>					
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT <b>Same as Report (SAR)</b>	18. NUMBER OF PAGES <b>4</b>	19a. NAME OF RESPONSIBLE PERSON
a. REPORT <b>unclassified</b>	b. ABSTRACT <b>unclassified</b>	c. THIS PAGE <b>unclassified</b>			

initiated, grain size distribution has not been completed; x-radiography and microscopic analyses indicate differential sediment disturbance by grass growth and burrowing. A significant amount of image analyses still to be done should help understand the pore water irrigation results (Furukawa). We have just begun identifying specific OM morphology and types; NRL's new high resolution TEM has just come on line.


## RESULTS

a. EMT modeling of permeability from North Key Harbor in the Dry Tortugas, Florida, on a representative section of carbonate matrix ( $k = 1.6 \times 10^{-4}$  cm/s) agreed well with bulk laboratory analyses of permeability coefficient ( $1.5 \times 10^{-4}$  cm/s) and in situ permeability coefficient of  $7.1 \times 10^{-4}$  cm/s. To model permeability, the microfabric was quantitatively characterized. The protocols developed are being used to model more complicated fabric of fine-grained coastal sediments. Serial SEM sections have been stacked to create a virtual 3D sediment section. Techniques to model 3D permeability are being developed. Image results from the Dry Tortugas experiment suggest that burrows (open or filled) result in differential fluid flow on the microscale (Figure 1).



***Figure 1. Carbonate sediments recovered from North Key Harbor, site of the first 13 C tracer experiment, indicate a number of burrows, both active and filled. Differential grain size results in differential pore distribution and permeability on the microscale.***

b. Sediments recovered from the St. Louis Bay experimental sites are heavily burrowed, and in two cases, disturbed by the growth of marsh grasses. The sediments are fine-grained clays (mineralogy not completed) of approximately 69 % porosity and permeability coefficient of  $6.1 \times 10^{-4}$  cm/s. The image analysis to date has concentrated on the effects of grass (mm-scale, x-radiography) on fluid flow. The grasses, modeled as a partially-filled tubular pore network give a bulk sediment permeability of  $6.8 \times 10^{-4}$  cm/s. In situ permeability measurements using NRL's falling-head field permeameter are planned as ground truth.



**Figure 2. a. x-ray image of sediment illustrating the pervasiveness of grasses throughout the section. b. The grasses are imaged as tubular networks in a binary image. c. Erosion-dilation, Euclidian Distance Mapping and other standard image analysis techniques are used to identify and quantify the required microfabric parameters (e.g., pore body and throat radii, connectivity, autocorrelation function, chord size distribution ) used as input for modeling permeability.**

## **IMPACT/APPLICATION**

A better understanding and mathematical description of biologically-enhanced transport, sediment fabric and particle surface chemistry during shallow diagenesis will allow us to better model and predict the fate and transport of particles and associated pollutants. By concentrating on fine-grained sediments over the next few years, we hope to make a significant contribution understanding harbor pollution solutions. In addition, by understanding the effect of fabric changes during diagenesis, we will be able to better predict sediment physical and geacoustic properties of interest to the MCM community, (predicting mine burial in shallow coastal regions) and the acoustic community for modeling acoustic propagation.

## **TRANSITIONS**

Techniques for quantitative characterization (2D and 3D) of sediment macro and microfabric will be transitioned to other ONR-funded programs including the High-frequency Sound Interaction in Ocean Sediments: Modeling Environmental Controls DRI. It is anticipated that results from this effort will contribute to applied environmental programs in the future.

## **RELATED PROJECTS**

This project has leveraged the NRL 6.1 core program (Microenvironmental Studies) for support, particularly for the field effort, and will continue to do so. Microfabric results have been used in modeling efforts to predict permeability and will undoubtedly continue to benefit other programs with similar requirements, e.g., the ONR High Frequency Sound Interaction DRI will require quantitative pore space and particle geometry data for prediction of permeability and porosity.

## REFERENCES/PUBLICATIONS

Furukawa, Yoko, Dawn Lavoie and Kevin Stephens, 1997, Early Diagenesis of Biologically Reworked Carbonate Sediments near the Dry Tortugas, Florida, GSA (poster).

Furukawa, Y., D. Lavoie, S. Bentley, P. Van Cappellen and A. Shiller. Early Diagenesis Driven by Biologically-enhanced Oxygen Transport in North Key Harbor, Dry Tortugas National Park, Florida. *Marine Geology* (in revision).

Lavoie, D., R. Anandarajah. and Kishore Mohanty, 1998. Microenvironmental Studies: Sediment Response to Biogeochemical Processes. "Microstructure and Sedimentary Environments as Related to Geotechnical Properties of Sediments" Workshop, Port and Harbor Commission, Yokosuka, Japan (invited paper)

Lavoie, D, Y. Furukawa, C. Vaughan, and A. Reed. 1999. Modeling shallow-water sediments off Panama City, Florida, using effective medium theory. Mississippi Academy of Science 1999 Annual Meeting (abstract).

Furukawa, Y., Lavoie, D., and Vaughan, C. 1999: Relationship between measured in situ permeability and diffusive transport of pore water constituents. Mississippi Academy of Science 1999 Annual Meeting (abstract).

Vaughan, C, D. Lavoie, Y. Furukawa and A. Reed. 1999. Predicting permeability as a function of biologically-induced diagenesis. Mississippi Academy of Science 1999 Annual Meeting (abstract).