

Modification of the Stratification and Velocity Profile within the Straits and Seas of the Indonesian Archipelago

Amy Ffield

Earth & Space Research, 290 Clausland Mountain Road, Upper Grandview, NY 10960-4113
phone: (845) 633-1383 email: ffield@esr.org

Award Number: N000141010316

<http://www.esr.org>

LONG-TERM GOALS

To understand the circulation and mixing within the Indonesian Seas associated with topographic configuration, monsoonal driven sea-air flux of momentum and buoyancy, and tides.

OBJECTIVES

1. To utilize existing in situ data (XBT/XCTD, CTD, LADCP, mooring time series, tide gauge records) and satellite data (TRMM, SAR, SST, scatterometer, altimeter) to advance the study of the spatial and temporal scales of topographically and tidally linked circulation, sea-air fluxes, and mixing and internal wave phenomenon within the Indonesian Seas.
2. To develop a program in collaboration with Indonesian marine agencies and universities to study regional variability of meso- and sub-mesoscale processes and ocean strait dynamics within the Indonesian Seas, centered upon field observations.

APPROACH

First to acquire existing in situ and satellite data and to assess the relevant regional scale, meso-scale and smaller processes within the Indonesian Seas, as well as to attend preliminary discussions in Indonesia for the development of a small-scale circulation and mixing program. Second, the data analysis will be refined and published and explicitly used to develop and define the Indonesian program design with Arnold Gordon (Lamont-Doherty Earth Observatory) and Indonesia into a formal research agreement.

WORK COMPLETED

Acquisition and quality control of existing in situ and satellite data with preliminary assessment of the relevance to regional scale circulation and mixing within the Indonesian Seas has been completed. Development of the preliminary plans for a collaborative oceanographic research program was completed during one month in Indonesia, which included presenting early results, "Increasing the Impact of Wind-Induced Mixing in the Indonesian seas: Upwelling, ENSO, Mindanao and Halmahera Eddies, and Internal Tides".

Report Documentation Page

Form Approved
OMB No. 0704-0188

Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.

1. REPORT DATE 2010		2. REPORT TYPE		3. DATES COVERED 00-00-2010 to 00-00-2010	
4. TITLE AND SUBTITLE Modification of the Stratification and Velocity Profile within the Straits and Seas of the Indonesian Archipelago				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Earth & Space Research, 290 Clausland Mountain Road, Upper Grandview, NY, 10960-4113				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT	18. NUMBER OF PAGES	19a. NAME OF RESPONSIBLE PERSON
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified			

RESULTS

The Seram Sea region has been identified as a key indicator of the processes and timing of the factors that impact the heat and freshwater inventories in the Indonesian Seas, and therefore the linkage by sea-air fluxes to the larger scale climate system. While a map of the Root-Mean-Squares (RMS) of February sea surface temperatures in the Indonesian Seas (Figure 1a) reveals the lowest RMS values in the Seram Sea, of less than 0.35 Celsius, the RMS values in the Seram Sea during August (Figure 1b) are highest within the Indonesian Seas, greater than 0.80 Celsius.

The map of the August average SST for the Indonesian Seas (Figure 2a) shows the overall cool SSTs in the eastern Indonesian Seas during August, but the August anomaly SSTs with the 4 coolest Augusts (1994, 1987, 1982, 2007) minus the 4 warmest Augusts (1998, 2000, 1996, 1988) again identifies the Seram Sea region, with larger than 2 Celsius anomaly, as a key location (Figure 2b). In addition, the ocean temperature profiles of the region reveal the same trends with little temperature variability in February, 1 Celsius or less (Figure 3a) in the western Seram Sea region, but significant temperature variability in August, up to 4 Celsius (Figure 3b) in the western Seram Sea region, when contrasting El Niño versus La Niña years. Therefore variability at both seasonal and interannual timescales is accentuated in the Seram Sea region of the Indonesian Seas identifying it as a key indicator of the processes and timing that impact heat and freshwater inventories and sea-air fluxes.




Figure 1a. A map of the Root-Mean-Squares (RMS) of February sea surface temperatures (Celsius) in the Indonesian Seas, calculated from the OI SST dataset. Low RMS, of less than 0.35 Celsius, is revealed in the Seram Sea (magenta region).




Figure 1b. A map of the Root-Mean-Squares (RMS) of August sea surface temperatures (Celsius) in the Indonesian Seas, calculated from the OI SST dataset. High RMS, greater than 0.80 Celsius, is revealed in the Seram Sea.




Figure 2. In panel (a) the map of the August average SST (Celsius) for the Indonesian Seas, and in panel (b) the August anomaly SST with the 4 coolest Augusts (1994, 1987, 1982, 2007) minus the 4 warmest Augusts (1998, 2000, 1996, 1988). In panel (b) the largest SST anomaly, larger than 2 Celsius, is revealed in the Seram Sea region.




Figure 3a. The February eastern Indonesian seas temperature profiles from XBT data revealing that there is little variability, 1 degree Celsius or less, in the thermocline during February when contrasting El Niño versus La Niña years.




Figure 3b. The August eastern Indonesian seas temperature profiles from XBT data revealing that there is large variability, up to 4 degrees Celsius, in the thermocline during August when contrasting El Niño versus La Niña years.

IMPACT/APPLICATIONS

The transfer of tropical water from the Pacific to the Indian Ocean through the complex archipelago of the Indonesian Seas, the Indonesian Throughflow [ITF] is considered to be a first order factor impacting the heat and freshwater inventories of those oceans, and as such is linked by sea-air fluxes to the larger scale climate system. Increased understanding of the circulation and mixing within the Indonesian Seas associated with topographic configuration, monsoonal driven sea-air flux of momentum and buoyancy, and tides will enable improved estimates of the factors impacting these inventories.

RELATED PROJECTS

None.