



 **Peer Reviewed**

Title:

Emergency Physician Estimation of Blood Loss

Author:

[Ashburn, Jeffrey C. MD](#), Madigan Army Medical Center, Department of Emergency Medicine, Tacoma, Washington

[Harrison, Tamara BS](#), University of Washington Medical Center, Department of Medicine, Seattle, Washington

[Ham, James J. MD](#), Madigan Army Medical Center, Department of Emergency Medicine, Tacoma, Washington

[Strote, Jared MD, MS](#), University of Washington Medical Center, Division of Emergency Medicine, Seattle, Washington

Publication Date:

12-06-2011

Publication Info:

Western Journal of Emergency Medicine Articles in Press, Department of Emergency Medicine, UC Irvine

Permalink:

<http://escholarship.org/uc/item/37p9w2x0>

DOI:

[DOI: 10.5811/westjem.2011.9.6669](https://doi.org/10.5811/westjem.2011.9.6669)

Keywords:

Emergency Medicine

License Statement:

This is an Open Access article distributed under the terms of the Creative Commons Non-Commercial Attribution License, which permits its use in any digital medium, provided the original work is properly cited and not altered. For details, please refer to <http://creativecommons.org/licenses/by-nc/3.0/>. Authors grant Western Journal of Emergency Medicine as well as the National Library of Medicine a nonexclusive license to publish the manuscript. Western Journal of Emergency Medicine is produced by the eScholarship Repository and bepress.

Report Documentation Page

Form Approved
OMB No. 0704-0188

Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.

1. REPORT DATE 2011		2. REPORT TYPE		3. DATES COVERED 00-00-2011 to 00-00-2011	
4. TITLE AND SUBTITLE Emergency Physician Estimation Of Blood Loss				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Madigan Army Medical Center, Department of Emergency Medicine, Tacoma, WA, 98466				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited					
13. SUPPLEMENTARY NOTES Western Journal of Emergency Medicine Articles in Press, Department of Emergency Medicine, Volume), No., Month 2011					
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT Same as Report (SAR)	18. NUMBER OF PAGES 6	19a. NAME OF RESPONSIBLE PERSON
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified			

Emergency Physician Estimation of Blood Loss

Jeffery C. Ashburn, MD*

Tamara Harrison, BS[†]

James J. Ham, MD*

Jared Strote, MD, MS[‡]

* Madigan Army Medical Center, Department of Emergency Medicine, Tacoma, Washington

[†] University of Washington Medical Center, Department of Medicine, Seattle, Washington

[‡] University of Washington Medical Center, Division of Emergency Medicine, Seattle, Washington

Supervising Section Editor: H. Bryant Nguyen, MD, MS

Submission history: Submitted November 9, 2010; Revision received June 13, 2011; Accepted September 19, 2011

Reprints available through open access at http://escholarship.org/uc/uciem_westjem

DOI: 10.5811/westjem.2011.9.6669

Introduction: Emergency physicians (EP) frequently estimate blood loss, which can have implications for clinical care. The objectives of this study were to examine EP accuracy in estimating blood loss on different surfaces and compare attending physician and resident performance.

Methods: A sample of 56 emergency department (ED) physicians (30 attending physicians and 26 residents) were asked to estimate the amount of moulage blood present in 4 scenarios: 500 mL spilled onto an ED cot; 25 mL spilled onto a 10-pack of 4 × 4-inch gauze; 100 mL on a T-shirt; and 150 mL in a commode filled with water. Standard estimate error (the absolute value of (estimated volume – actual volume)/actual volume × 100) was calculated for each estimate.

Results: The mean standard error for all estimates was 116% with a range of 0% to 1233%. Only 8% of estimates were within 20% of the true value. Estimates were most accurate for the sheet scenario and worst for the commode scenario. Residents and attending physicians did not perform significantly differently ($P > 0.05$).

Conclusion: Emergency department physicians do not estimate blood loss well in a variety of scenarios. Such estimates could potentially be misleading if used in clinical decision making. Clinical experience does not appear to improve estimation ability in this limited study. [West J Emerg Med. 2011;00(0):000–000.]

INTRODUCTION

Blood loss in the emergency department (ED) is common. Physicians are frequently required to estimate the loss from hemoptysis, epistaxis, rectal bleeding, vaginal bleeding, and traumatic injury as part of their physical examination and ongoing evaluation. Initial bedside hematocrit can often be a poor indicator of acute blood loss: children have a high physiologic reserve, and many cardiac medications, such as β -blockers and calcium channel blockers, affect the normal response to blood loss. These examples are just a few in which estimation of blood loss can alter the assessment of volume status.

Most prior research on blood loss estimation has evaluated practitioners from other specialties. Obstetricians, general

surgeons, trauma surgeons, nurses, and paramedics have all been found to be neither precise nor accurate.^{1–6} In a recent study, emergency physicians (EP) and paramedics were given vital signs and mechanism of injury then asked to visually estimate blood loss in trauma scenarios. As in other studies, estimations were again found to be both inaccurate and affected by the patient's presentation.⁷

The objective of our study was two-fold: (1) to examine ED physicians' accuracy when estimating blood loss on different surfaces in a broad and general set of clinically encountered situations; and (2) to perform a pilot study to compare the accuracy of attending and resident physicians in these tasks.

Table 1. Overall performance in blood estimation.

Scenario	T-shirt	Commode	Sheet	Gauze
Actual amount, mL	100	150	500	25
Standard error mean, %	93	160	63	149
Standard error IQ range, %	36–100	33–117	50–82	35–145
Overestimates,* %	34	43	13	34
Mean, mL/percentage error, [†] %	289/189	605/303	893/79	95/278
Underestimates, [‡] %	50	50	84	50
Mean, mL/percentage error, %	43/57	60/60	182/64	13/48
Within 20%, [§] %	17	11	11	25

IQ, interquartile; *standard error*, absolute value of ((estimated volume – actual volume)/actual volume × 100).

* Overestimates represent the percentage of respondents guessing above the actual amount.

[†] The percentage error is the (mean – actual amount)/average amount.

[‡] Underestimates represent the percentage of respondents guessing below the actual amount.

[§] Within 20% is the percentage of respondents who estimated within 20% of the actual amount.

METHODS

Study Design and Population

A prospective, single-blinded, observational design was used. Participants were emergency physician attending physicians and residents from a single residency program, working in academic hospitals in both urban and suburban settings. Participation was voluntary and a convenience sample of participants was chosen over 4 separate academic meetings.

Study Protocol

Moulage blood was used for all estimations (Ben Nye Stage Blood, Los Angeles, California). Four estimation stations were created: (1) 500 mL of blood spilled onto a standard, white bed sheet, which was spread over an ED bed; (2) 25 mL of blood spilled onto ten 4 × 4-inch gauze sponges; (3) 100 mL of blood spilled on a T-shirt and allowed to dry; and (4) 150 mL of blood added to a commode with 3 L of water. Volumes were chosen to represent a common range of blood loss seen in the ED. Data were collected on 4 separate occasions (grand rounds and faculty meetings) at varying times in order to maximize the number of participating physicians. There was no fundamental difference about the events or the participants at each event.

Physicians were asked to assess each blood loss scenario and estimate the amount of blood loss in milliliters. The same T-shirt and gauze were used for all days. The bed sheet and commode stations were made with fresh blood for each participation group. Subjects were aware of the amount of water that had been in the commode before blood addition. The first bed sheet station created was measured and photographed and subsequent stations were recreated to be as consistent as possible. One researcher interviewed all participants; he presented the scenarios and asked for estimates in the same way for each participant.

Demographic data on gender and years of clinical experience post-MD were also collected.

Verbal consent was obtained; the study was approved by

the human subjects division and qualified for exemption from federal regulations for the protection of human subjects at all participating institutions.

Data Analysis

Estimate error for each group was standardized by converting to percentage error (ie, the absolute value of: estimated volume – actual volume/actual volume × 100). Attending physicians' and residents' performances were compared by using the unpaired student *t* test (IBM SPSS Statistics, version 18, Chicago, Illinois).

RESULTS

A total of 56 subjects participated; all verbally consented and completed all 4 stations.

Of the participants, 64% were male (2 did not choose a gender). The median experience among the 30 attending physicians (54%) who participated was 8 years (first/third quartile = 6/21; 3 did not respond) and the median experience among the 26 residents was 2 years (first/third quartile = 0.5/2.5; 7 did not respond).

Overall standard error averages and interquartile ranges for each scenario are listed in Table 1. For all scenarios combined, mean standard error was 116% with a range from the rare correct answer to 1,233%. Only 8% of estimates were within 20% of the correct answer, while 70% of estimates were within 100% of the correct answer.

Other than for the sheet station, there was not a strong trend toward underestimation or overestimation; underestimates were 50% for the T-shirt station, 50% for the commode station, 84% for the sheet station, and 50% for the gauze station. Only 16% of all estimates were within 20% of the true value.

Table 2 compares attending physician and resident performance for each scenario. In general, attending physicians had less mean error and less extremes of error. Although no

Table 2. Comparison of attending physician and resident performance.

Scenario	T-shirt	Commode	Sheet	Gauze
Actual amount, mL	100	150	500	25
Attending mean std error, %	65	114	65	103
Resident mean std error, %	125	213	62	202
Attending std error IQ range, %	29–78	62–100	50–86	40–100
Resident std error IQ range, %	50–188	33–484	50–80	20–300
Std error mean difference,* %	59	99	–3	99
95% CI, %	0–119	–33–230	–17–11	–19–217
P value	0.05	0.12	0.67	0.10

CI, confidence interval; IQ, interquartile; *std error*, absolute value of ((estimated volume – actual volume)/actual volume × 100).

* Resident mean standard error – attending physician mean standard error.

significant difference was found, trends toward superior attending physician accuracy were seen in each scenario except the sheet station.

DISCUSSION

This study adds to the literature showing that ED physicians, like other specialists,^{1–7} do not estimate blood loss well on a variety of surfaces. The large range of answers seen suggests that physicians are imprecise in addition to being inaccurate. This is consistent with prior studies that have examined other specialties' estimates of blood loss in a variety of scenarios, including active patient blood loss,^{1–3} blood spilled on the ground,⁵ blood spilled onto drapes and sponges,⁴ blood placed in a commode,⁸ and blood loss estimated at the scene of trauma.⁷

Estimates ranged widely, averaging more than 100% off from the actual amounts. Prior studies have suggested that health professionals tend to overestimate small losses (<150 mL) and underestimate large ones (>150 mL).^{3,5–8} Our study found similar results. Comparing the larger blood loss on the bed sheet scenario (500 mL) to the smaller blood loss in the T-shirt scenario (100 mL), there were considerably more underestimates with the larger loss.

The commode and gauze scenarios present an additional challenge of estimating blood in a larger body of fluid and in a very small quantity, respectively; this led to a greater range of answers and more evenly distributed overestimates and underestimates. Patients with complaints of heavy vaginal bleeding or rectal bleeding frequently report blood loss while on the commode and occasionally ask physicians to evaluate losses. This may have clinical relevance because physicians may be just as inaccurate when making estimations in this scenario. Also, EPs are poor at estimating small losses and gauze saturation. Gauze is frequently used to absorb blood during epistaxis, and we observe that ongoing losses could significantly affect the estimation of cumulative loss. It is worthwhile for practitioners to be aware of the further decreased precision and accuracy in these types of situations.

The present study had similarities to a prior study that examined estimations of blood loss made by ED patients.⁹ Nearly identical commode, T-shirt, and gauze scenarios were presented. It is notable that resident and attending EPs had approximately one half and one quarter, respectively, of the average and standard deviation of standard error demonstrated by the patients. This difference makes intuitive sense, as making clinically relevant estimates and observing known quantities of various liquids is part of the day-to-day practice of the emergency physician; it is notably different from a prior study that showed no significant difference between physicians and patients estimating blood in a commode.⁸

No significant difference was found between resident and attending physician performance on each test, consistent with a prior study looking at paramedics with different levels of experience.⁵ There was, however, a strong trend toward superior attending physician performance, with residents demonstrating approximately double the mean error in all scenarios but the bed sheet. Also, there was no significant difference when comparing gender within the scenarios, although the small numbers used likely did not have the power to show a difference; other studies have not found a gender difference.⁹ Whether there would be utility in instituting some form of training is unclear. In the study by Moscatti et al⁵, the blood loss was revealed to the subjects in the study, then slide show presentations were used to train subjects on the appearance of various amounts of blood loss. The subjects were retested at a later date and found to have significantly improved. Given that small effort may lead to large estimation improvement, further research should consider training programs for physicians and how to incorporate these into emergency medicine (EM) or advanced trauma life support training.

The potential clinical relevance of misusing estimations of blood does exist, especially in early evaluation when the hematocrit is unreliable, when multiple different observers are making continued mistakes, in children whose total body blood volume is lower, and for the elderly or those taking medications

that may change the physiologic response to blood loss. Given these situations, we feel that there is value in creating awareness about common conditions causing overestimates and underestimates and the overall inaccuracy of EPs in making and using such estimates in clinical care.

LIMITATIONS

Our study has numerous limitations. Our results may not accurately reflect estimations of actual blood loss in the clinical setting. The study also does not reflect real-life situations in which mechanism of injury and vital signs can influence estimates. However, a prior study has shown physicians are no better when given this information.⁷

Because our study was unblinded to the investigators, bias could have been introduced in the way the study was presented to each patient. Also, physicians could not spread the blood or use other hands-on techniques for estimating blood loss because this would affect the scenario for other subjects. Scenarios that were created each day may have had small but important differences, affecting estimates in some way.

The physicians in this study came from 1 residency program in 1 geographic location in the United States; the study may not reflect the estimation skills of physicians in other academic programs, community settings, or other geographic areas.

Finally, this pilot study was not prospectively powered and the sample sizes used may not have been large enough to show a difference between attending physicians and residents.

CONCLUSION

Blood loss estimates are frequently made by EPs. Like all historical findings, such estimates could potentially be useful as 1 piece in an array of historical points, physical examination findings, and laboratory data results, in an effort to create a complete picture of the severity of the process. This and prior studies suggest that blood loss estimates are difficult for patients and healthcare providers alike, and interpretation of these numbers should therefore be made with extreme caution, if used at all. Further study may be warranted on the effect of blood estimation in clinical

decision making, as well as the value of adding estimation skills to EM training.

Address for Correspondence: Jared Strote, MD, MS, University of Washington Medical Center, Division of Emergency Medicine, Box 356123, 1959 NE Pacific St, Seattle, WA 98195. E-mail: strote@u.washington.edu.

Conflicts of Interest: By the WestJEM article submission agreement, all authors are required to disclose all affiliations, funding, sources, and financial or management relationships that could be perceived as potential sources of bias. The authors disclosed none.

REFERENCES

1. Brant HA. Precise estimation of postpartum haemorrhage: difficulties and importance. *Br Med J.* 1967;1:398–400.
2. Duthie SJ, Ven D, Yung GL, et al. Discrepancy between laboratory determination and visual estimation of blood loss during normal delivery. *Eur J Obstet Gynecol Reprod Biol.* 1991;38:119–124.
3. Larsson C, Saltvedt S, Wiklund I, et al. Estimation of blood loss after cesarean section and vaginal delivery has low validity with a tendency to exaggeration. *Acta Obstet Gynecol Scand.* 2006;85:1448–1452.
4. Meiser A, Casagrande O, Skipka G, et al. Quantification of blood loss. How precise is visual estimation and what does its accuracy depend on [article in German]? *Anaesthesist.* 2001;50:13–20.
5. Moscati R, Billittier AJ, Marshall B, et al. Blood loss estimation by out-of-hospital emergency care providers. *Prehosp Emerg Care.* 1999;3:239–242.
6. Razvi K, Chua S, Arulkumaran S, et al. A comparison between visual estimation and laboratory determination of blood loss during the third stage of labour. *Aust N Z J Obstet Gynaecol.* 1996;36:152–154.
7. Frank M, Schmucker U, Stengel D, et al. Proper estimation of blood loss on scene of trauma: tool or tale? *J Trauma.* 2010;69:1191–1195.
8. Simpson RR, Kennedy ML, Chew SB, et al. Do patients assess rectal bleeding accurately? *ANZ J Surg.* 2001;71:650–651.
9. Strote J, Mayo M, Townes D. ED patient estimation of blood loss. *Am J Emerg Med.* 2009;27:709–711.