



***Ultraviolet B Irradiance and Vitamin D  
Status are Inversely Associated  
With Incidence Rates of  
Pancreatic Cancer Worldwide***

***Sharif B. Mohr  
Cedric F. Garland  
Edward D. Gorham  
William B. Grant  
Frank C. Garland***



***Naval Health Research Center***

---

***Report No. 10-10***

***The views expressed in this article are those of the authors and do not necessarily reflect the official policy or position of the Department of the Navy, Department of Defense, nor the U. S. Government..  
Approved for public release, distribution unlimited.***

***Naval Health Research Center  
140 Sylvester Road  
San Diego, California 92106***

# Ultraviolet B Irradiance and Vitamin D Status are Inversely Associated With Incidence Rates of Pancreatic Cancer Worldwide

Sharif B. Mohr, MPH,\* Cedric F. Garland, DrPH,† Edward D. Gorham, MPH, PhD,\* William B. Grant, PhD,‡ and Frank C. Garland, PhD\*

**Objectives:** To determine if an inverse association exists between latitude, ultraviolet B (UVB) irradiance and incidence rates of pancreatic cancer worldwide.

**Methods:** Multiple linear regression was used to investigate the relationship and between UVB irradiance incidence rates of pancreatic cancer and while controlling for cigarette, alcohol and sugar consumption, and proportion overweight. Serum 25-hydroxyvitamin D [25(OH)D] levels were estimated, and their association with incidence rates also was analyzed.

**Results:** Incidence rates were higher at higher latitudes ( $R^2$  for latitude for men, 0.51;  $P < 0.001$ ;  $R^2$  for latitude for women, 0.32;  $P < 0.001$ ). Ultraviolet B irradiance also was independently inversely associated with incidence in men ( $P < 0.01$ ) and women ( $P = 0.02$ ). Alcohol ( $P < 0.0001$ ) and cigarette ( $P \leq 0.01$ ) consumption were positively associated with incidence in men ( $R^2$  for overall model for men, 0.76;  $P < 0.0001$ ). Alcohol ( $P < 0.0001$ ) and sugar ( $P = 0.001$ ) consumption were positively associated with incidence rates in women ( $R^2$  for overall model for women, 0.64;  $P < 0.0001$ ). Incidence rates were half as high in countries with estimated serum 25(OH)D  $>30$  ng/mL (75 nmol/L) than in those with  $\leq 30$  ng/mL.

**Conclusions:** Countries with lower UVB irradiance had higher incidence rates of pancreatic cancer in both hemispheres, with occasional exceptions.

**Key Words:** pancreatic neoplasms, incidence, vitamin D, alcohol, smoking, multiple regression

(*Pancreas* 2010;00: 00–00)

Approximately 232,000 new cases and 227,000 deaths occur worldwide annually from pancreatic cancer,<sup>1</sup> including 37,700 cases and 34,200 deaths in the United States.<sup>2</sup> The case-fatality rate is 98%.<sup>2</sup> Risk factors include cigarette smoking,<sup>3–6</sup> intake of alcohol<sup>6,7</sup> and red meat,<sup>6,8,9</sup> and history of type 2 diabetes,<sup>3–6</sup> although pancreatic cancer is enigmatic and most of the substantial worldwide differences in incidence cannot be

readily accounted for by known factors. The possibility that vitamin D might play a role in the etiology of pancreatic cancer was raised by studies showing that populations living at higher latitudes or having lower prediagnostic serum 25-hydroxyvitamin D [25(OH)D] levels have higher incidence rates of various cancers, including those of breast,<sup>10–14</sup> colon,<sup>14–19</sup> and ovary.<sup>20</sup>

While dietary and lifestyle factors are well established and highly relevant to risk of pancreatic cancer,<sup>3–9</sup> an inverse association between ultraviolet B (UVB) irradiance and pancreatic cancer mortality was identified in an ecological study in the United States by Grant.<sup>18</sup> The association persisted after controlling for per capita alcohol intake, a proxy estimate of cigarette consumption in the region, and several other covariates.<sup>21</sup> Two studies identified a positive association between latitude and pancreatic cancer mortality rates in Japan<sup>22,23</sup> and Spain.<sup>24</sup> These results suggest that exposure to solar UVB irradiance and resulting vitamin D photosynthesis may be associated with lower incidence of pancreatic cancer, although a worldwide analysis of incidence rates by latitude and UVB irradiance has not been performed.

To further investigate whether there may be an inverse association between UVB irradiance and incidence of pancreatic cancer, this study examined the associations of UVB irradiance, cloud cover, per capita cigarette consumption, intake of alcohol and sugar, and proportion of population overweight, with age-standardized incidence rates of pancreatic cancer in all countries of the world that had data available on all covariates.

## MATERIALS AND METHODS

### Data Sources

Age-standardized incidence rates of pancreatic cancer were obtained for 175 countries from the International Agency for Research on Cancer (IARC) Global Cancer (GLOBOCAN) database.<sup>1</sup> GLOBOCAN uses national registries and registration of vital events to estimate annual age-standardized incidence rates per 100,000 population in these countries. The latest year for which complete data were available was 2002. Data also were obtained on solar UVB irradiance at the top of the atmosphere at the vernal equinox, adjusted for cloud cover; alcohol, cigarette, and sugar consumption; and proportion of population overweight. Data on these covariates were available for 104 countries and were the basis for the multiple regression analyses in this study.

Total solar UVB irradiance at the top of the atmosphere on the vernal equinox was calculated using a standard algorithm.<sup>25</sup> The vernal equinox is March 21 in the Northern hemisphere and September 21 in the Southern hemisphere. The total noon solar irradiance at the top of the atmosphere for each country on the date of the vernal equinox was calculated using the formula ( $A' = A \times \cos x$ ) where  $x$  is the latitude of the country in degrees,  $A$  is the total solar radiation at the equator in  $W/m^2$ , and  $A'$  is the total solar radiation for the country on the date of the vernal equinox in  $W/m^2$ .<sup>25</sup> Because UVB irradiance is approximately

From the \*Naval Health Research Center, San Diego; †Department of Family and Preventive Medicine, University of California San Diego, La Jolla; and ‡Sunlight, Nutrition, and Health Research Center, San Francisco, CA. Received for publication February 19, 2009; accepted August 4, 2009.

Reprints: Cedric F. Garland, DrPH, Department of Family and Preventive Medicine, University of California San Diego, 0631C, 9500 Gilman Dr, La Jolla, CA 92093-0631 (e-mail: cgarland@ucsd.edu).

This research was supported by a Congressional allocation to the Penn State Hershey Cancer Institute of the Milton S. Hershey Medical Center, Hershey, Pennsylvania, through the Naval Health Research Center and the Bureau of Medicine and Surgery, Department of the Navy, under Work Unit No. 60126.

Supplemental digital contents are available for this article. Direct URL citations appear in the printed text and are provided in the HTML and PDF versions of this article on the journal's Web site ([www.pancreasjournal.com](http://www.pancreasjournal.com)).

Copyright © 2010 by Lippincott Williams & Wilkins


0.42% of the total solar irradiance, the total solar irradiance was multiplied by 0.0042 to obtain the estimated UVB irradiance.

To account for the influence of percentage of cloud cover on transmission of UVB through the atmosphere, solar UVB at the top of the atmosphere (TOA) was adjusted for mean December percentage cloud cover by multiplying the UVB at TOA by the following factor: (1 - mean percentage cloud cover). This was done for each country by using data from a NASA cloud cover measurement instrument, the International Satellite Cloud Climatology Project satellite that measures cloud cover from space.<sup>26</sup> Relative ground level UVB irradiance at solar noon on the equinox in each country also was evaluated using UVB data from Lubin et al.<sup>27</sup>


The latitude of the population centroid of each country was determined by the Columbia University Center for International Earth Science Information Network.<sup>25</sup> Intakes of alcohol and sugar were obtained from the United Nations Food and Agriculture Organization, as daily energy in kilocalories, for 132 countries.<sup>28</sup> Data on per capita cigarette consumption during

1980–1982 were obtained from the World Health Organization (WHO).<sup>29</sup> Data on the proportion of population of overweight (obese) individuals during 2002 also were obtained from the World Health Organization.<sup>30</sup>

Serum 25(OH)D was modeled using measured levels of serum 25(OH)D during winter obtained from 28 regions in 18 countries (Appendix Table 1, Supplemental Digital Content 1, <http://links.lww.com/MPA/A20>) as the dependent variable, and UVB irradiance adjusted for cloudiness and skin pigmentation levels in the areas where the studies were performed as the independent variables. The initial regression model that was based on the known values of all variables provided regression coefficients that were used in a separate multiple regression prediction equation. That equation used known UVB and skin pigmentation and the regression equation derived above to estimate mean winter serum 25(OH)D levels in countries where measurements were not available. The prediction equation included an empirically determined scaling constant. The measured 25(OH)D levels were used in the final calculations for



**FIGURE 1.** Incidence rates of pancreatic cancer per population of 100,000 men. Source: Data from GLOBOCAN.<sup>1</sup> 1. New Zealand; 2. Argentina; 3. Uruguay; 4. Chile; 5. South African Republic; 6. Swaziland; 7. Australia; 8. Paraguay; 9. Namibia; 10. Botswana; 11. Mauritius; 12. Madagascar; 13. Zimbabwe; 14. Fiji; 15. Mozambique; 16. Vanuatu; 17. Bolivia; 18. Polynesia; 19. Zambia; 20. Samoa; 21. Melanesia; 22. Angola; 23. Comoros; 24. Malawi; 25. Peru; 26. Brazil; 27. Solomon Islands; 28. Tanzania; 29. Indonesia; 30. Papua New Guinea; 32. Rwanda; 33. Congo Brazzaville; 34. Gabon; 36. Ecuador; 38. Kenya; 39. Uganda; 40. North Korea; 43. Suriname; 44. Brunei; 45. Guyana; 50. Sri Lanka; 53. Togo; 54. Venezuela; 56. Ethiopia; 57. Panama; 58. Costa Rica; 61. Guinea; 64. Djibouti; 65. Guinea-Bissau; 66. Philippines; 68. Cambodia; 69. Nicaragua; 72. Guam; 74. Senegal; 75. Guatemala; 78. Sudan; 80. Yemen; 83. Niger; 85. Viet Nam; 86. Belize; 87. Laos; 88. Jamaica; 89. Puerto Rico; 90. Dominican Republic; 91. Haiti; 93. Mauritania; 94. Cuba; 95. Myanmar; 96. Oman; 98. Bahamas; 99. Qatar; 100. Saudi Arabia; 101. United Arab Emirates; 102. Bahrain; 103. Bangladesh; 105. Libya; 109. Kuwait; 111. Pakistan; 112. Jordan; 113. Israel; 116. China; 117. Iraq; 118. Afghanistan; 121. Cyprus; 123. Malta; 124. Japan; 125. Slovakia; 126. United States of America; 128. Greece; 129. Turkey; 131. Armenia; 132. Korea, Republic of; 133. Spain; 134. Turkmenistan; 135. Azerbaijan; 136. Albania; 137. Kyrgyzstan; 138. Uzbekistan; 139. Macedonia; 141. Italy; 142. Bulgaria; 143. Serbia and Montenegro; 145. Croatia; 146. France; 147. Romania; 149. Singapore; 151. Hungary; 153. Mongolia; 154. Kazakhstan; 155. Sierra Leone; 157. Czech Republic; 158. Luxembourg; 160. Germany; 161. Poland; 162. Netherlands; 163. Ireland; 164. United Kingdom; 165. Belarus; 166. Denmark; 167. Lithuania; 168. Latvia; 169. Estonia; 170. Canada; 172. Norway; 173. Sweden; 174. Finland; 175. Iceland. Legend for countries not labeled on graph: 31. Burundi; 35. Congo; 37. Equatorial Guinea; 41. Malaysia; 42. Colombia; 46. Benin; 47. Cameroon; 48. Liberia; 49. Central African Republic; 51. Cote d'Ivoire; 52. Ghana; 55. Slovenia; 59. Nigeria; 60. Somalia; 62. Micronesia; 63. Trinidad and Tobago; 67. Burkina Faso; 70. Barbados; 71. Gambia; 73. El Salvador; 76. Chad; 77. Honduras; 79. Thailand; 81. Eritrea; 82. Cape Verde; 84. Mali; 92. India; 97. Mexico; 104. Egypt; 106. Bhutan; 107. Algeria; 108. Nepal; 110. Lesotho; 114. Iran; 115. Morocco; 119. Lebanon; 120. Tunisia; 122. Syria; 127. Tajikistan; 130. Portugal; 140. Georgia; 144. Bosnia Herzegovina; 148. Switzerland; 150. Moldova; 152. Austria; 156. Ukraine; 159. Belgium; 171. Russian Federation.



**TABLE 2.** Solar Ultraviolet B Irradiance and Other Covariates Associated With Pancreatic Cancer Incidence Rates; Multiple Regression, 104 Countries, Women, 2002

Variable	Regression Coefficient	SE	t	P
Solar UVB irradiance*	-0.1090	0.0470	-2.32	0.02
Alcohol intake <sup>†</sup>	0.0081	0.0020	4.08	<0.0001
Sugar intake <sup>†</sup>	0.0038	0.0011	3.35	<0.001
Cigarette consumption <sup>‡</sup>	0.0003	0.0002	1.60	0.11
Proportion of population overweight	-0.0101	0.0108	-0.93	0.35
Intercept	1.9810	0.6790	2.92	0.004

R<sup>2</sup> = 0.64, P < 0.0001.

\*Watts/m<sup>2</sup>, adjusted for cloud cover.

<sup>†</sup>Source: United Nations Food and Agriculture Organization.<sup>32</sup>

<sup>‡</sup>Source: United Nations Tobacco Atlas.<sup>33</sup>

§Source: United Nations, WHO Global InfoBase.<sup>34</sup>

individuals. The analyses were performed using SAS Version 9.1 and JMP Version 5.1.2 (SAS Institute, Cary, NC).

### RESULTS

Incidence rates of pancreatic cancer were higher at higher latitudes in men (R<sup>2</sup> = 0.51, P < 0.001; Fig. 1) and women (R<sup>2</sup> = 0.32, P < 0.001; Fig. 2), generally with a curvilinear gradient rising from a low in countries near the equator to a high in those most distant from it. There was an inverse dose-response relationship between modeled serum 25(OH)D and incidence rates of pancreatic cancer in both sexes combined (Fig. 3). Modeled serum 25(OH)D levels above 30 ng/mL (75 nmol/L) were associated with substantially lower incidence rates, with the exception of Belize, which is at a low latitude but had an unusually high incidence rate.

Ultraviolet B irradiance at ground level was independently inversely associated with incidence rates of pancreatic cancer in men (P < 0.01; Table 1). Per capita alcohol consumption (P < 0.0001) and cigarette consumption (P < 0.01) were positively associated with incidence rates (R<sup>2</sup> for full model in men = 0.76, P < 0.0001). In women, UVB irradiance at ground level was also independently inversely associated with incidence rates (P = 0.02; Table 2). Alcohol consumption (P < 0.0001) and sugar intake (P < 0.001) were positively associated with incidence rates (R<sup>2</sup> for full model in women = 0.64, P < 0.0001).

### DISCUSSION

This is the first report of the inverse association of incidence rates of pancreatic cancer with UVB irradiance by country, to our knowledge. Pancreatic cancer incidence rates were higher in countries located at latitudes distant from the equator, where UVB irradiance is low, than in countries closer to the equator, where it is high. Ultraviolet B irradiance varies inversely with latitude<sup>32</sup> and is the source of approximately 95% of circulating vitamin D and its metabolites.<sup>33</sup> The incidence curve corresponded approximately to the reciprocal of the cosine of the latitude, multiplied by a constant that may be different for each disease. The cosine of the latitude is the strongest single correlate of total solar irradiance at noon on the equinoxes. At the latitude of Boston, 42°, for example, the cosine of the latitude is 0.74 at the latitude of Helsinki, 60°, the cosine is 0.5.

The dose-response gradient between modeled serum 25(OH)D levels and incidence rates of pancreatic cancer fol-

lowed a sigmoid inverse dose-response curve (Fig. 3). There was a flat portion at the upper left that suggested that there is no effect of serum 25(OH)D level below approximately 30 ng/mL (75 nmol/L) on incidence rates of pancreatic cancer. The diagonal portion of the dose-response curve suggests that increasing increments in serum 25(OH)D levels in the range from approximately 30 to 36 ng/mL (75–90 nmol/L) were associated with incrementally lower incidence rates of pancreatic cancer. The ratio of the median of the highest incidence rates on the curve to the lowest points was approximately 2.3-fold.

Cigarette and alcohol consumption were positively associated with incidence rates, as expected from previous studies.<sup>3–7</sup> In women, sugar intake was positively associated with rates. This result is in agreement with previous studies that have found a positive association between sugar intake and the risk of pancreatic cancer.<sup>34</sup>

A cohort study by Giovannucci et al<sup>35</sup> reported that higher estimated serum 25(OH)D levels were associated with lower incidence rates of pancreatic cancer. Higher dietary intake of vitamin D also was associated with lower incidence rates of pancreatic cancer in the Harvard Health Professionals Follow-up Study<sup>36</sup> and the Nurses Health Study.<sup>37</sup> Despite these important findings, the literature on the association of vitamin D and 25(OH)D with pancreatic cancer is mixed because of a positive association in men with winter 25(OH)D in a cohort of Finnish smokers.<sup>38</sup> The Finnish population regularly consumed pickled and processed herring that may have been contaminated with environmental carcinogens and are known to contain carcinogenic nitrosamines.<sup>39</sup> This ecological analysis confirms that higher UVB irradiance, a correlate of serum 25(OH)D, is associated with lower incidence of pancreatic cancer. This finding is reassuring and is a new line of evidence that helps to confirm the validity of the inverse association of oral vitamin D intake and serum 25(OH)D levels with pancreatic cancer that has been found in previous research.<sup>36,37</sup>

### Strengths

This study had several strengths. It is the first, to our knowledge, to analyze incidence rates in a large number of countries located at widely different latitudes and to account for variation among countries using multiple linear regression. The latitude part of the study included virtually all countries of the world. The regression model accounted for 76% of the variation in age-standardized incidence rates of pancreatic cancer in men and 64% in women. The regression revealed that there was an independent inverse association of UVB irradiance with incidence rates of pancreatic cancer while controlling for several potential confounders.

The percentage of the variation among countries that was explained for pancreatic cancer was similar to that for cancers for which a role of vitamin D has been more fully established in observational studies of individuals, such as those of the breast (R<sup>2</sup> = 0.55, P < 0.0001),<sup>40</sup> colon (R<sup>2</sup> = 0.68, P < 0.0001),<sup>41</sup> and ovary (R<sup>2</sup> = 0.60, P < 0.0001).<sup>42</sup>

### Limitations

Studies of this type should be considered as hypothesis generating or confirmatory rather than as definitive in isolation. Findings that apply to aggregates, such as countries, may not apply to individuals.<sup>43</sup> For example, all individuals living in areas of high UVB irradiance may not have high exposure to it because of indoor urban lifestyle,<sup>44</sup> air pollution,<sup>45</sup> and regular use of chemical sunscreens.<sup>46</sup> On the other hand, regional solar UVB irradiance may affect a broad range of individuals, and the association reported here was present despite the possible

misclassification of exposure. This is important because non-differential misclassification of exposure generally obscures associations rather than creating them.<sup>47</sup>

The model does not account for all variables that have an impact on vitamin D production such as amount of clothing worn, resulting in some degree of misclassification. The main geophysical and pigmentary determinants of solar photosynthesis of vitamin D were included in the regression model for points that required estimation of 25(OH)D.<sup>31,48,49</sup> There are no accepted international measures of the proportion of skin surface area typically exposed to solar irradiance, so it was not possible to include this clothing factor in regression. The estimated 25(OH)D levels that were used in this model may have been overestimates in countries where cultural factors prevent exposure of most of the skin to solar irradiance, especially Muslim countries. In these countries, 25(OH)D levels, particularly in women wearing a hijab, may be quite low.<sup>50,51</sup> However, most women worldwide do not wear such clothing and the modeling is more directly applicable to them.

Despite the limitations of the ecological approach, the great differences that exist in UVB irradiance among countries worldwide provide a natural experiment on a global scale. Natural experiments have been of value in the past identifying potentially relevant etiological factors for disease.<sup>52</sup>

This study could not account for all possible confounders. For example, although a history of type 2 diabetes mellitus has been a mild risk factor for pancreatic cancer in previous studies,<sup>3–6</sup> the present study could not adjust for differences in prevalence among countries. No comprehensive database on type 2 diabetes prevalence by country is available.

Previous research has identified relatively mild associations between lower levels of physical activity and higher risk of pancreatic cancer<sup>53</sup> and possible associations with high consumption of animal fat<sup>54</sup> and red meat.<sup>6,8,9</sup> The present study did not attempt to control dietary factors other than alcohol and sugar consumption, the most prominent dietary risk factors. However, a covariate was included for proportion of the population overweight, which is a marker for risk of type 2 diabetes. No association was found in the present study between the percentage of population overweight and incidence rates of pancreatic cancer.

While there were departures from the parabolic least-squares trend line for many countries, none was as great as for Belize (17 degrees North), formerly British Honduras. Incidence rates in Belize could represent a chance fluctuation, but a similar association for brain cancer (S. B. Mohr, unpublished) suggests there is a carcinogen present in the environment or diet of Belize.

Further investigation is warranted to confirm the new line of evidence described in this study regarding UVB irradiance and vitamin D.

#### ACKNOWLEDGMENTS

*The views expressed in this report are those of the authors and do not represent an official position of the Department of the Navy, Department of Defense, or the US Government. The authors thank M. Ferlay and the staff of the GLOBOCAN database of the United Nations International Agency for Research on Cancer, Lyon, France.*

#### REFERENCES

1. Ferlay J, Bray F, Pisani P, et al. GLOBOCAN 2002: Cancer Incidence, Mortality and Prevalence Worldwide. IARC CancerBase No. 5. version 2.0. <http://www-dep.iarc.fr/>. Accessed 10 July 2008.
2. Jemal A, Siegel R, Ward E, et al. Cancer facts and figures, 2008. <http://www.cancer.org/downloads/STT/2008CAFFfinalsecured.pdf>. Accessed 6 July 2008.
3. Lowenfels AB, Maisonneuve P. Risk factors for pancreatic cancer. *J Cell Biochem*. 2005;95:649–656.
4. Michaud DS. Epidemiology of pancreatic cancer. *Minerva Chir*. 2004;59:99–111.
5. Larsson SC, Permert J, Hakansson N, et al. Overall obesity, abdominal adiposity, diabetes and cigarette smoking in relation to the risk of pancreatic cancer in two Swedish population-based cohorts. *Br J Cancer*. 2005;93:1310–1315.
6. Lu XH, Wang L, Li H, et al. Establishment of risk model for pancreatic cancer in Chinese Han population. *World J Gastroenterol*. 2006;12:2229–2234.
7. Go VL, Gukovskaya A, Pandolfi SJ. Alcohol and pancreatic cancer. *Alcohol*. 2005;35:205–211.
8. Nothlings U, Wilkens LR, Murphy SP, et al. Meat and fat intake as risk factors for pancreatic cancer: the multiethnic cohort study. *J Natl Cancer Inst*. 2005;97:1458–1465.
9. Larsson SC, Hakansson N, Permert J, et al. Meat, fish, poultry and egg consumption in relation to risk of pancreatic cancer: a prospective study. *Int J Cancer*. 2006;118:2866–2870.
10. Gorham E, Garland C, Garland F. Acid haze air pollution and breast and colon cancer in 20 Canadian cities. *Can J Publ Health*. 1989;80:96–100.
11. Gorham ED, Garland FC, Garland CF. Sunlight and breast cancer incidence in the USSR. *Int J Epidemiol*. 1990;19:820–824.
12. Garland F, Garland C, Gorham E, et al. Geographic variation in breast cancer mortality in the United States: a hypothesis involving exposure to solar radiation. *Prev Med*. 1990;19:614–622.
13. Grant WB. An ecologic study of dietary and solar ultraviolet-B links to breast carcinoma mortality rates. *Cancer*. 2002;94:272–281.
14. Freedman D, Dosemeci M, McGlynn K. Sunlight and mortality from breast, ovarian, colon, prostate, and non-melanoma skin cancer: a composite death certificate based case-control study. *Occup Environ Med*. 2002;59:257–262.
15. Garland C, Garland F. Do sunlight and vitamin D reduce the likelihood of colon cancer? *Int J Epidemiol*. 1980;9:227–231.
16. Tangrea J, Helzlsouer K, Pietinen P, et al. Serum levels of vitamin D metabolites and the subsequent risk of colon and rectal cancer in Finnish men. *Cancer Causes Control*. 1997;8:615–625.
17. Feskanich D, Ma J, Fuchs CS, et al. Plasma vitamin D metabolites and risk of colorectal cancer in women. *Cancer Epidemiol Biomarkers Prev*. 2004;13:1502–1508.
18. Grant WB. An estimate of premature cancer mortality in the U.S. due to inadequate doses of solar ultraviolet-B radiation. *Cancer*. 2002;94:1867–1875.
19. Mizoue T. Ecological studies of solar radiation and cancer mortality in Japan. *Health Phys*. 2004;87:532–538.
20. Lefkowitz ES, Garland CF. Sunlight, vitamin D, and ovarian cancer mortality rates in US women. *Int J Epidemiol*. 1994;23:1133–1136.
21. Grant W, Garland C. The association of solar ultraviolet B with reducing risk of cancer: multifactorial ecological analysis of geographic variation in age-adjusted cancer mortality rates. *Anticancer Res*. 2006;26:2687–2699.
22. Kato I, Tajima K, Kuroishi T, et al. Latitude and pancreatic cancer. *Jpn J Clin Oncol*. 1985;15:403–413.
23. Tominaga S, Kuroishi T. Epidemiology of pancreatic cancer. *Semin Surg Oncol*. 1998;15:3–7.
24. Grant WB. An ecologic study of cancer mortality rates in Spain with respect to indices of solar UVB irradiance and smoking. *Int J Cancer*. 2007;120:1123–1128.
25. Columbia University. Center for International Earth Science Information Network. <http://sedac.ciesin.columbia.edu/ozone/rtm/mval.html>. Accessed 10 July 2008. 2008.
26. National Aeronautics and Space Administration. International Satellite Cloud Climatology Project database. Available from: <http://isccpgisnasagov/products/browsed2html>, accessed 10 July 2008.

27. Lubin D, Jensen E, Gies P. Global surface ultraviolet radiation climatology from TOMS and ERBE data. *J Geophys Res.* 1998;103:26061–26091.
28. United Nations Food and Agriculture Organization. FAOSTAT Food and Agriculture database. Available from: <http://www.fao.org/geonetwork/srv/en/main.search>, accessed 10 July 2008.
29. Mackay J, Eriksen M. *The Tobacco Atlas*. In: Jeremy P, Lacey C, eds. Geneva: World Health Organization; 2002: 94–101.
30. World Health Organization. WHO Global InfoBase Online Available from: [http://www.who.int/ncd\\_surveillance/infobase/web/InfoBaseCommon/](http://www.who.int/ncd_surveillance/infobase/web/InfoBaseCommon/), accessed 10 July 2008.
31. Jablonski NG, Chaplin G. The evolution of human skin coloration. *J Hum Evol.* 2000;39:57–106.
32. Frederick J, Lubin D. The budget of biologically active ultraviolet radiation in the earth-atmosphere system. *J Geophys Res.* 1988;93:3825–3832.
33. Adams JS, Clemens TL, Parrish JA, et al. Vitamin-D synthesis and metabolism after ultraviolet irradiation of normal and vitamin-D-deficient subjects. *N Engl J Med.* 1982;306:722–725.
34. Hart AR, Kennedy H, Harvey I. Pancreatic cancer: a review of the evidence on causation. *Clin Gastroenterol Hepatol.* 2008;6:275–282.
35. Giovannucci E, Liu Y, Rimm EB, et al. Prospective study of predictors of vitamin D status and cancer incidence and mortality in men. *J Natl Cancer Inst.* 2006;98:451–459.
36. Giovannucci E. Vitamin D and cancer incidence in the Harvard cohorts. *Ann Epidemiol.* 2009;19:84–88.
37. Skinner HG, Michaud DS, Giovannucci E, et al. Vitamin D intake and the risk for pancreatic cancer in two cohort studies. *Cancer Epidemiol Biomarkers Prev.* 2006;15:1688–1695.
38. Stolzenberg-Solomon RZ, Vieth R, Azad A, et al. A prospective nested case-control study of vitamin D status and pancreatic cancer risk in male smokers. *Cancer Res.* 2006;66:10213–10219.
39. Knekt P, Jarvinen R, Dich J, et al. Risk of colorectal and other gastro-intestinal cancers after exposure to nitrate, nitrite and N-nitroso compounds: a follow-up study. *Int J Cancer.* 1999;80:852–856.
40. Mohr S, Garland C, Gorham E, et al. Mapping vitamin D deficiency, breast cancer, and colorectal cancer. *Proceedings of the ESRI International User Conference*. Redlands, CA: ESRI; 2005: 1468 Available at [www.esri.com](http://www.esri.com).
41. Gorham ED, Garland CF, Garland FC, et al. Vitamin D and prevention of colorectal cancer. *J Steroid Biochem Mol Biol.* 2005;97:179–194.
42. Garland CF, Mohr SB, Gorham ED, et al. Role of ultraviolet B irradiance and vitamin D in prevention of ovarian cancer. *Am J Prev Med.* 2006;31:512–514.
43. Robinson WS. Ecological correlations and the behavior of individuals. *Am Sociol Rev.* 1950;15:351–357.
44. Holick MF, Chen TC. Vitamin D deficiency: a worldwide problem with health consequences. *Am J Clin Nutr.* 2008;87:1080S–1086S.
45. Agarwal KS, Mughal MZ, Upadhyay P, et al. The impact of atmospheric pollution on vitamin D status of infants and toddlers in Delhi, India. *Arch Dis Child.* 2002;87:111–113.
46. Matsuoka LY, Ide L, Wortsman J, et al. Sunscreens suppress cutaneous vitamin D3 synthesis. *J Clin Endocrinol Metab.* 1987;64:1165–1168.
47. Szklo M, Nieto J. *Epidemiology: Beyond the basics*. Gaithersburg, MD: Aspen Publishers; 2000. p. 1141–1145.
48. Mohr S, Garland C, Gorham E, et al. Is ultraviolet B irradiance inversely associated with incidence rates of endometrial cancer: an ecological study of 107 countries. *Prev Med.* 2007;45:327–331.
49. Mohr S, Garland C, Gorham E, et al. Relationship between low ultraviolet B irradiance and higher breast cancer risk in 107 countries. *Breast J.* 2008;14:255–260.
50. Mishal AA. Effects of different dress styles on vitamin D levels in healthy young Jordanian women. *Osteoporos Int.* 2001;12: 931–935.
51. Al-Turki HA, Sadat-Ali M, Al-Elq AH, et al. 25-Hydroxyvitamin D levels among healthy Saudi Arabian women. *Saudi Med J.* 2008;29:1765–1768.
52. Dean HT, Arnold FA Jr, Jay P, et al. Studies on mass control of dental caries through fluoridation of the public water supply. *Public Health Rep.* 1950;65:1403–1408.
53. Calton BA, Stolzenberg-Solomon RZ, Moore SC, et al. A prospective study of physical activity and the risk of pancreatic cancer among women (United States). *BMC Cancer.* 2008;8:63.
54. Chan JM, Wang F, Holly EA. Pancreatic cancer, animal protein and dietary fat in a population-based study, San Francisco Bay Area, California. *Cancer Causes Control.* 2007;18:1153–1167.

# REPORT DOCUMENTATION PAGE

The public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing the burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB Control number. **PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.**

<b>1. Report Date (DD MM YY)</b> 03/15/2010	<b>2. Report Type</b> New	<b>3. DATES COVERED (from - to)</b> 1980- 2002
--	---------------------------	---

<b>4. TITLE AND SUBTITLE</b> Ultraviolet B irradiance and vitamin D status are inversely associated with incidence rates of pancreatic cancer worldwide	<b>5a. Contract Number:</b> <b>5b. Grant Number:</b> <b>5c. Program Element: 63706N</b> <b>5d. Project Number:</b> <b>5e. Task Number:</b> <b>5f. Work Unit Number: 60126</b>
--	--

<b>6. AUTHORS</b> Mohr, Sharif B; Garland, Cedric F; Gorham, Edward D; Grant, William B; Garland Frank C.	
--	--

<b>7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES)</b> Commanding Officer Naval Health Research Center 140 Sylvester Road San Diego, CA 92106-3521	<b>8. PERFORMING ORGANIZATION REPORT NUMBER</b>  10-10
---	--

<b>8. SPONSORING/MONITORING AGENCY NAMES(S) AND ADDRESS(ES)</b> <b>Commanding Officer</b> <b>Commander</b> <b>Naval Medical Research Center</b> <b>Navy Medicine Support Command</b> <b>503 Robert Grant Ave</b> <b>P O Box 140</b> <b>Silver Spring, MD 20910-7500</b> <b>Jacksonville, FL 32213-0140</b>	<b>10. Sponsor/Monitor's Acronyms(s)</b> BuMed/NHRC <b>11. Sponsor/Monitor's Report Number(s)</b>
--	---

**12 DISTRIBUTION/AVAILABILITY STATEMENT**  
Approved for public release; distribution unlimited.

**13. SUPPLEMENTARY NOTES** This research was supported by a Congressional allocation to the Penn State Hershey Cancer Institute of the Milton S. Hershey Medical Center, Hershey, PA through the Naval Health Research Center and the Department of the Navy, Bureau of Medicine and Surgery.

**14. ABSTRACT (maximum 200 words)**  
**A brief (approximately 200 words) factual summary of the most significant information.**

Approximately 232,000 new cases and 227,000 deaths occur worldwide annually from pancreatic cancer, including 37,700 cases and 34,200 deaths in the US alone. The possibility that Vitamin D might play a role in the etiology of pancreatic cancer was raised by studies showing that populations living at higher latitudes, or having lower prediagnostic serum 25-hydroxyvitamin D [25(OH)D] levels, have higher incidence rates of various cancers. Countries with lower UVB irradiance had higher incidence rates of pancreatic cancer in both hemispheres, with occasional exceptions.

**14. SUBJECT TERMS**  
Pancreatic neoplasms, incidence, vitamin D, alcohol, smoking, multiple regression

<b>16. SECURITY CLASSIFICATION OF:</b>			<b>17. LIMITATION OF ABSTRACT</b> UNCL	<b>18. NUMBER OF PAGES</b> 6	<b>18a. NAME OF RESPONSIBLE PERSON</b> Commanding Officer
<b>a. REPORT</b> UNCL	<b>b. ABSTRACT</b> UNCL	<b>b. THIS PAGE</b> UNCL			<b>18b. TELEPHONE NUMBER (INCLUDING AREA CODE)</b> COMM/DSN: (619) 553-8429