

Award Number:
W81XWH-10-1-0925

TITLE:
Neural and Behavioral Correlates of PTSD and Alcohol Use.

PRINCIPAL INVESTIGATOR:
Gina Forster, Ph.D.

CONTRACTING ORGANIZATION:
The University of South Dakota
Vermillion, SD, 57069

REPORT DATE:
October 2012

TYPE OF REPORT:
Annual

PREPARED FOR: U.S. Army Medical Research and Materiel Command
Fort Detrick, Maryland 21702-5012

DISTRIBUTION STATEMENT: (Check one)

- Approved for public release; distribution unlimited
- Distribution limited to U.S. Government agencies only;
report contains proprietary information

The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision unless so designated by other documentation.

REPORT DOCUMENTATION PAGE

Form Approved
OMB No. 0704-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number. **PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.**

1. REPORT DATE (DD-MM-YYYY) October2012		2. REPORT TYPE Annual		3. DATES COVERED (From - To) 24September2011-23September2012	
4. TITLE AND SUBTITLE Neural and Behavioral Correlates of PTSD and Alcohol Use.				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER W81XWH-10-1-0925	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S) Gina Forster, Ph.D				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) The University of South Dakota Vermillion, SD, 57069-2037				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) U.S. Army Medical Research and Materiel Command Fort Detrick, Maryland 21702-5012				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION / AVAILABILITY STATEMENT Approved for public release; distribution unlimited					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT Hazardous use of alcohol negatively impacts the treatment of PTSD. However, the mechanisms that underlie the association between PTSD and hazardous alcohol use in veterans are poorly understood. The current research takes a multi-level approach to study the psychological, behavioral, cognitive and neural relationships between PTSD and alcohol use. Progress to date has been focused on building the infrastructure to conduct neural imaging by training of personnel, setting up recruitment of veteran populations, and conducting test runs to ensure the recently purchased fMRI equipment and experimental procedures were suitable to test the study hypotheses with veteran participants. The ability to now conduct functional brain imaging at the University of South Dakota represents the first and only opportunity in the State of South Dakota to perform such research, and complements the strong emphasis and infrastructure already in place to conduct psychological, behavioral and cognitive assessment of South Dakota's underserved rural veteran population. Preliminary data collected in the current reporting period suggests that emotional stimuli used in the fMRI tasks elicit arousal responses and expected neural activity in non-veteran and veteran participants, thus validating the paradigms for testing the behavioral and neural association between PTSD and hazardous alcohol use. Research in the next funding period will build on recruitment efforts to increase subject numbers, thus allowing full analysis of the psychological, behavioral, cognitive and neural data to identify behavioral and neural predictors of poor psychological outcomes related to PTSD and hazardous alcohol use in South Dakota veterans. Overall, the current funding has built research infrastructure and tools to allow comprehensive studies regarding mental health and neurological issues specific to veterans in rural areas.					
15. SUBJECT TERMS Posttraumatic stress disorder; Alcohol; Neurobiology of anxiety disorders; Limbic System; Emotional Regulation.					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT UU	18. NUMBER OF PAGES 59	19a. NAME OF RESPONSIBLE PERSON USAMRMC
a. REPORT U	b. ABSTRACT U	c. THIS PAGE U			19b. TELEPHONE NUMBER (include area code)

Table of Contents

	<u>Page</u>
Introduction.....	4
Body.....	4
Key Research Accomplishments.....	16
Reportable Outcomes.....	16
Conclusion.....	16
References.....	16
Appendix.....	17

Introduction:

Posttraumatic stress disorder (PTSD) is a signature impairment of US soldiers arising from Operation Iraqi Freedom (OIF) and Operation Enduring Freedom (OEF) (Kennedy et al., 2007; Gaylord et al., 2008; Hoge et al., 2008). Furthermore, 25-35% of returning OEF/OIF veterans report some alcohol misuse (i.e., using more than intended) (Hoge et al., 2004), and approximately 15% of veterans screen positive for both PTSD and hazardous drinking (Jacobson et al., 2008; McDevitt-Murphy et al., 2010). The association between PTSD and alcohol use has clear therapeutic implications given that impairment associated with PTSD may be compounded by alcohol use (Dao-Castellana et al., 1998; Gilman and Homer, 2008). Furthermore, patients with PTSD and comorbid substance use disorder are least likely to benefit from treatment and more likely to relapse when compared to patients who have substance use disorder alone (Ouimette, et al., 2007; Schafer and Najavits, 2007). Therefore, it is important to study the behavioral and neural relationships between PTSD and alcohol use.

Objective 1) *To test a model linking PTSD symptoms and alcohol-related problems in a population of recently returned soldiers.*

Associations between PTSD symptoms and alcohol use disorder likely represent complex reciprocal relationships. However, longitudinal and experimental research indicates that PTSD symptomatology increases risk for alcohol use disorders. This objective will focus on enhancing understanding of the potential mechanisms by which PTSD may increase risk for alcohol use disorder. Recently returned veterans will be assessed for PTSD symptoms and other mental health issues, alcohol use and dependence issues, and factors related to stress coping and behavioral disinhibition. Specifically, we will test a model linking PTSD symptoms and alcohol use disorder via behavioral disinhibition and affective lability. Behavioral disinhibition is posited to link PTSD symptoms and alcohol-related negative consequences (e.g., interpersonal aggression), whereas affect lability is posited to link PTSD and symptoms of alcohol dependence. Secondary analyses will examine whether hyperarousal and reexperiencing symptoms represent core symptoms underlying the link between PTSD and alcohol use disorder.

Objective 2) *To test a neural circuitry model linking trauma, alcohol consumption, and PTSD symptoms.*

This objective will focus on enhancing understanding of potential neural mechanisms by which traumatic experience and excessive alcohol use may contribute to PTSD symptoms in recently returned veterans. We will test a model linking trauma and alcohol consumption with PTSD symptoms via altered functioning in the anterior cingulate cortex and amygdala. The non-recursive model will incorporate bi-directional effects linking alcohol consumption, neural functioning, and PTSD symptoms. This will be achieved by conducting an fMRI study to examine activity of the amygdala during a masked emotional faces paradigm and to examine the activity of anterior cingulate cortex during the emotional counting Stroop task. The central hypothesis of this objective predicts that trauma and heavy alcohol use negatively impact the activity of and balance between ACC and amygdala function, thus increasing dysfunction associated with PTSD.

A further purpose of this funding was to enhance the capacity of USD to perform fMRI research at USD's health care partner, Avera Sacred Heart Hospital. Therefore, the technical objectives of this project are:

- 1) To substantially increase the ability of The University of South Dakota and its health care partners to research the neural bases of psychological health impairment, diagnosis and treatment by initiating fMRI research.
- 2) To conduct a short-term initial feasibility study to bring together the components of recruitment, screening, imaging, testing, and data analysis; all critical for subsequent fMRI longitudinal studies.

Body:

Objectives and Milestones:

The scientific objectives for this research project were:

- 1) To test a model linking PTSD symptoms and alcohol-related problems in a population of recently returned soldiers.

2) To test a neural circuitry model linking trauma, alcohol consumption, and PTSD symptoms.

Given a year no-cost extension and the delays incurred in the first year of the award, the originally approved Milestone timeline was no longer relevant. Table 1 and Figure 1 below show the updated timeline for each task, which were presented at the Product Line Review meeting on June 12th 2012 and included in the June 2012 quarterly report.

Table 1: Project Objectives, Milestones and Timeline

Tasks/Milestones	Start Date	Duration (days)	End Date
Objective 1: Testing a model linking PTSD symptoms and alcohol-use.			
Task 1: Obtain IRB approval.	9/24/2010	447	12/15/2011
Task 2: Hiring of personnel.	10/1/2010	304	8/1/2011
Task 3: Advertising & recruitment.	12/15/2011	502	4/30/2013
Task 4: Interview participants.	1/1/2012	486	5/1/2013
Task 5: Analysis/statistical modeling of assessment data.	4/30/2012	426	6/30/2013
Task 6: Manuscript/s for assessment data prepared for publication.	5/30/2013	116	9/23/2013
Objective 2: Testing a neurocircuitry model linking trauma, alcohol consumption and PTSD symptoms.			
Task 7: Purchase, set-up & test infrastructure for fMRI study.	9/24/2010	495	2/1/2012
Task 8: Conduct fMRI study.	2/1/2012	469	5/15/2013
Task 9: Analysis of images.	4/30/2012	426	6/30/2013
Task 10: Statistical analysis/modeling of fMRI & behavioral data.	7/1/2013	45	8/15/2013
Task 11: Manuscript/s for fMRI study prepared for publication.	5/30/2013	116	9/23/2013




Figure 1: Gantt chart illustrating the timeline for the various tasks listed in Table 1.

As outlined by Table 1 and Figure 1, the major objectives for the second year of the award included work towards:

- Objective 1 Task 1: Obtain IRB Approval
- Objective 1 Task 3: Advertising and Recruitment
- Objective 1 Task 4: Interview Participants
- Objective 1 Task 5: Analysis/Statistical Modeling of Assessment Data
- Objective 2 Task 7: Infrastructure for fMRI
- Objective 2 Task 8: Conduct fMRI Study
- Objective 2 Task 9: Analysis of Images

Work Towards Objective 1 Task 1: Obtain IRB Approval:

Detailed information regarding the delays and progression of IRB approval over the previous reporting periods was supplied in the 2011 annual report. At that stage (September 2011), the University of South Dakota IRB had approved our human subjects protocol in June 2011, but the results of the review by the DoD Office of

Research Protection (ORP) had not been made available to the PI. In October 2011, reviews from the DoD ORP were sent to the PI, and minor revisions required. The revisions were made, and sent to the University of South Dakota (USD) IRB and to the DoD ORP. The final version of the protocol was approved by the USD IRB on December 12th 2011, and by the DoD ORP on December 14th 2011. Therefore, this task has been completed.

Work Towards Objective 1 Task 3: Advertising and Recruitment:

The investigators generated several forms of advertising materials targeting veterans, including fliers, letters, email scripts, and newspaper advertisements. These were approved in their final form by the DoD ORP and USD IRB in mid-December 2011, and these materials were emailed or mailed to Iraq/Afghanistan veterans in South Dakota directly, in addition to fliers being posted on notice boards on University campuses, National Guard offices and local clinics. Fliers were also distributed during meetings of the USD veterans association. Furthermore, newspaper advertisements in daily and weekly newspapers within the southeast corner of South Dakota and adjacent Nebraska towns were placed. These efforts resulted in the first participants being scheduled for the study in the first week of January 2012. To date, we have recruited 46 veterans who met the selection criteria for phase I of the study (psychological and alcohol use assessment). Of these, 22 participants met the MRI screening criteria and agreed to participate in phase II of the study (fMRI and emotional suppression tasks). See Table 3 for more details on participant recruitment. Work towards this task will continue in year three of this award to ensure sufficient participant numbers (at least 20 per group) for appropriate analysis of the data.

Work Towards Objective 1 Task 4: Interview Participants:

The psychological assessment interview represents phase I of the study. Participants were invited to an interview session with the one of the Clinical Psychology investigators in the location of the participant’s choice (Sioux Falls, Yankton or Vermillion). The participants filled in on-line validated assessment tools probing combat exposure, social support, PTSD symptoms, other mental health and health symptoms, alcohol use and misuse, stress coping and personality traits. They then underwent a clinical interview to confirm mental health status and cognitive status. Finally, screening for MRI contraindications was performed to assist in determining eligibility for phase II of the study. Each interview session was 2-3 hours in length. Forty-six participants were interviewed for phase I of the study. Work towards this task will continue in year three of this award.

Work Towards Objective 1 Task 5: Analysis/Statistical Modeling of Assessment Data:

Clinical interview data is being collated by the Clinical Psychologist investigators, and at this stage there are not enough participants to begin the statistical modeling of this data. However, individual subject information is analyzed on a weekly basis as it relates to MRI compatibility, PTSD symptoms and alcohol use, and is made available by a confidential online process (PsychData) to the Neuroscience investigators of this project, so that they can quickly determine whether the participant could be invited to participate in phase II of this study (Objective 2 Task 8). Participants that do not have contraindications to MRI and that fall into one of the following groups qualify for phase II of the study: 1) low/no alcohol use and low/no PTSD symptoms; 2) low/no alcohol use and PTSD symptoms; 3) high alcohol use and low/no PTSD symptoms; 4) high alcohol use and high PTSD symptoms (Table 2). Table 3 shows the number of participants in each group to date.

Table 2: Criteria Used for Allocating Participants to PTSD and Alcohol Use Groups

	Low-Moderate Alcohol Use	Hazardous Alcohol Use
No PTSD	< score of 29 on PCL < 7/14 drinks/week (women/men) < score of 8 on AUDIT	< score of 29 on PCL ≥ 7/14 drinks/week (women/men) ≥ score of 8 on AUDIT
PTSD	≥ score of 29 on PCL < 7/14 drinks/week (women/men) < score of 8 on AUDIT	≥ score of 29 on PCL ≥ 7/14 drinks/week (women/men) ≥ score of 8 on AUDIT

PCL = PTSD checklist, is used clinically to diagnose PTSD, with a score of 29 or more indicating PTSD symptoms.

The number of drinks (shown for both men and women) is based on the National Institute for Alcohol Abuse and Alcoholism (NIAAA) guidelines for problem drinking within a week, over the last 30 days.

AUDIT = Alcohol Use Disorders Identification Test, a score of 8 or more is used to diagnose alcohol use disorder (questions relate to problems associated with alcohol use in the last year).

Table 3: Number of Participants Tested in Phase I (Psychological Assessment) and Phase II (fMRI) within each Experimental Group

	Low-Moderate Alcohol Use	Hazardous Alcohol Use
No PTSD	Phase I = 6/ Phase II = 4	Phase I = 8/ Phase II = 4
PTSD	Phase I = 15/ Phase II = 6	Phase I = 15/ Phase II = 8

Phase I total = 46 participants, Phase II total = 22 participants.

Work Towards Objective 2 Task 7: Infrastructure for fMRI:

As detailed in the first annual report for this award, the 1.5 T MRI unit at Avera Sacred Heart Hospital was replaced with a 3 T scanner, and outfitted for research (e.g. presenting stimuli, recording subject responses) during the second half of Year 1. The new scanner (Siemens 3T Open Bore Magnetom Skyra) and the Esys fMRI system (stimulus presentation-response system with E-Prime stimulus generation software) required optimization during the current reporting period, prior to being used to image veteran populations. The PI and study personnel received training on the use of the fMRI equipment and software in September and October 2011, and obtained IRB approval to conduct test experiments with student subjects. During December 2011-Feberurary 2012, the PI and co-investigator Dr. Vince Magnotta (Director of MR Research Facility at the University of Iowa) conducted studies with test subjects recruited from USD student groups. We optimized the parameters for basic visual and motor tasks, and for the emotional tasks that were to be used in the scanner with the veteran populations. One problem did arise with the equipment when presenting the emotional tasks which involved the rate at which stimuli needed to be presented was not being supported by the Esys stimulus system. This was solved by the investigators and Esys technical staff with modifications to the stimulus design and mode of presentation. Also in this current reporting year, a DICOM server with XNAT interface was set up between Avera Sacred Heart Hospital and USD, with password protected access being given to Dr. Magnotta so that he can download and process images. This system was tested with the data collected from the student participants in the fMRI optimization studies, and was a very successful means by which to transfer images to Dr. Magnotta at the University of Iowa. The fMRI infrastructure for this study was fully functional by March 2012, allowing the fMRI phase of the current study with veteran subjects to begin. Therefore, this task is complete.

Work Towards Objective 2 Task 8: Conduct fMRI Study:

Following psychological assessment as outlined in Objective 1 Task 4 above, qualifying veteran participants were invited to Avera Sacred Heart Hospital in Yankton for the fMRI component of the study. Participants were re-screened for MRI compatibility, tested for blood-alcohol levels, and then within the scanner, passively viewed masked emotional faces and performed an emotional stroop counting task (a task of emotional suppression – see Figure 3 for more details regarding each task). After the tasks in the scanner were complete, participants then filled out a self-assessment manikin to rate their arousal levels upon viewing the images that comprised the two tasks in the scanner (Figure 2). Although low participant numbers at this stage preclude any statistical analysis of the arousal rating data, preliminary data suggests that participants rate fearful faces higher than neutral and happy faces, and rate combat and negative words higher than neutral words. This suggests that the negatively-valanced emotive stimuli used in the fMRI-based tasks appear to elicit a greater state of arousal in participants. The entire phase II study is approximately 2 hours long, with participants in the scanner for 45 minutes to 1 hour. Twenty-two veteran participants have been scanned in phase II of the study to date. Work towards this task will continue in year three of this award to increase participant numbers to a total of 20 participants each group.






Figure 2: Very preliminary arousal ratings for stimuli used in fMRI tasks (4-8 participants per group, see Table 3). Following the scan, participants were asked to rate their arousal level (scale 1-9 with 9 being high arousal, 1 being very calm) using the self-assessment manikin (SAM). Participants were presented with each stimulus used in the **A**) masked faces paradigm (fearful, neutral, or happy face) and in the **B**) emotional stroop task (neutral, negative or combat word) for 3 sec each, and given 8 sec to rate each stimulus. See Figure 3 for more details regarding the stimuli. While very preliminary, there is a trend for participants to rate fearful faces higher than neutral and happy faces, and there is also a trend for participants to rate combat and negative words higher than neutral words.

Work Towards Objective 2 Task 9: Analysis of Images:

Preliminary fMRI analysis of the first 12 participants from phase II was performed by Dr. Magnotta in June 2012 to determine whether the stimuli and tasks performed in the scanner elicited the expected neural activity. The data from the first 12 participants analyzed was sufficient to determine activation in the regions of interest (amygdala and anterior cingulate cortex) with the planned contrasts that will be employed when performing between-group analyses (Figure 3). Specifically, masked presentation of fearful faces elicited increased activation within the amygdala (Figure 3A) as has been observed in other studies using Vietnam veterans and non-combat participants (e.g., Shin et al., 2005). Furthermore, robust activation of the anterior cingulate cortex was observed during the combat-word condition of the emotional counting stroop task (Figure 3B), as has been observed by previous studies with Vietnam veterans (using Vietnam war-related words; e.g. Shin et al., 2001). Therefore, the paradigms and stimuli used in the current study were validated by this preliminary analysis.

A Masked Fearful Faces



B Emotional Counting Stroop Task




Figure 3: Preliminary analysis ($n = 12$) of fMRI data collected during the masked faces paradigm and the emotional counting stroop task. **A) Masked Faces Paradigm:** Used to elicit processing of fearful information in the brain. Each emotional face (happy or fearful) is backward masked by the neutral expression, as illustrated by the composite stimulus above (500 ms total). Stimuli are derived from the Ekman-Friesen stimulus set, and comprise 8 individuals (5 male, 3 female). Each fearful or happy block comprises 56 presentations of masked fearful or masked happy faces respectively. There are three blocks of each within the 10 block run. Two runs are conducted, with the presentation of fearful and happy blocks counterbalanced between runs. Significant activation in the region of interest – the amygdala (circled in red) was observed in fearful vs. fixation conditions. With an increase in subject number, we expect to see more robust activation of the amygdala. **B) Emotional Counting Stroop Task:** Used to assess suppression of emotional information. Each word (in the table above) is presented 1-4 times on the screen as illustrated above, and the subjects are required to count the number of words and report that number via button press. Words from neutral, negative and combat conditions are matched for length, number of syllables and frequency of use in American-English. Each word condition block comprises 20 word presentations (each 1450 msec duration with 50 msec between each), and each block is presented twice within the run. Two runs are conducted with the presentation of neutral, negative and combat blocks counterbalanced between runs. Robust significant bilateral activation of the region of interest, the rostral anterior cingulate cortex, was observed (circled in green) in combat vs. negative word conditions.

In September 2012, Dr. Magnotta performed a preliminary group-based analysis of the fMRI data collected from twenty participants scanned in phase II of the study. Since there were not sufficient subject numbers to analyze the data across all four groups (see Table 3), within-group analyses were performed for each of the task conditions for no-PTSD groups, the PTSD with no hazardous alcohol use group, and the PTSD with hazardous alcohol use group. Analyses were conducted with a threshold of 0.01 and a cluster size of 50 voxels. The results are summarized in Tables 4-5. While very preliminary due to low subject numbers, the vast majority of regions identified within each group during the masked faces paradigm (Table 4) are regions known to be activated during the processing of emotional faces or images. For example, changes in activation in the lingual gyrus, superior temporal gyrus, anterior and posterior cingulate, cuneus, fusiform gyrus, and parahippocampal gyrus are associated with processing of positively-valanced images while changes in activation of the middle frontal gyrus, parahippocampal gyrus, precuneus, cingulate, superior temporal gyrus, middle temporal gyrus, fusiform gyrus are associated with processing of negatively-valanced images (Kensinger and Schacter, 2006). Furthermore, changes in activity of the middle frontal gyrus, parahippocampal gyrus, fusiform gyrus and middle temporal gyrus are specifically associated with emotional face processing (Shin et al., 2005; Sabatinelli et al. 2011). Similarly, the majority of regions identify within each group during the emotional counting stroop task (Table 5) are consistent with those known to be activated during the processing of negatively-valanced words. It is known that changes in activity within the middle frontal gyrus, inferior temporal lobule, superior temporal gyrus, middle temporal gyrus, precuneus, and cingulate are associated with processing of negative words (Kensinger and Schacter, 2006). Furthermore, combat-related words are specifically associated with regional changes in the middle frontal gyrus, precentral gyrus, anterior cingulate, precuneus, insula, superior temporal gyrus, middle temporal gyrus, lingual gyrus and claustrum (Shin et al., 2001). Therefore, the preliminary within-group analysis of our fMRI data for both paradigms suggests activation in expected regions of the brain associated with processing of emotional content.

Table 4: Summary of Regions in which Changes in BOLD Signal was Observed during the Masked Faces Paradigm.

Group	Task Condition	Region
No PTSD	Happy Faces	Left Lingual Gyrus Right Parahippocampal Gyrus Left Parahippocampal Gyrus Right Superior Temporal Gyrus Right Insula Right Cingulate Gyrus Right Cuneus
PTSD	Happy Faces	Right Lingual Gyrus Left Lingual Gyrus Left Fusiform Gyrus Right Inferior Frontal Gyrus
PTSD with Hazardous Alcohol Use	Happy Faces	Left Lingual Gyrus Right Caudate Tail Left Caudate Tail Left Parahippocampal Gyrus Right Superior Temporal Gyrus Right Insula Right Cingulate Gyrus Right Cuneus
No PTSD	Fearful Faces	Right Parahippocampal Gyrus Left Parahippocampal Gyrus Left Precuneus Right Precuneus Right Cuneus Right Insula Right Precentral Gyrus Right Postcentral Gyrus Right Cingulate Gyrus Right Superior Temporal Gyrus Left Posterior Cingulate Gyrus
PTSD	Fearful Faces	Right Fusiform Gyrus

		Left Fusiform Gyrus Right Middle Frontal Gyrus Left Precentral Gyrus
PTSD with Hazardous Alcohol Use	Fearful Faces	Right Parahippocampal Gyrus Left Parahippocampal Gyrus Right Insula Right Cuneus Right Precuneus Left Precuneus Right Precentral Gyrus Right Postcentral Gyrus Right Cingulate Gyrus Left Posterior Cingulate Gyrus Right Superior Temporal Gyrus
No PTSD	Fearful – Happy Faces	Left Lingual Gyrus
PTSD	Fearful – Happy Faces	None
PTSD with Hazardous Alcohol Use	Fearful – Happy Faces	Right Middle Temporal Gyrus Right Anterior Cingulate Gyrus

The summary presented in Table 4 also suggest activity of many of the same regions in participants within no-PTSD and PTSD groups within the happy faces or fearful faces conditions, as would be expected with similar regions processing emotional content across participants. However, the fearful-happy contrast analysis (Table 4; Figure 4) suggests differential activation of regions in those participants reporting hazardous alcohol use and PTSD when compared to PTSD alone, specifically related to emotional processing (e.g. middle temporal gyrus and anterior cingulate). Like the masked faces paradigm, the presentation of negatively-valenced words in the emotional counting stroop task (Table 5) elicited activity in many of the same brain regions across PTSD and no-PTSD groups. However, some notable differences were observed – for example, combat words elicited changes in regions involved in top-down control of emotional/attentional processing (such as the claustrum and anterior cingulate gyrus) in the PTSD with hazardous alcohol use group only (Table 5, Figure 5). Increasing subject numbers in each group over the next year will determine whether these findings are confirmed.





Figure 4: Preliminary analysis ($n = 8$) of fMRI data collected during the fearful faces condition in the masked faces paradigm from participants in the PTSD with hazardous alcohol use group. These participants show


reduced activation of the anterior cingulate and increased activation of the middle temporal gyrus when viewing masked fearful faces.

Table 5: Summary of Regions in which Changes in BOLD Signal was Observed during the Emotional Counting Stroop Task.

Group	Task Condition	Region
No PTSD	General Negative Words	Left Lingual Gyrus Left Posterior Cingulate Gyrus Left Inferior Parietal Lobule Left Inferior Parietal Lobule Right Middle Temporal Gyrus Left Middle Temporal Gyrus Left Precentral Gyrus Right Insula Right Middle Frontal Gyrus
PTSD	General Negative Words	Right Lingual Gyrus Left Lingual Gyrus Right Precentral Gyrus Left Postcentral Gyrus Right Fusiform Gyrus
PTSD with Hazardous Alcohol Use	General Negative Words	Left Lingual Gyrus Left Posterior Cingulate Gyrus Right Inferior Parietal Lobule Left Inferior Parietal Lobule Right Middle Temporal Gyrus Left Middle Temporal Gyrus Left Precentral Gyrus Left Postcentral Gyrus Right Insula Right Middle Frontal Gyrus
No PTSD	Combat Words	Left Precuneus Left Precentral Gyrus Left Insula Left Superior Temporal Gyrus Right Cingulate Gyrus
PTSD	Combat Words	Left Middle Frontal Gyrus
PTSD with Hazardous Alcohol Use	Combat Words	Right Anterior Cingulate Gyrus Right Cingulate Gyrus Left Cingulate Gyrus Left Lingual Gyrus Left Precentral Gyrus Left Postcentral Gyrus Left Insula Right Middle Temporal Gyrus Right Middle Frontal Gyrus Left Inferior Temporal Lobule Right Claustrum



Anterior Cingulate Gyrus



Claustrum


Figure 5: Preliminary analysis (n = 8) of fMRI data collected during the combat word condition in the emotional counting stroop task from participants in the PTSD with hazardous alcohol use group. These participants show increased activation of the anterior cingulate and claustrum when viewing combat-related words.

Furthermore, subject numbers allowed contrasts between-groups for PTSD vs. No PTSD and PTSD vs. PTSD with hazardous alcohol use for each of the task conditions. These contrasts were performed with a threshold of 0.01 and 20 voxel clusters and are thus quite preliminary. The results are summarized in Tables 6-7 and illustrated by Figures 6-8. Of relevance to the planned comparisons when the full data set is available, we observed regional differences in precuneus and cuneus areas as well as the middle temporal gyrus when PTSD groups were contrasted with the PTSD and hazardous alcohol group in the fearful-happy faces contrast condition of the masked faces paradigm (Table 6, Figure 6). Furthermore, changes in activity of the postcentral gyrus and middle temporal gyrus were noted when PTSD groups were contrasted with no-PTSD groups, and in the precentral gyrus when PTSD groups were contrasted with the PTSD and hazardous alcohol group in the combat word condition of the emotional counting stroop task (Table 7, Figures 7-8). Changes in these regions within PTSD populations during the combat word condition of the counting emotional stroop task have been noted previously (Shin et al., 2001), with the current data adding the variable of hazardous alcohol use. Work towards this task will continue in year three of this award, with greater subject numbers allowing full within-group and between-group contrasts for each task condition in the future.


Table 6: Differences in BOLD Signal Between Experimental Groups during the Masked Faces Paradigm.

Group Contrast	Task Condition	Region
PTSD - No PTSD	Happy Faces	Right Insula Right Cingulate Gyrus Left Posterior Cingulate
PTSD – PTSD & Hazardous Alcohol	Happy Faces	Right Putamen Right Cingulate Gyrus Left Posterior Cingulate
PTSD – No PTSD	Fearful Faces	Right Postcentral Gyrus Right Cingulate Gyrus
PTSD – PTSD & Hazardous Alcohol	Fearful Faces	Right Precentral Gyrus


PTSD - No PTSD	Fearful – Happy Faces	Right Postcentral Gyrus Right Cingulate Gyrus Left Cingulate Gyrus Right Anterior Cingulate Right Insula Right Inferior Frontal Gyrus Right Precuneus Left Precuneus
PTSD – PTSD & Hazardous Alcohol	Fearful – Happy Faces	Right Middle Temporal Gyrus Right Precuneus Right Cuneus Left Cuneus Right Middle Temporal Gyrus



Precuneus



Cuneus




Middle Temporal Gyrus


Figure 6: Preliminary analysis (n = 12) of fMRI data collected during the masked faces paradigm from participants in the PTSD vs. PTSD with hazardous alcohol use groups. Data represents fearful vs. happy faces conditions. Participants with PTSD show increased activation of the precuneus and reduced activation of the cuneus and middle temporal gyrus as compared to participants with PTSD and hazardous alcohol use when activity elicited by masked fearful faces is contrasted with masked happy faces.

Table 7: Differences in BOLD Signal Between Experimental Groups during the Emotional Counting Stroop Task.

Group Contrast	Task Condition	Region
PTSD - No PTSD	General Negative Words	Right Precentral Gyrus
PTSD – PTSD & Hazardous Alcohol	General Negative Words	Right Precentral Gyrus
PTSD – No PTSD	Combat Words	Left Postcentral Gyrus
		Left Middle Temporal Gyrus
PTSD – PTSD & Hazardous Alcohol	Combat Words	Right Precentral Gyrus




Postcentral Gyrus



Middle Temporal Gyrus

Figure 7: Preliminary analysis ($n = 20$) of fMRI data collected during the combat word condition in the emotional counting stroop task from participants in the PTSD vs. no-PTSD groups. Participants with PTSD show decreased activation of the postcentral gyrus and middle temporal gyrus as compared to participants without PTSD when viewing combat-related words.



Precentral Gyrus

Figure 8: Preliminary analysis ($n = 12$) of fMRI data collected during the combat word condition in the emotional counting stroop task from participants in the PTSD vs. PTSD with hazardous alcohol use groups. Participants with PTSD show decreased activation of the precentral gyrus as compared to participants with PTSD and hazardous alcohol use when viewing combat-related words.

Key Research Accomplishments:

In relation to the objectives and tasks in Table 1, key accomplishments for Year 2 were:

- Obtaining IRB approval and setting up the infrastructure for fMRI research with veterans – the first of its kind in South Dakota.
- Recruitment of mid-west Iraq/Afghanistan war veterans to participate in a study assessing the relationship between alcohol use, PTSD symptoms and neural processing of emotions.
- Preliminary analysis validating that the emotional stimuli used in the fMRI tasks elicit arousal responses and expected neural activity in non-veteran and veteran participants.

Reportable Outcomes:

- Review Article in Press (see Appendix):
Forster GL, Novick AM, Scholl JL, Watt MJ. (2012). The role of the amygdala in anxiety disorders. In B. Ferry (Ed.), *Amygdala*. Intech (in press).
- Conference Presentation of Preliminary Data:
Olson, D., Hansen, J., Baugh, L., Gaher, R., Simons, J, Magnotta, V and Forster, G. (2012). Neural Imaging: The Effects of Alcohol Use on Emotional Processing in Posttraumatic Stress Disorder (PTSD). Presented at the *University of South Dakota Health Affairs Research Symposium*, Vermillion, SD, April 2012.
- Conference Presentation of Preliminary Data:
Olson, D., Hansen, J., Baugh, L., Gaher, R., Simons, J., Magnotta, V. and Forster, G. (2012). Neural Imaging: The Effects of Alcohol Use on Emotional Processing in Posttraumatic Stress Disorder (PTSD). Presented at the *Sanford School of Medicine Research Symposium*, Vermillion SD, May 2012.

Conclusions:

Hazardous use of alcohol negatively impacts the treatment of PTSD. However, the mechanisms that underlie the association between PTSD and hazardous alcohol use in veterans are poorly understood. The current research takes a multi-level approach to study the psychological, behavioral, cognitive and neural relationships between PTSD and alcohol use. Progress to date has been focused on building the infrastructure to conduct neural imaging by training of personnel, setting up recruitment of veteran populations, and conducting test runs to ensure the recently purchased fMRI equipment and experimental procedures were suitable to test the study hypotheses with veteran participants. The ability to now conduct functional brain imaging at the University of South Dakota represents the first and only opportunity in the State of South Dakota to perform such research, and complements the strong emphasis and infrastructure already in place to conduct psychological, behavioral and cognitive assessment of South Dakota's underserved rural veteran population. Preliminary data collected in the current reporting period suggests that emotional stimuli used in the fMRI tasks elicit arousal responses and expected neural activity in non-veteran and veteran participants, thus validating the paradigms for testing the behavioral and neural association between PTSD and hazardous alcohol use. Research in the next funding period will build on recruitment efforts to increase subject numbers, thus allowing full analysis of the psychological, behavioral, cognitive and neural data to identify behavioral and neural predictors of poor psychological outcomes related to PTSD and hazardous alcohol use in South Dakota veterans. Overall, the current funding has built research infrastructure and tools to allow comprehensive studies regarding mental health and neurological issues specific to veterans in rural areas.

References:

Dao-Castellana, M.H., Samson, Y., Legault, F., Martinot, J.L., Aubin, H.J., Crouzel, C., Feldman, L., Barrucand, D., Rancurel, G., Feline, A. & Syrota, A. (1998) Frontal dysfunction in neurologically normal chronic alcoholic subjects: metabolic and neuropsychological findings. *Psychological Medicine*, **28**, 1039-1048.

- Gilman, J.M. & Hommer, D.W. (2008) Modulation of brain response to emotional images by alcohol cues in alcohol-dependent patients. *Addiction Biology*, **13**, 423-434.
- Hoge, C. W., Castro, C. A., Messer, S. C., McGurk, D., Cotting, D. I., & Koffman, R. L. (2004). Combat Duty in Iraq and Afghanistan, Mental Health Problems, and Barriers to Care. *New England Journal of Medicine*, **351**(1), 13-22.
- Hoge, C.W., McGurk, D., Thomas, J.L., Cox, A.L., Engel, C.C. & Castro, C.A. (2008) Mild traumatic brain injury in U.S. Soldiers returning from Iraq. *The New England Journal of Medicine*, **358**, 453-463.
- Jacobsen, L. K., Southwick, S. M., & Kosten, T. R. (2001). Substance use disorders in patients with posttraumatic stress disorder: A review of the literature. *American Journal of Psychiatry*, **158**(8), 1184-1190.
- Kensinger EA, Schacter DL (2006) Processing emotional pictures and words: effects of valence and arousal. *Cognitive, Affective & Behavioral Neuroscience* **6**:110-126.
- McDevitt-Murphy, M. E., Williams, J. L., Bracken, K. L., Fields, J. A., et al., (2010). PTSD symptoms, hazardous drinking, and functioning among U. S. OEF and OIF Veterans present to primary care. *Journal of Traumatic Stress*, **23** (1), 108-111.
- Ouimette. P., Coolhart, D., Schum Funderburk, J., Wade, M., & Brown P. J. (2007). Precipitants of first substance use in recently abstinent substance use disorder patients with PTSD. *Addictive Behaviors*, **32**, 1719–1727.
- Sabatinelli D, Fortune EE, Li Q, Siddiqui A, Krafft C, Oliver WT, Beck S, Jeffries J (2011) Emotional perception: meta-analyses of face and natural scene processing. *NeuroImage* **54**:2524-2533.
- Schafer, I., & Najavits, L. M. (2007). Clinical challenges in the treatment of patients with posttraumatic stress disorder and substance abuse. *Current Opinion in Psychiatry*, **20**(6), 614-618.
- Shin LM, Whalen PJ, Pitman RK, Bush G, Macklin ML, Lasko NB, Orr SP, McInerney SC, Rauch SL (2001) An fMRI study of anterior cingulate function in posttraumatic stress disorder. *Biological Psychiatry* **50**:932-942.
- Shin LM, Wright CI, Cannistraro PA, Wedig MM, McMullin K, Martis B, Macklin ML, Lasko NB, Cavanagh SR, Krangel TS, Orr SP, Pitman RK, Whalen PJ, Rauch SL (2005) A functional magnetic resonance imaging study of amygdala and medial prefrontal cortex responses to overtly presented fearful faces in posttraumatic stress disorder. *Archives of General Psychiatry* **62**:273-281.

Appendix:

- Forster GL, Novick AM, Scholl JL, Watt MJ. (2012). The role of the amygdala in anxiety disorders. In B. Ferry (Ed.), *Amygdala*. Intech (in press).

1 The Role of the Amygdala in Anxiety Disorders

2 Gina L. Forster, Andrew M. Novick, Jamie L. Scholl and Michael J. Watt

3 Additional information is available at the end of the chapter

4 <http://dx.doi.org/10.5772/50323>

5 1. Introduction

6 1.1. Defining anxiety and fear

7 Anxiety is a term often used to encompass feelings of apprehension, dread, unease or
8 similarly unpleasant emotions. Trait anxiety defines the affect of an organism over time and
9 across situations, whereas state anxiety is the response or adaptation to a given situation [1].
10 Anxiety can be differentiated from fear, both biologically and behaviorally [see 1 for an
11 extensive review]. Converging theories and evidence from clinical psychology and
12 comparative neuroscience suggest that fear can be considered a negatively-valenced
13 emotion that is brief, focused on the present, occurs in situations of specific threat, and aids
14 in avoidance or escape [1,2]. Anxiety, on the other hand, is a negatively-valenced emotion
15 that is characterized by sustained hyperarousal in response to uncertainty, is thus future-
16 focused, and aids in defensive approach or risk assessment [1,2]. Both anxiety and fear are
17 emotions experienced by all individuals and can serve to be adaptive in shaping decisions
18 and behaviors related to survival of an organism [1,3]. However, when excessive, or
19 pathological, or triggered inappropriately, fear and anxiety form the basis of a variety of
20 anxiety disorders [3,4,5; Table 1]. As illustrated by Table 1, some anxiety disorders such as
21 generalized anxiety disorder (GAD) or obsessive-compulsive disorder (OCD) are
22 characterized by excessive anxiety as defined above [1]. However, other anxiety disorders
23 are characterized, at least in part, by excessive and inappropriate fear, such as posttraumatic
24 stress disorder (PTSD), specific phobias and social anxiety disorder [1,3; Table 1]. Thus, it is
25 important to understand the neurobiology of both anxiety and fear to obtain a
26 comprehensive picture of the physiological basis of anxiety disorders.

27 1.2. Anxiety disorders

28 One in three people will develop one of the anxiety disorders outlined by Table 1 within
29 their life-time, with the life-time prevalence at least two times more likely for women [5,6].

1 Furthermore, individuals may present with one or more comorbid anxiety disorders, and
 2 anxiety disorders are highly likely to be comorbid with other psychiatric illnesses, such as
 3 major depressive disorder, psychosis, mania, and substance abuse disorder [4-6]. Several
 4 non-psychiatric disorders are also associated with anxiety disorders, and these include
 5 hyperthyroidism, Cushing's disease and mitral value prolapse [4,5]. Thus, anxiety disorders
 6 are one of the most prevalent psychiatric disorders, posing great personal, economic, and
 7 societal burdens [4-6].
 8

Generalized Anxiety Disorder (GAD)
Excessive worry occurring more days than not over at least a 6 month period, accompanied by restlessness, fatigue, sleep disturbances, muscle tension or irritability.
Posttraumatic Stress Disorder (PTSD)
Characterized by a history of trauma and symptoms related to avoidance, re-experiencing, and physiological hyperarousal in the face of triggering cue.
Obsessive-Compulsive Disorder (OCD)
Compulsions (repeated actions) produced to reduce anxiety associated with obsessions (unwanted, intrusive thoughts).
Panic Disorder
Characterized by panic attacks; a period of intense fear or discomfort accompanied by a variety of physiological symptoms (e.g. sweating, trembling, chest pains, tachycardia).
Agoraphobia
Fear and avoidance of situations from which escape would be difficult in the event of having panic-like symptoms.
Specific Phobia
Excessive or unreasonable fear in anticipation or in response to a specific object or situation.
Social Anxiety Disorder (Social Phobia)
Excessive/unreasonable fear and avoidance of social situations (including performances) in which the person is exposed to unfamiliar people or possible scrutiny by others.

9 **Table 1.** Major Classes of Anxiety Disorders [4,5,7]

10 **1.3. Goals of the current review**

11 The neurobiological bases of anxiety and fear appear to be very similar across species [1],
 12 thus complementary findings from both animal models (most often rodents) and human
 13 studies can contribute to theories of the neurobiological basis of anxiety disorders. State fear
 14 within animal models is most often studied by measures of freezing and fear-potentiated
 15 startle, both acquired via classical conditioning of rodents [1,8]. State anxiety, on the other
 16 hand, is most often studied using apparatus such as an open field, elevated plus maze, or
 17 light-dark box, which all take advantage of the rodent's preference for familiar, dark, and/or
 18 enclosed areas [1,9]. Notably, these paradigms do not rely on the processes underlying
 19 classical conditioning, although McNaughton and Corr [2] caution against defining fear
 20 verses anxiety as conditioned versus unconditioned responses. While trait fear is not well-

1 defined by animal studies [1], trait anxiety is often examined in animal models by the use of
2 selective breeding, resulting in high- and low-anxiety strains and lines of rodents [for
3 example, see 1, 10]. However, one can argue that experimental manipulations (such as early-
4 life stress or amphetamine withdrawal) that drive a group of animals towards greater fear-
5 and anxiety-like phenotypes also examine the underlying basis of trait fear or anxiety [e.g.
6 11, 12]. As noted by Sylver et al [1] clinical studies most often examine trait anxiety, whereas
7 experiments involving animal models most often focus on state anxiety and fear, and then
8 relate these findings to concepts associated with trait anxiety. Regardless, both human and
9 animal studies suggest an important role for the amygdala, and subregions within, in
10 mediating fear and anxiety, and in the manifestation of anxiety disorders (Sections 2 and 3).
11 Therefore, the goals of this review are to first evaluate and integrate classical and recent
12 findings from human studies and relevant animal models that reveal the specific role the
13 amygdala plays in fear and anxiety, and then to elucidate how anxiolytic drugs may affect
14 the amygdala function to ameliorate heightened fear and/or anxiety. This is important,
15 given that traditional drug and cognitive behavioral therapy (CBT) are effective in reducing
16 symptoms of the various anxiety disorders for many individuals, but often do not provide
17 long-term relief, and relapse is a common post-treatment outcome [as reviewed by 3].
18 Therefore, the final goal of the current review is to identify future potential therapeutic
19 targets for the treatment of anxiety disorders.

20 **2. Human imaging studies: Amygdala hyperfunction and anxiety** 21 **disorders**

22 **2.1. Amygdala reactivity and anxiogenic or fearful stimuli**

23 Human imaging studies that explore the neurobiological bases of anxiety or fear processing
24 typically use functional magnetic resonance imaging (fMRI) or positron emission tomography
25 (PET) as measures of neural activity or cerebral blood flow. Imaging experiments that are
26 designed to study neural reactivity to fearful stimuli utilize either conditioned fear paradigms
27 similar to those used in animal models, or involve the presentation of unconditioned stimuli
28 such as fearful faces [1]. It has become clear that masked stimuli can elicit conditioned and
29 unconditioned fear responses from human subjects, suggesting unconscious, implicit
30 processing of these cues [as reviewed by 1]. Similarly, increased activity of the amygdala is
31 observed in response to both conditioned and unconditioned fearful stimuli, independent of
32 whether the subject is aware of the stimulus [1,13-16].

33 Comparable studies that have examined neural correlates of anxiety in healthy controls are
34 limited. One of the reasons for this is that many studies use fearful stimuli, such as the
35 fearful faces or conditioned fear paradigms [1], blurring the distinction between fear and
36 anxiety. Therefore, conclusions regarding neural bases of anxiety are better drawn from
37 studies that include trait anxiety as a variable while utilizing fearful stimuli, or those fewer
38 studies in which an anxiogenic situation is created within the experimental design. Like for
39 studies of fear processing, the majority of these studies show a relationship between trait
40 anxiety and greater amygdala reactivity [as reviewed by 17]. For example, a study of healthy

1 subjects found that reactivity of the amygdala was positively correlated with anticipatory
2 anxiety, and when the anticipated event was imminent, amygdala activation positively
3 correlated with the degree of trait anxiety [18]. Furthermore, college students who scored in
4 the upper 15th percentile for trait anxiety show greater amygdala reactivity to emotional
5 faces as compared to students who scored in the normative range, suggesting that anxiety-
6 prone individuals have greater amygdala reactivity [19]. A similar hyperactivity of the
7 amygdala in high trait anxiety participants is noted when a masked emotional faces or
8 unattended faces paradigm are used [20,21], suggesting the individual does not need to be
9 aware of the stimulus to exhibit heightened amygdala activity. Interestingly, Etkin et al., [21]
10 differentiate between different subregions of the amygdala (see Section 3.1 for more details
11 on amygdala subregions), with the basolateral amygdala activated during masked
12 presentations of emotional faces while the dorsal/central amygdala was activated during
13 unmasked presentations. Thus, there may be subregion specificity within the amygdala
14 when processing unconscious versus conscious emotionally-valenced stimuli.

15 When gender has been examined as a factor in populations of healthy subjects, higher trait
16 anxiety is associated with greater amygdala responses to unattended fearful faces in female
17 but not male participants [22]. A further factor potentially mediating the relationship
18 between trait anxiety and amygdala reactivity appears to be perceived social support. To
19 illustrate, Hyde et al. [17] show a positive correlation between the degree of trait anxiety and
20 amygdala reactivity to fearful faces in subjects that report below-average social support, but
21 not in those who report above average support. Related, it is also thought that the degree of
22 social anxiety rather than trait anxiety may be more closely related to amygdala reactivity to
23 emotional faces [23]. These factors, and other similar considerations, may explain why some,
24 but not all, studies show a positive correlation between trait anxiety and amygdala
25 reactivity in non-patient populations [18-21,23].

26 **2.2. Amygdala reactivity in anxiety disorders**

27 Hyperactivity of the amygdala in response to negatively-valenced stimuli also appears to be
28 a common finding from a variety of clinical anxiety populations [16]. For example,
29 individuals suffering from social anxiety disorder show heightened amygdala responses to
30 both social and non-social highly emotive stimuli as compared to healthy control groups,
31 with the degree of social anxiety positively correlated with amygdala reactivity [24-27].
32 Furthermore, activation of the amygdala by non-social stimuli has been correlated with trait
33 anxiety in social anxiety disorder, leading to the conclusion that social anxiety disorder is
34 characterized by a more general dysfunction in emotional processing in addition to altered
35 processing of social stimuli and situations [26]. Importantly, reduced symptoms in a public
36 speaking situation following either CBT or antidepressant treatment was associated with
37 reduced amygdala reactivity [24], further suggesting a tight link between symptomology
38 and amygdala reactivity in social anxiety disorder.

39 Like social anxiety disorder, a commonly replicated finding from various PTSD populations
40 is hyperactivity of the amygdala in response to masked fearful faces or trauma-related

1 stimuli [3,28,29]. This manifests as higher amygdala reactivity as compared to non-PTSD
2 groups and/or a positive correlation between severity of PTSD symptoms and amygdala
3 reactivity [28,30-33]. Furthermore, in a group of unmedicated acute PTSD subjects (1 month
4 post trauma), the degree of PTSD symptoms also positively correlated with activity of the
5 amygdala in response to masked fearful faces [34]. Thus, amygdala hyperactivity observed
6 in chronic PTSD appears early in the disorder. However, it should be noted that in these
7 same individuals, the degree of PTSD symptoms negatively correlated with activity in the
8 amygdala in response to unmasked fearful faces [34]. This suggests amygdala hypoactivity
9 in response to consciously-processed fearful stimuli in the early stages of PTSD, further
10 implying a dissociation in amygdala activity in response to consciously-processed versus
11 unconsciously-processed fearful stimuli. Interestingly, activity of the amygdala in response
12 to fearful stimuli might not only be characteristic of PTSD, but might predict treatment
13 outcome. Bryant et al [33] show that individuals diagnosed with PTSD that do not respond
14 to CBT (8 one weekly sessions) show significantly greater pre-treatment amygdala
15 activation in response to masked fearful faces as compared to those PTSD subjects who did
16 respond to CBT, as defined by a 50% or more reduction in scores on the Clinician-
17 Administered PTSD Scale (CAPS). Therefore, hyper-function of the amygdala might provide
18 a useful tool for future selections of treatment options for PTSD.

19 Similar to PTSD and social anxiety disorder, amygdala hyperactivity as a result of highly
20 emotional stimuli presentation or symptom provocation has been observed in specific
21 phobia, panic disorder, and OCD [35-38]. Given the prevalence of GAD, it is surprising that
22 few studies have assessed amygdala reactivity in GAD participants. Somewhat more
23 surprising is that of those studies that have determined amygdala activity in response to
24 emotive stimuli in adult GAD populations, a lack of amygdala hyperactivity has been
25 observed [27,39,40]. This stands in contrast to findings from pediatric GAD, where
26 hyperactivity of the amygdala is apparent in response to emotional stimuli and positively
27 correlated with symptom severity [41,42]. However, recent findings examining amygdala
28 function within paradigms that elicit anticipatory anxiety or emotional conflict have
29 implicated a role for amygdala hyper-reactivity in adult GAD populations. For example,
30 Nitschke et al. [43] report greater anticipatory amygdala activation in response to both
31 emotional and neutral images in adult GAD subjects. Furthermore, Etkin et al [44] found
32 that adult participants with GAD exhibited poor performance on a task that involved
33 emotional conflict (incongruent visual emotional stimuli), accompanied by a failure of the
34 frontal cortex to exert negative top-down control of amygdala activity (see Section 3.1 for
35 more on top-down control of the amygdala). Therefore, amygdala hypofunction in adult
36 GAD might be better revealed by imaging studies that create anxiogenic or conflict
37 situations, rather than the standard presentation of fearful stimuli. While this conclusion
38 requires direct testing, the findings that anxiogenic but not fearful stimuli reveal
39 hypofunction of the amygdala in GAD, whereas fearful stimuli consistently elicit amygdala
40 hyper-reactivity in other anxiety disorders (such as social anxiety disorder, PTSD and also
41 pediatric GAD), suggests a neural dichotomy between GAD and other anxiety disorders on
42 the anxiety to fear continuum.

1 In summary, there appears to be reasonable overlap across various experimental paradigms
2 and study populations to conclude that the amygdala is reactive to fearful stimuli and
3 anxiogenic situations, and exhibits hyper-function to emotive stimuli, anxiogenic situations
4 and/or symptom provocation in anxiety disorders. However, which neurotransmitters and
5 subregions of the amygdala mediate these responses is often better answered by animal
6 studies, where spatial and neurochemical resolution is greatly improved over human
7 imaging studies.

8 **3. Amygdala subregions, connectivity, neurotransmission and** 9 **fear/anxiety**

10 **3.1. The role of amygdala subregions in mediating fear and anxiety**

11 As discussed above, hyper-function of the amygdala appears to be a key component of
12 human anxiety disorders. However, the contribution of particular amygdalar subregions in
13 the development and maintenance of this hyperactive state in humans is still being
14 established. Only very recently have refinements in the acquisition and analysis of fMRI
15 data allowed subregion function to be segregated effectively during emotional tasks such as
16 avoidance learning [45] and facial expression recognition [21,46]. Similarly, effective
17 structural identification of human amygdalar subregions and assessment of their functional
18 connectivity using imaging techniques is still fairly new [for example, see 47-51]. Therefore,
19 most of our understanding of causal neurochemical pathways in amygdalar circuitry related
20 to fear and anxiety has derived from extensive studies using rodent and non-human primate
21 models [for example, see 9,52-58].

22 Anatomical arrangement of the mammalian amygdala appears to have been evolutionarily
23 conserved, with particular subregions being connected to homologous brain structures
24 across species [as reviewed by 59]. The lateral (LA) nucleus of the amygdala is reciprocally
25 connected with the auditory, somatosensory and visual sensory association centers in the
26 temporal and insular cortices [59], and in rats also receives further auditory information via
27 projections from the posterior thalamus [59,60]. The medial amygdala (MeA) is reciprocally
28 connected with the accessory olfactory bulb and many hypothalamic and preoptic nuclei
29 [59,61], creating a locus for assimilation of olfactory stimuli and information regarding
30 internal hormonal state [62,63]. Information summated within the LA and MeA is then
31 conveyed to the adjacent basal (B) and accessory basal (AB) nuclei [64], which also receive
32 projections from the CA1 and subiculum areas of the ventral hippocampus [65-67]. The
33 B/AB nuclei send excitatory and inhibitory projections back to the LA and MeA [64,68],
34 creating a localized circuit that may assist in fine-tuning the filtering of sensory input into
35 these regions [64]. Excitatory projections from this basolateral (BLA) complex target the
36 central nucleus of the amygdala (CeA) either directly or via a series of GABAergic
37 interneurons known as intercalated (ITC) cells located between the BLA and CeA [69],
38 providing an effective means of gating CeA activity and output through a combination of
39 direct excitation and feed-forward inhibition [64,70,71]. The CeA itself, principally the medial
40 sector, sends GABAergic projections to brainstem, hypothalamic and basal forebrain regions

1 that control expression of autonomic, hormonal and behavioral responses to emotive
2 situations [72,73]. It should also be noted that in addition to activating the CeA, the BLA
3 projects to the adjacent bed nucleus of the stria terminalis (BNST), which in turn targets many
4 of the same regions as the CeA to produce similar behavioral and physiological responses [73].
5 The MeA is also able to regulate these responses not only via its influence on hypothalamic
6 nuclei and brainstem targets, but by modulating activity in the BNST and CeA [61,64].

7 The functional connectivity between the BLA, MeA and CeA ensures that sensory and
8 contextual information associated with emotional situations, such as fearful or anxiogenic
9 circumstances, is channeled to effector regions to produce appropriate responses necessary
10 for survival. The BLA and CeA, unlike the MeA, do not appear necessary for expression of
11 unconditioned fear responses to olfactory stimuli in rodents, e.g., to novel presentation of
12 predator odor [74-76], although the BLA does appear to play a role in responses to other
13 types of unconditioned stimuli [77,78]. However, the functional arrangement of the BLA and
14 CeA with other regions facilitates learning about the situation, such that appropriate
15 reactions are maintained if cues associated with initial exposure are experienced again. The
16 BLA in particular appears to play a crucial role in encoding positive or negative salience to
17 relevant stimuli for future reference, as indicated by numerous studies showing that the
18 BLA is required for fear learning and acquisition of conditioned fear responses [see 56,60].
19 Once fear conditioning is acquired, the CeA is necessary for expression of the conditioned
20 response [56,60], the magnitude of which will be influenced by BLA gating of CeA activity
21 and output. Similarly, the BLA is needed for acquisition and expression of fear extinction
22 [79,80], which requires a subject to learn that expression of a previously conditioned fear
23 response is no longer necessary when the conditioned stimulus no longer predicts an
24 aversive event [57,81]. To achieve this, the BLA must integrate new sensory information
25 (absence of the unconditioned aversive stimulus) that will result in a dampening of CeA
26 excitation. This may result from increased BLA excitation of ITC cells during fear extinction
27 acquisition to enhance feed-forward inhibition of the CeA [79,82,83], followed by structural
28 remodeling within the BLA during consolidation of the extinction memory to inhibit later
29 BLA output [79]. However, while the roles of the BLA and CeA in fear behaviors are well
30 established, their contribution to anxiety is less clear, especially for the CeA. Animal studies
31 suggest that changes in BLA and CeA activity can alter state anxiety [9; also see Section 3.2.].
32 However, most investigations have focused on the BLA with the exact role of the CeA
33 remaining ill-defined [for example, see 84,85], although it appears that BLA to CeA circuitry
34 can directly regulate anxiety-like behavior as measured on the elevated plus maze [EPM,
35 86]. This direct control is thought to result from BLA excitation of GABAergic neurons in the
36 lateral CeA to induce feed-forward inhibition of output from the medial CeA [86], similar to
37 that induced by BLA excitation of ITC cells during fear extinction. Thus, suppression of CeA
38 output may be equally important for mediating expression of both fear and anxiety.
39 Alternatively, some studies have suggested that it is BLA activation of the BNST, not of the
40 CeA, that is responsible for mediating anxiety-like behavior as measured using light-
41 potentiated startle responses in rodents [56,87,88]. Startle responses are also potentiated by
42 corticotropin releasing factor (CRF) infused into the BNST [56]. This effect is presumed to
43 result through facilitation of glutamate release from BLA afferents by CRF neurons that

1 originate in the lateral CeA [88,89], implying that even if BNST is the principal output center
2 for certain types of anxiety-like behaviors, the CeA may still play some modulatory role.
3 Furthermore, the MeA has been strongly implicated in animal models of state anxiety [for
4 example, see 90-93 and see Section 3.2], but whether its effects involve modulation of CeA
5 activity is unknown. To direct translational research into the neurological underpinning of
6 anxiety disorders more effectively, animal studies employing as wide a range of state
7 anxiety paradigms as possible, along with animal models that generate trait anxiety, are
8 required to establish the exact nature of CeA involvement and of amygdala subregion
9 interplay in mediating anxiety-like behavior.

10 It is important to remember that while the amygdala can mediate fear and anxiety-like
11 behavior, other brain regions play a major role in expression of these states, presumably by
12 influencing activity in particular amygdalar subregions to alter the balance of output from
13 the CeA. For example, input from the ventral hippocampus to the B/AB nuclei within the
14 BLA is required for expression of conditioned fear responses to contextual cues in rodents
15 and humans [60,94,95], and so receipt of this information presumably increases BLA
16 activity, to in turn enhance CeA output in the aversive context. In rodents, the ventromedial
17 prefrontal cortex (vmPFC) also appears to be crucial in regulating amygdalar activity,
18 especially during fearful experiences [79]. The prelimbic (PL) subregion of the vmPFC can
19 enhance conditioned fear expression via excitatory projections to the BLA and CeA [96-98].
20 In contrast, expression of conditioned fear appears to be decreased by activation of the
21 infralimbic (IL) subregion of the vmPFC [99, but see 100]. The IL cortex is also required for
22 effective consolidation and recall of fear extinction memories [79,98]. Both decreased
23 conditioned fear responding and fear extinction require suppression of CeA output, which
24 is thought to result in part via IL cortex stimulation of the series of inhibitory ITC cells that
25 project to the CeA [71,79,96,101]. The bidirectional roles of the PL and IL cortices in
26 regulating conditioned fear through opposing influences on CeA activity and output imply
27 that imbalance in the influence of either cortical structure could contribute to amygdala
28 hyperactivity seen in anxiety disorders characterized by excessive and inappropriate fear
29 (see Table 1). This is supported by fMRI studies investigating neural correlates of impaired
30 fear extinction in PTSD patients, who compared to healthy subjects show hyperactivity of
31 the amygdala during extinction learning [102]. This enhanced amygdala function in PTSD
32 patients is accompanied by greater activation of the dorsal anterior cingulate cortex (dACC,
33 functionally equivalent to the rodent PL cortex, [3,57], which is also present during recall of
34 the extinction memory [102]. This is in line with rodent studies demonstrating potentiated
35 fear conditioning upon PL cortex activation [98]. However, PTSD individuals exhibit
36 hypoactivation of the ventral portion of the vmPFC (equivalent to rodent IL cortex, [3,57])
37 during extinction learning and recall [102,103]. Human imaging studies also suggest that
38 impaired regulation of amygdala activity by the ventral vmPFC may contribute to anxiety
39 disorders characterized by hypervigilance in the absence of conditioned stimuli, such as in
40 GAD. Specifically, the strength of the connection between the vmPFC and the amygdala, as
41 measured using diffusion tensor imaging, predicts levels of self-reported trait anxiety, such
42 that weaker connections are seen in more anxious individuals [104]. As mentioned earlier
43 (Section 2.2), participants with GAD exhibited a failure of the vmPFC to exert negative top-

1 down control of amygdala activity during a task that involved emotional conflict [44].
2 Further, resting state fMRI revealed that in anxious individuals, vmPFC activity was
3 negatively correlated with amygdala activity, while a positive relationship was observed for
4 low anxious subjects [105]. The combination of animal and human studies strongly indicates
5 that inadequate suppression by the ventral portion of the vmPFC, most likely of the CeA, is
6 a key factor in amygdala hyperactivity underlying the emergence of excessive fear and
7 anxiety states.

8 **3.2. Monoaminergic neurotransmission in the amygdala: Relation to fear and** 9 **anxiety**

10 The monoamine neurotransmitters (serotonin, dopamine and norepinephrine) have long
11 been associated with fear and anxiety, and drugs that alter monoaminergic function are
12 often effective across the range of anxiety disorders [8, 9, 52, 55]. Animal studies suggest a
13 variety of anxiogenic stressors or fearful stimuli increase monoamine levels in the amygdala.
14 To illustrate, increased serotonin (5-HT) release or increased activity of 5-HT neurons in the
15 amygdala have been observed in response to restraint or footshock, or in association with
16 expression of conditioned fear behavior [106-110]. Similarly, dopamine (DA) and
17 norepinephrine (NE) levels in the amygdala are increased following restraint, handling
18 stress, footshock or during the expression of conditioned fear behavior [107,111-118]. The
19 source of monoamines to the amygdala arise from monoaminergic cell body regions in the
20 brainstem. Specifically, the dorsal raphe nucleus (dRN) provides 5-HT innervation to the
21 amygdala, while NE and DA innervation of the amygdala arise from the locus coeruleus
22 (LC) and ventral tegmental area (VTA) respectively [55,119,120]. Regulation of
23 monoaminergic activity in the amygdala thus can occur at the level of these brainstem cell
24 body regions, or within the terminal regions of the amygdala.

25 One of the important mediators of amygdala monoaminergic activity in response to
26 anxiogenic or fearful stimuli is CRF. A strong body of evidence implicates central CRF in
27 mediating fear and anxiety [12,121-128], and recent clinical studies suggest an important
28 role for CRF in anxiety disorders [129]. Like anxiogenic and fearful stimuli, central infusion
29 of CRF or CRF receptor agonists increases 5-HT, NE and DA levels in the amygdala [130-
30 133], and stress-induced increases in monoamine levels in the amygdala are prevented by
31 CRF receptor antagonists [108,111]. It is thought that CRF regulation of monoaminergic
32 activity in the amygdala occurs at the level of the monoaminergic cell bodies. The
33 monoaminergic cell body regions receive CRF innervation from the CeA and BNST, and
34 CRF type 1 and 2 (CRF₁ and CRF₂) receptors are localized to the dRN, LC and VTA [134-
35 140]. Direct infusion of CRF or CRF receptor agonists into the dRN stimulates 5-HT release
36 in the CeA or BLA [131-133]. Interestingly, CRF-induced 5-HT release in the amygdala
37 appears to be dependent on CRF₂ receptor activation in the dRN [131,133], and CRF₂
38 receptors are known to increase 5-HT neuronal firing rates in the dRN [141]. Importantly,
39 increased neuronal surface expression of CRF₂ receptors occurs in the dRN as a result of
40 stress [142], and increased expression of CRF₂ receptors in the dRN has been observed in rat
41 models of high anxiety [11,128,137,143]. Furthermore, CRF₂ receptor antagonists infused

1 directly into the dRN reduce heightened anxiety-like behavior in rat models of
2 amphetamine withdrawal or early life stress [12,128]. Combined, these findings suggest that
3 CRF₂ receptor modulation of 5-HT activity in the amygdala may play an important role in
4 heightened anxiety. While similar studies have not been performed to elucidate the role of
5 CRF receptors in the LC and VTA in mediating NE and DA activity in the amygdala and
6 anxiety states, some indirect evidence suggests an important role for CRF receptors in the
7 LC and VTA stress responses [136,138,144]. Overall, it is clear that further investigations are
8 needed to ascertain the role of CRF receptors in mediating NE and DA activity in the
9 amygdala and how CRF modulation of this activity could relate to fear or anxiety.

10 Studies demonstrating increased monoamine activity in the amygdala in response to
11 anxiogenic or fearful stimuli, and CRF modulation of these responses (as described above)
12 do not allow conclusions to be made about the specific role of each monoamine in mediating
13 anxiety or fear. Direct manipulation of monoaminergic activity within the amygdala or
14 specific amygdala subregions, and the measurement of resultant anxiety-like or fear-related
15 behaviors, have gone some way to providing a picture of how monoamine function in the
16 amygdala might translate to anxiety or fear. Table 2 summarizes such studies directly
17 manipulating 5-HT levels or 5-HT receptor activity in the amygdala. When 5-HT or 5-HT
18 activity is decreased in the entire amygdala [145,146], a consistent increase in anxiety-like
19 behavior is observed (Table 2). This would suggest that increased 5-HT activity in the
20 amygdala would thus be associated with decreased anxiety, implying an anxiolytic role of 5-
21 HT. However, this does not appear to be supported by experiments that directly manipulate
22 5-HT receptor activity in the amygdala with 5-HT receptor ligands (Table 2). For example,
23 activation of postsynaptic excitatory 5-HT₂ or 5-HT₃ receptors in the amygdala decreases
24 social interaction and increases anxiety-like behavior, whereas antagonism of 5-HT₃
25 receptors in particular increases social interaction and decreases anxiety-like behaviors,
26 suggesting that 5-HT actions on postsynaptic receptors is anxiogenic (Table 2), although, see
27 [147] for an exception to this pattern. Similarly, activation of excitatory 5HT₂ receptors in the
28 BLA generally increases anxiety-like behavior (Table 2), suggesting an anxiogenic role for
29 postsynaptic 5-HT receptors in the BLA (although an exception to this is observed, [148]). In
30 contrast, inhibitors of 5-HT₂ receptors in the MeA increase anxiety-like behavior while
31 activation of these receptors increases social interaction and decreases anxiety behavior
32 (Table 2). Thus like the some findings from the amygdala as a whole (Table 2), 5-HT activity
33 in the MeA appears to play an anxiolytic role. The role of 5-HT or 5-HT receptors has not
34 been well studied in the CeA. However, rats undergoing amphetamine withdrawal that
35 exhibit greater anxiety-like behavior have greater 5-HT release in the CeA [12,133],
36 suggesting a similar anxiogenic relationship between 5-HT and anxiety as for the BLA.
37 Future work should determine whether 5-HT in the CeA reduces anxiety-like behaviors as is
38 suggestive for the MeA, or in contrast, increases anxiety-like behaviors as appears to be the
39 case for the BLA. Overall, the findings summarized in Table 2 suggest a dichotomy in the
40 potential role of 5-HT in the amygdala in mediating anxiety depending on whether the
41 entire amygdala or a specific subregion is targeted. Potential confounds in comparing the
42 studies listed in Table 2 could be the different paradigms used to measure anxiety-like

1 behaviors and the relative selectivity of 5-HT receptor ligands across different experiments.
 2 Future studies directly comparing the effects of 5-HT manipulations within the different
 3 amygdala subregions across several well-validated tests of anxiety-like behaviors will better
 4 elucidate the role of amygdala 5-HT in mediating anxiety.
 5

Amygdala Subregion	Monoamine or Receptor Involvement	Behavioral Outcome	Citation
<i>Anxiety-like Behavior</i>			
Amygdala	Decreased 5-HT (induced by MDMA)	Increased anxiety behavior	Faria et al. [145]
Amygdala	Decreased 5-HIAA (induced by stress)	Increased anxiety behavior	Niwa et al. [146]
Amygdala	5-HT _{1A} agonist	No change in anxiety behavior	Zangrossi and Graeff [149]
Amygdala	5-HT _{2B/2C} agonist	Increased anxiety behavior	Cornelio and Nunes-De-Souza [150]
Amygdala	5-HT ₃ agonist	Decreased social interaction	Higgans et al. [151]
Amygdala	5-HT ₃ agonist	Decreased anxiety behavior	Costall et al. [147]
Amygdala	5-HT ₃ antagonist	Increased social interaction	Higgans et al. [151]
Amygdala	5-HT ₃ antagonist	Decreased anxiety behavior	Costall et al. [147]
Amygdala	5-HT ₃ antagonist	Decreased anxiety behavior	Tomkins et al. [152]
BLA	5-HT _{1A} agonist	Decreased social interaction	Gonzalez et al. [153]
BLA	5-HT _{1A} agonist	No change in anxiety behavior	Gonzalez et al. [153]
BLA	5-HT _{2A} agonist	Increased anxiety behavior	Zangrossi and Graeff [149]
BLA	5-HT _{2A/2C} agonist	No change in anxiety behavior	Cruz et al [148]
BLA	5-HT _{2C} agonist	Increased anxiety behavior	Vincente et al. [154]
MeA	5-HT _{2A} antagonist	Increased anxiety behavior	Zangrossi and Graeff [149]
MeA	5-HT ₂ agonist	No change in anxiety behavior	Duxon et al. [155]
MeA	5-HT _{2B} agonist	Increased social interaction	Duxon et al. [156]
MeA	5-HT _{2B} agonist	Decreased anxiety behavior	Duxon et al. [155]

Amygdala Subregion	Monoamine or Receptor Involvement	Behavioral Outcome	Citation
MeA	5-HT _{2B/2C} agonist	No change in anxiety behavior	Duxon et al. [155]
<i>Fear-related Behavior</i>			
CeA	Increased 5-HT	Increased unconditioned freezing	Forster et al. [132]
BLA	Increased 5-HT	Decreased conditioned freezing	Inoue et al. [157]
BLA	Increased 5-HT	Decreased unconditioned tonic immobility	Leite-Panissi et al. [158]
BLA	5-HT _{1A} agonist	Decreased conditioned freezing	Li et al. [159]
BLA	5-HT _{1A} agonist	Decreased acquisition and expression of conditioned defeat	Morrison et al. [160]
BLA	5-HT _{1A/2} agonist	Decreased unconditioned tonic immobility	Leite-Panissi et al. [158]

1 Abbreviations: 5-HIAA = 5-Hydroxyindoleacetic acid (5-HT metabolite); 5-HT = serotonin; BLA = basolateral
 2 amygdala; CeA = central nucleus of the amygdala; MDMA = 3,4-methylenedioxy-N-methylamphetamine; MeA =
 3 medial amygdala.

4 **Table 2.** The Role of Serotonin in Anxiety-Like and Fear-Related Behaviors

5 Determining the role of amygdala 5-HT in fear-related behavior has mainly utilized studies
 6 of freezing or immobility responses in rodents, and of 5-HT manipulation in the BLA (Table
 7 2). From these studies, it seems clear that 5-HT in the BLA decreases the expression of
 8 unconditioned and conditioned fear responses, likely via activation of the inhibitory
 9 postsynaptic 5-HT_{1A} receptor (Table 2). Thus, it has been suggested that 5-HT in the
 10 BLA/amygdala ameliorates fear [8]. This conclusion is in contrast to the apparent role for
 11 BLA 5-HT in enhancing anxiety (Table 2), suggesting a fear versus anxiety dissociation for
 12 the role of 5-HT in the BLA. This dissociation, if upheld by more in-depth future work,
 13 could prove important information for the development of treatment strategies for the
 14 various anxiety disorders that differ in the degree of anxiety-like and fear-like
 15 symptomology (as discussed in Section 1.1).

16 A role for amygdala DA in anxiety has not been as well explored as for 5-HT. However, a
 17 summary of studies that have manipulated DA function in the amygdala provides a
 18 consistent picture of the role of amygdala DA in mediating anxiety in animal models (Table
 19 3). Indirect evidence suggests that decreased DA in the amygdala leads to increased anxiety,
 20 and this is supported by direct manipulation of the CeA (Table 3). For example, decreased
 21 DA or DA receptor antagonism within the CeA all increase anxiety-like behavior (Table 3),

1 suggesting that DA activity in the CeA is anxiolytic. This role for DA in the CeA is in direct
 2 contrast to the BLA, where converging evidence suggests that decreased DA function in the
 3 BLA decreases anxiety-like behaviors while increased DA receptor activity in the BLA
 4 increases anxiety (Table 3). Thus, DA activity in the BLA is anxiogenic, revealing an
 5 opposite role for DA activity in the CeA and BLA in mediating anxiety-like behaviors in
 6 animal models.
 7

Amygdala Subregion	Monoamine or Receptor Involvement	Behavioral Outcome	Citation
<i>Anxiety-like Behavior</i>			
Amygdala	Decreased DA	Decreased rearing in open field indicative of increased anxiety behavior	Summavielle et al. [163]
CeA	Decreased DA	Decreased voluntary activity indicative of increased anxiety behavior	Izumo et al. [164]
CeA	D ₁ antagonist	Increased anxiety behavior	Rezayof et al. [165]
CeA	D _{2/3} antagonist	Increased anxiety behavior	de la Mora et al. [166]
BLA	DA depletion	Decreased anxiety in males but not females	Sullivan et al. [167]
BLA	D ₁ agonist	Increased anxiety behavior	Banaej et al. [168]
BLA	D ₂ agonist	Increased anxiety behavior	Banaej et al. [168]
BLA	D ₁ antagonist	Decreased anxiety behavior	Banaej et al. [168]
BLA	D ₁ antagonist	Decreased anxiety behavior	de la Mora et al. [169]
BLA	D ₂ antagonist	Decreased anxiety behavior	Banaej et al. [168]
<i>Fear-related Behavior</i>			
Amygdala	D ₂ antagonist	Decreased acquisition and retention of fear conditioning	Greba et al. [170]
CeA	D ₁ agonist	Increased conditioned fear behavior	Guarraci et al. [171]
CeA	D ₁ antagonist	Inhibited conditioned fear behavior	Guarraci et al. [171]
CeA	D ₂ antagonist	Decreased	Guarraci et al. [172]

Amygdala Subregion	Monoamine or Receptor Involvement	Behavioral Outcome	Citation
		conditioned fear behavior	
BLA	DA depletion	Decreased fear conditioning	Seldon et al. [173]
BLA	D ₁ antagonist	Inhibited acquisition of fear conditioning	Greba and Kokkinidis [174]
BLA	D ₂ antagonist	Inhibited fear potentiated startle	De Oliveira et al. [175]

1 Abbreviations: BLA = basolateral amygdala; CeA = central nucleus of the amygdala; DA = dopamine.

2 **Table 3.** The Role of Dopamine in Anxiety-Like and Fear-Related Behaviors

3 In contrast, the role of DA in mediating fear-related behaviors does not appear to differ
 4 based on amygdala subregion (Table 3). Reducing DA function in the amygdala reduces or
 5 inhibits processes associated with fear conditioning, while increasing DA receptor activity
 6 increases conditioned fear (Table 3). Thus, DA in the amygdala is required for fear
 7 conditioning, and enhanced DA levels in the amygdala as elicited by fearful stimuli and
 8 conditioned cues [107,112] would thus facilitate fear conditioning. It should be noted that
 9 the studies summarized by Table 3 indicate a role for both excitatory D₁ receptors and
 10 inhibitory D₂ receptors. Dopamine D₂ receptors are localized both pre- and post-
 11 synaptically, with pre-synaptic D₂ autoreceptors limiting DA neuronal activity and DA
 12 release [161,162]. Thus, antagonism of presynaptic D₂ receptors would actually increase DA
 13 within the amygdala. Since the effects of D₂ receptor antagonism on fear-related behaviors is
 14 characteristic of reduced, not enhanced, DA function in the amygdala, it may be concluded
 15 that the results of D₂ receptor antagonism summarized by Table 3 are due to postsynaptic D₂
 16 receptor effects. However, this conclusion requires direct testing.

17 Very few studies have examined the role of amygdala NE in mediating anxiety-like
 18 behavior in animal models, surprising given that anxiogenic stimuli increase NE in this
 19 region [for example, see 111,115,116] and drugs that alter NE neurotransmission are used to
 20 treat anxiety disorders [8]. There appears to be little role for NE receptors in the CeA in
 21 mediating anxiety-like behavior, although infusion of a α_1 antagonist can increase social
 22 interaction following an anxiogenic stimulus [restraint; 176; Table 4]. It is clear that more
 23 experiments are required to delineate the role of amygdala NE in mediating anxiety.

24 Studies determining the role of NE in fear-related behaviors have concentrated on the BLA,
 25 due to the importance of this amygdala subregion in conditioned fear responses (see Section
 26 3.1.). The major focus of the studies summarized by Table 4 has been on the role of NE in
 27 fear conditioning and reconsolidation of fear memories in conditioned fear paradigms.
 28 Taken as a whole, findings suggest that NE in the BLA facilitates fear conditioning and fear
 29 memory, via activation of adrenergic β receptors (Table 4). Recent evidence suggests a role
 30 for α_1 receptors in the BLA in mediating fear memory, in this case, activation of α_1 receptors
 31 by NE would appear to decrease fear memory (Table 4). Thus, it is possible that NE in the

1 BLA could have opposing effects on reconsolidation of fear memory based on the balance of
 2 α_1 versus β receptor activity – a hypothesis that requires direct testing. The role of NE in the
 3 BLA (and β receptors in particular) in fear memory has generated interest in targeting this
 4 NE system for the treatment of anxiety disorders where enhancement in fear memory is
 5 apparent, such as PTSD [for example, see 177]. Whether NE within the BLA plays a role in
 6 other aspects of fear processing (e.g. unconditioned fear responses to non-olfactory based
 7 stimuli) or NE within other amygdala subregions mediate fear should be subjects of future
 8 investigations to fully elucidate the role of amygdala NE in fear.

9

Amygdala Subregion	Monoamine or Receptor Involvement	Behavioral Outcome	Citation
<i>Anxiety-like Behavior</i>			
CeA	α_1 antagonist	Increased social interaction	Cecchi et al. [176]
CeA	α_1 antagonist	No effect on anxiety behavior	Cecchi et al. [176]
CeA	$\beta_{1/2}$ antagonist	No effect on social interaction	Cecchi et al. [176]
CeA	$\beta_{1/2}$ antagonist	No effect on anxiety behavior	Cecchi et al. [176]
<i>Fear-related Behavior</i>			
BLA	Increased NE	Increased memory and retention of fear conditioning	LaLumiere et al. [178]
BLA	Decreased NE	Impaired fear conditioning	Seldon et al. [173]
BLA	Decreased NE	Impaired fear memory	Debiec and LeDoux [177]
BLA	α_1 antagonist	Increased fear memory	Lazzaro et al. [179]
BLA	$\beta_{1/2}$ antagonist	Impaired of fear memory	Debiec and LeDoux [180]
BLA	β_1 antagonist	Impaired fear memory (as enhanced by glucocorticoids)	Roosendaal et al. [181]

10 Abbreviations: BLA = basolateral amygdala; CeA = central nucleus of the amygdala; NE = norepinephrine.

11 **Table 4.** The Role of Norepinephrine in Anxiety-Like and Fear-Related Behaviors

12 In summary, it is clear that more work is required to fully understand the role of amygdala
 13 monoamines in mediating fear and anxiety. However, several patterns of interest emerge
 14 from the current literature, namely that there are distinct subregion differences in the role

1 each monoamine plays in mediating anxiety and fear, with the one monoamine possibly
2 playing opposing roles depending on subregion or depending on whether anxiety or fear
3 measures are employed. Therefore, these findings suggest neurochemical dissociations
4 between amygdala subregions and monoamines in mediating fear or anxiety.

5 **4. The amygdala as a potential site of anxiolytic drug action**

6 Psychopharmacological management of anxiety disorders includes the benzodiazepines,
7 antidepressants, 5-HT_{1A} agonists and various “off-label” drugs such as β -blockers, mood
8 stabilizers and antipsychotics. The mechanism by which these drugs produce anti-anxiety
9 effects has yet to be definitively established and represents a frequently updated field of
10 research. Because these drugs bind to target receptors throughout the brain, it is unlikely
11 that their efficacy can be attributed to action in one particular region. However, given the
12 role that the amygdala plays in fear and anxiety, modification of amygdala function by
13 pharmacological agents represents a likely mechanism of action as well as a target to guide
14 future drug development. The evidence for amygdala involvement in anxiolytic action
15 comes from both human imaging studies as well as work in animal models.

16 **4.1. Human imaging studies: Effects of anxiolytics on amygdala activity and** 17 **emotion**

18 Given the highly complex and subjective nature of anxiolytic drug response in humans,
19 neuroimaging represents an invaluable tool for drug evaluation and discovery.

20 *Benzodiazepines:* Benzodiazepines exert their anxiolytic action through binding to GABA_A
21 receptors, which leads to enhanced GABA activity and a subsequent increase in inhibitory
22 tone. Despite the long history and current prevalence of benzodiazepine use for anxiety
23 disorders [182,183], there is a paucity of human neuroimaging studies utilizing this class of
24 drug, especially compared to those using antidepressants. This may have to do with eclipse
25 of benzodiazepines by antidepressants as first line agents for many anxiety disorders [182].

26 Various studies have utilized healthy volunteers undergoing experimental challenges in an
27 attempt to elucidate the neurobiology underlying the anxiolytic effect of benzodiazepines.
28 These studies have found that benzodiazepines have the ability to impair functions related
29 to amygdala activity including fear conditioning [184-186], recognition of fearful emotional
30 faces [187], and memory for emotional stimuli relative to neutral stimuli [188,189].

31 Neuroimaging work appears to support a role for the amygdala in benzodiazepine action,
32 although this may be dependent upon the nature of the accompanying neuropsychological
33 challenge. Specifically, lorazepam was found to decrease amygdala activation during an
34 emotional face assessment task without modifying baseline levels of anxiety or task
35 recognition [190]. A similar finding was found with diazepam, which decreased amygdala
36 response to fearful faces, and also impaired fearful face recognition [191]. However, during
37 anticipation of aversive electrical stimulation, lorazepam failed to produce changes in
38 amygdala activity [192]. Thus, while there is support for benzodiazepine induced

1 modulation of the amygdala during processing of threatening/emotional stimuli, further
2 studies are needed to clarify the neural correlates of benzodiazepine-induced anxiolysis.

3 *β-Blockers:* The β-blocker propranolol has a substantial history of being utilized to reduce
4 somatic symptoms of fear and anxiety in situations such as stage fright [193] and acute panic
5 [194-195]. More recently, research on the role of amygdala NE and β-receptors in facilitating
6 emotional memory formation (see Table 4 and associated text) has caused much excitement
7 and controversy about the use of propranolol to prevent PTSD [196-198]. Thus far, initial
8 trials have demonstrated limited efficacy [199,200]. Despite lack of success in the application
9 of propranolol to PTSD, neuroimaging studies in healthy human subjects have confirmed
10 the ability of propranolol to modulate amygdala activation to emotional stimuli.
11 Propranolol was found to decrease amygdala activation to emotional faces irrespective of
12 emotional valence [201]. Furthermore, supporting a role for the amygdala NE in the
13 encoding and consolidation of emotional stimuli, a separate study found that propranolol
14 was able to decrease amygdala reactivity to emotional pictures of high valence as well as
15 decrease the subject's memory for them [202].

16 *Selective Serotonin Re-uptake Inhibitors:* Antidepressant drugs, and selective serotonin re-
17 uptake inhibitors (SSRIs) in particular, have become first line drugs for many of the anxiety
18 disorders [182,203]. As such, there has been comparatively more work investigating these
19 drugs in humans using advanced imaging techniques.

20 Most antidepressants are unique from benzodiazepines and β-blockers in that a time lag
21 exists between initial treatment and onset of anxiolytic effects. In line with a potential
22 anxiogenic role of serotonin in the amygdala (see Table 2 and associated text), some patients
23 have reported an initial exacerbation of anxiety upon acute dosing of SSRIs [203]. In studies
24 on healthy subjects, acute dosing of the SSRI citalopram can enhance recognition of fearful
25 faces as well as increase emotion-potentiated startle response [204-206]. These effects are
26 reversed when citalopram treatment is continued for 7 days [207,208].

27 Attempts to correlate the acute versus sub-chronic effects of SSRIs with neural activation
28 have resulted in unexpected findings. On one hand, sub-chronic citalopram treatment was
29 found to decrease amygdala activation to unconscious fearful stimuli [209], suggesting a
30 relationship between repeated SSRI treatment, changes in emotional processing, and
31 decreased amygdala activity. However, acute doses of citalopram have also been found to
32 decrease amygdala activation to fearful faces [208,210,211]. Divergent effects of acute versus
33 sub-chronic citalopram on emotional recognition but similar effects on amygdala response
34 could suggest that the amygdala does not play a core role in acute SSRI-induced anxiety or
35 chronic SSRI-induced anxiolysis. However, it has been emphasized that the effects of
36 serotonergic challenge on fear recognition and amygdala activation appear to be dependent
37 upon the individual's baseline sensitivity to threat [212], gender [213] and genotype [214].
38 Thus differences in subject profiles both between and within studies could have confounded
39 results.

40 Overall, it appears that pharmacotherapeutics commonly used to treat anxiety disorders may
41 modulate amygdala function. In particular, it appears that anxiolytics can reduce amygdala

1 reactivity to highly emotive or fearful stimuli. Given that amygdala hyper-reactivity to
2 similar stimuli is the most common finding across all anxiety disorders (with the exception
3 of adult GAD – see Section 2.2), it is possible that the anxiolytic effects of these drugs may be
4 in part, mediated by dampening amygdala function.

5 **4.2. Evidence delineating effects of anxiolytic drugs on amygdala function in** 6 **animal models of anxiety states**

7 *Benzodiazepines:* While benzodiazepine receptors exist throughout the brain, there is a
8 particularly high density in amygdala regions [215,216]. There is much evidence from
9 animal models to suggest that it is the action of benzodiazepines in the amygdala that
10 mediates their anxiolytic effect. For example, early evidence demonstrated that local
11 amygdala infusion of benzodiazepines produces anxiolytic-like effects in conflict models of
12 anxiety [217-220]. These effects can be reversed by systemic [217,219] or direct amygdala
13 administration of benzodiazepine antagonists [220]. Anti-conflict effects are most apparent
14 when the benzodiazepines are injected into the BLA, and are absent when injected into the
15 CeA [219,220]. While anti-conflict effects of benzodiazepines have been observed in the CeA,
16 these were with substantially higher doses [221]. Further studies suggest that the BLA and
17 not the CeA is essential for the anxiolytic effects of benzodiazepines in the EPM
18 [149,222,223]. However, with regards to the shock probe burying test, it appears that the
19 CeA is responsible for benzodiazepine-induced impairment of passive avoidance [223].
20 Although contradictory results exist on the role of benzodiazepines in the BLA versus CeA,
21 particularly when animals are tested on the EPM [9,84,224] have suggested that distinct
22 benzodiazepine receptor subtypes located within subregions of the amygdala may
23 differentially alter avoidance responses to “potential threat” (EPM and BLA) versus
24 “discrete, unambiguous threat” (shock probe burying and CeA).

25 As discussed in the human studies in Section 4.1 above [184,188,189], a key aspect of
26 benzodiazepine action may be the ability to modulate emotional memory. Here the BLA
27 once again appears to be a main site of benzodiazepine action. Lesions of the BLA, but not
28 the CeA, block the benzodiazepine induced deficits in inhibitory avoidance memory
29 [225,226]. Similar impairments were seen by direct injection of benzodiazepine into the BLA
30 and not the CeA [227]. Enhancement of memory consolidation could be induced by BLA
31 infusion of a benzodiazepine antagonist [228]. Given that individuals with anxiety disorders
32 may be hypervigilant to cues associated with threatening stimuli and biased to form
33 memories regarding such stimuli [229,230], the pro-amnestic effects of benzodiazepines in
34 the BLA may represent a putative mechanism of action.

35 *β-Blockers:* The evaluation of β-blockers (with propranolol being the prototypical agent) in
36 animal models has revolved mainly around their utility in models of memory and fear
37 conditioning. Within the BLA, stress hormone elicited increases in norepinephrine have
38 been found to enhance the consolidation of emotionally relevant memories [231,232]. This
39 appears to be particularly true with contextual fear conditioning [178] and reconsolidation
40 of fear memory following extinction [180,197,233; Table 4]. In particular, local infusions of

1 propranolol are able to block reconsolidation of fear [180,233]. Recently, it has been
2 demonstrated that β -adrenoreceptor activation within the BLA decreases surface expression
3 of GABA_A receptors, and this phenomenon is necessary for the reinstatement of fear
4 following extinction [234]. It is proposed that propranolol, through blocking the decrease in
5 GABA_A receptor surface expression, prevents fear reinstatement by maintaining feed
6 forward inhibition from BLA interneurons and thus dampening activity of BLA projections
7 [234]. This finding is noteworthy as it suggests that hyperactive noradrenergic activity in
8 PTSD [235,236] may lead to reduced GABA_A availability, explaining a potential mechanism
9 for the relative ineffectiveness of benzodiazepines in PTSD populations [237,238].

10 Despite the action of β -blockers within the amygdala to modulate fear conditioning (see
11 Table 4), attempts at testing propranolol in other animal models of PTSD have met with
12 mixed results, echoing the mixed efficacy seen thus far in humans [199,200,239,240]. One
13 such model is exposure to predator odor in rodents, which produces long lasting increases
14 in anxiety like behavior [241-243]. The increases in anxiety like behavior following exposure
15 to predator odor is influenced by a long lasting potentiation in BLA activity [243],
16 supporting the role of the amygdala in mediating the consequences of fear and trauma.
17 Propranolol administered 1 minute following exposure to predator odor to rats blocks the
18 development of anxiogenesis in various tests, including the EPM, one week later [241].
19 However, when propranolol administration is delayed to 1 hour following predator odor
20 exposure, no effects are seen when rats are subsequently tested on the EPM 30 days later
21 [242]. These results highlight once again a potential key role of timing if propranolol is to be
22 effectively implemented in clinical patients. Similarly, findings that propranolol seems most
23 effective in blocking the *reconsolidation* of fearful memories [233, 180, 197] (also see Table 4)
24 suggests that future work should be aimed at establishing protocols for the integration of
25 propranolol during exposure therapy, in which extinction and reconsolidation processes are
26 most active. Specifically, it would seem important that propranolol not be administered
27 shortly after exposure therapy, as this might interfere reconsolidation processes within the
28 amygdala. On the other hand, propranolol would likely have utility when PTSD patients
29 encounter aversive stimuli outside the context of therapy which could potentially
30 undermine the therapeutic process and lead to reinstatement.

31 *Selective Serotonin Re-uptake Inhibitors:* Similar to human studies, animal models of anxiety-
32 like behavior demonstrate divergent behavioral effects of acute versus chronic SSRI
33 administration. Increased anxiety-like behavior with acute treatment of SSRIs and its
34 reversal with chronic treatment has been found in novelty-suppressed feeding [244], EPM
35 testing [245,246], and the social interaction test [247]. While a large percentage of studies
36 reveal acute anxiogenic effects and chronic anxiolytic effect, there are exceptions (for review,
37 see [248]).

38 Much evidence suggests that enhanced activity at 5-HT_{2C} within the BLA by SSRIs produces
39 acute anxiogenic effects, while the eventual downregulation of these receptors by chronic
40 treatment leads to eventual anxiolysis. For example, amygdala or BLA 5-HT_{2C} receptors
41 have been found to produce anxiety-like responses in a variety of tests [249,250] (see Table
42 2). Blockade of 5-HT_{2C} receptors within the BLA prevents the acute anxiogenic effect of the

1 SSRI fluoxetine on the Vogel conflict test [251]. Systemic 5-HT_{2c} antagonism also prevents
2 the increase in fear conditioning [252], decrease in social interaction [247,253], and escape
3 response to airjet [254] following acute SSRI treatment. Following chronic treatment with
4 SSRIs, 5-HT_{2c} agonists have attenuated anxiogenic effects on the exacerbation of OCD
5 symptoms in humans [255,256], on social interaction [257] and hyperlocomotion [258],
6 suggesting down-regulation of the ability to 5-HT_{2c} receptors in the amygdala to produce
7 anxiogenic responses following chronic SSRI treatment. Thus, the amygdala (BLA in
8 particular) may be an important locus of action for the long-term effects of SSRIs on anxiety.

9 **4.3. Future potential anxiolytic targets**

10 The literature reviewed above suggests that in part, the effects of anxiolytic drugs may be
11 mediated by altering amygdala function – either global dampening of the amygdala by
12 benzodiazepines, or specific actions on 5-HT and NE receptors within particular amygdala
13 subregions. However, to improve therapeutic efficacy and reduce relapse, several aspects of
14 amygdala pharmacology discussed above might provide useful potential anxiolytic targets
15 in the future.

16 Findings suggesting down-regulation of anxiogenic 5-HT_{2c} receptors in the amygdala
17 following chronic SSRI treatment (Section 4.2.) present a potential strategy of reducing onset
18 latency of SSRIs as well as enhancing their effects. Specifically, blocking 5-HT_{2c} receptors at
19 the initiation of SSRI treatment would be expected to produce a faster onset of anxiolytic
20 action. Currently, there are no selective 5-HT_{2c} antagonists available for human use.
21 However, atypical antipsychotics [259] as well as atypical antidepressants such as
22 mirtazapine [260] possess 5-HT_{2c} antagonist activity. While there is evidence that
23 antipsychotic augmentation of SSRIs may improve anxiolytic efficacy, their use has been
24 limited by poor tolerability [for review see 261]. Although research is lacking, mirtazapine
25 and the melatonin receptor agonist/5-HT_{2c} receptor antagonist agomelatine [262] may
26 provide the advantage of targeting anxiogenic 5-HT_{2c} in the amygdala with less side effects.

27 Furthermore, the recent observation that β -adrenoreceptor activation within the BLA results
28 in decreased of GABA_A receptor surface expression necessary for fear reinstatement [234]
29 (and see Section 4.2.) suggests that the combination of propranolol and a benzodiazepine
30 may have unique benefit for PTSD. By blocking β -adrenoreceptors with propranolol, one
31 might be able to enhance benzodiazepine receptor availability, and increase
32 benzodiazepine-induced inhibition of fear circuits within the amygdala. While currently
33 speculative, the use of propranolol to enhance benzodiazepine action in the amygdala may
34 represent a potential creative treatment strategy in a population that is traditionally
35 refractory to benzodiazepine treatment.

36 While current pharmacotherapeutic strategies for the treatment of anxiety disorders target
37 monoamine function, this has predominantly been related to altering 5-HT or NE levels or
38 receptor activity [8]. However, Table 3 clearly shows a role for DA in the amygdala in
39 mediating both fear and anxiety, and the role for DA and both D₁ and D₂ receptors in
40 acquisition and retention of conditioned fear in particular appears quite robust. Thus,

1 reducing DA function might serve as means by which to treat anxiety disorders in which
2 fear plays a major component. The obvious disadvantage of dopaminergic-based
3 pharmacotherapeutics is potential for major cognitive and motoric side-effects, limiting the
4 treatment options with the currently available dopaminergic agents. Atypical antipsychotic
5 drugs incorporate DA receptor blocking activity while avoiding many of the motoric and
6 cognitive issues of traditional agents. There is evidence that atypical agents possess
7 anxiolytic activity [261], but metabolic side effects make them poorly tolerated. Furthermore,
8 because atypical antipsychotics also have high affinity for 5-HT receptors, the contribution
9 of DA modulation to their anxiolytic effects in humans is currently unknown. One potential
10 strategy may be the use of partial agonists to reduce DA activity in the amygdala via
11 activation of inhibitory presynaptic D₂ autoreceptors. While non-selective for DA, the D₂
12 partial agonist aripiprazole has demonstrated anxiolytic efficacy similar to other atypical
13 antipsychotic drugs [263]. In the future, more selective DA partial agonists may have
14 additional benefit without unwanted side-effects.

15 Finally, CRF has been identified as an important neuropeptide in the regulation of
16 monoaminergic activity in the amygdala in response to anxiogenic or fearful stimuli (Section
17 3.2). Furthermore, CRF and its receptors (CRF₁ and CRF₂) are implicated in fear and anxiety
18 within animal models and in the development of anxiety disorders [12,121-129]. Upon the
19 development of non-peptide CRF₁ receptor antagonists that cross the blood-brain barrier,
20 there was great interest in the use of CRF₁ receptor antagonist in the treatment of anxiety
21 disorders. To date, there have been limited phase II clinical trials published regarding the
22 use of CRF₁ receptor antagonists in anxiety disorders [264]. Of those, preliminary findings
23 suggest the CRF₁ receptor antagonist-treated groups did not differ from placebo-treated
24 groups in anxiety symptomology in both social anxiety disorder and GAD [264]. However,
25 it has been suggested that efficacious concentrations have not been established for the
26 various CRF₁ receptor antagonists, and it is clear that further clinical trials are necessary.
27 One potential promising area in the treatment of anxiety disorders may actually lie in CRF₂
28 receptor antagonists. As outlined in Section 3.2, CRF₂ receptors mediate 5-HT activity in the
29 amygdala, are up-regulated in animal models of anxiety, and an antagonist of this receptor
30 reduces heightened anxiety in rats [11,12,127,128,131,132,137]. The challenge lies in
31 developing non-peptide CRF₂ receptor antagonists that cross the blood-brain barrier, so that
32 the efficacy of such ligands can be determined for anxiety disorders.

33 5. Conclusion

34 Human imaging studies in non-patient populations suggest amygdala activation in
35 response to fearful stimuli, and that the magnitude of this response is positively correlated
36 with trait anxiety. Furthermore, individuals suffering from an anxiety disorder (with the
37 possible exception of adult GAD) show exaggerated amygdala responses to fearful or
38 emotive stimuli, which again is positively correlated with the severity of symptoms.
39 Moreover, reactivity of the amygdala to fearful stimuli is reduced by anxiolytic drugs in
40 healthy subjects, and long-term pharmacotherapy or CBT reduces amygdala hyper-
41 reactivity in anxiety disorders. Animal studies corroborate an important role for the

1 amygdala in fear and anxiety, with specific subregions mediating acquisition and expression
2 of fear, fear memories and anxiety, and the monoamines within each of these regions often
3 playing a very specific role in facilitating or attenuating fear or anxiety. Both human and
4 animal studies suggest dysfunction of the amygdala might arise in part, from inadequate
5 top-down control by regions such as the medial prefrontal cortex, and in part, from altered
6 neuropeptide regulation of amygdala monoaminergic systems. Overall, the amygdala plays
7 a critical role in anxiety disorders, and understanding the function of this region in fear and
8 anxiety states and how dysfunction of the amygdala results in anxiety disorders is critical to
9 improving long-term treatment outcomes.

10 **Author details**

11 Gina L. Forster*, Andrew M. Novick, Jamie L. Scholl and Michael J. Watt
12 *Division of Basic Biomedical Sciences, Sanford School of Medicine, University of South Dakota,*
13 *Vermillion, SD, USA*

14 **Acknowledgement**

15 This work was supported by National Institutes of Health grant R01 DA019921, and
16 Department of Defense grants W81XWH-10-1-0925 and W81XWH-10-1-0578.

17 **6. References**

- 18 [1] Sylvers P, Lilienfeld SO, LaPrairie JL (2011) Differences between Trait Fear and Trait
19 Anxiety: Implications for Psychopathology. *Clin Psychol Rev.* 31, 122-137. S0272-
20 7358(10)00131-5 [pii] 10.1016/j.cpr.2010.08.004.
- 21 [2] McNaughton N, Corr PJ (2004) A Two-Dimensional Neuropsychology of Defense:
22 Fear/Anxiety and Defensive Distance. *Neuroscience & Biobehavioral Reviews.* 28, 285-
23 305. 10.1016/j.neubiorev.2004.03.005.
- 24 [3] Graham BM, Milad MR (2011) The Study of Fear Extinction: Implications for Anxiety
25 Disorders. *Am J Psychiatry.* 168, 1255-1265.
26 10.1176/appi.ajp.2011.11040557 appi.ajp.2011.11040557 [pii].
- 27 [4] Ninan PT (2001) Recent Perspectives on the Diagnosis and Treatment of Generalized
28 Anxiety Disorder. *Am J Manag Care.* 7, S367-376.
- 29 [5] Alexander JL, Dennerstein L, Kotz K, Richardson G (2007) Women, Anxiety and Mood:
30 A Review of Nomenclature, Comorbidity and Epidemiology. *Expert Rev Neurother.* 7,
31 S45-58.
- 32 [6] Katzman MA (2009) Current Considerations in the Treatment of Generalized Anxiety
33 Disorder. *CNS Drugs.* 23, 103-120.
- 34 [7] American Psychiatric Association. *Diagnostic and Statistical Manual of Mental*
35 *Disorders.* 4th ed. Washington, DC: American Psychiatric Association; 1994.

* Corresponding Author

- 1 [8] Inoue T, Kitaichi Y, Koyama T (2011) SSRIs and Conditioned Fear. *Progress in Neuro-*
2 *Psychopharmacology and Biological Psychiatry.* 35, 1810-1819.
3 10.1016/j.pnpbp.2011.09.002.
- 4 [9] Engin E, Treit D (2008) The Effects of Intra-Cerebral Drug Infusions on Animals'
5 Unconditioned Fear Reactions: A Systematic Review. *Prog Neuropsychopharmacol Biol*
6 *Psychiatry.* 32, 1399-1419. 10.1016/j.pnpbp.2008.03.020.
- 7 [10] Keck ME, Sartori SB, Welt T, Müller MB, Ohl F, Holsboer F, Landgraf R, Singewald N
8 (2005) Differences in Serotonergic Neurotransmission between Rats Displaying High or
9 Low Anxiety/Depression-Like Behaviour: Effects of Chronic Paroxetine Treatment. *J*
10 *Neurochem.* 92, 1170-1179. 10.1111/j.1471-4159.2004.02953.x.
- 11 [11] Lukkes JL, Mokin MV, Scholl JL, Forster GL (2009) Adult Rats Exposed to Early-Life
12 Social Isolation Exhibit Increased Anxiety and Conditioned Fear Behavior, and Altered
13 Hormonal Stress Responses. *Horm Behav.* 55, 248-256.
- 14 [12] Vuong SM, Oliver HA, Scholl JL, Oliver KM, Forster GL (2010) Increased Anxiety-Like
15 Behavior of Rats During Amphetamine Withdrawal Is Reversed by Crf2 Receptor
16 Antagonism. *Behavioural Brain Research.* 208, 278-281.
- 17 [13] Furmark T, Fischer H, Wik G, Larsson M, Fredrikson M (1997) The Amygdala and
18 Individual Differences in Human Fear Conditioning. *Neuroreport.* 8, 3957-3960.
- 19 [14] Whalen PJ, Bush G, McNally RJ, Wilhelm S, McInerney SC, Jenike MA, Rauch SL (1998)
20 The Emotional Counting Stroop Paradigm: A Functional Magnetic Resonance Imaging
21 Probe of the Anterior Cingulate Affective Division. *Biol Psychiatry.* 44, 1219-1228.
- 22 [15] Liddell BJ, Brown KJ, Kemp AH, Barton MJ, Das P, Peduto A, Gordon E, Williams LM
23 (2005) A Direct Brainstem–Amygdala–Cortical ‘Alarm’ System for Subliminal Signals of
24 Fear. *Neuroimage.* 24, 235-243. 10.1016/j.neuroimage.2004.08.016.
- 25 [16] Etkin A, Wager TD (2007) Functional Neuroimaging of Anxiety: A Meta-Analysis of
26 Emotional Processing in Ptsd, Social Anxiety Disorder, and Specific Phobia. *Am J*
27 *Psychiatry.* 164, 1476-1488. 10.1176/appi.ajp.2007.07030504.
- 28 [17] Hyde LW, Gorka A, Manuck SB, Hariri AR (2011) Perceived Social Support Moderates
29 the Link between Threat-Related Amygdala Reactivity and Trait Anxiety.
30 *Neuropsychologia.* 49, 651-656. 10.1016/j.neuropsychologia.2010.08.025.
- 31 [18] Carlson JM, Greenberg T, Rubin D, Mujica-Parodi LR (2011) Feeling Anxious:
32 Anticipatory Amygdalo-Insular Response Predicts the Feeling of Anxious Anticipation.
33 *Soc Cogn Affect Neurosci.* 6, 74-81. 10.1093/scan/nsq017.
- 34 [19] Stein MB, Simmons AN, Feinstein JS, Paulus MP (2007) Increased Amygdala and Insula
35 Activation During Emotion Processing in Anxiety-Prone Subjects. *Am J Psychiatry.* 164,
36 318-327. 10.1176/appi.ajp.164.2.318.
- 37 [20] Bishop S, Duncan J, Brett M, Lawrence AD (2004) Prefrontal Cortical Function and
38 Anxiety: Controlling Attention to Threat-Related Stimuli. *Nat Neurosci.* 7, 184-188.
- 39 [21] Etkin A, Klemenhagen KC, Dudman JT, Rogan MT, Hen R, Kandel ER, Hirsch J (2004)
40 Individual Differences in Trait Anxiety Predict the Response of the Basolateral
41 Amygdala to Unconsciously Processed Fearful Faces. *Neuron.* 44, 1043-1055.
42 10.1016/j.neuron.2004.12.006.

- 1 [22] Dickie EW, Armony JL (2008) Amygdala Responses to Unattended Fearful Faces:
2 Interaction between Sex and Trait Anxiety. *Psychiatry Research: Neuroimaging*. 162, 51-
3 57. 10.1016/j.psychres.2007.08.002.
- 4 [23] Ball TM, Sullivan S, Flagan T, Hitchcock CA, Simmons A, Paulus MP, Stein MB (2012)
5 Selective Effects of Social Anxiety, Anxiety Sensitivity, and Negative Affectivity on the
6 Neural Bases of Emotional Face Processing. *Neuroimage*. 59, 1879-1887.
7 10.1016/j.neuroimage.2011.08.074.
- 8 [24] Furmark T, Tillfors M, Marteinsdottir I, Fischer H, Pissiota A, Langstrom B, Fredrikson
9 M (2002) Common Changes in Cerebral Blood Flow in Patients with Social Phobia
10 Treated with Citalopram or Cognitive-Behavioral Therapy. *Arch Gen Psychiatry*. 59,
11 425-433.
- 12 [25] Shah SG, Klumpp H, Angstadt M, Nathan PJ, Phan KL (2009) Amygdala and Insula
13 Response to Emotional Images in Patients with Generalized Social Anxiety Disorder.
14 *Journal of Psychiatry & Neuroscience*. 34, 296-302.
- 15 [26] Brühl AB, Rufer M, Delsignore A, Kaffenberger T, Jäncke L, Herwig U (2011) Neural
16 Correlates of Altered General Emotion Processing in Social Anxiety Disorder. *Brain Res*.
17 1378, 72-83. 10.1016/j.brainres.2010.12.084.
- 18 [27] Blair K, Shaywitz J, Smith BW, Rhodes R, Geraci M, Jones M, McCaffrey D, Vythilingam
19 M, Finger E, Mondillo K, Jacobs M, Charney DS, Blair RJ, Drevets WC, Pine DS (2008)
20 Response to Emotional Expressions in Generalized Social Phobia and Generalized
21 Anxiety Disorder: Evidence for Separate Disorders. *Am J Psychiatry*. 165, 1193-1202.
22 10.1176/appi.ajp.2008.07071060.
- 23 [28] Rauch SL, Shin LM, Phelps EA (2006) Neurocircuitry Models of Posttraumatic Stress
24 Disorder and Extinction: Human Neuroimaging Research--Past, Present, and Future.
25 *Biol Psychiatry*. 60, 376-382.
- 26 [29] Jovanovic T, Norrholm SD (2011) Neural Mechanisms of Impaired Fear Inhibition in
27 Posttraumatic Stress Disorder. *Frontiers in Behavioral Neuroscience*. 5,
28 10.3389/fnbeh.2011.00044.
- 29 [30] Liberzon I, Abelson JL, Flagel SB, Raz J, Young EA (1999) Neuroendocrine and
30 Psychophysiologic Responses in Ptsd: A Symptom Provocation Study.
31 *Neuropsychopharmacology*. 21, 40-50.
- 32 [31] Shin LM, Orr SP, Carson MA, Rauch SL, Macklin ML, Lasko NB, Peters PM, Metzger LJ,
33 Dougherty DD, Cannistraro PA, Alpert NM, Fischman AJ, Pitman RK (2004) Regional
34 Cerebral Blood Flow in the Amygdala and Medial Prefrontal Cortex During Traumatic
35 Imagery in Male and Female Vietnam Veterans with Ptsd. *Arch Gen Psychiatry*. 61, 168-
36 176.
- 37 [32] Shin LM, Wright CI, Cannistraro PA, Wedig MM, McMullin K, Martis B, Macklin ML,
38 Lasko NB, Cavanagh SR, Krangel TS, Orr SP, Pitman RK, Whalen PJ, Rauch SL (2005) A
39 Functional Magnetic Resonance Imaging Study of Amygdala and Medial Prefrontal
40 Cortex Responses to Overtly Presented Fearful Faces in Posttraumatic Stress Disorder.
41 *Arch Gen Psychiatry*. 62, 273-281.

- 1 [33] Bryant RA, Felmingham K, Kemp A, Das P, Hughes G, Peduto A, Williams L (2008)
2 Amygdala and Ventral Anterior Cingulate Activation Predicts Treatment Response to
3 Cognitive Behaviour Therapy for Post-Traumatic Stress Disorder. *Psychological*
4 *Medicine*. 38, 555-561. doi:10.1017/S0033291707002231.
- 5 [34] Armony JL, Corbo V, Clement MH, Brunet A (2005) Amygdala Response in Patients
6 with Acute Ptsd to Masked and Unmasked Emotional Facial Expressions. *Am J*
7 *Psychiatry*. 162, 1961-1963.
- 8 [35] Dilger S, Straube T, Mentzel H-J, Fitzek C, Reichenbach JR, Hecht H, Krieschel S,
9 Gutberlet I, Miltner WHR (2003) Brain Activation to Phobia-Related Pictures in Spider
10 Phobic Humans: An Event-Related Functional Magnetic Resonance Imaging Study.
11 *Neurosci Lett*. 348, 29-32. 10.1016/s0304-3940(03)00647-5.
- 12 [36] Schienle A, Schäfer A, Walter B, Stark R, Vaitl D (2005) Brain Activation of Spider
13 Phobics Towards Disorder-Relevant, Generally Disgust- and Fear-Inducing Pictures.
14 *Neurosci Lett*. 388, 1-6. 10.1016/j.neulet.2005.06.025.
- 15 [37] van den Heuvel OA, Veltman DJ, Groenewegen HJ, Witter MP, Merkelbach J, Cath DC,
16 van Balkom AJLM, van Oppen P, van Dyck R (2005) Disorder-Specific Neuroanatomical
17 Correlates of Attentional Bias in Obsessive-Compulsive Disorder, Panic Disorder, and
18 Hypochondriasis. *Arch Gen Psychiatry*. 62, 922-933. 10.1001/archpsyc.62.8.922.
- 19 [38] Straube T, Mentzel H-J, Miltner WHR (2006) Neural Mechanisms of Automatic and
20 Direct Processing of Phobogenic Stimuli in Specific Phobia. *Biol Psychiatry*. 59, 162-170.
21 10.1016/j.biopsych.2005.06.013.
- 22 [39] Whalen PJ, Johnstone T, Somerville LH, Nitschke JB, Polis S, Alexander AL, Davidson
23 RJ, Kalin NH (2008) A Functional Magnetic Resonance Imaging Predictor of Treatment
24 Response to Venlafaxine in Generalized Anxiety Disorder. *Biol Psychiatry*. 63, 858-863.
25 10.1016/j.biopsych.2007.08.019.
- 26 [40] Palm ME, Elliott R, McKie S, Deakin JFW, Anderson IM (2011) Attenuated Responses to
27 Emotional Expressions in Women with Generalized Anxiety Disorder. *Psychological*
28 *Medicine*. 41, 1009-1018. doi:10.1017/S0033291710001455.
- 29 [41] McClure EB, Adler A, Monk CS, Cameron J, Smith S, Nelson EE, Leibenluft E, Ernst M,
30 Pine DS (2007) fMRI Predictors of Treatment Outcome in Pediatric Anxiety Disorders.
31 *Psychopharmacology (Berl)*. 191, 97-105. 10.1007/s00213-006-0542-9.
- 32 [42] Monk CS, Telzer EH, Mogg K, Bradley BP, Mai X, Louro HM, Chen G, McClure-Tone
33 EB, Ernst M, Pine DS (2008) Amygdala and Ventrolateral Prefrontal Cortex Activation
34 to Masked Angry Faces in Children and Adolescents with Generalized Anxiety
35 Disorder. *Arch Gen Psychiatry*. 65, 568-576. 10.1001/archpsyc.65.5.568.
- 36 [43] Nitschke JB, Sarinopoulos I, Oathes DJ, Johnstone T, Whalen PJ, Davidson RJ, Kalin NH
37 (2009) Anticipatory Activation in the Amygdala and Anterior Cingulate in Generalized
38 Anxiety Disorder and Prediction of Treatment Response. *Am J Psychiatry*. 166, 302-310.
39 *appi.ajp*.2008.07101682 [pii] 10.1176/appi.ajp.2008.07101682.
- 40 [44] Etkin A, Prater KE, Hoefl F, Menon V, Schatzberg AF (2010) Failure of Anterior
41 Cingulate Activation and Connectivity with the Amygdala During Implicit Regulation

- 1 of Emotional Processing in Generalized Anxiety Disorder. *Am J Psychiatry*. 167, 545-
2 554. 10.1176/appi.ajp.2009.09070931.
- 3 [45] Prevost C, McCabe JA, Jessup RK, Bossaerts P, O'Doherty JP (2011) Differentiable
4 Contributions of Human Amygdalar Subregions in the Computations Underlying
5 Reward and Avoidance Learning. *Eur J Neurosci*. 34, 134-145. 10.1111/j.1460-
6 9568.2011.07686.x.
- 7 [46] Boll S, Gamer M, Kalisch R, Buchel C (2011) Processing of Facial Expressions and Their
8 Significance for the Observer in Subregions of the Human Amygdala. *Neuroimage*. 56,
9 299-306. S1053-8119(11)00160-1 [pii] 10.1016/j.neuroimage.2011.02.021.
- 10 [47] Ball T, Rahm B, Eickhoff SB, Schulze-Bonhage A, Speck O, Mutschler I (2007) Response
11 Properties of Human Amygdala Subregions: Evidence Based on Functional MRI
12 Combined with Probabilistic Anatomical Maps. *PLoS One*. 2,
13 e307. 10.1371/journal.pone.0000307.
- 14 [48] Ball T, Derix J, Wentlandt J, Wieckhorst B, Speck O, Schulze-Bonhage A, Mutschler I
15 (2009) Anatomical Specificity of Functional Amygdala Imaging of Responses to Stimuli
16 with Positive and Negative Emotional Valence. *J Neurosci Methods*. 180, 57-70. S0165-
17 0270(09)00144-7 [pii] 10.1016/j.jneumeth.2009.02.022.
- 18 [49] Etkin A, Prater KE, Schatzberg AF, Menon V, Greicius MD (2009) Disrupted Amygdalar
19 Subregion Functional Connectivity and Evidence of a Compensatory Network in
20 Generalized Anxiety Disorder. *Arch Gen Psychiatry*. 66, 1361-1372. 66/12/1361 [pii]
21 10.1001/archgenpsychiatry.2009.104.
- 22 [50] Entis JJ, Doerga P, Barrett LF, Dickerson BC (2012) A Reliable Protocol for the Manual
23 Segmentation of the Human Amygdala and Its Subregions Using Ultra-High Resolution
24 MRI. *Neuroimage*. 60, 1226-1235. S1053-8119(12)00002-X
25 [pii] 10.1016/j.neuroimage.2011.12.073.
- 26 [51] Kim N, Kim HJ, Hwang J, Yoon SJ, Cho HB, Renshaw PF, Lyoo IK, Kim JE (2011)
27 Amygdalar Shape Analysis Method Using Surface Contour Aligning, Spherical
28 Mapping, and Probabilistic Subregional Segmentation. *Neurosci Lett*. 488, 65-69. S0304-
29 3940(10)01446-1 [pii] 10.1016/j.neulet.2010.11.005.
- 30 [52] Millan MJ (2003) The Neurobiology and Control of Anxious States. *Prog Neurobiol*. 70,
31 83-244. S030100820300087X [pii].
- 32 [53] Kalin NH, Shelton SE (2003) Nonhuman Primate Models to Study Anxiety, Emotion
33 Regulation, and Psychopathology. *Annals of the New York Academy of Sciences*. 1008,
34 189-200. 10.1196/annals.1301.021.
- 35 [54] Kalin NH, Shelton SE, Davidson RJ (2004) The Role of the Central Nucleus of the
36 Amygdala in Mediating Fear and Anxiety in the Primate. *J Neurosci*. 24, 5506-5515.
37 10.1523/JNEUROSCI.0292-04.2004 24/24/5506 [pii].
- 38 [55] Lowry CA, Johnson PL, Hay-Schmidt A, Mikkelsen J, Shekhar A (2005) Modulation of
39 Anxiety Circuits by Serotonergic Systems. *Stress*. 8, 233-246.
- 40 [56] Davis M (2006) Neural Systems Involved in Fear and Anxiety Measured with Fear-
41 Potentiated Startle. *Am Psychol*. 61, 741-756. 2006-21079-027 [pii] 10.1037/0003-
42 066X.61.8.741.

- 1 [57] Milad MR, Rauch SL, Pitman RK, Quirk GJ (2006) Fear Extinction in Rats: Implications
2 for Human Brain Imaging and Anxiety Disorders. *Biol Psychol.* 73, 61-71.
- 3 [58] Davis M, Antoniadis EA, Amaral DG, Winslow JT (2008) Acoustic Startle Reflex in
4 Rhesus Monkeys: A Review. *Rev Neurosci.* 19, 171-185.
- 5 [59] Price JL (2003) Comparative Aspects of Amygdala Connectivity. *Ann N Y Acad Sci.* 985,
6 50-58.
- 7 [60] LeDoux JE (2000) Emotion Circuits in the Brain. *Annu Rev Neurosci.* 23, 155-184.
8 10.1146/annurev.neuro.23.1.155.
- 9 [61] Canteras NS, Simerly RB, Swanson LW (1995) Organization of Projections from the
10 Medial Nucleus of the Amygdala: A Phal Study in the Rat. *J Comp Neurol.* 360, 213-245.
11 10.1002/cne.903600203.
- 12 [62] Newman SW (1999) The Medial Extended Amygdala in Male Reproductive Behavior a
13 Node in the Mammalian Social Behavior Network. *Annals of the New York Academy*
14 *of Sciences.* 877, 242-257. 10.1111/j.1749-6632.1999.tb09271.x.
- 15 [63] Herman JP, Ostrander MM, Mueller NK, Figueiredo H (2005) Limbic System
16 Mechanisms of Stress Regulation: Hypothalamo-Pituitary-Adrenocortical Axis.
17 *Progress in Neuro-Psychopharmacology and Biological Psychiatry.* 29, 1201-1213.
18 10.1016/j.pnpbp.2005.08.006.
- 19 [64] Pitkanen A, Savander V, LeDoux JE (1997) Organization of Intra-Amygdaloid
20 Circuitries in the Rat: An Emerging Framework for Understanding Functions of the
21 Amygdala. *Trends Neurosci.* 20, 517-523. S0166223697011259 [pii].
- 22 [65] Canteras NS, Swanson LW (1992) Projections of the Ventral Subiculum to the
23 Amygdala, Septum, and Hypothalamus: A Phal Anterograde Tract-Tracing Study in the
24 Rat. *J Comp Neurol.* 324, 180-194. 10.1002/cne.903240204.
- 25 [66] Canteras NS, Simerly RB, Swanson LW (1992) Connections of the Posterior Nucleus of
26 the Amygdala. *J Comp Neurol.* 324, 143-179. 10.1002/cne.903240203.
- 27 [67] Richter-Levin G, Akirav I (2000) Amygdala-Hippocampus Dynamic Interaction in
28 Relation to Memory. *Mol Neurobiol.* 22, 11-20. MN:22:1-3:011 [pii] 10.1385/MN:22:1-
29 3:011.
- 30 [68] Savander V, Miettinen R, Ledoux JE, Pitkanen A (1997) Lateral Nucleus of the Rat
31 Amygdala Is Reciprocally Connected with Basal and Accessory Basal Nuclei: A Light
32 and Electron Microscopic Study. *Neuroscience.* 77, 767-781. S0306-4522(96)00513-1 [pii].
- 33 [69] Pare D, Smith Y (1993) The Intercalated Cell Masses Project to the Central and Medial
34 Nuclei of the Amygdala in Cats. *Neuroscience.* 57, 1077-1090. 0306-4522(93)90050-P
35 [pii].
- 36 [70] Royer S, Martina M, Pare D (1999) An Inhibitory Interface Gates Impulse Traffic
37 between the Input and Output Stations of the Amygdala. *J Neurosci.* 19, 10575-10583.
- 38 [71] Pare D, Quirk GJ, Ledoux JE (2004) New Vistas on Amygdala Networks in Conditioned
39 Fear. *J Neurophysiol.* 92, 1-9. 10.1152/jn.00153.2004 92/1/1 [pii].
- 40 [72] LeDoux JE, Iwata J, Cicchetti P, Reis DJ (1988) Different Projections of the Central
41 Amygdaloid Nucleus Mediate Autonomic and Behavioral Correlates of Conditioned
42 Fear. *J Neurosci.* 8, 2517-2529.

- 1 [73] Davis M, Whalen PJ (2001) The Amygdala: Vigilance and Emotion. *Mol Psychiatry*. 6,
2 13-34.
- 3 [74] Dielenberg RA, McGregor IS (2001) Defensive Behavior in Rats Towards Predatory
4 Odors: A Review. *Neuroscience & Biobehavioral Reviews*. 25, 597-609.
5 10.1016/s0149-7634(01)00044-6.
- 6 [75] Müller M, Fendt M (2006) Temporary Inactivation of the Medial and Basolateral
7 Amygdala Differentially Affects Tmt-Induced Fear Behavior in Rats. *Behavioural Brain*
8 *Research*. 167, 57-62. 10.1016/j.bbr.2005.08.016.
- 9 [76] Rosen JB, Pagani JH, Rolla KL, Davis C (2008) Analysis of Behavioral Constraints and
10 the Neuroanatomy of Fear to the Predator Odor Trimethylthiazoline: A Model for
11 Animal Phobias. *Neurosci Biobehav Rev*. 32, 1267-1276. S0149-7634(08)00068-7 [pii]
12 10.1016/j.neubiorev.2008.05.006.
- 13 [77] Ramos C, Leite-Panissi R, Monassi R, Menescal-De-Oliveira L (1999) Role of the
14 Amygdaloid Nuclei in the Modulation of Tonic Immobility in Guinea Pigs. *Physiol*
15 *Behav*. 67, 717-724. S003193849900133X [pii].
- 16 [78] Macedo CE, Martinez RC, Brandao ML (2006) Conditioned and Unconditioned Fear
17 Organized in the Inferior Colliculus Are Differentially Sensitive to Injections of
18 Muscimol into the Basolateral Nucleus of the Amygdala. *Behav Neurosci*. 120, 625-631.
19 2006-07279-013 [pii] 10.1037/0735-7044.120.3.625.
- 20 [79] Quirk GJ, Mueller D (2008) Neural Mechanisms of Extinction Learning and Retrieval.
21 *Neuropsychopharmacology*. 33, 56-72.
- 22 [80] Zimmerman JM, Maren S (2010) Nmda Receptor Antagonism in the Basolateral but Not
23 Central Amygdala Blocks the Extinction of Pavlovian Fear Conditioning in Rats. *Eur J*
24 *Neurosci*. 31, 1664-1670. EJN7223 [pii] 10.1111/j.1460-9568.2010.07223.x.
- 25 [81] Sotres-Bayon F, Cain CK, LeDoux JE (2006) Brain Mechanisms of Fear Extinction:
26 Historical Perspectives on the Contribution of Prefrontal Cortex. *Biol Psychiatry*. 60,
27 329-336.
- 28 [82] Likhtik E, Popa D, Apergis-Schoute J, Fidacaro GA, Pare D (2008) Amygdala
29 Intercalated Neurons Are Required for Expression of Fear Extinction. *Nature*. 454, 642-
30 645. nature07167 [pii] 10.1038/nature07167.
- 31 [83] Amano T, Unal CT, Pare D (2010) Synaptic Correlates of Fear Extinction in the
32 Amygdala. *Nat Neurosci*. 13, 489-494. nn.2499 [pii] 10.1038/nn.2499.
- 33 [84] Moreira CM, Masson S, Carvalho MC, Brandao ML (2007) Exploratory Behaviour of
34 Rats in the Elevated Plus-Maze Is Differentially Sensitive to Inactivation of the
35 Basolateral and Central Amygdaloid Nuclei. *Brain Res Bull*. 71, 466-474. S0361-
36 9230(06)00305-4 [pii] 10.1016/j.brainresbull.2006.10.004.
- 37 [85] Zarrindast MR, Solati J, Oryan S, Parivar K (2008) Effect of Intra-Amygdala Injection of
38 Nicotine and GABA Receptor Agents on Anxiety-Like Behaviour in Rats.
39 *Pharmacology*. 82, 276-284. 000161129 [pii] 10.1159/000161129.
- 40 [86] Tye KM, Prakash R, Kim SY, Fenno LE, Grosenick L, Zarabi H, Thompson KR,
41 Gradinaru V, Ramakrishnan C, Deisseroth K (2011) Amygdala Circuitry Mediating

- 1 Reversible and Bidirectional Control of Anxiety. *Nature*. 471, 358-362. nature09820 [pii]
2 10.1038/nature09820.
- 3 [87] Walker DL, Toufexis DJ, Davis M (2003) Role of the Bed Nucleus of the Stria Terminalis
4 Versus the Amygdala in Fear, Stress, and Anxiety. *Eur J Pharmacol*. 463, 199-216.
5 S0014299903012822 [pii].
- 6 [88] Walker DL, Miles LA, Davis M (2009) Selective Participation of the Bed Nucleus of the
7 Stria Terminalis and Crf in Sustained Anxiety-Like Versus Phasic Fear-Like Responses.
8 *Prog Neuropsychopharmacol Biol Psychiatry*. 33, 1291-1308. S0278-5846(09)00212-7 [pii]
9 10.1016/j.pnpbp.2009.06.022.
- 10 [89] Walker DL, Davis M (2008) Role of the Extended Amygdala in Short-Duration Versus
11 Sustained Fear: A Tribute to Dr. Lennart Heimer. *Brain Struct Funct*. 213, 29-42.
12 10.1007/s00429-008-0183-3.
- 13 [90] Duncan GE, Knapp DJ, Breese GR (1996) Neuroanatomical Characterization of Fos
14 Induction in Rat Behavioral Models of Anxiety. *Brain Res*. 713, 79-91. 0006-
15 8993(95)01486-1 [pii].
- 16 [91] Herdade KC, Strauss CV, Zangrossi Junior H, Viana MB (2006) Effects of Medial
17 Amygdala Inactivation on a Panic-Related Behavior. *Behav Brain Res*. 172, 316-323.
18 S0166-4328(06)00292-0 [pii] 10.1016/j.bbr.2006.05.021.
- 19 [92] Troakes C, Ingram CD (2009) Anxiety Behaviour of the Male Rat on the Elevated Plus
20 Maze: Associated Regional Increase in C-Fos mRNA Expression and Modulation by
21 Early Maternal Separation. *Stress*. 12, 362-369. 906308344
22 [pii] 10.1080/10253890802506391.
- 23 [93] Vinkers CH, Bijlsma EY, Houtepen LC, Westphal KG, Veening JG, Groenink L, Olivier
24 B (2010) Medial Amygdala Lesions Differentially Influence Stress Responsivity and
25 Sensorimotor Gating in Rats. *Physiol Behav*. 99, 395-401. S0031-9384(09)00392-8 [pii]
26 10.1016/j.physbeh.2009.12.006.
- 27 [94] Marschner A, Kalisch R, Vervliet B, Vansteenwegen D, Buchel C (2008) Dissociable
28 Roles for the Hippocampus and the Amygdala in Human Cued Versus Context Fear
29 Conditioning. *J Neurosci*. 28, 9030-9036. 28/36/9030 [pii] 10.1523/JNEUROSCI.1651-
30 08.2008.
- 31 [95] Onishi BK, Xavier GF (2010) Contextual, but Not Auditory, Fear Conditioning Is
32 Disrupted by Neurotoxic Selective Lesion of the Basal Nucleus of Amygdala in Rats.
33 *Neurobiol Learn Mem*. 93, 165-174. S1074-7427(09)00197-X
34 [pii] 10.1016/j.nlm.2009.09.007.
- 35 [96] McDonald AJ, Mascagni F, Guo L (1996) Projections of the Medial and Lateral
36 Prefrontal Cortices to the Amygdala: A Phaseolus Vulgaris Leucoagglutinin Study in
37 the Rat. *Neuroscience*. 71, 55-75. 0306-4522(95)00417-3 [pii].
- 38 [97] Vertes RP (2004) Differential Projections of the Infralimbic and Prelimbic Cortex in the
39 Rat. *Synapse*. 51, 32-58. 10.1002/syn.10279.
- 40 [98] Corcoran KA, Quirk GJ (2007) Recalling Safety: Cooperative Functions of the
41 Ventromedial Prefrontal Cortex and the Hippocampus in Extinction. *CNS Spectr*. 12,
42 200-206.

- 1 [99] Vidal-Gonzalez I, Vidal-Gonzalez B, Rauch SL, Quirk GJ (2006) Microstimulation
2 Reveals Opposing Influences of Prelimbic and Infralimbic Cortex on the Expression of
3 Conditioned Fear. *Learn Mem.* 13, 728-733. 13/6/728 [pii] 10.1101/lm.306106.
- 4 [100] Sierra-Mercado D, Padilla-Coreano N, Quirk GJ (2011) Dissociable Roles of Prelimbic
5 and Infralimbic Cortices, Ventral Hippocampus, and Basolateral Amygdala in the
6 Expression and Extinction of Conditioned Fear. *Neuropsychopharmacology.* 36, 529-
7 538. npp2010184 [pii] 10.1038/npp.2010.184.
- 8 [101] Li G, Amano T, Pare D, Nair SS (2011) Impact of Infralimbic Inputs on Intercalated
9 Amygdala Neurons: A Biophysical Modeling Study. *Learn Mem.* 18, 226-240.
10 10.1101/lm.1938011 18/4/226 [pii].
- 11 [102] Milad MR, Pitman RK, Ellis CB, Gold AL, Shin LM, Lasko NB, Zeidan MA,
12 Handwerker K, Orr SP, Rauch SL (2009) Neurobiological Basis of Failure to Recall
13 Extinction Memory in Posttraumatic Stress Disorder. *Biol Psychiatry.* 66, 1075-1082.
14 S0006-3223(09)00896-8 [pii] 10.1016/j.biopsych.2009.06.026.
- 15 [103] Rougemont-Bucking A, Linnman C, Zeffiro TA, Zeidan MA, Lebron-Milad K,
16 Rodriguez-Romaguera J, Rauch SL, Pitman RK, Milad MR (2011) Altered Processing of
17 Contextual Information During Fear Extinction in PTSD: An fMRI Study. *CNS Neurosci*
18 *Ther.* 17, 227-236. 10.1111/j.1755-5949.2010.00152.x CNS152 [pii].
- 19 [104] Kim MJ, Whalen PJ (2009) The Structural Integrity of an Amygdala-Prefrontal
20 Pathway Predicts Trait Anxiety. *J Neurosci.* 29, 11614-11618. 29/37/11614 [pii]
21 10.1523/JNEUROSCI.2335-09.2009.
- 22 [105] Kim MJ, Gee DG, Loucks RA, Davis FC, Whalen PJ (2011) Anxiety Dissociates Dorsal
23 and Ventral Medial Prefrontal Cortex Functional Connectivity with the Amygdala at
24 Rest. *Cereb Cortex.* 21, 1667-1673. bhq237 [pii] 10.1093/cercor/bhq237.
- 25 [106] Inoue T, Koyama T, Yamashita I (1993) Effect of Conditioned Fear Stress on Serotonin
26 Metabolism in the Rat Brain. *Pharmacol Biochem Behav.* 44, 371-374. 0091-
27 3057(93)90476-A [pii].
- 28 [107] Yokoyama M, Suzuki E, Sato T, Maruta S, Watanabe S, Miyaoka H (2005) Amygdalic
29 Levels of Dopamine and Serotonin Rise Upon Exposure to Conditioned Fear Stress
30 without Elevation of Glutamate. *Neurosci Lett.* 379, 37-41. S0304-3940(04)01584-8 [pii]
31 10.1016/j.neulet.2004.12.047.
- 32 [108] Mo B, Feng N, Renner K, Forster G (2008) Restraint Stress Increases Serotonin Release
33 in the Central Nucleus of the Amygdala Via Activation of Corticotropin-Releasing
34 Factor Receptors. *Brain Res Bull.* 76, 493-498. S0361-9230(08)00039-7 [pii]
35 10.1016/j.brainresbull.2008.02.011.
- 36 [109] Zanoveli JM, Carvalho MC, Cunha JM, Brandão ML (2009) Extracellular Serotonin
37 Level in the Basolateral Nucleus of the Amygdala and Dorsal Periaqueductal Gray
38 under Unconditioned and Conditioned Fear States: An in Vivo Microdialysis Study.
39 *Brain Res.* 1294, 106-115. S0006-8993(09)01537-6 [pii] 10.1016/j.brainres.2009.07.074.
- 40 [110] Spannuth BM, Hale MW, Evans AK, Lukkes JL, Campeau S, Lowry CA (2011)
41 Investigation of a Central Nucleus of the Amygdala/Dorsal Raphe Nucleus Serotonergic

- 1 Circuit Implicated in Fear-Potentiated Startle. *Neuroscience*. 179, 104-119. S0306-
2 4522(11)00065-0 [pii] 10.1016/j.neuroscience.2011.01.042.
- 3 [111] Emoto H, Koga C, Ishii H, Yokoo H, Yoshida M, Tanaka M (1993) A CRF Antagonist
4 Attenuates Stress-Induced Increases in NA Turnover in Extended Brain Regions in Rats.
5 *Brain Res.* 627, 171-176. 0006-8993(93)90762-C [pii].
- 6 [112] Dunn AJ (1988) Stress-Related Activation of Cerebral Dopaminergic Systems. *Ann N Y*
7 *Acad Sci.* 537, 188-205.
- 8 [113] Quirarte, GL, Galvez, R, Roozendaal B, McGaugh JL. (1998) Norepinephrine Release
9 in the Amygdala in Response to Footshock and Opioid Peptidergic Drugs. *Brain Res.*
10 808, 134-140.
- 11 [114] Inglis FM, Moghaddam B (1999) Dopaminergic Innervation of the Amygdala Is Highly
12 Responsive to Stress. *J Neurochem.* 72, 1088-1094.
- 13 [115] Morilak DA, Barrera G, Echevarria DJ, Garcia AS, Hernandez A, Ma S, Petre CO (2005)
14 Role of Brain Norepinephrine in the Behavioral Response to Stress. *Prog*
15 *Neuropsychopharmacol Biol Psychiatry.* 29, 1214-1224. S0278-5846(05)00270-8 [pii]
16 10.1016/j.pnpbp.2005.08.007.
- 17 [116] Ma S, Morilak DA (2005) Norepinephrine Release in Medial Amygdala Facilitates
18 Activation of the Hypothalamic-Pituitary-Adrenal Axis in Response to Acute
19 Immobilisation Stress. *J Neuroendocrinol.* 17, 22-28. JNE1279 [pii] 10.1111/j.1365-
20 2826.2005.01279.x.
- 21 [117] Mitsushima D, Yamada K, Takase K, Funabashi T, Kimura F (2006) Sex Differences in
22 the Basolateral Amygdala: The Extracellular Levels of Serotonin and Dopamine, and
23 Their Responses to Restraint Stress in Rats. *Eur J Neurosci.* 24, 3245-3254. EJN5214 [pii]
24 10.1111/j.1460-9568.2006.05214.x.
- 25 [118] McIntyre CK, Hatfield T, McGaugh, JL (2002) Amygdala Norepinephrine Levels after
26 Training Predict Inhibitory Avoidance Retention Performance in Rats. *Eur J Neurosci.*
27 16, 1223-1226.
- 28 [119] Carrasco GA, Van de Kar LD (2003) Neuroendocrine Pharmacology of Stress. *Eur J*
29 *Pharmacol.* 463, 235-272.
- 30 [120] Feltenstein MW, See RE. (2008) The Neurocircuitry of Addiction: An Overview. *British*
31 *Journal of Pharmacology* 154, 261-274.
- 32 [121] Butler PD, Weiss JM, Stout JC, Nemeroff CB (1990) Corticotropin-Releasing Factor
33 Produces Fear-Enhancing and Behavioral Activating Effects Following Infusion into the
34 Locus Coeruleus. *J Neurosci.* 10, 176-183.
- 35 [122] Basso AM, Spina M, Rivier J, Vale W, Koob GF (1999) Corticotropin-Releasing Factor
36 Antagonist Attenuates the "Anxiogenic-Like" Effect in the Defensive Burying Paradigm
37 but Not in the Elevated Plus-Maze Following Chronic Cocaine in Rats.
38 *Psychopharmacology (Berl).* 145, 21-30.
- 39 [123] Radulovic J, Ruhmann A, Liepold T, Spiess J (1999) Modulation of Learning and
40 Anxiety by Corticotropin-Releasing Factor (CRF) and Stress: Differential Roles of CRF
41 Receptors 1 and 2. *J Neurosci.* 19, 5016-5025.

- 1 [124] Kikusui T, Takeuchi Y, Mori Y (2000) Involvement of Corticotropin-Releasing Factor in
2 the Retrieval Process of Fear-Conditioned Ultrasonic Vocalization in Rats. *Physiol*
3 *Behav.* 71, 323-328.
- 4 [125] Takahashi LK (2001) Role of CRF(1) and CRF(2) Receptors in Fear and Anxiety.
5 *Neurosci Biobehav Rev.* 25, 627-636.
- 6 [126] Bale TL (2005) Sensitivity to Stress: Dysregulation of CRF Pathways and Disease
7 Development. *Horm Behav.* 48, 1-10.
- 8 [127] Lukkes J, Vuong S, Scholl J, Oliver H, Forster G (2009) Corticotropin-Releasing Factor
9 Receptor Antagonism within the Dorsal Raphe Nucleus Reduces Social Anxiety-Like
10 Behavior after Early-Life Social Isolation. *The Journal of Neuroscience.* 29, 9955-9960.
11 10.1523/jneurosci.0854-09.2009.
- 12 [128] Bledsoe AC, Oliver KM, Scholl JL, Forster GL (2011) Anxiety States Induced by Post-
13 Weaning Social Isolation Are Mediated by CRF Receptors in the Dorsal Raphe Nucleus.
14 *Brain Res Bull.* 85, 117-122. 10.1016/j.brainresbull.2011.03.003.
- 15 [129] Risbrough VB, Stein MB (2006) Role of Corticotropin Releasing Factor in Anxiety
16 Disorders: A Translational Research Perspective. *Horm Behav.* 50, 550-561.
- 17 [130] Matsuzaki I, Takamatsu Y, Moroji T (1989) The Effects of Intracerebroventricularly
18 Injected Corticotropin-Releasing Factor (CRF) on the Central Nervous System:
19 Behavioural and Biochemical Studies. *Neuropeptides.* 13, 147-155.
- 20 [131] Amat J, Tamblin JP, Paul ED, Bland ST, Amat P, Foster AC, Watkins LR, Maier SF
21 (2004) Microinjection of Urocortin 2 into the Dorsal Raphe Nucleus Activates
22 Serotonergic Neurons and Increases Extracellular Serotonin in the Basolateral
23 Amygdala. *Neuroscience.* 129, 509-519. S0306-4522(04)00718-3
24 [pii] 10.1016/j.neuroscience.2004.07.052.
- 25 [132] Forster GL, Feng N, Watt MJ, Korzan WJ, Mouw NJ, Summers CH, Renner KJ (2006)
26 Corticotropin-Releasing Factor in the Dorsal Raphe Elicits Temporally Distinct
27 Serotonergic Responses in the Limbic System in Relation to Fear Behavior.
28 *Neuroscience.* 141, 1047-1055. S0306-4522(06)00490-8
29 [pii] 10.1016/j.neuroscience.2006.04.006.
- 30 [133] Scholl JL, Vuong SM, Forster GL (2010) Chronic Amphetamine Treatment Enhances
31 Corticotropin-Releasing Factor-Induced Serotonin Release in the Amygdala. *Eur J*
32 *Pharmacol.* 644, 80-87. S0014-2999(10)00705-3 [pii] 10.1016/j.ejphar.2010.07.008.
- 33 [134] Gray TS (1993) Amygdaloid CRF Pathways. Role in Autonomic, Neuroendocrine, and
34 Behavioral Responses to Stress. *Ann N Y Acad Sci.* 697, 53-60.
- 35 [135] Day HE, Greenwood BN, Hammack SE, Watkins LR, Fleshner M, Maier SF, Campeau S
36 (2004) Differential Expression of 5HT-1a, Alpha 1b Adrenergic, CRF-R1, and CFR-R2
37 Receptor mRNA in Serotonergic, Gamma-Aminobutyric Acidergic, and
38 Catecholaminergic Cells of the Rat Dorsal Raphe Nucleus. *J Comp Neurol.* 474, 364-378.
- 39 [136] Wang B, You ZB, Rice KC, Wise RA (2007) Stress-Induced Relapse to Cocaine Seeking:
40 Roles for the CRF(2) Receptor and CRF-Binding Protein in the Ventral Tegmental Area
41 of the Rat. *Psychopharmacology (Berl).* 193, 283-294.

- 1 [137] Pringle RB, Mouw NJ, Lukkes JL, Forster GL (2008) Amphetamine Treatment
2 Increases Corticotropin-Releasing Factor Receptors in the Dorsal Raphe Nucleus.
3 *Neurosci Res.* 62, 62-65.
- 4 [138] Reyes BA, Valentino RJ, Van Bockstaele EJ (2008) Stress-Induced Intracellular
5 Trafficking of Corticotropin-Releasing Factor Receptors in Rat Locus Coeruleus
6 Neurons. *Endocrinology.* 149, 122-130.
- 7 [139] Wise RA, Morales M (2010) A Ventral Tegmental CRF–Glutamate–Dopamine
8 Interaction in Addiction. *Brain Res.* 1314, 38-43. 10.1016/j.brainres.2009.09.101.
- 9 [140] Tagliaferro P, Morales M (2008) Synapses between Corticotropin-Releasing Factor-
10 Containing Axon Terminals and Dopaminergic Neurons in the Ventral Tegmental Area
11 Are Predominantly Glutamatergic. *J Comp Neurol.* 506, 616-626. 10.1002/cne.21576.
- 12 [141] Pernar L, Curtis AL, Vale WW, Rivier JE, Valentino RJ (2004) Selective Activation of
13 Corticotropin-Releasing Factor-2 Receptors on Neurochemically Identified Neurons in
14 the Rat Dorsal Raphe Nucleus Reveals Dual Actions. *J Neurosci.* 24, 1305-1311.
- 15 [142] Waselus M, Nazzaro C, Valentino RJ, Van Bockstaele EJ (2009) Stress-Induced
16 Redistribution of Corticotropin-Releasing Factor Receptor Subtypes in the Dorsal Raphe
17 Nucleus. *Biol Psychiatry.* 66, 76-83.
- 18 [143] Lukkes JL, Summers CH, Scholl JL, Renner KJ, Forster GL (2009) Early Life Social
19 Isolation Alters Corticotropin-Releasing Factor Responses in Adult Rats. *Neuroscience.*
20 158, 845-855.
- 21 [144] Boyson C, Miguel T, Quadros I, DeBold J, Miczek K (2011) Prevention of Social Stress-
22 Escalated Cocaine Self-Administration by CRF-R1 Antagonist in the Rat Vta.
23 *Psychopharmacology (Berl).* 1-13. 10.1007/s00213-011-2266-8.
- 24 [145] Faria R, Magalhães A, Monteiro PR, Gomes-Da-Silva J, Amélia Tavares M,
25 Summavielle T (2006) MDMA in Adolescent Male Rats: Decreased Serotonin in the
26 Amygdala and Behavioral Effects in the Elevated Plus-Maze Test. *Ann N Y Acad Sci.*
27 1074, 643-649. 1074/1/643 [pii] 10.1196/annals.1369.062.
- 28 [146] Niwa M, Matsumoto Y, Mouri A, Ozaki N, Nabeshima T (2011) Vulnerability in Early
29 Life to Changes in the Rearing Environment Plays a Crucial Role in the Aetiopathology
30 of Psychiatric Disorders. *Int J Neuropsychopharmacol.* 14, 459-477. S1461145710001239
31 [pii] 10.1017/S1461145710001239.
- 32 [147] Costall B, Jones BJ, Kelly ME, Naylor RJ, Oakley NR, Onaivi ES, Tyers MB (1989) The
33 Effects of Ondansetron (Gr38032f) in Rats and Mice Treated Subchronically with
34 Diazepam. *Pharmacol Biochem Behav.* 34, 769-778.
- 35 [148] Cruz APM, Pinheiro G, Alves SH, Ferreira G, Mendes M, Faria L, et al. (2005)
36 Behavioral effects of systemically administered MK-212 are prevented by ritanserin
37 microinfusion into the basolateral amygdala of rats exposed to the elevated plus-maze.
38 *Psychopharmacology.* 182:345–54.
- 39 [149] Zangrossi H, Graeff FG (1994) Behavioral effects of intraamygdala injections of GABA
40 and 5-HT acting drugs in the elevated plus-maze. *Braz. J Med Biol Res.* 27, 2453–6.

- 1 [150] Cornelio AM, Nunes-de-Souza RL (2007) Anxiogenic-Like Effects of MCP
2 Microinfusions into the Amygdala (but not Dorsal or Ventral Hippocampus) in Mice
3 Exposed to Elevated Plus-Maze. *Behav Brain Res.* 178, 82-89. 10.1016/j.bbr.2006.12.003.
- 4 [151] Higgins GA, Jones BJ, Oakley NR, Tyers MB (1991) Evidence that the Amygdala Is
5 Involved in the Disinhibitory Effects of 5-HT₃ Receptor Antagonists.
6 *Psychopharmacology (Berl).* 104, 545-551.
- 7 [152] Tomkins DM, Costall B, Kelly ME. (1990) Release of Suppressed Behavior of rat on the
8 Elevated X-maze by 5-HT₃ Receptor Antagonists injected into the Basolateral
9 Amygdala. *J Psychopharmacol.* 4, 203-5.
- 10 [153] Gonzalez LE, Andrews N, File SE (1996) 5-HT_{1a} and Benzodiazepine Receptors in the
11 Basolateral Amygdala Modulate Anxiety in the Social Interaction Test, but Not in the
12 Elevated Plus-Maze. *Brain Res.* 732, 145-153.
- 13 [154] Vicente MA, Zangrossi H (2011) Serotonin-2c Receptors in the Basolateral Nucleus of
14 the Amygdala Mediate the Anxiogenic Effect of Acute Imipramine and Fluoxetine
15 Administration. *Int J Neuropsychopharmacol.* 1-12. S1461145711000873
16 [pii] 10.1017/S1461145711000873.
- 17 [155] Duxon MS, Beckett SR, Baxter GS, Blackburn TP, Fone KCF. (1995). Intraamygdala
18 Injection of the 5-HT_{2B} Receptor Agonist BW-72386 Produces Anxiolysis on the
19 Elevated Plus-Maze in the Rat. *Br J Pharmacol* 116, P331-P331.
- 20 [156] Duxon MS, Kennett GA, Lightowler S, Blackburn TP, Fone KC (1997) Activation of 5-
21 HT_{2b} Receptors in the Medial Amygdala Causes Anxiolysis in the Social Interaction
22 Test in the Rat. *Neuropharmacology.* 36, 601-608.
- 23 [157] Inoue T, Li XB, Abekawa T, Kitaichi Y, Izumi T, Nakagawa S, Koyama T (2004)
24 Selective Serotonin Reuptake Inhibitor Reduces Conditioned Fear through Its Effect in
25 the Amygdala. *Eur J Pharmacol.* 497, 311-316. S0014-2999(04)00702-2 [pii]
26 10.1016/j.ejphar.2004.06.061.
- 27 [158] Leite-Panissi CR, Ferrarese AA, Terzian AL, Menescal-de-Oliveira L (2006)
28 Serotonergic Activation of the Basolateral Amygdala and Modulation of Tonic
29 Immobility in Guinea Pig. *Brain Res Bull.* 69, 356-364. S0361-9230(06)00051-7 [pii]
30 10.1016/j.brainresbull.2006.02.007.
- 31 [159] Li X, Inoue T, Abekawa T, Weng S, Nakagawa S, Izumi T, Koyama T (2006) 5-Ht_{1a}
32 Receptor Agonist Affects Fear Conditioning through Stimulation of the Postsynaptic 5-
33 Ht_{1a} Receptors in the Hippocampus and Amygdala. *Eur J Pharmacol.* 532, 74-80. S0014-
34 2999(05)01310-5 [pii] 10.1016/j.ejphar.2005.12.008.
- 35 [160] Morrison KE, Cooper MA (2012) A Role for 5-Ht_{1a} Receptors in the Basolateral
36 Amygdala in the Development of Conditioned Defeat in Syrian Hamsters. *Pharmacol*
37 *Biochem Behav.* 100, 592-600. S0091-3057(11)00312-1 [pii] 10.1016/j.pbb.2011.09.005.
- 38 [161] Centonze D, Usiello A, Gubellini P, Pisani A, Borrelli E, Bernardi G, Calabresi P (2002)
39 Dopamine D₂ Receptor-Mediated Inhibition of Dopaminergic Neurons in Mice Lacking
40 D_{2l} Receptors. *Neuropsychopharmacology.* 27, 723-726.

- 1 [162] Rouge-Pont F, Usiello A, Benoit-Marand M, Gonon F, Piazza PV, Borrelli E (2002)
2 Changes in Extracellular Dopamine Induced by Morphine and Cocaine: Crucial Control
3 by D2 Receptors. *J Neurosci.* 22, 3293-3301.
- 4 [163] Summavielle T, Magalhães A, Castro-Vale I, de Sousa L, Tavares MA (2002) Neonatal
5 Exposure to Cocaine: Altered Dopamine Levels in the Amygdala and Behavioral
6 Outcomes in the Developing Rat. *Ann N Y Acad Sci.* 965, 515-521.
- 7 [164] Izumo N, Ishibashi Y, Ohba M, Morikawa T, Manabe T (2012) Decreased Voluntary
8 Activity and Amygdala Levels of Serotonin and Dopamine in Ovariectomized Rats.
9 *Behav Brain Res.* 227, 1-6. S0166-4328(11)00764-9 [pii] 10.1016/j.bbr.2011.10.031.
- 10 [165] Rezaayof A, Hosseini SS, Zarrindast MR (2009) Effects of Morphine on Rat Behaviour in
11 the Elevated Plus Maze: The Role of Central Amygdala Dopamine Receptors. *Behav*
12 *Brain Res.* 202, 171-178. S0166-4328(09)00193-4 [pii] 10.1016/j.bbr.2009.03.030.
- 13 [166] de la Mora MP, Gallegos-Cari A, Crespo-Ramirez M, Marcellino D, Hansson AC, Fuxe
14 K (2012) Distribution of Dopamine D(2)-Like Receptors in the Rat Amygdala and Their
15 Role in the Modulation of Unconditioned Fear and Anxiety. *Neuroscience.* 201, 252-266.
16 S0306-4522(11)01238-3 [pii] 10.1016/j.neuroscience.2011.10.045.
- 17 [167] Sullivan RM, Duchesne A, Hussain D, Waldron J, Laplante F (2009) Effects of
18 Unilateral Amygdala Dopamine Depletion on Behaviour in the Elevated Plus Maze:
19 Role of Sex, Hemisphere and Retesting. *Behav Brain Res.* 205, 115-122. S0166-
20 4328(09)00440-9 [pii] 10.1016/j.bbr.2009.07.023.
- 21 [168] Bananej M, Karimi-Sori A, Zarrindast MR, Ahmadi S (2011) D1 and D2 Dopaminergic
22 Systems in the Rat Basolateral Amygdala Are Involved in Anxiogenic-Like Effects
23 Induced by Histamine. *J Psychopharmacol.* 0269881111405556
24 [pii] 10.1177/02698811111405556.
- 25 [169] de la Mora MP, Cardenas-Cachon L, Vazquez-Garcia M, Crespo-Ramirez M, Jacobsen
26 K, Hoistad M, Agnati L, Fuxe K (2005) Anxiolytic Effects of Intra-Amygdaloid Injection
27 of the D1 Antagonist SCH23390 in the Rat. *Neurosci Lett.* 377, 101-105.
28 10.1016/j.neulet.2004.11.079.
- 29 [170] Greba Q, Gifkins A, Kokkinidis L (2001) Inhibition of Amygdaloid Dopamine D2
30 Receptors Impairs Emotional Learning Measured with Fear-Potentiated Startle. *Brain*
31 *Res.* 899, 218-226. S0006-8993(01)02243-0 [pii].
- 32 [171] Guarraci FA, Frohardt RJ, Kapp BS (1999) Amygdaloid D1 Dopamine Receptor
33 Involvement in Pavlovian Fear Conditioning. *Brain Res.* 827, 28-40. S0006-
34 8993(99)01291-3 [pii].
- 35 [172] Guarraci FA, Frohardt RJ, Falls WA, Kapp BS (2000) The Effects of Intra-Amygdaloid
36 Infusions of a D2 Dopamine Receptor Antagonist on Pavlovian Fear Conditioning.
37 *Behav Neurosci.* 114, 647-651.
- 38 [173] Selden NR, Everitt BJ, Jarrard LE, Robbins TW (1991) Complementary Roles for the
39 Amygdala and Hippocampus in Aversive Conditioning to Explicit and Contextual
40 Cues. *Neuroscience.* 42, 335-350. 0306-4522(91)90379-3 [pii].
- 41 [174] Greba Q, Kokkinidis L (2000) Peripheral and Intraamygdalar Administration of the
42 Dopamine D1 Receptor Antagonist Sch 23390 Blocks Fear-Potentiated Startle but Not

- 1 Shock Reactivity or the Shock Sensitization of Acoustic Startle. *Behav Neurosci.* 114,
2 262-272.
- 3 [175] de Oliveira AR, Reimer AE, de Macedo CE, de Carvalho MC, Silva MA, Brandão ML
4 (2011) Conditioned Fear Is Modulated by D2 Receptor Pathway Connecting the Ventral
5 Tegmental Area and Basolateral Amygdala. *Neurobiol Learn Mem.* 95, 37-45. S1074-
6 7427(10)00172-3 [pii] 10.1016/j.nlm.2010.10.005.
- 7 [176] Cecchi M, Khoshbouei H, Morilak DA (2002) Modulatory Effects of Norepinephrine,
8 Acting on Alpha 1 Receptors in the Central Nucleus of the Amygdala, on Behavioral
9 and Neuroendocrine Responses to Acute Immobilization Stress. *Neuropharmacology.*
10 43, 1139-1147. S0028390802002927 [pii].
- 11 [177] Debiec J, LeDoux JE (2006) Noradrenergic Signaling in the Amygdala Contributes to
12 the Reconsolidation of Fear Memory: Treatment Implications for PTSD. *Ann N Y Acad*
13 *Sci.* 1071, 521-524. 1071/1/521 [pii] 10.1196/annals.1364.056.
- 14 [178] LaLumiere RT, Buen TV, McGaugh JL (2003) Post-Training Intra-Basolateral
15 Amygdala Infusions of Norepinephrine Enhance Consolidation of Memory for
16 Contextual Fear Conditioning. *J Neurosci.* 23, 6754-6758. 23/17/6754 [pii].
- 17 [179] Lazzaro SC, Hou M, Cunha C, LeDoux JE, Cain CK (2010) Antagonism of Lateral
18 Amygdala Alpha1-Adrenergic Receptors Facilitates Fear Conditioning and Long-Term
19 Potentiation. *Learn Mem.* 17, 489-493. 17/10/489 [pii] 10.1101/lm.1918210.
- 20 [180] Debiec J, Ledoux JE (2004) Disruption of Reconsolidation but Not Consolidation of
21 Auditory Fear Conditioning by Noradrenergic Blockade in the Amygdala.
22 *Neuroscience.* 129, 267-272. S0306-4522(04)00745-6
23 [pii] 10.1016/j.neuroscience.2004.08.018.
- 24 [181] Roozendaal B, Hui GK, Hui IR, Berlau DJ, McGaugh JL, Weinberger NM (2006)
25 Basolateral Amygdala Noradrenergic Activity Mediates Corticosterone-Induced
26 Enhancement of Auditory Fear Conditioning. *Neurobiol Learn Mem.* 86, 249-255.
27 S1074-7427(06)00032-3 [pii] 10.1016/j.nlm.2006.03.003.
- 28 [182] Cloos J-M, Ferreira V (2009) Current Use of Benzodiazepines in Anxiety Disorders.
29 *Curr Opin Psychiatry.* 22, 90-95.
- 30 [183] Ravindran LN, Stein MB (2010) The Pharmacologic Treatment of Anxiety Disorders: A
31 Review of Progress. *J Clin Psychiatry.* 71, 839-854.
- 32 [184] Brignell CM, Curran HV (2006) Drugs, Sweat and Fears: A Comparison of the Effects
33 of Diazepam and Methylphenidate on Fear Conditioning. *Psychopharmacol.* 186, 504-
34 516. 10.1007/s00213-006-0363-x.
- 35 [185] Hellewell JS, Guimaraes FS, Wang M, Deakin JF (1999) Comparison of Buspirone with
36 Diazepam and Fluvoxamine on Aversive Classical Conditioning in Humans. *J*
37 *Psychopharmacol.* 13, 122-127.
- 38 [186] Scaife JC, Hou RH, Samuels ER, Baqui F, Langley RW, Bradshaw CM, Szabadi E (2007)
39 Diazepam-Induced Disruption of Classically-Conditioned Fear-Potentiation of Late-
40 Latency Auditory Evoked Potentials Is Prevented by Flumazenil Given before, but Not
41 after, CS/US Pairing. *J Psychopharmacol.* 21, 93-101.

- 1 [187] Zangara A, Blair RJR, Curran HV (2002) A Comparison of the Effects of a Beta-
2 Adrenergic Blocker and a Benzodiazepine Upon the Recognition of Human Facial
3 Expressions. *Psychopharmacology (Berl)*. 163, 36-41.
- 4 [188] Brignell CM, Rosenthal J, Curran HV (2007) Pharmacological Manipulations of
5 Arousal and Memory for Emotional Material: Effects of a Single Dose of
6 Methylphenidate or Lorazepam. *J Psychopharmacol*. 21, 673-683.
7 10.1177/0269881107077351.
- 8 [189] Buchanan TW, Karafin MS, Adolphs R (2003) Selective Effects of Triazolam on
9 Memory for Emotional, Relative to Neutral, Stimuli: Differential Effects on Gist Versus
10 Detail. *Behav Neurosci*. 117, 517-525.
- 11 [190] Paulus MP, Feinstein JS, Castillo G, Simmons AN, Stein MB (2005) Dose-Dependent
12 Decrease of Activation in Bilateral Amygdala and Insula by Lorazepam During
13 Emotion Processing. *Arch Gen Psychiatry*. 62, 282-288.
- 14 [191] Del-Ben CM, Ferreira CA, Sanchez TA, Alves-Neto WC, Guapo VG, de Araujo DB,
15 Graeff FG (2010) Effects of Diazepam on Bold Activation During the Processing of
16 Aversive Faces. *J Psychopharmacol*.
- 17 [192] Schunck T, Mathis A, Erb G, Namer IJ, Demazieres A, Luthringer R (2010) Effects of
18 Lorazepam on Brain Activity Pattern During an Anxiety Symptom Provocation
19 Challenge. *J Psychopharmacol*. 24, 701-708.
- 20 [193] Brantigan CO, Brantigan TA, Joseph N (1979) The Effect of Beta Blockade on Stage
21 Fright. A Controlled Study. *Rocky Mt Med J*. 76, 227-233.
- 22 [194] Heiser JF, Defrancisco D (1976) The Treatment of Pathological Panic States with
23 Propranolol. *Am J Psychiatry*. 133, 1389-1394.
- 24 [195] Tyrer P (1988) Current Status of Beta-Blocking Drugs in the Treatment of Anxiety
25 Disorders. *Drugs*. 36, 773-783.
- 26 [196] Bell J (2008) Propranolol, Post-Traumatic Stress Disorder and Narrative Identity. *J Med*
27 *Ethics*. 34, e23. 10.1136/jme.2008.024752.
- 28 [197] Dębiec J, Bush DEA, LeDoux JE (2011) Noradrenergic Enhancement of
29 Reconsolidation in the Amygdala Impairs Extinction of Conditioned Fear in Rats--a
30 Possible Mechanism for the Persistence of Traumatic Memories in PTSD. *Depress*
31 *Anxiety*. 28, 186-193.
- 32 [198] Fletcher S, Creamer M, Forbes D (2010) Preventing Post Traumatic Stress Disorder:
33 Are Drugs the Answer? *The Australian and New Zealand Journal of Psychiatry*. 44,
34 1064-1071.
- 35 [199] McGhee LL, Maani CV, Garza TH, Desocio PA, Gaylord KM, Black IH (2009) The
36 Effect of Propranolol on Posttraumatic Stress Disorder in Burned Service Members.
37 *Journal of Burn Care & Research*, 30, 92-97.
- 38 [200] Stein MB, Kerridge C, Dimsdale JE, Hoyt DB (2007) Pharmacotherapy to Prevent Ptsd:
39 Results from a Randomized Controlled Proof-of-Concept Trial in Physically Injured
40 Patients. *Journal of Traumatic Stress*. 20, 923-932.
- 41 [201] Hurlemann R, Walter H, Rehme AK, Kukolja J, Santoro SC, Schmidt C, Schnell K,
42 Musshoff F, Keyzers C, Maier W, Kendrick KM, Onur OA (2010) Human Amygdala

- 1 Reactivity Is Diminished by the B-Noradrenergic Antagonist Propranolol. *Psychological*
2 *Medicine.* 40, 1839-1848.
- 3 [202] van Stegeren AH, Goekoop R, Everaerd W, Scheltens P, Barkhof F, Kuijper JPA,
4 Rombouts SARB (2005) Noradrenaline Mediates Amygdala Activation in Men and
5 Women During Encoding of Emotional Material. *Neuroimage.* 24, 898-909.
- 6 [203] Kent JM, Coplan JD, Gorman JM (1998) Clinical Utility of the Selective Serotonin
7 Reuptake Inhibitors in the Spectrum of Anxiety. *Biol Psychiatry.* 44, 812-824.
- 8 [204] Browning M, Reid C, Cowen PJ, Goodwin GM, Harmer CJ (2007) A Single Dose of
9 Citalopram Increases Fear Recognition in Healthy Subjects. *J Psychopharmacol.* 21, 684-
10 690. 10.1177/0269881106074062.
- 11 [205] Grillon C, Levenson J, Pine DS (2007) A Single Dose of the Selective Serotonin
12 Reuptake Inhibitor Citalopram Exacerbates Anxiety in Humans: A Fear-Potentiated
13 Startle Study. *Neuropsychopharmacology.* 32, 225-231.
- 14 [206] Harmer CJ, Bhagwagar Z, Perrett DI, Völlm BA, Cowen PJ, Goodwin GM (2003) Acute
15 Ssri Administration Affects the Processing of Social Cues in Healthy Volunteers.
16 *Neuropsychopharmacology.* 28, 148-152.
- 17 [207] Harmer CJ, Shelley NC, Cowen PJ, Goodwin GM (2004) Increased Positive Versus
18 Negative Affective Perception and Memory in Healthy Volunteers Following Selective
19 Serotonin and Norepinephrine Reuptake Inhibition. *Am J Psychiatry.* 161, 1256-1263.
- 20 [208] Murphy SE, Yiend J, Lester KJ, Cowen PJ, Harmer CJ (2009) Short-Term Serotonergic
21 but Not Noradrenergic Antidepressant Administration Reduces Attentional Vigilance
22 to Threat in Healthy Volunteers. *The International Journal of*
23 *Neuropsychopharmacology.* 12, 169-179.
- 24 [209] Harmer CJ, Mackay CE, Reid CB, Cowen PJ, Goodwin GM (2006) Antidepressant
25 Drug Treatment Modifies the Neural Processing of Nonconscious Threat Cues. *Biol*
26 *Psychiatry.* 59, 816-820.
- 27 [210] Anderson IM, Del-Ben CM, McKie S, Richardson P, Williams SR, Elliott R, Deakin JFW
28 (2007) Citalopram Modulation of Neuronal Responses to Aversive Face Emotions: A
29 Functional MRI Study. *Neuroreport.* 18, 1351-1355.
- 30 [211] Del-Ben CM, Deakin JFW, McKie S, Delvai NA, Williams SR, Elliott R, Dolan M,
31 Anderson IM (2005) The Effect of Citalopram Pretreatment on Neuronal Responses to
32 Neuropsychological Tasks in Normal Volunteers: An fMRI Study.
33 *Neuropsychopharmacology.* 30, 1724-1734.
- 34 [212] Cools R, Calder AJ, Lawrence AD, Clark L, Bullmore E, Robbins TW (2005) Individual
35 Differences in Threat Sensitivity Predict Serotonergic Modulation of Amygdala
36 Response to Fearful Faces. *Psychopharmacology (Berl).* 180, 670-679.
- 37 [213] Attenburrow MJ, Williams C, Odontiadis J, Reed A, Powell J, Cowen PJ, Harmer CJ
38 (2003) Acute Administration of Nutritionally Sourced Tryptophan Increases Fear
39 Recognition. *Psychopharmacology (Berl).* 169, 104-107. 10.1007/s00213-003-1479-x.
- 40 [214] Rhodes RA, Murthy NV, Dresner MA, Selvaraj S, Stavrakakis N, Babar S, Cowen PJ,
41 Grasby PM (2007) Human 5-HT Transporter Availability Predicts Amygdala Reactivity
42 in Vivo. *The Journal of Neuroscience.* 27, 9233-9237.

- 1 [215] Niehoff DL, Kuhar MJ (1983) Benzodiazepine Receptors: Localization in Rat
2 Amygdala. *The Journal of Neuroscience*. 3, 2091-2097.
- 3 [216] Young WS, 3rd, Kuhar MJ (1980) Radiohistochemical Localization of Benzodiazepine
4 Receptors in Rat Brain. *J Pharmacol Exp Ther*. 212, 337-346.
- 5 [217] Hodges H, Green S, Glenn B (1987) Evidence That the Amygdala Is Involved in
6 Benzodiazepine and Serotonergic Effects on Punished Responding but Not on
7 Discrimination. *Psychopharmacology (Berl)*. 92, 491-504.
- 8 [218] Nagy J, Zámbo K, Decsi L (1979) Anti-Anxiety Action of Diazepam after Intra-
9 Amygdaloid Application in the Rat. *Neuropharmacology*. 18, 573-576.
- 10 [219] Petersen EN, Braestrup C, Scheel-Krüger J (1985) Evidence That the Anticonflict Effect
11 of Midazolam in Amygdala Is Mediated by the Specific Benzodiazepine Receptors.
12 *Neurosci Lett*. 53, 285-288.
- 13 [220] Scheel-Krüger J, Petersen EN (1982) Anticonflict Effect of the Benzodiazepines
14 Mediated by a Gabaergic Mechanism in the Amygdala. *Eur J Pharmacol*. 82, 115-116.
- 15 [221] Shibata K, Kataoka Y, Gomita Y, Ueki S (1982) Localization of the Site of the
16 Anticonflict Action of Benzodiazepines in the Amygdaloid Nucleus of Rats. *Brain Res*.
17 234, 442-446.
- 18 [222] Green S, Vale AL (1992) Role of Amygdaloid Nuclei in the Anxiolytic Effects of
19 Benzodiazepines in Rats. *Behavioural Pharmacology*. 3, 261-264.
- 20 [223] Pesold C, Treit D (1995) The Central and Basolateral Amygdala Differentially Mediate
21 the Anxiolytic Effects of Benzodiazepines. *Brain Res*. 671, 213-221.
- 22 [224] Carvalho MC, Moreira CM, Zanoveli JM, Brandão ML (2012) Central, but Not
23 Basolateral, Amygdala Involvement in the Anxiolytic-Like Effects of Midazolam in Rats
24 in the Elevated Plus Maze. *J Psychopharmacol*. 26, 543-54.
- 25 [225] Tomaz C, Dickinson-Anson H, McGaugh JL (1992) Basolateral Amygdala Lesions
26 Block Diazepam-Induced Anterograde Amnesia in an Inhibitory Avoidance Task. *Proc*
27 *Natl Acad Sci U S A*. 89, 3615-3619.
- 28 [226] Tomaz C, Dickinson-Anson H, McGaugh JL, Souza-Silva MA, Viana MB, Graeff FG
29 (1993) Localization in the Amygdala of the Amnestic Action of Diazepam on Emotional
30 Memory. *Behav Brain Res*. 58, 99-105.
- 31 [227] Silva MA, Tomaz C (1995) Amnesia after Diazepam Infusion into Basolateral but Not
32 Central Amygdala of *Rattus Norvegicus*. *Neuropsychobiology*. 32, 31-36.
- 33 [228] Da Cunha C, Roozendaal B, Vazdarjanova A, McGaugh JL (1999) Microinfusions of
34 Flumazenil into the Basolateral but Not the Central Nucleus of the Amygdala Enhance
35 Memory Consolidation in Rats. *Neurobiology of Learning and Memory*. 72, 1-7.
- 36 [229] Coles ME, Heimberg RG (2002) Memory Biases in the Anxiety Disorders: Current
37 Status. *Clin Psychol Rev*. 22, 587-627.
- 38 [230] Coles ME, Turk CL, Heimberg RG (2007) Memory Bias for Threat in Generalized
39 Anxiety Disorder: The Potential Importance of Stimulus Relevance. *Cognitive*
40 *Behaviour Therapy*. 36, 65-73.
- 41 [231] Ferry B, McGaugh JL (2000) Role of Amygdala Norepinephrine in Mediating Stress
42 Hormone Regulation of Memory Storage. *Acta Pharmacol Sin*. 21, 481-493.

- 1 [232] Ferry B, Roozendaal B, McGaugh J (1999) Role of Norepinephrine in Mediating Stress
2 Hormone Regulation of Long-term Memory Storage: A Critical Involvement of the
3 Amygdala. *Biol Psych*. 46, 1140-1152. S0006322399001572 [pii].
- 4 [233] Berlau DJ, McGaugh JL (2006) Enhancement of Extinction Memory Consolidation: The
5 Role of the Noradrenergic and Gabaergic Systems within the Basolateral Amygdala.
6 *Neurobiology of Learning and Memory*. 86, 123-132.
- 7 [234] Lin H-C, Tseng Y-C, Mao S-C, Chen P-S, Gean P-W (2011) Gabaa Receptor Endocytosis
8 in the Basolateral Amygdala Is Critical to the Reinstatement of Fear Memory Measured
9 by Fear-Potentiated Startle. *The Journal of Neuroscience*. 31, 8851-8861.
- 10 [235] Geraciotti TD, Jr., Baker DG, Ekhtator NN, West SA, Hill KK, Bruce AB, Schmidt D,
11 Rounds-Kugler B, Yehuda R, Keck PE, Jr., Kasckow JW (2001) CSF Norepinephrine
12 Concentrations in Posttraumatic Stress Disorder. *Am J Psychiatry*. 158, 1227-1230.
- 13 [236] Strawn JR, Geraciotti TD, Jr. (2008) Noradrenergic Dysfunction and the
14 Psychopharmacology of Posttraumatic Stress Disorder. *Depress Anxiety*. 25, 260-271.
- 15 [237] Asnis GM, Kohn SR, Henderson M, Brown NL (2004) Sris Versus Non-Sris in Post-
16 Traumatic Stress Disorder: An Update with Recommendations. *Drugs*. 64, 383-404.
- 17 [238] Braun P, Greenberg D, Dasberg H, Lerer B (1990) Core Symptoms of Posttraumatic
18 Stress Disorder Unimproved by Alprazolam Treatment. *J Clin Psychiatry*. 51, 236-238.
- 19 [239] Pitman RK, Sanders KM, Zusman RM, Healy AR, Cheema F, Lasko NB, Cahill L, Orr
20 SP (2002) Pilot Study of Secondary Prevention of Posttraumatic Stress Disorder with
21 Propranolol. *Biol Psychiatry*. 51, 189-192.
- 22 [240] Vaiva G, Ducrocq F, Jezequel K, Averland B, Lestavel P, Brunet A, Marmar CR (2003)
23 Immediate Treatment with Propranolol Decreases Posttraumatic Stress Disorder Two
24 Months after Trauma. *Biol Psychiatry*. 54, 947-949.
- 25 [241] Adamec R, Muir C, Grimes M, Pearcey K (2007) Involvement of Noradrenergic and
26 Corticoid Receptors in the Consolidation of the Lasting Anxiogenic Effects of Predator
27 Stress. *Behav Brain Res*. 179, 192-207. 10.1016/j.bbr.2007.02.001.
- 28 [242] Cohen H, Kaplan Z, Koresh O, Matar MA, Geva AB, Zohar J (2011) Early Post-Stressor
29 Intervention with Propranolol Is Ineffective in Preventing Posttraumatic Stress
30 Responses in an Animal Model for PTSD. *European Neuropsychopharmacology*. 21,
31 230-240.
- 32 [243] Adamec RE, Blundell J, Collins A (2001) Neural Plasticity and Stress Induced Changes
33 in Defense in the Rat. *Neurosci Biobehav Rev*. 25, 721-744.
- 34 [244] Bodnoff SR, Suranyi-Cadotte B, Quirion R, Meaney MJ (1989) A Comparison of the
35 Effects of Diazepam Versus Several Typical and Atypical Anti-Depressant Drugs in an
36 Animal Model of Anxiety. *Psychopharmacology (Berl)*. 97, 277-279.
- 37 [245] Griebel G, Moreau JL, Jenck F, Misslin R, Martin JR (1994) Acute and Chronic
38 Treatment with 5-HT Reuptake Inhibitors Differentially Modulate Emotional Responses
39 in Anxiety Models in Rodents. *Psychopharmacology (Berl)*. 113, 463-470.
- 40 [246] Kurt M, Arik AC, Celik S (2000) The Effects of Sertraline and Fluoxetine on Anxiety in
41 the Elevated Plus-Maze Test in Mice. *Journal of Basic and Clinical Physiology and
42 Pharmacology*. 11, 173-180.

- 1 [247] Dekeyne A, Denorme B, Monneyron S, Millan MJ (2000) Citalopram Reduces Social
2 Interaction in Rats by Activation of Serotonin (5-HT)_{2c} Receptors.
3 *Neuropharmacology*. 39, 1114-1117.
- 4 [248] Griebel G (1995) 5-Hydroxytryptamine-Interacting Drugs in Animal Models of
5 Anxiety Disorders: More Than 30 Years of Research. *Pharmacology & Therapeutics*. 65,
6 319-395.
- 7 [249] Campbell BM, Merchant KM (2003) Serotonin 2c Receptors within the Basolateral
8 Amygdala Induce Acute Fear-Like Responses in an Open-Field Environment. *Brain Res*.
9 993, 1-9.
- 10 [250] Christianson JP, Ragole T, Amat J, Greenwood BN, Strong PV, Paul ED, Fleshner M,
11 Watkins LR, Maier SF (2010) 5-Hydroxytryptamine 2c Receptors in the Basolateral
12 Amygdala Are Involved in the Expression of Anxiety after Uncontrollable Traumatic
13 Stress. *Biol Psychiatry*. 67, 339-345.
- 14 [251] Vicente MA, Zangrossi H (2011) Serotonin-2c Receptors in the Basolateral Nucleus of
15 the Amygdala Mediate the Anxiogenic Effect of Acute Imipramine and Fluoxetine
16 Administration. *The International Journal of Neuropsychopharmacology*. 1-12.
- 17 [252] Burghardt NS, Bush DEA, McEwen BS, LeDoux JE (2007) Acute Selective Serotonin
18 Reuptake Inhibitors Increase Conditioned Fear Expression: Blockade with a 5-HT_{2c}
19 Receptor Antagonist. *Biol Psychiatry*. 62, 1111-1118.
- 20 [253] Bagdy G, Graf M, Anheuer ZE, Modos EA, Kantor S (2001) Anxiety-Like Effects
21 Induced by Acute Fluoxetine, Sertraline or M-CPP Treatment Are Reversed by
22 Pretreatment with the 5-HT_{2c} Receptor Antagonist Sb-242084 but not the 5-HT_{1a}
23 Receptor Antagonist Way-100635. *Int J Neuropsychopharmacol*. 4, 399-408.
24 doi:10.1017/S1461145701002632.
- 25 [254] Salchner P, Singewald N (2006) 5-HT Receptor Subtypes Involved in the Anxiogenic-
26 Like Action and Associated Fos Response of Acute Fluoxetine Treatment in Rats.
27 *Psychopharmacology (Berl)*. 185, 282-288.
- 28 [255] Hollander E, DeCaria C, Gully R, Nitescu A, Suckow RF, Gorman JM, Klein DF,
29 Liebowitz MR (1991) Effects of Chronic Fluoxetine Treatment on Behavioral and
30 Neuroendocrine Responses to Meta-Chlorophenylpiperazine in Obsessive-Compulsive
31 Disorder. *Psychiatry Res*. 36, 1-17.
- 32 [256] Zohar J, Insel TR, Zohar-Kadouch RC, Hill JL, Murphy DL (1988) Serotonergic
33 Responsivity in Obsessive-Compulsive Disorder. Effects of Chronic Clomipramine
34 Treatment. *Archives of General Psychiatry*. 45, 167-172.
- 35 [257] Kennedy AJ, Gibson EL, O'Connell MT, Curzon G (1993) Effects of Housing, Restraint
36 and Chronic Treatments with MCPP and Sertraline on Behavioural Responses to MCPP.
37 *Psychopharmacology (Berl)*. 113, 262-268.
- 38 [258] Yamauchi M, Tatebayashi T, Nagase K, Kojima M, Imanishi T (2004) Chronic
39 Treatment with Fluvoxamine Desensitizes 5-HT_{2c} Receptor-Mediated Hypolocomotion
40 in Rats. *Pharmacol Biochem Behav*. 78, 683-689.
- 41 [259] Seeman P. (2002) Atypical antipsychotics: mechanism of action. *Can J Psychiatry*, 47,
42 27-38.

- 1 [260] de Boet T. (1996) The pharmacological profile of mirtazapine. *J Clin Psychiatry*. 57,
2 19-25.
- 3 [261] Ravindran LN, Stein MB. (2010) The pharmacologic treatment of anxiety disorders: a
4 review of progress. *J Clin Psychiatry*. 71, 839-54.
- 5 [262] Rainer Q, Xia L, Guilloux JP, Gabriel C, Mocaër E, Hen R, Enhamre E, Gardier AM,
6 David DJ. (2010) Beneficial Behavioural and Neurogenic effects of Agomelatine in a
7 Model of Depression/Anxiety. *Int J Neuropsychopharmacol*. 8, 1-15.
- 8 [263] Katzman MA. (2011) Aripiprazole: a clinical review of its use for the treatment of
9 anxiety disorders and anxiety as a comorbidity in mental illness. *J Affect Disord*. 128,
10 S11-20.
- 11 [264] Kehne JH, Cain CK (2010) Therapeutic Utility of Non-Peptidic CRF1 Receptor
12 Antagonists in Anxiety, Depression, and Stress-Related Disorders: Evidence from
13 Animal Models. *Pharmacology & Therapeutics*. 128, 460-487.
14 10.1016/j.pharmthera.2010.08.011.