

Evolution of particle size in turbid discharge plumes

Paul S. Hill
Department of Oceanography
Dalhousie University
Halifax, Nova Scotia, CANADA B3H 4J1
phone: (902) 494-2266 fax: (902) 494-3877 e-mail: paul.hill@dal.ca
Award #N00014-97-1-0160
<http://www.phys.ocean.dal.ca/~phill>

LONG-TERM GOALS

The goals of this research are to develop greater understanding of the fate of fine sediment delivered to the continental shelf by the Eel River during floods and to refine methods for interpreting quantitatively the environmental record stored in grain size distributions of fine sediments on the shelf.

SCIENTIFIC OBJECTIVES

The proposed research has three objectives.

- Develop and test energy-based parameterization of flocculation settling velocity.
- Monitor patterns of post-event, cross-shelf transport of flood sediment in the bottom boundary layer (BBL).
- Refine and test models for quantitative process-based interpretation of disaggregated inorganic grain size (DIGS) in suspension and in the seabed.

APPROACH

The field effort employs a rapid response strategy, in which preset streamflow levels in the Eel River elicit a mobilization of people and gear. Within 24 hours of a “trigger,” an instrument package comprising a CTD, optical backscatter sensor, silhouette flocc camera (SFC), and 2 depth-actuated Niskin bottles is ready to be deployed from a Coast Guard helicopter. In 1999 two additional samplers were added to the instrument package: a bottom-actuated syringe for collecting samples of suspension within the bottom boundary layer and a bottom-actuated micro-grab sampler for collecting samples of surficial sediment. The instrument package is lowered on a wire into the water on a grid of 9-12 stations in the Eel River plume. Stations extend from just seaward of the surf zone to 40 meters water depth. Stations are occupied during the flood and for several days afterward, allowing examination of the post-flood distribution of suspended sediment on the shelf.

Analysis of DIGS focuses on process-based parameterization of fine-sediment grain size distributions. The parameterization divides a deposit into flocc-delivered and single-grain delivered fractions. These distinctions are more valuable from a dynamic standpoint than typical sand-silt-clay ratios because they relate directly to mode of transport. This parameterization is being

Report Documentation Page

Form Approved
OMB No. 0704-0188

Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.

1. REPORT DATE 30 SEP 1999		2. REPORT TYPE		3. DATES COVERED 00-00-1999 to 00-00-1999	
4. TITLE AND SUBTITLE Evolution of particle size in turbid discharge plumes				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Dalhousie University, Department of Oceanography, Halifax, Nova Scotia B3H 4J1, Canada,				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT	18. NUMBER OF PAGES	19a. NAME OF RESPONSIBLE PERSON
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified			

applied to closely spaced samples of x-radiographed slabs from box cores taken by Rob Wheatcroft on the Eel shelf.

All work is being conducted collaboratively with Tim Milligan of Bedford Institute of Oceanography. Milligan takes primary responsibility for equipment design, data acquisition, and particle size analysis. Hill takes primary responsibility for modeling, data analysis, and communication of results.

WORK COMPLETED

In winter 1999 particle size distributions were monitored during and after one flood event on 4 helicopter missions. SFC photos for each mission have been digitized. Water samples have been analyzed for suspended particulate mass, and disaggregated grain size distributions have been generated with a Coulter Multisizer. Also, disaggregated size distributions have been generated for a variety of cores on the Eel shelf.

A manuscript exploring the controls on effective settling velocity of sediment in the Eel plume was prepared and submitted in July, 1999 to a *Continental Shelf Research* special volume on event sedimentation on the continental shelf. This manuscript made use of sediment concentration, salinity, and floc size data gathered by Milligan, Hill, and Rocky Geyer in the 1997 and 1998 flood seasons. It also made use of velocity data gathered on moorings.

Contributions were made to a manuscript on the structure of the Eel River plume put together by Geyer and submitted to the same special volume in September, 1999. This manuscript explored the controls on plume structure and sediment fate.

A former CoOp student named Kristian Curran returned as a technician to work on the STRATAFORM project. In addition to assisting in data collection, he performed all image analysis and submitted a manuscript describing a flume study, which was carried out during his previous work term at Dalhousie. This experiment was designed to explore the evolution of disaggregated grain size distribution in a flowing suspension.

RESULTS

Bulk effective settling velocities required to explain sinking losses from the Eel River flood plume off the coast of northern California are of order 0.1 mm s^{-1} for five different helicopter-based sampling surveys conducted in January and February 1998. These effective settling velocities exceed those expected by single-grain sinking by a factor of two and implicate flocculation as an important mechanism for speeding the removal of sediment from the Eel River plume. The relative constancy of effective clearance rates despite widely varying winds, waves, and currents is consistent with photographs in the plume that show little variability in floc size with day, depth, or across- and along-shelf position. These observations of floc size contrast with those made in winter 1997 during the exceptionally large New Year's flood. During that event, increases of floc size with depth are evident. In 1997, higher sediment concentrations associated with the significantly larger discharge allowed flocs to grow substantially as they sank through the plume, whereas in 1998 low concentrations precluded significant increases in floc size with depth. These observations do not support the hypothesis that concentration controls maximal floc size; rather they indicate that the time required to form large flocs scales inversely with concentration. Using a published relationship between floc size and settling velocity for the Eel shelf suggests that approximately one half of the sediment in the plume was packaged as flocs during the 1998 floods. These data provide unique quantification of the role of flocculation in determining clearance rates

from riverine discharge plumes, thus advancing efforts to develop a process-based understanding of flood deposits on continental shelves.

In a laboratory flume, size-specific clearance rates suggest that sediment deposits primarily within small “microflocs” that sink at fractions of a millimeter per second. The dominance of microflocs over large, fast-sinking macroflocs is expected given the highly turbulent environment within the flume and its impellor-driven flow-return section. Methods developed in this study will be used by Curran, now a Masters student at Dalhousie, to constrain further the controls on sinking from the Eel River plume.

IMPACT/APPLICATION

Observations are helping to refine understanding of modes of delivery of flood sediment to the Eel River Shelf. These observations will allow quantitative constraint of aggregation time scales that has been lacking in the past.

TRANSITIONS

No transitions have occurred to date.

RELATED PROJECTS

Similar observations of in situ and disaggregated grain size were made with ONR support in a turbid discharge plume in Yakutat Bay, Alaska. Collaborators are James Syvitski (U. Colorado), Ellen Cowan (Appalachian St. U.) and Ross Powell (N. Illinois U.).

Control of floc size by turbulent kinetic energy is being investigated under the aegis of ONR’s Coastal Mixing and Optics program. Collaborators are George Voulgaris and John Trowbridge (WHOI).

Controls on floc size in tidal rivers in South Carolina are being investigated by Milligan in collaboration with Gail Kineke (Boston College) and Clark Alexander (SKIO). Hill is assisting with data analysis and interpretation.

PUBLICATIONS

- Curran, K. J., P. S. Hill, and T. G. Milligan. The role of particle aggregation in size dependent deposition of drill wastes. *Geo-Marine Letters*, submitted August, 1999.
- Geyer, W. R., P. S. Hill, T. G. Milligan, and P. Traykovski. The structure of the Eel River plume during floods. *Continental Shelf Research*, submitted September, 1999.
- Gonzalez, E. A. and P. S. Hill (1999). An investigation of Bremer et al.’s aggregation time. *Colloids and Surfaces A: Physicochemical and Engineering Aspects*, **155**: 113–116.
- Gonzalez, E. A. and P. S. Hill (1998). A method for estimating the flocculation time of monodispersed sediment suspensions. *Deep-Sea Research I*, **54**: 1931-1954.
- Hill, P. S., T. G. Milligan, and W. R. Geyer. Controls on effective settling velocity in the Eel River flood plume, *Continental Shelf Research*, submitted July, 1999.