

UNCLASSIFIED

AD NUMBER
ADB258969
NEW LIMITATION CHANGE
TO Approved for public release, distribution unlimited
FROM Distribution authorized to U.S. Gov't. agencies only; Proprietary Info.; Jun 2000. Other requests shall be referred to U.S. Army Medical Research and Materiel Command, 504 Scott St., Fort Detrick, MD 21702-5012.
AUTHORITY
USAMRMC ltr, dtd 15 May 2003

THIS PAGE IS UNCLASSIFIED

REPORT DOCUMENTATION PAGE

Form Approved
OMB No. 074-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503

1. AGENCY USE ONLY (Leave blank)	2. REPORT DATE June 2000	3. REPORT TYPE AND DATES COVERED Annual Summary (1 Jun 99 - 31 May 00)
----------------------------------	-----------------------------	---

4. TITLE AND SUBTITLE Matrix Dependent Mechanisms Involved in Tumor Promotion in Initiated Human Mammary Epithelium by Reactive Stroma	5. FUNDING NUMBERS DAMD17-99-1-9113
---	--

6. AUTHOR(S)
Bryony Wiseman, Ph.D.
Zena Werb, Ph.D.

7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES)
University of California
San Francisco, California 94143-0962

E-MAIL:
bryony@itsa.ucsf.edu

8. PERFORMING ORGANIZATION REPORT NUMBER

9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES)
U.S. Army Medical Research and Materiel Command
Fort Detrick, Maryland 21702-5012

10. SPONSORING / MONITORING AGENCY REPORT NUMBER

11. SUPPLEMENTARY NOTES

12a. DISTRIBUTION / AVAILABILITY STATEMENT
Distribution authorized to U.S. Government agencies only (proprietary information, Jun 00). Other requests for this document shall be referred to U.S. Army Medical Research and Materiel Command, 504 Scott Street, Fort Detrick, Maryland 21702-5012.

12b. DISTRIBUTION CODE

13. ABSTRACT (Maximum 200 Words)

Matrix metalloproteinases (MMPs) are thought to play important regulatory roles in normal development and disease. In addition to their established role in tumor invasion and metastasis, recent studies using mice that overexpress or lack specific MMPs or their inhibitors indicate that several MMPs can influence early tumor development. Our data indicate that stromal MMPs play an important role in tumor establishment. The coinjection of mouse embryo fibroblasts (MEFs) fosters human breast carcinoma cell growth in nude mice, whereas isogenic MEFs lacking either MMP-2 (gelatinase-A) or MMP-3 (stromelysin-1), but not MMP-9 (gelatinase-B), are deficient in their ability to foster this growth. This occurs even though the tumor cells and the murine hosts retain the ability to express these genes. This supports the emerging thesis that stromal signals and enzymes are an important factor in early tumor development. In addition, MMP-2 and MMP-3 are upregulated and activated during normal pubertal branching morphogenesis of the murine mammary gland. Transgenic mice that overexpress MMP-3 in mammary gland show increased ductal branching and mice null for MMP-3 show significantly diminished branching, whereas mice lacking MMP-2 have retarded ductal elongation. Thus MMP-2 and MMP-3 play critical roles in both tumor development and normal mammary gland morphogenesis.

14. SUBJECT TERMS
Breast Cancer, Matrix Metalloproteinases, Stromal-Epithelial Interactions

15. NUMBER OF PAGES
16

16. PRICE CODE

17. SECURITY CLASSIFICATION OF REPORT
Unclassified

18. SECURITY CLASSIFICATION OF THIS PAGE
Unclassified

19. SECURITY CLASSIFICATION OF ABSTRACT
Unclassified

20. LIMITATION OF ABSTRACT
Unlimited

AD _____

Award Number: DAMD17-99-1-9113

TITLE: Matrix Dependent Mechanisms Involved in Tumor Promotion in
Initiated Human Mammary Epithelium by Reactive Stroma

PRINCIPAL INVESTIGATOR: Bryony Wiseman, Ph.D.
Zena Werb, Ph.D.

CONTRACTING ORGANIZATION: University of California
San Francisco, California 94143-0962

REPORT DATE: June 2000

TYPE OF REPORT: Annual Summary

PREPARED FOR: U.S. Army Medical Research and Materiel Command
Fort Detrick, Maryland 21702-5012

DISTRIBUTION STATEMENT: Distribution authorized to U.S. Government
agencies only (proprietary information, Jun 00). Other requests
for this document shall be referred to U.S. Army Medical Research
and Materiel Command, 504 Scott Street, Fort Detrick, Maryland
21702-5012.

The views, opinions and/or findings contained in this report are
those of the author(s) and should not be construed as an official
Department of the Army position, policy or decision unless so
designated by other documentation.

NOTICE

USING GOVERNMENT DRAWINGS, SPECIFICATIONS, OR OTHER DATA INCLUDED IN THIS DOCUMENT FOR ANY PURPOSE OTHER THAN GOVERNMENT PROCUREMENT DOES NOT IN ANY WAY OBLIGATE THE U.S. GOVERNMENT. THE FACT THAT THE GOVERNMENT FORMULATED OR SUPPLIED THE DRAWINGS, SPECIFICATIONS, OR OTHER DATA DOES NOT LICENSE THE HOLDER OR ANY OTHER PERSON OR CORPORATION; OR CONVEY ANY RIGHTS OR PERMISSION TO MANUFACTURE, USE, OR SELL ANY PATENTED INVENTION THAT MAY RELATE TO THEM.

LIMITED RIGHTS LEGEND

Award Number: DAMD17-99-1-9113
Organization: University of California
Location of Limited Rights Data (Pages):

Those portions of the technical data contained in this report marked as limited rights data shall not, without the written permission of the above contractor, be (a) released or disclosed outside the government, (b) used by the Government for manufacture or, in the case of computer software documentation, for preparing the same or similar computer software, or (c) used by a party other than the Government, except that the Government may release or disclose technical data to persons outside the Government, or permit the use of technical data by such persons, if (i) such release, disclosure, or use is necessary for emergency repair or overhaul or (ii) is a release or disclosure of technical data (other than detailed manufacturing or process data) to, or use of such data by, a foreign government that is in the interest of the Government and is required for evaluational or informational purposes, provided in either case that such release, disclosure or use is made subject to a prohibition that the person to whom the data is released or disclosed may not further use, release or disclose such data, and the contractor or subcontractor or subcontractor asserting the restriction is notified of such release, disclosure or use. This legend, together with the indications of the portions of this data which are subject to such limitations, shall be included on any reproduction hereof which includes any part of the portions subject to such limitations.

THIS TECHNICAL REPORT HAS BEEN REVIEWED AND IS APPROVED FOR PUBLICATION.

Kath More 10/12/00

FOREWORD

Opinions, interpretations, conclusions and recommendations are those of the author and are not necessarily endorsed by the U.S. Army.

 Where copyrighted material is quoted, permission has been obtained to use such material.

 Where material from documents designated for limited distribution is quoted, permission has been obtained to use the material.

 Citations of commercial organizations and trade names in this report do not constitute an official Department of Army endorsement or approval of the products or services of these organizations.

N/A In conducting research using animals, the investigator(s) adhered to the "Guide for the Care and Use of Laboratory Animals," prepared by the Committee on Care and use of Laboratory Animals of the Institute of Laboratory Resources, national Research Council (NIH Publication No. 86-23, Revised 1985).

X For the protection of human subjects, the investigator(s) adhered to policies of applicable Federal Law 45 CFR 46.

N/A In conducting research utilizing recombinant DNA technology, the investigator(s) adhered to current guidelines promulgated by the National Institutes of Health.

N/A In the conduct of research utilizing recombinant DNA, the investigator(s) adhered to the NIH Guidelines for Research Involving Recombinant DNA Molecules.

N/A In the conduct of research involving hazardous organisms, the investigator(s) adhered to the CDC-NIH Guide for Biosafety in Microbiological and Biomedical Laboratories.

B. Wiseman

B. WISEMAN 06.27.00.

Table of Contents

Cover.....	
SF298.....	2
Foreword.....	3
Table of Contents.....	4
Introduction.....	5
Body.....	6-9
Key Research Accomplishments.....	10
Reportable Outcomes.....	11

Appended Abstracts:

B.S. Wiseman, M.D. Sternlicht, M. Sciabica & Z.Werb (2000). Roles for Matrix Metalloproteinases in Mammary Tumorigenesis and Morphogenesis. VIIIth International Congress of the Metastasis Research Society.

B.S. Wiseman, M.D. Sternlicht, M. Sciabica & Z.Werb (2000). Roles for Matrix Metalloproteinases in Mammary Morphogenesis and Tumorigenesis. Hannah Symposium 2000: The Natural History of Breast Cancer

M.D.Sternlicht, **B.S. Wiseman**, M.Simian, M.J.Bissell, Z.Werb (2000). Stromal Metalloproteinases in Morphogenesis, Remodelling and Cancer Proc. ENDO 2000 The Endocrine Society 82nd Annual Meeting, 17 (# 57)

J.L. Rinkenberger, M.D. Sternlicht, **B.S. Wiseman**, M.Sciabica, K. Holmbeck, J. Wiesen, Z. Werb (2000). Matrix Metalloproteinases and Tissue Inhibitors of Matrix Metalloproteinases in the Stroma Regulate Mammary Ductal Morphogenesis. Era of Hope, Department of Defense Breast Cancer Research Program Meeting, Abstract # Z-10.

Introduction

Breast tumors mostly arise in epithelial cells as carcinomas, however there is a growing body of evidence that reciprocal signals between carcinoma cells and adjacent stromal cells are likely to influence tumor evolution. Human conditions such as ulcerative colitis in which inflammation causes the development of a highly reactive stroma or juvenile polyposis syndrome, in which expansion of the stromal compartment results in hamartomatous polyps of the colon, can result in carcinomas. Experimentally similar phenomena have been demonstrated. Stromal fibroblasts are able to promote non-tumorigenic but immortalized epithelial cell-lines or barely tumorigenic carcinoma derived cell-lines to form significant tumors when recombinants are grown subcutaneously or as a graft under the kidney capsule of immunocompromised mice. By using these methods and also utilizing *in vitro* systems we hope to demonstrate that the stroma does have a role in breast tumor promotion and to dissect what factor/s it may contribute. Matrix metalloproteinases (MMPs), stromally expressed metzincins, are thought to play important regulatory roles in normal development and disease through their ability to alter the cellular microenvironment. Accumulating evidence from transgenic and knockout mice suggests that these stromal enzymes influence early carcinogenesis in addition to their established role in invasion and metastasis. Mice null or overexpressing specific MMPs are respectively more resistant or sensitive to tumor promoting agents. Furthermore, overexpression of MMP-3 in the mammary glands of transgenic mice predisposes these mice to mammary carcinomas in the absence of tumor promoters. In addition, MMPs are thought to have a role in normal mammary development. MMP-2 and MMP-3 are upregulated and activated during normal pubertal branching morphogenesis of the murine mammary gland. We have demonstrated that these proteases have distinct roles in ductal development of the mammary gland.

Body

Stromal Factors Can Promote to Tumor Formation

One goal for this research proposal was to develop *in vivo* and *in vitro* systems to study the promotion of mammary epithelial tumorigenesis by stromal factors. Our initial approach has been to use a mammary carcinoma cell-line, MCF7 that is reportedly poorly tumorigenic *in vivo*, but whose tumorigenicity increases if grown in the presence of fibroblasts from a variety of origins. We have shown that murine embryo fibroblasts (MEFs) are able to promote the tumorigenic potential of MCF7 cells when injected subcutaneously into immunocompromised nude mice. After 6 weeks, MCF7 cells formed average tumors of 53 mg if grown alone in the presence of matrigel, in contrast to MCF7 cells combined with MEFs in matrigel which formed tumors with an average mass of 110 mg ($p < 0.03$). This system should provide a good model with which to investigate the stromal influence on tumor formation. Initial data indicates a very significant increase in MCF7 cell proliferation (through assessment of the cell proliferation marker PCNA) in very early tumors (6 days) with the presence of MEFs compared to those without ($p < 0.0001$). In addition, 6 day old MCF7 cell recombinants in the presence of MEFs show faint signs of angiogenesis through immunohistochemical analysis of the endothelial cell marker, PECAM or CD31, whereas this marker is not evident in implants containing only MCF7 cells. Work is continuing to elucidate differences between these tumors that may provide clues to the mechanisms promoting tumor growth, including assessing different endothelial cell markers, markers of differentiation, cell death rates and *in situ* zymography at a variety of time points.

A Requirement for MMPs in the Stromal Promotion of Tumorigenesis

As mentioned above, candidate effectors of tumor promotion may be the matrix metalloproteinases (MMPs). Hence we utilized our laboratory's transgenic mice which are

Note: This report contains unpublished data

either null for specific MMPs or overexpress tissue inhibitor of MMPs-1 (TIMP-1) to generate MEFs in order to find out whether MMPs indeed have a role in the stromal promotion of tumorigenesis. We are currently in the process of subcutaneously implanting knockout MEFs (or isogenic wild type(wt) MEFs) with MCF7 cells to discover whether the presence of particular MMPs is required for tumor promotion. Our initial results look promising. MEFs lacking MMP-3 or MMP-2 were significantly retarded in their abilities to promote tumor take as compared to isogenic wtMEFs (with p values of 0.02 and 0.01 respectively). In each case the tumors that did form were not significantly different in size than those that grew with wtMEFs. Thus it appears that stromally expressed MMPs may be important in the establishment of tumors. Interestingly, we see this effect even though the murine host and the MCF7 cells both are able to express the MMP that is lacking in the MEFs. In addition, host stromal cells enter into the tumor plug (as evidenced by immunohistochemical analysis for the mesenchymal marker, vimentin), but this does not appear to be sufficient for tumor establishment. This effect on tumor establishment may be due to a lack of early filtration of endogenous fibroblasts, thus far we have only looked at 6 days. Alternatively, mouse embryo fibroblasts may have different properties from those found in adult mouse skin or may exclude endogenous fibroblasts. We are currently attempting to distinguish between host and transplanted fibroblasts by their different HLA complexes within the tumors to see whether host fibroblasts are specifically excluded from the MEF/MCF7 tumors. We also plan to use fibroblasts of different origins such as skin and mammary gland to determine whether fibroblast origin is an important factor in tumor promotion. These experiments are in their early stages and need to be repeated with different batches of MEFs. In addition, to test whether the effect of MEFs and the requirement for MMPs therein, is a common phenomenon in mammary cell tumor development, the response of other mammary cell-lines to MEFs will be tested. Currently we plan to test the very poorly tumorigenic MCF10, BT-10 and T47D human cell-lines, MBA-MD-231 cells which are highly tumorigenic and EPH4 cells which are a murine mammary cell-line which has been shown to

increase in tumorigenicity in the presence of exogenous MMP-3. Moreover, to establish that the MMP deficiency of the MEFs is responsible for the difference in tumor promotion we will attempt to rescue the tumor promoting effects of the wtMEFs by re-expressing the deficient MMP in the null MEFs.

Establishment of a Co-culture System

To allow us to investigate the mechanisms of stromal promotion of tumorigenesis in more detail we wish to re-capitulate the effects we see *in vivo* in the context of an *in vitro* co-cultivation system. For some studies we have used a direct co-culture system in which fibroblasts are labeled with a fluorescent dye to distinguish them from the epithelial cells both visually and through fluorescent activated cell sorting (FACS). We are currently looking for a marker of transformation that correlates with the ability of MEFs to promote tumorigenesis. Possibilities are proliferative capacity; anchorage independent growth; the ability to migrate and the ability to invade through a basement membrane. DNA synthesis assays are currently being optimized to allow differentiation between epithelial cells and fibroblasts undergoing S-phase using the fluorescent dye and fluorescently labeling incorporated BrdU. We have not observed any increase in the anchorage independent ability of MCF7 cells to grow in soft agar in the presence of MEFs. However fibroblasts signals may be transmitted through breakdown or release of matrix bound proteins in the basement membrane and we would see an effect using matrigel or collagen as a substrate. Those experiments are currently being performed, as are invasion and migration assays. In addition, we have shown *in vivo* that proteases are an important factor in the ability of MEFs to promote tumorigenesis. Co-incubation of epithelial cells with fibroblasts alters protease expression and activation in comparison with these cells grown alone. We have seen interesting differences in protease expression through gelatin and casein zymography when MEFs are directly co-cultured with MCF7 or T47D cells in different proportions. Thus the cells are able to signal and respond to signals from the other cell type. We are still working to identify the specific proteases involved through the use of inhibitors specific to certain classes of

proteases, western blot and by using fibroblasts null for specific MMPs. Thus both cell types are able to signal and respond to signals from the other cell type.

Distinct Roles for MMP-2 and MMP-3 in Mammary Gland Morphogenesis

In order to understand the role of MMPs in breast cancer, it is also important to understand their role in normal breast development. To this end, mammary development in mice deficient in MMPs has been examined in collaboration with others in the laboratory. We have shown that overexpression of Timp-1 in transgenic mice or treatment of mice with synthetic MMP inhibitors retards ductal elongation and branching in the pubescent mammary gland. In addition, through examination of mice null for specific MMPs, the specific requirement of a single MMP for each of these properties has been elucidated. MMP-3 has no effect on ductal penetration but its overexpression increases and its absence diminishes branching of the ducts. Conversely MMP-2 is required for ductal penetration. MMP-2 null mice up to 50 days old have shorter ducts than heterozygotes, but normal branching patterns. However by 10 weeks of age MMP-2 deficient glands have filled the fat pad indicating that these mice can overcome the lack of MMP-2 by an alternative, as yet unknown mechanism. Interestingly, MMP-9 null mice have normal ductal development indicating that the requirements for MMP-2 and MMP-3 are specific. Thus we have separated two distinct processes in mammary gland ductal development and described a requirement for distinct proteases in these processes.

Key Research Accomplishments.

- Development of *in vivo* system for studying stromal promotion of tumor growth
- Demonstrated fostering of tumor take and growth by mouse embryo fibroblasts (MEFs) with subcutaneously injected MCF-7 cell recombinants.
- Observed a significant increase in cell proliferation in early MCF-7 recombinants in the presence of MEFs over recombinants with MCF-7 cells alone.
- Observed more early angiogenesis in MCF-7 recombinants in the presence of MEFs over recombinants with MCF-7 cells alone.
- Identified a requirement for MMP-3 and MMP-2, but not MMP-9 in fibroblasts, and thus the promotion of tumor take by the stroma.
- Development of *in vitro* co-cultivation systems
- Identified differences in secreted gelatinolytic and caseinolytic activities when breast carcinoma epithelial cells are co-cultured with fibroblasts
- Identified requirement for gelatinase A in ductal invasion of the pubescent murine mammary gland

Reportable Outcomes

Published Abstracts (For platform presentations the presenting author is underlined)

B.S. Wiseman, M.D. Sternlicht, M. Sciabica & Z.Werb. Roles for Matrix Metalloproteinases in Mammary Tumorigenesis and Morphogenesis. VIIIth International Congress of the Metastasis Research Society. Abstract accepted. September 24-27 2000.

B.S. Wiseman, M.D. Sternlicht, M. Sciabica & Z.Werb Roles for Matrix Metalloproteinases in Mammary Morphogenesis and Tumorigenesis.. Hannah Symposium 2000:The Natural History of Breast Cancer. Abstract accepted. September 6-8 2000.

M.D.Sternlicht, **B.S. Wiseman**, M.Simian, M.J.Bissell, Z.Werb. Stromal Metalloproteinases in Morphogenesis, Remodelling and Cancer. Proc. ENDO 2000 The Endocrine Society 82nd Annual Meeting, 17 (# 57) June 23, 2000.

J.L. Rinkenberger, M.D. Sternlicht, **B.S.Wiseman**, M.Sciabica, K. Holmbeck, J. Wiesen, Z. Werb. Matrix Metalloproteinases and Tissue Inhibitors of Matrix Metalloproteinases in the Stroma Regulate Mammary Ductal Morphogenesis.. Era of Hope, Department of Defense Breast Cancer Research Program Meeting, Abstract # Z-10. June 8, 2000.

Training

- UCSF Committee on Animal Research/Animal Care Facility Training and Certification Program
- Animal Care Facility Training Workshop on animal handling and aseptic techniques
- Laboratory for Cytometric Analysis Flow Cytometry Course

Roles for Stromal Matrix Metalloproteinases in Mammary Tumorigenesis and Morphogenesis

B.S. Wiseman, M.D. Sternlicht, M. Sciabica & Z. Werb

Matrix metalloproteinases (MMPs) are thought to play important regulatory roles in normal development and disease. In addition to their established role in tumor invasion and metastasis, recent studies using mice that overexpress or lack specific MMPs or their inhibitors indicate that several MMPs can influence early tumor development. Although most tumors are epithelial in origin, the majority of MMPs are expressed by the stroma and alter the cellular microenvironment. Our data indicate that stromal MMPs play an important role in tumor establishment. The coinjection of mouse embryo fibroblasts (MEFs) fosters human breast carcinoma cell growth in nude mice, whereas isogenic MEFs lacking either MMP-2 (gelatinase-A) or MMP-3 (stromelysin-1), but not MMP-9 (gelatinase-B), are deficient in their ability to foster this growth. This occurs even though the tumor cells and the murine hosts retain the ability to express these genes. This supports the emerging thesis that stromal signals and enzymes are an important factor in early tumor development. In addition, MMP-2 and MMP-3 are upregulated and activated during normal pubertal branching morphogenesis of the murine mammary gland. Transgenic mice that overexpress MMP-3 in mammary gland show increased ductal branching and mice null for MMP-3 show significantly diminished branching, whereas mice lacking MMP-2 have retarded ductal elongation. Thus MMP-2 and MMP-3 play critical roles in both tumor development and normal mammary gland morphogenesis.

Roles for Matrix Metalloproteinases in Mammary Morphogenesis and Tumorigenesis

B.S. Wiseman, M.D. Sternlicht, M. Sciabica & Z. Werb

Dept of Anatomy, Box 0452, University of California, San Francisco, CA 94143, USA.

Matrix metalloproteinases (MMPs) are thought to play important regulatory roles in normal development and disease. MMP-2 (gelatinase-A) and MMP-3 (stromelysin-1) are upregulated and activated during normal pubertal branching morphogenesis of the murine mammary gland. Transgenic mice that overexpress MMP-3 in mammary gland show increased ductal branching and mice null for MMP-3 show significantly diminished branching, whereas mice lacking MMP-2 have retarded ductal elongation. MMP-9 (gelatinase-B) is not upregulated during mammary development, nor does its absence alter ductal development. Thus proper development of the mammary gland requires coordinated action of at least two, and possibly more, members of the MMP family. In addition to their important roles in development, MMPs also have an established role in tumor invasion and metastasis. Moreover, recent studies using mice that overexpress or lack specific MMPs or their inhibitors indicate that several MMPs can also influence early tumor development. Most tumors are epithelial in origin and yet the majority of MMPs are expressed by the stroma and alter the cellular microenvironment. Thus reciprocal signals between carcinoma cells and adjacent stromal cells are likely to influence tumor evolution. Our data indicate that stromal MMPs play an important role in tumor establishment. The coinjection of mouse embryo fibroblasts (MEFs) fosters human breast carcinoma cell growth in nude mice, whereas isogenic MEFs lacking either MMP-2 or MMP-3, but not MMP-9, are deficient in their ability to foster this growth. This occurs even though the tumor cells and the murine hosts retain the ability to express these genes. This supports the emerging thesis that stromal signals and enzymes are an important factor in early tumor development. Thus MMP-2 and MMP-3 play critical roles in both tumor development and normal mammary gland morphogenesis.

Please select Print from the file menu to print your Abstract.

ENDO 2000 The Endocrine Society 82nd Annual Meeting

Invited Speaker Submission

Filename: 9938

Corresponding Author: Mark D Sternlicht Ph.D.

Department/Institution: Dept. of Anatomy, Univ of California-San Francisco

Address: HSW 1320/1321 University of California, 513 Parnassus Avenue, San Francisco, CA, 94143-0452

Phone: 415 476 4758 **Fax:** 415 476 4565 **E-Mail:** sternli@itsa.ucsf.edu

Keywords: Mammary Development, Matrix Metalloproteinases, Mammary Cancer

STROMAL METALLOPROTEINASES IN MORPHOGENESIS, REMODELING AND CANCER.

Mark D. Sternlicht¹, Bryony S. Wiseman¹, Marina Simian², Mina J. Bissell² and Zena Werb¹. ¹Dept. of Anatomy, University of California-San Francisco, San Francisco, CA; and ²Life Sciences Div., Lawrence Berkeley National Laboratory, Berkeley, CA.

At least 19 matrix metalloproteinases (MMPs) influence various aspects of mammalian development, health and disease. In the mammary gland, MMP2/gelatinase A (GelA) and MMP3/stromelysin-1 (Str1) are upregulated and activated during pubertal branching morphogenesis and postlactational involution. MMPs are required for branching of cultured mammary organoids and recombinant Str1 suffices to promote branching in the absence of growth factors. Analysis of transgenic and knockout mice reveals that Str1 indeed regulates mammary branching in vivo, whereas GelA regulates ductal penetration and elongation within the mammary fat pad. MMP9/gelatinase B is not normally upregulated at puberty and, as expected, its absence in null animals has no effect on mammary development. Transgenic data also suggest that Str1 plays a key role in the apoptosis and remodeling seen during involution. Stromal MMPs are also consistently upregulated in malignant disease. Indeed, many MMPs were cloned as cancer-specific genes and are key agonists in angiogenesis, invasion and metastasis. In addition, recent studies in MMP-overexpressing and -deficient mice indicate that several MMPs can influence early cancer development, but most such studies have required carcinogens or pre-existing mutations to initiate tumorigenesis. We, however, have observed spontaneous premalignant and malignant lesions in the mammary glands of mice with a mammary-targeted Str1 transgene. These changes were absent in nontransgenic littermates and were quenched by the co-expression of a tissue inhibitor of metalloproteinases-1 transgene. Moreover, mouse embryonic fibroblasts foster human breast cancer cell growth in vivo, yet Str1-null fibroblasts do not; and an autoactivating Str1 can convert functionally normal mammary epithelial cells into invasive and tumorigenic mesenchymal-like cells, and can modulate the expression of genes that control cancer development. Thus by altering the cellular microenvironment, Str1 can act as a natural tumor promoter and can enhance cancer susceptibility.

Disclosure: There is no disclosure information to present

Presenting Author Signature: (Mark D. Sternlicht)



Date:

2/28/00

Return To Thank You

Note: Use the buttons on the left to move between screens or the "Reload" button if you encounter an error message.

Questions about the Online Abstract Submission process?
Contact [Marathon Multimedia](mailto:support@marathonmultimedia.com) at support@marathonmultimedia.com

Questions about the The Endocrine Society 82nd Annual Meeting?
Contact [The Endocrine Society](mailto:endostaff@endo-society.org) at endostaff@endo-society.org

Online Submission[®] is a product of

MATRIX METALLOPROTEINASES AND TISSUE INHIBITORS OF MATRIX METALLOPROTEINASES IN THE STROMA REGULATE MAMMARY DUCTAL MORPHOGENESIS.

J. L. Rinkenberger[^], M. D. Sternlicht[^], B. S. Wiseman[^], M. Sciabica[^], K. Holmbeck^{*}, J. Wiesen[^], H. Birkedal-Hansen^{*}, Z. Werb[^].

[^]University of California, Department of Anatomy, HSW 1301, 513 Parnassus Avenue, San Francisco, CA 94143-0452. ^{*}MMP Unit, National Institute of Dental and Craniofacial Research, Bethesda, MD 20892.

rinken@itsa.ucsf.edu.

Mammary gland ductal morphogenesis begins in late fetal development, continues slowly during the early post-natal period and accelerates in response to ovarian hormones at the onset of puberty. It is well known that ductal epithelial cell morphogenesis is regulated by stromal factors in the surrounding fat pad. Here we show that stromal matrix metalloproteinases (MMPs) and their natural inhibitors, the tissue inhibitors of matrix metalloproteinases (TIMPs), influence ductal elongation and branching by regulating extracellular matrix remodeling. Transgenic mice that constitutively over-express TIMP-1 show diminished ductal elongation and branching, and branching of cultured mammary organoids is blocked by synthetic MMP inhibitors in vitro. However, these models do not indicate which MMPs are responsible for ductal elongation and branching. MMP-3/stromelysin-1 and MMP-2/gelatinase A are both upregulated at the mRNA and protein levels in the stroma surrounding the developing ducts making them particularly good candidates. In order to determine which MMPs are involved in ductal elongation and branching we have examined mammary gland development in specific MMP over-expressing and null mouse models. MMP-3 has no effect on ductal penetration but its over-expression increases, and its absence significantly decreases, both dichotomous and lateral branching. On the other hand, the MMP-2 null mice exhibit reduced ductal penetration of the fat pad without changes in branching. MMP-9/gelatinase B is not normally upregulated at puberty and, not surprisingly, its absence in MMP-9 null mice has no effect on ductal development. The transmembrane MMP, MMP-14, regulates MMP-2 activation at the cell surface. Consequently, deletion of the MMP-14 protease from the mammary gland may also inhibit ductal development through regulation of MMP-2 activation. These animals exhibit retarded adipocyte differentiation within the mammary fat pad and we are currently evaluating whether epithelial alterations exist as well. Although we have shown that MMP-2 and MMP-3 have distinct roles during ductal development, the absence of either MMP-2 or MMP-3 fails to ablate ductal morphogenesis entirely indicating that other MMP family members are likely to have a role in these processes.

The U.S. Army Medical Research and Materiel Command under DAMD17-97-7324 supported this work.



DEPARTMENT OF THE ARMY
US ARMY MEDICAL RESEARCH AND MATERIEL COMMAND
504 SCOTT STREET
FORT DETRICK, MD 21702-5012

REPLY TO
ATTENTION OF

MCMR-RMI-S (70-1y)

15 May 03

MEMORANDUM FOR Administrator, Defense Technical Information Center (DTIC-OCA), 8725 John J. Kingman Road, Fort Belvoir, VA 22060-6218

SUBJECT: Request Change in Distribution Statement

1. The U.S. Army Medical Research and Materiel Command has reexamined the need for the limitation assigned to technical reports written for this Command. Request the limited distribution statement for the enclosed accession numbers be changed to "Approved for public release; distribution unlimited." These reports should be released to the National Technical Information Service.
2. Point of contact for this request is Ms. Kristin Morrow at DSN 343-7327 or by e-mail at Kristin.Morrow@det.amedd.army.mil.

FOR THE COMMANDER:

Encl


PHYLIS M. RINEHART
Deputy Chief of Staff for
Information Management

ADB266022	ADB265793
ADB260153	ADB281613
ADB272842	ADB284934
ADB283918	ADB263442
ADB282576	ADB284977
ADB282300	ADB263437
ADB285053	ADB265310
ADB262444	ADB281573
ADB282296	ADB250216
ADB258969	ADB258699
ADB269117	ADB274387
ADB283887	ADB285530
ADB263560	
ADB262487	
ADB277417	
ADB285857	
ADB270847	
ADB283780	
ADB262079	
ADB279651	
ADB253401	
ADB264625	
ADB279639	
ADB263763	
ADB283958	
ADB262379	
ADB283894	
ADB283063	
ADB261795	
ADB263454	
ADB281633	
ADB283877	
ADB284034	
ADB283924	
ADB284320	
ADB284135	
ADB259954	
ADB258194	
ADB266157	
ADB279641	
ADB244802	
ADB257340	
ADB244688	
ADB283789	
ADB258856	
ADB270749	
ADB258933	