

UNCLASSIFIED

AD NUMBER
ADB275851
NEW LIMITATION CHANGE
TO Approved for public release, distribution unlimited
FROM Distribution authorized to U.S. Gov't. agencies only; Proprietary Info.; Oct 2001. Other requests shall be referred to U.S. Army Medical Research and Materiel Command, 504 Scott St., Ft. Detrick, MD 21702-5012.
AUTHORITY
USAMRMC ltr, 28 Aug 2002

THIS PAGE IS UNCLASSIFIED

AD _____

Award Number: DAMD17-00-1-0684

TITLE: Role of Sulfation Pharmacogenetics in Breast Cancer Treatment with 2-Methoxyestradiol

PRINCIPAL INVESTIGATOR: Araba A. Adjei, Ph.D.

CONTRACTING ORGANIZATION: Mayo Clinic Foundation
Rochester, Minnesota 55905

REPORT DATE: October 2001

TYPE OF REPORT: Final

PREPARED FOR: U.S. Army Medical Research and Materiel Command
Fort Detrick, Maryland 21702-5012

DISTRIBUTION STATEMENT: Distribution authorized to U.S. Government agencies only (proprietary information, Oct 01). Other requests for this document shall be referred to U.S. Army Medical Research and Materiel Command, 504 Scott Street, Fort Detrick, Maryland 21702-5012.

The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision unless so designated by other documentation.

20020225 061

NOTICE

USING GOVERNMENT DRAWINGS, SPECIFICATIONS, OR OTHER DATA INCLUDED IN THIS DOCUMENT FOR ANY PURPOSE OTHER THAN GOVERNMENT PROCUREMENT DOES NOT IN ANY WAY OBLIGATE THE U.S. GOVERNMENT. THE FACT THAT THE GOVERNMENT FORMULATED OR SUPPLIED THE DRAWINGS, SPECIFICATIONS, OR OTHER DATA DOES NOT LICENSE THE HOLDER OR ANY OTHER PERSON OR CORPORATION; OR CONVEY ANY RIGHTS OR PERMISSION TO MANUFACTURE, USE, OR SELL ANY PATENTED INVENTION THAT MAY RELATE TO THEM.

LIMITED RIGHTS LEGEND

Award Number: DAMD17-00-1-0684
Organization: Mayo Clinic Rochester

Those portions of the technical data contained in this report marked as limited rights data shall not, without the written permission of the above contractor, be (a) released or disclosed outside the government, (b) used by the Government for manufacture or, in the case of computer software documentation, for preparing the same or similar computer software, or (c) used by a party other than the Government, except that the Government may release or disclose technical data to persons outside the Government, or permit the use of technical data by such persons, if (i) such release, disclosure, or use is necessary for emergency repair or overhaul or (ii) is a release or disclosure of technical data (other than detailed manufacturing or process data) to, or use of such data by, a foreign government that is in the interest of the Government and is required for evaluational or informational purposes, provided in either case that such release, disclosure or use is made subject to a prohibition that the person to whom the data is released or disclosed may not further use, release or disclose such data, and the contractor or subcontractor or subcontractor asserting the restriction is notified of such release, disclosure or use. This legend, together with the indications of the portions of this data which are subject to such limitations, shall be included on any reproduction hereof which includes any part of the portions subject to such limitations.

THIS TECHNICAL REPORT HAS BEEN REVIEWED AND IS APPROVED FOR PUBLICATION.

Hath More 215702

REPORT DOCUMENTATION PAGE

Form Approved
OMB No. 074-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503

1. AGENCY USE ONLY (Leave blank)		2. REPORT DATE October 2001	3. REPORT TYPE AND DATES COVERED Final (1 Oct 00 - 30 Sep 01)	
4. TITLE AND SUBTITLE Role of Sulfation Pharmacogenetics in Breast Cancer Treatment with 2-Methoxyestradiol			5. FUNDING NUMBERS DAMD17-00-1-0684	
6. AUTHOR(S) Araba A. Adjei, Ph.D.				
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Mayo Clinic Foundation Rochester, Minnesota 55905 E-mail: adjei.araba@mayo.edu			8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) U.S. Army Medical Research and Materiel Command Fort Detrick, Maryland 21702-5012			10. SPONSORING / MONITORING AGENCY REPORT NUMBER	
11. SUPPLEMENTARY NOTES				
12a. DISTRIBUTION / AVAILABILITY STATEMENT Distribution authorized to U.S. Government agencies only (proprietary information, Oct 01). Other requests for this document shall be referred to U.S. Army Medical Research and Materiel Command, 504 Scott Street, Fort Detrick, Maryland 21702-5012.			12b. DISTRIBUTION CODE	
13. ABSTRACT (Maximum 200 Words) <p>2-ME₂ is an endogenous estrogen metabolite that inhibits the proliferation of breast and other human cancer cell lines. 2-ME₂ also has potent anti-angiogenic and anti-tubulin properties, and it may inhibit estrogen-induced carcinogenesis in the mammary gland. We set out to test the hypothesis that 2-ME₂ might be a substrate for sulfate conjugation and, therefore, that individual variations in the sulfation of 2-ME₂ might contribute to individual differences in its metabolism, pharmacokinetics and therapeutic efficacy. As a first step, we tested 2-ME₂ as a substrate for 7 human sulfotransferase (SULT) isoforms -- as well as all of the common allozymes for SULT1A1 and 1A2. Substrate kinetic studies were conducted in two stages -- starting with concentrations over 5 orders of magnitude, followed by determination of K_m values over a narrow concentration range. 2-ME₂ was a sulfate acceptor substrate for SULT1A1*1, *2, *3; 1A2*1, *2, *3; 1A3; 1E1; 2A1; 2B1a and 2B1b, with apparent K_m values of 2.5, 5.2, 1.6; 4.2, 111, 5.3; 91; 0.067; 8.3; 4.1 and 4.1 μM, respectively. These results suggest that individual pharmacogenetic variation in sulfate conjugation might contribute to individual differences in 2-ME₂ pharmacokinetics and therapeutic effect.</p>				
14. SUBJECT TERMS Anti-angiogenic, anti-tubulin, estrogen-induced carcinogenesis, Pharmacogenetic variation, Sulfation, Sulfotransferases, 2-Methoxyestradiol			15. NUMBER OF PAGES 12	
			16. PRICE CODE	
17. SECURITY CLASSIFICATION OF REPORT Unclassified	18. SECURITY CLASSIFICATION OF THIS PAGE Unclassified	19. SECURITY CLASSIFICATION OF ABSTRACT Unclassified	20. LIMITATION OF ABSTRACT Unlimited	

TABLE OF CONTENTS

FRONT COVER..... 1

STANDARD FORM (SF) 298, REPORT DOCUMENTATION PAGE 2

TABLE OF CONTENTS..... 3

INTRODUCTION..... 4

BODY 4

KEY RESEARCH ACCOMPLISHMENTS 5

REPORTABLE OUTCOMES 5

CONCLUSIONS..... 5

REFERENCES 6

APPENDICES 8

INTRODUCTION

The risk of estrogen-induced breast cancer is affected by the balance between activities of several enzymes involved in the biotransformation of estrogen and its metabolites. 2-methoxyestradiol (2-ME), an endogenous estrogen metabolite that inhibits the proliferation of many human cell lines in vitro and in vivo, is being developed for clinical testing as an anticancer agent. 2-ME has unique biological properties not shared with the other estrogen metabolites. It has potent anti-angiogenic activity in vitro and in vivo as well as antitubulin properties. While the exact mechanism of antiproliferative activity of 2-ME is unknown, emerging evidence suggests that 2-ME may inhibit estrogen-induced carcinogenesis in target tissues such as the mammary gland. Conjugation of 2-ME, catalyzed by sulfotransferase (SULT) enzymes may alter its anti-tumorigenic effects in the treatment of breast cancer. Many human SULTs are genetically polymorphic therefore, individual variations in SULT enzyme activity imply variations in the inactivation of 2-ME, with subsequent variations in antitumorigenic activity. We therefore hypothesized that SULT enzyme activities may play an active and important role in the response of breast cancer patients to treatment with 2-ME, through changes in catabolism. Hence, sulfation, may play a role in the therapeutic response of individuals to the treatment of breast cancer with 2-ME .

BODY

The main task was to characterize the SULT enzymes involved in the catabolism of 2-ME₂. This task was completed.

To initiate this work, cDNAs for human *SULT1A1*1*, *SULT1A1*2*, *SULT1A1*3*; *SULT1A2*1*, *SULT1A2*2*, *SULT1A2*3*; *SULT1A3*; *SULT1B1*; *SULT1C1*; *SULT1E1*, *SULT2A1*, *SULT2B1a*; *SULT2B1b* and *SULT4A1* were each ligated into either the eukaryotic expression vector pCR3.1 or p91023B. Sequences of the cDNA inserts were confirmed by DNA sequencing prior to transfection of COS-1 cells using the DEAE-dextran or the Transfast method. Cytosol from transfected COS-1 cells served as a source of recombinant protein.

The resulting recombinant SULT proteins were used for the biochemical characterization of 2-ME₂. Substrate kinetic studies for the sulfation of 2-ME₂ was performed using the modified assay method of Foldes and Meek for sulfotransferases. Because of profound substrate inhibition displayed by SULTs, two sets of experiments were performed for each enzyme. The K_m and V_{max} values were then calculated using the method by Cleland. See appendix.

The long term goal of this project would be to identify the existence of functionally significant polymorphism(s) in the SULT isoform(s) responsible for catabolism of 2-ME₂ in the target tissue. Genotyping of patients prior to treatment with 2-ME₂ would be expected to predict response and/or toxicity, and allow for the tailoring of drug doses to individual patients.

In the appendix are :

Figure 1: Scheme showing the sulfate conjugation of 2-ME₂.

Figure 2: Substrate curves and double inverse plot for 2-ME₂ catalyzed by SULT1E1.

Table 1 : Substrate Kinetic results obtained from this study.

KEY RESEARCH ACCOMPLISHMENTS

The proposed task/work indicated in the concept was completed.

A poster with this work was presented at the 102nd Annual meeting of the American Society for Clinical Pharmacology and Therapeutics (ASCPT), in March 2001, at Orlando, FL.

REPORTABLE OUTCOMES

Adjei, A.A., Wood, T.C. and Weinshilboum, R.M. (2001). 2-Methoxyestradiol (2-ME₂) Sulfation: Possible metabolic pathway. *Clinical Pharmacology and Therapeutics* 69:75. The abstract is attached in the appendix.

CONCLUSIONS

- 2-ME₂ is an endogenous estrogen metabolite formed *in vivo* by the O-methylation of 2-hydroxyestradiol, a reaction catalyzed by COMT.
- 2-ME₂ is being tested as an antineoplastic agent because of its anti-proliferative, anti-angiogenic and anti-tubulin properties.
- Sulfate conjugation is one potential metabolic pathway for 2-ME₂.
- We found that seven of the ten known human SULT isoforms can catalyze the sulfation of 2-ME₂.
- Of the isoforms studied, SULT1E1 had the lowest apparent K_m value for 2-ME₂.
- The common allozymes for SULT1A1 also catalyzed the sulfation of 2-ME₂. Therefore, if this isoform contributes significantly to 2-ME₂ biotransformation *in vivo*, genetic variation in SULT1A1 might contribute to individual differences in 2-ME₂ metabolism, pharmacokinetics and therapeutic efficacy.
- The next step in these studies will require a determination of the relative importance of sulfate conjugation in the metabolism of 2-ME₂ when this agent is administered in a clinical setting.

"SO WHAT"

As a result of these studies, we have evidence that SULTs metabolize the anti-tumorigenic drug, 2-ME₂. Since many of the human SULTs are genetically polymorphic, genotyping patients prior to treatment, perhaps, may predict response and/or toxicity and allow for tailoring of drug doses to individual patients.

REFERENCES

Bradford, M.M. A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal. Biochem.* 72:248-254, 1976.

Bradlow, H.L., Miichnovicz, J.J., Telang, N.T., and Osborne, M.P. Effects of dietary indole-3-carbinol on estradiol metabolism and spontaneous mammary tumors in mice. *Carcinogenesis* 12:1571-1574, 1991.

Campbell, N.R.C., Van Loon, J.A. and Weinshilboum, R.M. Human liver phenol sulfotransferase: assay conditions, biochemical properties and partial purification of isozymes of the thermostable form. *Biochem. Pharmacol.* 36:1435-1446, 1987.

Cleland, W.W. Computer programmes for processing enzyme kinetic data. *Nature (Lond.)* 198:463-465, 1963.

Foldes, A. and Meek, J.L. Rat brain phenolsulfotransferase - partial purification and some properties. *Biochim. Biophys. Acta* 327:365-374, 1973.

Cushman, M., He, H.-M., Katzenellenbogen, J.A., Lin, C.M. and Hamel, E. Synthesis, antitubulin and antimetabolic activity, and cytotoxicity of analogs of 2-methoxyestradiol, an endogenous mammalian metabolite of estradiol that inhibits tubulin polymerization by binding to the colchicine binding site. *J. Med. Chem.* 38:2041-2049, 1995.

Cushman, M., He, H.-M., Katzenellenbogen, J.A., Varma, R.K. and Hamel, E., Lin, C.M., Ram, S. and Sachdeva, Y. P. Synthesis of analogs of 2-methoxyestradiol with enhanced inhibitory effects on tubulin polymerization and cancer cell growth. *J. Med. Chem.* 40:2323- 2334, 1997.

D'Amato, R.J., Lin, C.M., Flynn, E., Folkman, J. and Hamel, E. 2-Methoxyestradiol, an endogenous mammalian metabolite, inhibits tubulin polymerization by interacting at the colchicine site. *Proc. Natl Acad. Sci. USA*, 91:3964 -3968, 1994.

Fotsis, T., Zhang, Y., Pepper, M.S., Adlercreutz, H., Montesano, R., Nawroth, P.P. and Schweigerer, L. The endogenous oestrogen metabolite 2-methoxyoestradiol inhibits angiogenesis and suppresses tumour growth. *Nature* 368:237-239, 1994.

Hernández, J.S., Watson, R.W.G., Wood, T.C. and Weinshilboum, R.M. Sulfation of estrone and 17 β -estradiol in human liver: catalysis by thermostable phenol sulfotransferase and by dehydroepiandrosterone sulfotransferase. *Drug Met. Dispos.* 20:413-422, 1992.

Klauber, N., Parangi, S., Flynn, E., Hamel, E. and D'Amato, R.J. Inhibition of angiogenesis and breast cancer in mice by the microtubule inhibitors 2-methoxyestradiol and taxol. *Cancer Res.* 57:81-86, 1997.

Lottering, M.-L., Haag, M. and Seegers, J.C. Effects of 17 β -estradiol metabolites on cell cycle events in MCF-7 cells. *Cancer Res.* 52:5926 -5932, 1992.

Osborne, M.P., Telang, N.T., Kaur, S. and Bradlow, H.L. Influence of chemopreventive agents on estradiol metabolism and mammary preneoplasia in the C3H mouse. *Steroids* 55:114-119, 1990.

Raftogianis, R.B., Wood, T.C., Otterness, D.M., Van Loon, J.A. and Weinshilboum, R.M. Phenol sulfotransferase pharmacogenetics in humans: association of common *SULT1A1* alleles with TS PST phenotype. *Biochem. Biophys. Res. Commun.* 239:298-304, 1997.

Raftogianis, R.B., Wood, T.C. and Weinshilboum, R.M. Human phenol sulfotransferases *SULT1A2* and *SULT1A1*: genetic polymorphisms, allozyme properties and human liver genotype-phenotype correlations. *Biochem. Pharmacol.* 58:605-610, 1999.

Wilkinson, G.N. Statistical estimations in enzyme kinetics. *Biochem. J.* 80:324-332, 1961.

Yager, J.D. and Liehr, J.G. Molecular mechanisms of estrogen carcinogenesis. *Ann. Rev. Pharmacol. Toxicol.* 36:203-232, 1996.

Zhu, B.T. and Liehr, J.G. Quercetin increases the severity of estradiol-induced tumorigenesis in hamster kidney. *Toxicol. Appl. Pharmacol.* 125:149-158, 1994.

Zhu, B.T. and Liehr, J.G. Inhibition of catechol-O-methyltransferase-catalyzed O-methylation of 2 and 4-hydroxyestradiol by quercetin: possible role in estradiol-induced tumorigenesis. *J. Biol. Chem.* 271:1357-1363, 1996.

Zhu, B.T. and Conney, A.H. Is 2-methoxyestradiol an endogenous estrogen metabolite that inhibits mammary carcinogenesis? *Cancer Res.* 58:2269-2277, 1998.

Zheng, W., Xie, D.W., Cerhan, J.R., Sellers, T.A., Wen, W.Q. and Folsom, A.R. Sulfotransferase 1A1 (*SULT1A1*) polymorphism, endogenous estrogen exposure, well-done meat intake, and breast cancer risk. *Cancer Epi, Biomarker Prevent.* In press, 2001.

APPENDICES

Appendix I :

2-METHOXYESTRADIOL (2-ME₂) SULFATION: POSSIBLE METABOLIC PATHWAY. A.A. Adjei, PhD*, T.C. Wood, B.A.* and R.M. Weinshilbourn, MD. Mayo Clinic-Mayo Foundation, Rochester, MN

2-ME₂ is an endogenous estrogen metabolite that inhibits the proliferation of breast and other human cancer cell lines. 2-ME₂ also has potent anti-angiogenic and anti-tubulin properties, and it may inhibit estrogen-induced carcinogenesis in the mammary gland. We set out to test the hypothesis that 2-ME₂ might be a substrate for sulfate conjugation and, therefore, that individual variations in the sulfation of 2-ME₂ might contribute to individual differences in its metabolism, pharmacokinetics and therapeutic efficacy. As a first step, we tested 2-ME₂ as a substrate for 7 human sulfotransferase (SULT) isoforms -- as well as all of the common allozymes for SULT1A1 and 1A2. Substrate kinetic studies were conducted in two stages -- starting with concentrations over 5 orders of magnitude, followed by determination of K_m values over a narrow concentration range. 2-ME₂ was a sulfate acceptor substrate for SULT1A1*1, *2, *3; 1A2*1, *2, *3; 1A3; 1E1; 2A1; 2B1a and 2B1b, with apparent K_m values of 2.5, 5.2, 1.6; 4.2, 111, 5.3; 91; 0.067; 8.3; 4.1 and 4.1 μM, respectively. These results suggest that individual pharmacogenetic variation in sulfate conjugation might contribute to individual differences in 2-ME₂ pharmacokinetics and therapeutic effect.


[Supported by DAMD Grant DAMD17-00-1-0684]

Appendix II : See attached Figures and Table on pages 9-11.

N.B.

FIGURES AND TABLE IN APPENDIX II ARE PROPRIETARY DATA.

SULT Catalyzed 2-Methoxyestradiol Sulfation



2-Methoxyestradiol

2-Methoxyestradiol-3-O-Sulfate

Sulfation of 2-Methoxyestradiol by SULT1E1




Table 1.

**SUBSTRATE KINETICS FOR SULT ISOFORMS :
REACTION WITH 2-METHOXYESTRADIOL**

Recombinant SULT Isoforms	Apparent Km Value (uM \pm S.E)	Vmax Units/B-Gal units	V/K(x 10 ³)
1A1*1	2.5 \pm 0.1	2.97	1188
1A1*2	5.2 \pm 0.4	3.79	729
1A1*3	1.6 \pm 0.2	1.13	707
1A2*1	4.2 \pm 0.3	1.56	372
1A2*2	111 \pm 0.5	1.51	14
1A2*3	5.3 \pm 0.3	0.34	65
1A3	91.4 \pm 23.0	0.97	11
1B1	ND	ND	ND
1C1	ND	ND	ND
1E1	0.067 \pm 0.0	1.65	24591
2A1	8.3 \pm 0.9	0.21	25
2B1a	4.1 \pm 0.1	0.51	124
2B1b	4.1 \pm 0.3	0.71	173
4A1	ND	ND	ND

ND : No detectable signal



DEPARTMENT OF THE ARMY
US ARMY MEDICAL RESEARCH AND MATERIEL COMMAND
504 SCOTT STREET
FORT DETRICK, MARYLAND 21702-5012

REPLY TO
ATTENTION OF:

MCMR-RMI-S (70-1y)

28 Aug 02

MEMORANDUM FOR Administrator, Defense Technical Information Center (DTIC-OCA), 8725 John J. Kingman Road, Fort Belvoir, VA 22060-6218

SUBJECT: Request Change in Distribution Statement

1. The U.S. Army Medical Research and Materiel Command has reexamined the need for the limitation assigned to technical reports written for this Command. Request the limited distribution statement for the enclosed accession numbers be changed to "Approved for public release; distribution unlimited." These reports should be released to the National Technical Information Service.

2. Point of contact for this request is Ms. Kristin Morrow at DSN 343-7327 or by e-mail at Kristin.Morrow@det.amedd.army.mil.

FOR THE COMMANDER:

Encl


PHYLIS M. RINEHART
Deputy Chief of Staff for
Information Management

ADB231838
ADB240253
ADB251610
ADB275099
ADB253637
ADB261538
ADB275186
ADB264648
ADB275102
ADB241899
ADB259033
ADB266113
ADB275663
ADB254489
ADB262700
ADB276708
ADB274345
ADB274844
ADB275154
ADB275535
ADB275101
ADB275451
ADB274597
ADB273871
ADB275145
ADB274505
ADB275851
ADB274459
ADB277942
ADB277404
ADB277494
ADB277536