

UNCLASSIFIED

Defense Technical Information Center
Compilation Part Notice

ADP010922

TITLE: The Riemann Non-Differentiable Function
and Identities for the Gaussian Sums

DISTRIBUTION: Approved for public release, distribution unlimited

This paper is part of the following report:

TITLE: Paradigms of Complexity. Fractals and
Structures in the Sciences

To order the complete compilation report, use: ADA392358

The component part is provided here to allow users access to individually authored sections of proceedings, annals, symposia, ect. However, the component should be considered within the context of the overall compilation report and not as a stand-alone technical report.

The following component part numbers comprise the compilation report:

ADP010895 thru ADP010929

UNCLASSIFIED

THE RIEMANN NON-DIFFERENTIABLE FUNCTION AND IDENTITIES FOR THE GAUSS SUMS

B. T. MATKARIMOV

Institute of Mathematics, Pushkin str. 125, Almaty 480100, Kazakhstan

E-mail: matkarimov@itte.kz

Riemann's example of a continuous, non-differentiable function is given by the sum $\sum_{n=1}^{\infty} \sin(n^2x)/n^2$. This function is sufficiently irregular and its graph is fractal. Hardy proved ¹ that Riemann's non-differentiable function is not differentiable at any irrational point because of a square root singularity at these points. Careful investigation of the differentiability of Riemann's non-differentiable function was carried out by Gerver, who showed ² that this function has derivative equal to $-1/2$ at every rational point of a special type (forming the orbit of the point 1 under the theta-modular group³). Different proofs of this surprising fact was given by other authors^{3,4,5}, providing also a close relation between Riemann's non-differentiable function and classical θ -function and Gauss sums. Duistermaat obtained ³ an exact functional equations for this function under transformations of the theta-modular group. In this article we use functional equations for Riemann's non-differentiable function under theta-modular transformations to derive functional equations on Gauss sums generalizing Genocchi-Schaar identity.

A Gauss sum is a sum of the form

$$S(p, q) = \sum_{n=0}^{q-1} \exp(\pi i n^2 p/q),$$

where p and q are relatively prime integers of opposite parity, i.e. one is odd and the other is even. The Genocchi-Schaar identity on Gauss sums is the following: for positive integers p and q of opposite parity,

$$\frac{1}{\sqrt{q}} \sum_{n=0}^{q-1} \exp(\pi i n^2 p/q) = \frac{\exp(\pi i/4)}{\sqrt{p}} \sum_{n=0}^{p-1} \exp(-\pi i n^2 q/p).$$

This identity can be interpreted as the transformation of the Gauss sum under the change $\sigma : z \rightarrow -1/z$ where $z = p/q$.

The modular group Γ is a group of fractional linear transformations $\gamma : z \rightarrow (az + b)/(cz + d)$ where $a, b, c, d \in Z$ and $ad - bc = 1$. Theta-modular group Γ_{θ} is a sub-group of modular group generated by the following mappings: $\tau : z \rightarrow z + 2$ and $\sigma : z \rightarrow -1/z$. For any element of the theta-modular group the following are valid: $ab \equiv 0 \pmod{2}$ and $cd \equiv 0 \pmod{2}$. Every fractional transformation $\gamma \in \Gamma_{\theta}$ has a simple pole at the point $z = -d/c$. The rational points $x = p/q$ with integers p, q of opposite parity and infinity constitute theta-modular orbit of point 0.

Riemann's non-differentiable function has the following local estimations at the point x : $f(x \pm \epsilon) = f(x) + R_{\pm} \epsilon^{\delta}$, where $0 < \delta \leq 1$. Using the functional equations for $f(x)$ it is possible to find functional equations on the functions R_{\pm} . These functions are known only at the rational points, where they coincide with the Gauss sums. The following theorems describe these functional equations:

Theorem: Let r and s be integers of which one is even and the other is odd, and s is positive, $\gamma \in \Gamma_\theta$ be an element of the theta-modular group, $\gamma : z \rightarrow (az + b)/(cz + d)$, and suppose c is positive and $cr + ds \neq 0$. Define $r' = ar + bs$ and $s' = cr + ds$. Then the following formula for Gauss sums is valid:

$$\frac{\exp(\pi i \text{Sign}(s')/4)}{\sqrt{|s'|}} S(r', s') \frac{1}{\sqrt{c}} S(-d, c) = \frac{1}{\sqrt{s}} S(r, s) \quad (1)$$

Proof: The proof is based on the differentiability properties of Riemann's non-differentiable function. Let us consider the following function $\phi(z) = \sum_{n=1}^{\infty} \exp(\pi i n^2 z)/\pi i n^2$. Using technique of papers^{4,5} we calculate the following estimation for the function $\phi(z)$ at the point $x = u/v$, where u and v are relatively prime integers of opposite parity, $(u, v) = 1$ and $uv \equiv 0 \pmod{2}$:

$$\phi(x + h^2) - \phi(x - h^2) = h^2 \sum_{n=-\infty}^{+\infty} \exp(\pi i n^2 x) \varphi(nh) - h^2 = \quad (2)$$

$$h^2 \sum_{t=0}^{|v|-1} \exp(\pi i t^2 x) \sum_{k=-\infty}^{+\infty} \varphi(k|v|h + th) - h^2 = 2^{1/2} S(u, v) h/|v| + O(h^2).$$

Here $\varphi(x) = \sin(\pi x^2)/\pi x^2$ if $x \neq 0$ and $\varphi(x) = 1$ if $x = 0$, and we write $n = k|v| + t$, $(0 \leq t \leq |v|)$ and use that $\exp(\pi i n^2 x) = \exp(\pi i t^2 x)$, since $uv \equiv 0 \pmod{2}$. The function $\phi(z)$ obey a functional equation under the action of theta-modular group³. Let $\gamma \in \Gamma_\theta$ be an element of theta modular group, then the function $\psi(z) = \phi(z) - \gamma'(z)^{-1} \mu_\gamma(z) \phi(\gamma(z))$ is differentiable, and analytical function μ_γ is given by:

$$\mu_\gamma(z) = e^{(\pi i/4)} c^{-1} (z + d/c)^{-1/2} S(-d, c). \quad (3)$$

Eq. (2) is valid for any point of theta-modular orbit of $x = r/s$, except infinity. Calculating estimation (2) for the differentiable function $\psi(x)$ at the point $x = r/s$ and supposing $\gamma(r/s) \neq \infty$, we find that

$$\frac{S(r, s)}{s} - \gamma'(r/s)^{-1/2} \mu_\gamma(r/s) \frac{S(ar + bs, cr + ds)}{|cr + ds|} = 0. \quad (4)$$

Note, that $r' = ar + bs$ and $s' = cr + ds$ are relatively prime integers. Substituting the expression of $\mu_\gamma(z)$ in terms of Gauss sums (Eq. 3) we obtain the desired result.

Q.E.D

The Genocchi-Schaar identity corresponds to the case of $a = 0$, $b = -1$, $c = 1$ and $d = 0$. Functional equation (Eq. 1) describe the theta-modular transformations of Gauss sums and can be used to derive the values of the Gauss sums.

References

1. G. H. Hardy, *Trans. AMS* **17**, 301 (1916)
2. J. Gerver, *Amer. J. Math.* **92**, 33 (1970)
3. J. J. Duistermaat, *Nieuw Archief voor Wiskunde* **9**, 303 (1991)
4. A. Smith, *Proc. AMS* **34**, 463 (1972)
5. W. Luther, *J. Approx. Theory* **48**, 303 (1986)