


AD 699 127

PERFORMANCE OF MONKEYS AFTER PARTIAL  
BODY IRRADIATION

J. W. Thorp, et al

Armed Forces Radiobiology Research Institute  
Bethesda, Maryland

November 1969



This document has been approved for public release and sale.


AFRRI SR69-18  
November 1969

PERFORMANCE OF MONKEYS AFTER PARTIAL BODY IRRADIATION

J. W. THORP  
J. E. GERMAS



R. E. GEORGE  
Commander, MSC, USN  
Chairman  
Radiation Biology Department



HUGH B. MITCHELL  
Colonel, USAF, MC  
Director



W. F. DAVIS, JR.  
Acting Chairman  
Behavioral Sciences Department

ARMED FORCES RADIOBIOLOGY RESEARCH INSTITUTE  
Defense Atomic Support Agency  
Bethesda, Maryland

---

Distribution of this document is unlimited

#### ACKNOWLEDGMENT

The authors wish to acknowledge the valuable assistance of Q. H. Shelton and J. H. Haines who helped train and prepare the subjects. W. F. Pfeiffer contributed greatly in providing the extensive dosimetry measurements.

TABLE OF CONTENTS

	Page
Foreword (Nontechnical summary) . . . . .	iii
Abstract . . . . .	v
I. Introduction . . . . .	1
II. Procedure . . . . .	1
III. Results . . . . .	3
Two-Hour Test Period . . . . .	3
Overall Performance and Survival . . . . .	5
IV. Discussion . . . . .	6
References . . . . .	9

LIST OF TABLES

Table I. Performance of Unshielded Monkeys after a 4500-rad Pulse of Mixed Gamma-Neutron Radiations . . . . .	4
Table II. Performance of Head-Shielded Monkeys after a 4500-rad Pulse of Mixed Gamma-Neutron Radiations . . . . .	4
Table III. Performance of Trunk-Shielded Monkeys after a 4500-rad Pulse of Mixed Gamma-Neutron Radiations . . . . .	5

FOREWORD  
(Nontechnical summary)

The results of several studies have shown that, for a broad range of supra-lethal doses of ionizing radiation, postirradiation performance of trained monkeys has a rather well-defined pattern. Almost immediately after irradiation there is a period of transient performance decrement; within 2 hours, ability to perform effectively usually returns to a functional level. Later there is a period of permanent complete incapacitation, and death occurs shortly thereafter.

The main purpose of this research was to determine whether head shielding or trunk shielding would alter the pattern of postirradiation performance. Another purpose was to assess the relative importance of head and trunk structures in postirradiation performance decrement.

Fifteen male and fifteen female monkeys (Macaca mulatta) were trained by shock avoidance conditioning to work a visual discrimination problem. Each animal received a 4500-rad midline tissue dose of pulsed mixed gamma-neutron radiations; of each sex, five were head-shielded, five were trunk-shielded, and five were not shielded. The midline tissue dose behind the shield was less than 8 percent of the midline tissue dose to the same point without the shield in place. Postirradiation performance was evaluated.

The results indicate that, after irradiation under the conditions of this study, there is a high probability that severe early performance decrement will occur in unshielded monkeys. Either trunk shielding or head shielding will decrease the probability of early performance decrement. However, if a shielded animal does suffer

an early performance decrement, the head-shielded monkey will recover sooner than the trunk-shielded monkey.

The results also indicate that, if a monkey survives the first 2 hours postirradiation, it will recover to perform well for some time before death. Since both types of shielding decrease the probability of an early death, both will increase the chance that a shielded subject can work effectively at some time after irradiation, even though early transient incapacitation might occur.

#### ABSTRACT

Fifteen male and fifteen female monkeys (Macaca mulatta) were trained by shock avoidance conditioning to work a visual discrimination problem. Each animal received a 4500-rad midline tissue dose of pulsed mixed gamma-neutron radiations; of each sex, five were head-shielded, five were trunk-shielded, and five were not shielded. The midline tissue dose behind the shield was less than 8 percent of the midline tissue dose to the same point without the shield in place. Postirradiation performance was evaluated. Early performance decrement occurred after irradiation in all unshielded monkeys. Either trunk shielding or head shielding decreased the probability of performance decrement. However, among the shielded monkeys that did suffer an early decrement, the head-shielded subjects recovered sooner than the trunk-shielded subjects.

## I. INTRODUCTION

The results of several studies<sup>2-4</sup> have shown that, for a broad range of supra-lethal doses of ionizing radiation, the postirradiation performance of trained monkeys has a rather well-defined pattern. Almost immediately after irradiation there is a period of transient performance decrement; within 2 hours or less, ability to perform effectively usually returns to a functional level. Later there is a period of permanent complete incapacitation, and death occurs shortly thereafter.

The main purpose of this research was to determine whether head shielding or trunk shielding would alter the pattern of postirradiation performance. Another purpose was to assess the relative importance of head and trunk structures in postirradiation performance decrement.

## II. PROCEDURE

The subjects were male and female monkeys (Macaca mulatta) that weighed between 3 and 5 kg. The monkeys were trained by shock avoidance conditioning to work a visual discrimination problem. They were continuously restrained in primate chairs for about 25 days, i.e., the period required for training, exposure, and postexposure testing. Each chaired subject was kept in an isolation box throughout the experimental period.

Fifteen males and fifteen females were irradiated individually with the AFRRITRIGA reactor operated in the pulsed mode.<sup>5</sup> All received approximately the same dose of pulsed mixed gamma-neutron radiations except that five of each sex were head-shielded, five were trunk-shielded, and five were not shielded. Each subject



faced away from the reactor core with its center line about 80 cm from the vertical core center line. The methods of shielding have been described.<sup>6</sup>

Midline tissue doses to the head and chest of each monkey were calculated as follows. First, the midline tissue doses to the head and chest of a Lucite phantom were measured with miniature, tissue-equivalent ionization chambers. Simultaneously, the reactor power output was monitored with sulfur tablets. Factors were then derived to describe the relationship of sulfur-monitored neutron fluence from the reactor and midline tissue doses in the phantom.

During each animal exposure, the reactor power was again monitored with sulfur tablets; doses were calculated by using the appropriate factors. All animals received midline tissue doses of about 4500 rads to the unshielded parts of the body. The midline tissue dose behind the shield was less than 8 percent of the midline tissue dose to the same point without the shield in place.

The discrimination problem consisted of a circle and square presented simultaneously by illuminating two keys on a console in front of the monkey. The relative positions of the circle and square were switched in a random fashion. The correct response was to press the key illuminated with the square.

Each trial lasted 10 seconds. It was initiated by simultaneous illumination of a house light in the isolation box and the stimulus keys on the console. The monkey had 5 seconds to respond. For a correct response, both house light and stimuli extinguished for the duration of the 10 seconds. However, for an incorrect response or for no response, the stimuli extinguished, the house light remained on, a tone was initiated, and the animal received a brief electrical shock. Before irradiation,

each subject was trained to a proficiency of 90 or more correct responses per 100 presentations.

Each test period consisted of 100 trials presented at 10-second intervals and lasted 16.7 minutes. Each test period was followed by a rest period with the monkey in complete darkness. One test period started at the time of the reactor pulse and others started at 20, 40, 60, 80, and 100 minutes postexposure. The subject was then moved from the exposure room to another area for later testing. One test period was presented each time beginning at 3, 4, 5, 6, 7, and 8 hours postexposure. Testing was then continued at 2-hour intervals until death.

### III. RESULTS

The term "performance decrement" is used to indicate fewer than 90 correct responses in a 100-trial test period. Among all subjects, incorrect responses were infrequent, i.e., the subject usually responded correctly or not at all. Results are presented in Tables I, II, and III.

Two-Hour Test Period. All of the unshielded subjects suffered performance decrement after irradiation and within this 2-hour test period. Eight of the ten monkeys failed to recover completely within 2 hours; in fact, five died within that time.

Among the head-shielded animals, three females and one male exhibited no early performance decrement. Only one female suffered any performance decrement beyond 20 minutes postexposure, and the decrement was very slight. All other performance decrements occurred very early, and the monkeys recovered to work at acceptable levels within 20 minutes postexposure.

Table I. Performance of Unshielded Monkeys after a 4500-rad Pulse of Mixed Gamma-Neutron Radiation :

	Performance during first 2 hours after exposure						Survival time (hours)	Performance from 2 hours postirradiation until death		
	Time testing began relative to pulse (minutes)							Last response (hours after irradiation)	Number of test periods with 90 or more correct	Number of test periods with any correct
	0	20	40	60	80	100				
	Number of correct responses (100 possible)									
<b>Males</b>										
1	17	91	41	0	1	76	79	28.3	14	17
2	18	3	96	98	97	98	20.7	14.3	9	9
3	66	86	55	64	71	71	27	24.3	8	14
4	73	93	74	80	68	38	151	127	3	52
5	71	51	1	0	0	0	1.2	.7	--	--
<b>Females</b>										
6	17	0	0	0	0	0	.3	.1	--	--
7	31	99	0	0	0	0	1.9	.6	--	--
8	27	98	23	14	99	99	33	29.2	14	14
9	10	0	0	0	0	0	1	<.1	--	--
10	4	0	0	0	0	0	.3	<.1	--	--

Table II. Performance of Head-Shielded Monkeys after a 4500-rad Pulse of Mixed Gamma-Neutron Radiations

	Performance during first 2 hours after exposure						Survival time (hours)	Performance from 2 hours postirradiation until death		
	Time testing began relative to pulse (minutes)							Last response (hours after irradiation)	Number of test periods with 90 or more correct	Number of test periods with any correct
	0	20	40	60	80	100				
	Number of correct responses (100 possible)									
<b>Males</b>										
11	95	99	100	98	100	90	42	40.2	21	22
12	55	99	97	99	97	99	29.7	26.2	14	15
13	14	91	98	98	98	95	18	16	3	6
14	74	98	100	99	99	99	26.5	26	12	15
15	55	91	94	98	98	99	37	35.1	20	21
<b>Females</b>										
16	94	99	98	97	99	97	4.5	2	--	--
17	49	88	79	93	91	78	21.5	20	11	12
18	91	91	100	98	99	99	24.3	22	10	13
19	39	99	97	99	99	92	18	17	10	11
20	93	99	100	99	99	92	20.5	18	10	10

Table III. Performance of Trunk-Shielded Monkeys after a 4500-rad Pulse of Mixed Gamma-Neutron Radiations

	Performance during first 2 hours after exposure						Survival time (hours)	Performance from 2 hours postirradiation until death		
	Time testing began relative to pulse (minutes)							Last response (hours after irradiation)	Number of test periods with 90 or more correct	Number of test periods with any correct
	0	20	40	60	80	100				
	Number of correct responses (100 possible)									
<b>Males</b>										
21	92	92	87	90	86	85	19	16	4*	10
22	98	98	86	96	96	96	78	51	17	28
23	97	99	99	100	100	98	36	30.5	16	17
24	13	0	96	100	100	100	35	32	18	18
25	13	97	77	21	100	100	49	43.5	22	24
<b>Females</b>										
26	45	67	88	96	96	99	38	28	13	16
27	42	75	65	82	20	6	50	32	15	19
28	98	76	50	43	40	16	41	22	3	8
29	76	98	100	100	100	100	21.5	21	11	12
30	40	96	94	94	90	91	29.5	20.3	11	12

\* Shock mechanism became defective after 3 hours postexposure so monkey was not reinforced for poor performance. Condition was corrected about 7 hours postexposure.

Three of the trunk-shielded males showed slight or no performance decrement in the first 2 hours. Two females had early serious decrements but recovered fully within 20 minutes. Four subjects (two males and two females) suffered serious early performance decrements and required more than 20 minutes for recovery. One female had no decrement in the first 20 minutes but did show a continuous deterioration of performance through the remainder of the first 2 hours.

Overall Performance and Survival. Survival times of unshielded monkeys ranged from 17 minutes to 151 hours. One male and four females survived less than 2 hours. None died in the interval of 2 to 21 hours postirradiation.

The survival time of shielded monkeys was less variable than with unshielded animals. All shielded subjects lived more than 2 hours after irradiation but one

head-shielded female lived for only 4.5 hours. All other shielded subjects lived for at least 18 hours, and most lived for 20 hours or more. In both shielded groups, the mean survival time was slightly less for females than for males. The mean survival time for trunk-shielded subjects was slightly longer than for head-shielded subjects.

Except for the head-shielded female that died in 4.5 hours, all animals which survived the first 2 hours performed at acceptable levels for some time after the first 2 hours postexposure. This was true whether the monkey was unshielded, head-shielded, or trunk-shielded, and whether or not there was early performance decrement. Each monkey usually made its last response about 3 or 4 hours before death.

#### IV. DISCUSSION

The results indicate that, after irradiation under the conditions of this study, there is a high probability that early performance decrement will occur in unshielded monkeys. Either trunk shielding or head shielding will decrease the probability of early performance decrement. However, if a shielded animal does suffer an early performance decrement, the head-shielded monkey will recover sooner than the trunk-shielded monkey. Chapman and Young<sup>1</sup> found a similar response in monkeys that were required to perform a discrete avoidance task after a 6250-rad pulsed dose of fission spectrum radiations.

It was not possible to completely separate the importance of head and trunk effects. All of the monkeys did receive more than 350 rads to the shielded part of the body. That dose, about equal to the LD<sub>50/60</sub> for monkeys,<sup>7</sup> may have been

adequate (when combined with the higher dose to the unshielded body) to elicit some of the effects that are normally mediated through the structures that were shielded. Furthermore, the mixed gamma-neutron radiation could not be well collimated due to the large source dimensions and the short source-to-animal distance. Therefore, a dose gradient existed in the animals from the position corresponding to that of the dosimeter in the phantom to the midline position at the edge of the shield. Until these problems can be resolved, it is suggested that both head and trunk structures are implicated in the early performance decrement that occurs after irradiation of trained monkeys.

The results of the currently reported study indicate that, if an animal survives the first 2 hours postirradiation, it will recover to perform well for some time before death. Since both types of shielding decrease the probability of an early death, both increase the probability that the shielded subject can work several hours after irradiation, even though early transient incapacitation might occur. However, the unshielded subjects that escaped the early death worked and survived nearly as long as the shielded subjects.

Trunk shielding did extend survival time when compared to unshielded and head-shielded subjects ( $P < 0.05$ ). However, the differences were small, especially if the unshielded animals that survived less than 2 hours are omitted. The extended survival of trunk-shielded animals contrasts to earlier results where head shielding extended the survival of beagles receiving pulsed mixed gamma-neutron radiations, but trunk shielding was not beneficial.<sup>6</sup> However, the beagles received doses of 19,000 rads or more compared to 4500 rads for the monkeys. This leads one to

speculate that, at relatively lower whole-body supralethal doses, the primary cause of death is damage to trunk structures, and survival time does not change rapidly with increased trunk doses. At higher doses, cephalic structures become relatively more important, and survival time decreases rapidly as the head dose is increased. However, since the monkeys were chaired and working, caution is required in comparing them to the dogs that were unrestrained (except for about 20 minutes during exposure) and not required to work.

## REFERENCES

1. Chapman, P. H. and Young, R. J. Effect of head versus trunk fission-spectrum radiation on learned behavior in the monkey. Brooks Air Force Base, Texas, U. S. Air Force School of Aerospace Medicine Report TR68-80, 1968.
2. de Haan, H. J. and Germas, J. E. Visual discrimination performance in the monkey (Macaca mulatta): A technique and assessment of 5000 rads gamma-neutron irradiation. Bethesda, Maryland, Armed Forces Radiobiology Research Institute Scientific Report SR68-16, 1968.
3. de Haan, H. J., Kaplan, S. J. and Germas, J. E. Visual discrimination performance in the monkey following a 5,000-rad pulse of mixed gamma-neutron radiation. Bethesda, Maryland, Armed Forces Radiobiology Research Institute Scientific Report SR69-1, 1969.
4. Germas, J. E., Kaplan, S. J. and de Haan, H. J. Visual discrimination performance in the monkey following a 10,000-rad pulse of mixed gamma-neutron radiation. Bethesda, Maryland, Armed Forces Radiobiology Research Institute Scientific Report SR69-2, 1969.
5. Pitchford, T. L. Beagle incapacitation and survival time after pulsed mixed gamma-neutron irradiation. Bethesda, Maryland, Armed Forces Radiobiology Research Institute Scientific Report SR68-24, 1968.
6. Thorp, J. W. Head shielding protection for beagles exposed to supralethal doses of pulsed mixed gamma-neutron radiations. Bethesda, Maryland, Armed Forces Radiobiology Research Institute Scientific Report SR68-26, 1968.
7. Wise, D. and Turbyfill, C. L. The acute mortality response of monkeys (Macaca mulatta) to pulsed mixed gamma-neutron radiations. Bethesda, Maryland, Armed Forces Radiobiology Research Institute Scientific Report SR68-17, 1968.



UNCLASSIFIED

Security Classification

DOCUMENT CONTROL DATA - R&D		
<i>(Security classification of title, body of abstract and indexing annotation must be entered when the overall report is classified)</i>		
1. ORIGINATING ACTIVITY (Corporate author)		2a. REPORT SECURITY CLASSIFICATION
Armed Forces Radiobiology Research Institute Defense Atomic Support Agency Bethesda, Maryland 20014		UNCLASSIFIED
		2b. GROUP N/A
3. REPORT TITLE		
PERFORMANCE OF MONKEYS AFTER PARTIAL BODY IRRADIATION		
4. DESCRIPTIVE NOTES (Type of report and inclusive dates)		
5. AUTHOR(S) (Last name, first name, initial)		
Thorp, J. W. and Germas, J. E.		
6. REPORT DATE	7a. TOTAL NO OF PAGES	7b. NO OF REFS
November 1969	17	7
8a. CONTRACT OR GRANT NO.		9a. ORIGINATOR'S REPORT NUMBER(S)
A. PROJECT NO.		AFRRI SR69-18
c. MA 1 90406		9b. OTHER REPORT NO(S) (Any other numbers that may be assigned this report)
d.		
10. AVAILABILITY/LIMITATION NOTICES		
Distribution of this document is unlimited		
11. SUPPLEMENTARY NOTES		12. SPONSORING MILITARY ACTIVITY
		Defense Atomic Support Agency Washington, D. C. 20305
13. ABSTRACT		
<p>Fifteen male and fifteen female monkeys (<u>Macaca mulatta</u>) were trained by shock avoidance conditioning to work a visual discrimination problem. Each animal received a 4500-rad midline tissue dose of pulsed mixed gamma-neutron radiations; of each sex, five were head-shielded, five were trunk-shielded, and five were not shielded. The midline tissue dose behind the shield was less than 8 percent of the midline tissue dose to the same point without the shield in place. Postirradiation performance was evaluated. Early performance decrement occurred after irradiation in all unshielded monkeys. Either trunk shielding or head shielding decreased the probability of performance decrement. However, among the shielded monkeys that did suffer an early decrement, the head-shielded subjects recovered sooner than the trunk-shielded subjects.</p>		

DD FORM 1473  
1 JAN 66UNCLASSIFIED  
Security Classification

14. KEY WORDS	LINK A		LINK B		LINK C	
	ROLE	WT	ROLE	WT	ROLE	WT

INSTRUCTIONS

1. **ORIGINATING ACTIVITY:** Enter the name and address of the contractor, subcontractor, grantee, Department of Defense activity or other organization (corporate author) issuing the report.
- 2a. **REPORT SECURITY CLASSIFICATION:** Enter the overall security classification of the report. Indicate whether "Restricted Data" is included. Marking is to be in accordance with appropriate security regulations.
- 2b. **GROUP:** Automatic downgrading is specified in DoD Directive 5200.10 and Armed Forces Industrial Manual. Enter the group number. Also, when applicable, show that optional markings have been used for Group 3 and Group 4 as authorized.
3. **REPORT TITLE:** Enter the complete report title in all capital letters. Titles in all cases should be unclassified. If a meaningful title cannot be selected without classification, show title classification in all capitals in parentheses immediately following the title.
4. **DESCRIPTIVE NOTES:** If appropriate, enter the type of report, e.g., interim, progress, summary, annual, or final. Give the inclusive dates when a specific reporting period is covered.
5. **AUTHOR(S):** Enter the name(s) of author(s) as shown on or in the report. Enter last name, first name, middle initial. If military, show rank and branch of service. The name of the principal author is an absolute minimum requirement.
6. **REPORT DATE:** Enter the date of the report as day, month, year, or month, year. If more than one date appears on the report, use date of publication.
- 7a. **TOTAL NUMBER OF PAGES:** The total page count should follow normal pagination procedures, i.e., enter the number of pages containing information.
- 7b. **NUMBER OF REFERENCES:** Enter the total number of references cited in the report.
- 8a. **CONTRACT OR GRANT NUMBER:** If appropriate, enter the applicable number of the contract or grant under which the report was written.
- 8b, 8c, & 8d. **PROJECT NUMBER:** Enter the appropriate military department identification, such as project number, subproject number, system numbers, task number, etc.
- 9a. **ORIGINATOR'S REPORT NUMBER(S):** Enter the official report number by which the document will be identified and controlled by the originating activity. This number must be unique to this report.
- 9b. **OTHER REPORT NUMBER(S):** If the report has been assigned any other report numbers (either by the originator or by the sponsor), also enter this number(s).

10. **AVAILABILITY/LIMITATION NOTICES:** Enter any limitations on further dissemination of the report, other than those imposed by security classification, using standard statements such as:
  - (1) "Qualified requesters may obtain copies of this report from DDC."
  - (2) "Foreign announcement and dissemination of this report by DDC is not authorized."
  - (3) "U. S. Government agencies may obtain copies of this report directly from DDC. Other qualified DDC users shall request through \_\_\_\_\_."
  - (4) "U. S. military agencies may obtain copies of this report directly from DDC. Other qualified users shall request through \_\_\_\_\_."
  - (5) "All distribution of this report is controlled. Qualified DDC users shall request through \_\_\_\_\_."

If the report has been furnished to the Office of Technical Services, Department of Commerce, for sale to the public, indicate this fact and enter the price, if known.

11. **SUPPLEMENTARY NOTE:** Use for additional explanatory notes.
12. **SPONSORING MILITARY ACTIVITY:** Enter the name of the departmental project office or laboratory sponsoring (paying for) the research and development. Include address.
13. **ABSTRACT:** Enter an abstract giving a brief and factual summary of the document indicative of the report, even though it may also appear elsewhere in the body of the technical report. If additional space is required, a continuation sheet shall be attached.

It is highly desirable that the abstract of classified reports be unclassified. Each paragraph of the abstract shall end with an indication of the military security classification of the information in the paragraph, represented as (TS), (S), (C), or (U).

There is no limitation on the length of the abstract. However, the suggested length is from 150 to 225 words.

14. **KEY WORDS:** Key words are technically meaningful terms or short phrases that characterize a report and may be used as index entries for cataloging the report. Key words must be selected so that no security classification is required. Identifiers, such as equipment model designation, trade name, military project code name, geographic location, may be used as key words but will be followed by an indication of technical context. The assignment of lines, rules, and weights is optional.