



DEPARTMENT OF THE ARMY
 WATERWAYS EXPERIMENT STATION, CORPS OF ENGINEERS
 P.O. BOX 631
 VICKSBURG, MISSISSIPPI 39180-0631

REPLY TO
 ATTENTION OF

CEWES-EV-I

18 August 1989

Environmental Effects of Dredging
 Technical Note
 EEDP-01-15 (May 1989)

SDTIC
 ELECTE
 MAR 29 1995
 C D

BIOACCUMULATION OF CHLORINATED CONTAMINANTS AND CONCOMITANT
 SUBLETHAL EFFECTS IN MARINE ANIMALS: AN ASSESSMENT
 OF THE CURRENT LITERATURE

Technical Note EEDP-01-15 (May 1989) was published without the complete
 References listing. It has been reprinted to correct this error; please
 discard your other copy.

Approved for public release
 Dissem. Unlimited

19950327 124

DEPT OF ARMY



Environmental Effects of Dredging Technical Notes



BIOACCUMULATION OF CHLORINATED CONTAMINANTS AND CONCOMITANT SUBLETHAL EFFECTS IN MARINE ANIMALS: AN ASSESSMENT OF THE CURRENT LITERATURE

PURPOSE: This note focuses on studies evaluating the sublethal effects of chlorinated organic contaminants on marine and estuarine organisms. Its objective is twofold: (1) to survey the literature for papers reporting both the sublethal effects of organohalogens and the corresponding body burdens in marine fish and invertebrates and (2) to provide a source of information for Corps field elements who have site-specific concerns (e.g., reproductive effects in a particular organism exposed to a specific organohalogen).

BACKGROUND: The US Army Corps of Engineers has the responsibility to ensure that contaminated sediments are dredged and disposed of in a manner that will not have an unacceptable adverse impact on the environment. The aquatic disposal of dredged material is regulated under two Federal statutes: Section 404(b)(1) of the Federal Water Pollution Control Act, as amended (PL 92-500) and Section 103 of the Marine Protection, Research, and Sanctuaries Act, as amended (PL 92-532). The regulations implementing these laws often require an evaluation of sediment toxicity and bioaccumulation potential prior to dredging and aquatic disposal.

Approximately 370 million cu m of sediments are dredged every year in the United States (Engler 1980). Approximately half of that volume is placed in open water. In most instances, dredged material is not acutely toxic to aquatic organisms. Therefore, decision-makers have had to rely less on toxicity data and more heavily on the results of bioaccumulation tests to evaluate potential impacts on the environment. There is very little interpretive guidance to assist in this evaluation (Peddicord and Hansen 1983). This report, produced under Work Unit 31773, Environmental Interpretation of Consequences from Bioaccumulation, of the Long-term Effects of Dredging Operations (LEDO) Program was designed, in part, to help provide that interpretive guidance.

ADDITIONAL INFORMATION: Contact one of the authors, Ms. Alfreda B. Gibson, (601) 634-4027, or Dr. Thomas M. Dillon (601) 634-3922, or the manager of the Environmental Effects of Dredging Programs, Dr. Robert M. Engler, (601) 634-3624.

Approach

A literature search was performed for information on the sublethal effects of organohalogenated contaminants on marine and estuarine animals. Only those investigations which examined organismic endpoints (growth, reproduction, behavior, morphology, osmoregulation, and metabolism) were considered. The reasons for evaluating organismic sublethal endpoints are discussed in Dillon (1984). The scope of this literature review was large. More than 50 technical journals were individually reviewed (Table 1). Ten data base literature search services were also used to identify any additional papers (Table 1).

For every paper included in this review, the following information was recorded: contaminant, test animal, exposure time, contaminant exposure concentration, reported tissue concentration, and any observed biological effects. The test animal was identified by common name and/or phylogenetic group. Tissue concentrations were expressed on a wet-weight basis. Exposure concentrations were all reported as micrograms per litre (parts per billion) unless noted otherwise.

Analysis

Approximately 1,200 published papers reporting the sublethal effects of chlorinated contaminants on marine and estuarine animals were identified in the literature. Of these, only 37 papers (3 percent) contained both sublethal effects data and contaminant tissue concentrations (Table 2).

Growth and behavior were the most frequently examined sublethal endpoints, while metabolism and osmoregulation were the least examined. Effects on reproduction and morphology appeared to be intermediate choices. The test organisms used by most investigators were fish and arthropods. They appeared in 51 percent and 36 percent of the papers, respectively. The environmental contaminants most frequently tested were kepone (30 percent) and polychlorinated biphenyls (PCBs) (24 percent). Exposure to contaminants was mainly via aqueous solutions (73 percent) or food (27 percent). None of the residue-effects papers involved contaminated sediment.

Because only 3 percent of the sublethal effects investigations considered published concomitant tissue residue information, the data base for

establishing quantitative residue-effects relationships is very limited. Variations due to interspecific differences, exposure regimes, and analytical capabilities diminish the ability to generate quantitative contaminant-specific guidance. However, a very broad generalization can be made based on data contained in Table 2. For marine and estuarine organisms with whole body tissue residues of chlorinated organic contaminants at or near steady-state, the level of concern associated with potential adverse sublethal effects is:

- LOW for tissue concentrations $<0.1 \mu\text{g/g}$ wet weight
- MEDIUM for tissue concentrations $0.1-1.0 \mu\text{g/g}$ wet weight
- HIGH for tissue concentrations $>1.0 \mu\text{g/g}$ wet weight

These are not discrete thresholds, pass-fail values, or numerical criteria. Rather they are heuristic and are the only general guidance the data will allow.

| | |
|------------------------------|-------------------------|
| Accession For | |
| NTIS | CRA&I |
| DTIC | TAB |
| Unannounced Justification | |
| Per Form 50 | |
| Availability Code | |
| Dist | Final and/or Special |
| A-1 | |

Discussion

This review and assessment of the literature has shown that few laboratory investigations (3 percent) report both sublethal effects of chlorinated organic contaminants and tissue residue data. In an earlier review which included biochemical and cellular endpoints as well as organismic responses (Dillon 1984), a similar frequency of residue-effects in the published literature was noted (6 percent). This paucity of residue-effects information hampers the ability to generate contaminant-specific guidance for interpreting results of bioaccumulation tests. This does not mean, however, that evaluative techniques are nonexistent. There are several.

It is often desirable to make relative comparisons among bioaccumulation data rather than to infer specific effects from absolute tissue concentrations. For example, bioassay data from a specific project sediment can be compared to a reference value. This reference value may be generated in the laboratory by exposing bioassay organisms to sediment collected at or near the aquatic disposal site. The resultant tissue concentration is then the comparative standard. Exposure to reference sediment is carried out concurrently with project sediment bioassays. A reference value may also emerge from a consensus process in which tissue concentrations, representing indigenous

organisms in a spatially discrete area, are identified (e.g., New York District matrix values). In both instances, bioassay results are interpreted from the standpoint of "no further degradation." It is important to note that although statistically significant differences may be observed in the laboratory, they do not necessarily imply that unacceptable adverse impacts in the environment are imminent or even inevitable.

In addition to ecological effects, tissue residue data may be interpreted in terms of human health issues. This can be done directly when the bioassay organism (or appropriate surrogate) is one commonly ingested by man. Numerical guidance for assessing contaminated seafood has been developed by agencies such as the US Food and Drug Administration and the Australian National Health and Medical Research Council. A summary of these data can be found in Peddicord et al. (1986). Local guidance in the form of action levels for seafood may also be available from state officials and US Environmental Protection Agency (USEPA) Regional Offices. If the bioassay does not involve an organism normally consumed by man, human health impacts can still be evaluated, albeit in a more circuitous manner. This is done by examining the potential for trophic transfer in the marine food web. Transfer may include the phenomenon of biomagnification, but this process is not common for many contaminants when trophic levels are strictly aquatic (Kay 1984). Biomagnification can become very important quantitatively when the trophic transfer process exits the aquatic environment. An in-depth technical discussion of this subject can be found in Biddinger and Gloss (1984).

When interpreting bioassay results, one must assess not only individual contaminants but also the impacts of multiple contaminants within the same tissue matrix. The first step in this analysis is, "How many and how much?" This approach can be quite useful in initial evaluations. For example, one would be very concerned if 10 out of 12 compounds were taken up in substantial amounts. The concern would lessen if only a few contaminants were accumulated and/or the magnitude of uptake was small. If only 1 out of 12 was accumulated to levels just above control or reference concentrations, the level of concern would be lower still.

Once a significant potential for bioaccumulation is established, the toxicological importance of the different contaminants must be considered. The potential for unacceptable adverse effects is elevated when toxic contaminants such as mercury, cadmium, and PCBs are accumulated. In contrast, concern

lessens if less toxic compounds such as phthalates are found in the tissues of biota. How does one gauge relative toxicity? One of the better sources of information is the numerous toxicity tests conducted by the USEPA as part of their program to develop Water Quality Criteria (USEPA 1980). Additional guidance for determining the toxicological importance of various environmental contaminants can be found in Peddicord et al. (1986).

One question often asked when reviewing tissue residue data is, "How do interactions among the contaminants (e.g., synergism or antagonism) affect the organism?" All interactions that may (or may not) be occurring are expressed in the acute toxicity data. Therefore, this question is somewhat irrelevant for sediment bioassays. To determine the interactive effects among specific contaminants of concern for a particular marine organism, additional laboratory experiments would have to be conducted.

Summary

A review of the literature has shown that about 3 percent of studies investigating the sublethal effects of chlorinated organic contaminants on marine organisms contain both effects and concomitant tissue residue data.

The residue-effects information that is available (Table 2) can be very useful for interpreting the results of project-specific bioassays. It is believed that they also represent heuristic guidance, not to be confused with pass-fail or threshold criteria. For marine and estuarine organisms with whole body tissue residues of chlorinated organic contaminants at or near steady-state, the level of concern associated with adverse sublethal effects is generally:

LOW for tissue concentrations $<0.1 \mu\text{g/g}$ wet weight

MEDIUM for tissue concentrations $0.1-1.0 \mu\text{g/g}$ wet weight

HIGH for tissue concentrations $>1.0 \mu\text{g/g}$ wet weight

Although the paucity of residue-effects information hampers contaminant-specific guidance, other evaluative techniques are available for interpreting the biological importance of bioaccumulation. For evaluating potential ecological effects, comparisons to reference values derived either in the laboratory or by consensus agreements can be carried out. To assess the potential

for human health impacts, bioaccumulation results can be compared directly to previously developed numerical guidance for contaminated seafood. Human health effects can also be evaluated indirectly by examining trophic transfer potential of contaminants in the marine food web. Finally, tissue residue information can be evaluated by determining the number of contaminants showing mobility, the magnitude of uptake relative to control and/or reference values, and the toxicological importance of contaminants that are bioaccumulated.

References

- Armstrong, D. A., Buchanan, D. V., Mallon, M. H., Caldwell, R. S., and Millemann, R. E. 1976. "Toxicity of the Insecticide Methoxychlor to the Dungeness Crab *Cancer magister*," *Marine Biology*, Vol 38, pp 239-252.
- Biddinger, G. R., and Gloss, S. P. 1984. "The Importance of Trophic Transfer in the Bioaccumulation of Chemical Contaminants in Aquatic Ecosystems," *Residues Reviews*, Vol 91, pp 104-130.
- Bookhout, C. G., Costlow, J. D., Jr., and Monroe, R. 1976. "Effects of Methoxychlor on Larval Development of Mud-Crab and Blue Crab," *Water, Air, and Soil Pollution*, Vol 5, pp 349-365.
- Caldwell, R. S. 1974. "Osmotic and Ionic Regulation in Decapod Crustacean Exposed to Methoxychlor," In: *Pollution and Physiology of Marine Organisms*, F. J. Vernberg and W. B. Vernberg (eds.), Academic Press, New York, pp 197-223.
- Couch, J. A., and Courtney, L. 1977. "Interaction of Chemical Pollutants and Virus in a Crustacean: A Novel Bioassay System," *Annals, New York Academy of Science*, Vol 298, pp 497-504.
- Dillon, T. M. 1984. "Biological Consequences of Bioaccumulation in Aquatic Animals: An Assessment of the Current Literature," Technical Report D-84-2, US Army Engineer Waterways Experiment Station, Vicksburg, MS.
- Emanuelson, M., Lincer, J. L., and Rifkin, E. 1978. "The Residue Uptake and Histology of American Oysters (*Crassostrea virginica* Gmelin) Exposed to Dieldrin," *Bulletin of Environmental Contamination and Toxicology*, Vol 19, pp 121-129.
- Engler, R. M. 1980. "Prediction of Pollution Potential through Geochemical and Biological Procedure: Development of Regulation Guidelines and Criteria for the Discharge of Dredged and Fill Material," In: *Contaminants and Sediments*, Vol 1, *Fate and Transport, Case Studies, Modeling Toxicity*, R. A. Balner (ed.), Ann Arbor Science Publishers, Ann Arbor, MI, pp 143-169.
- Fisher, D. J., Bender, M. E., and Roberts, M. H. 1983. "Effects of Ingestion of Kepone Contaminated Food by Juvenile Blue Crabs (*Callinectes sapidus* Rathbun)," *Aquatic Toxicology*, Vol 4, pp 219-234.
- Freeman, H. C., Sangalang, G., and Flemming, B. 1982. "The Sublethal Effects of a Polychlorinated Biphenyl (Aroclor 1254) Diet on the Atlantic Cod (*Gadus morhua*)," *Science Total Environment*, Vol 24, pp 1-11.
- Goodman, L. R., Hansen, D. J., Manning, C. S., and Faas, L. F. 1982. "Effects of Kepone on the Sheepshead Minnow in an Entire Life-Cycle Toxicity Test," *Archives of Environmental Contamination and Toxicology*, Vol 11, pp 335-342.
- Hansen, D. J., Goodman, L. R., and Wilson, A. J., Jr. 1977. Kepone: "Chronic Effects on Embryo, Fry, Juvenile, and Adult Sheepshead Minnows (*Cyprinodon variegatus*)," *Chesapeake Science*, Vol 18, pp 227-232.
- Hansen, D. J., Schimmel, S. C., and Forester, J. 1973. "Aroclor 1254 in Eggs of Sheepshead Minnows: Effect on Fertilization Success and Survival of Embryos and Fry," In: *Proceedings of the Twenty-Seventh Annual Conference Southeastern Association of Game and Fish Commissioners*, October 14-17, 1973, Hot Springs, AR.
- Hansen, D. J., Schimmel, S. C., and Forester, J. 1975. "Effects of Aroclor 1016 on Embryos, Fry, Juveniles, and Adults of Sheepshead Minnows (*Cyprinodon variegatus*)," *Transactions, American Fisheries Society*, Vol 104, pp 584-588.
- Hansen, D. J., Schimmel, S. C., and Forester, J. 1977. "Endrin: Effects on the Entire Life Cycle of a Saltwater Fish, *Cyprinodon variegatus*," *Journal of Toxicology and Environmental Health*, Vol 3, pp 721-733.
- Hansen, P.-D., von Westernhagen, H., and Rosenthal, H. 1985. "Chlorinated Hydrocarbons and Hatching Success in Baltic Herring Spring Spawners," *Marine Environmental Research*, Vol 15, pp 59-76.

- Kay, S. H. 1984. "Potential for Biomagnification of Contaminants within Marine and Freshwater Food Webs," Technical Report D-84-7, US Army Engineer Waterways Experiment Station, Vicksburg, MS.
- Neff, J. M., and Giam, C. S. 1977. "Effects of Aroclor 1016 and Halowax 1099 on Juvenile Horseshoe Crabs *Limulus polyphemus*," In: *Physiological Responses of Marine Biota to Pollutants*, E. J. Vernberg, A. Calabrese, F. P. Thurberg, and W. B. Vernberg (eds.), Academic Press, New York, pp 21-35.
- Neufeld, G. J., and Pritchard, J. B. 1979. "Osmoregulation and Gill Na, K-ATPase in the Rock Crab, *Cancer irroratus*: Response to DDT," *Comparative Biochemistry and Physiology*, Vol 62C, pp 165-172.
- Olofsson, S., and Lindahl, P. E. 1979. "Decreased Fitness of Cod (*Gadus morhua* L.) from Polluted Waters," *Marine Environmental Research*, Vol 2, pp 33-45.
- Peddicord, R. K., and Hansen, J. C. 1983. "Technical Implementation of the Regulations Governing Ocean Disposal of Dredged Material," In: *Wastes in the Ocean*, Vol II, *Dredged Material Disposal in the Ocean*, D. R. Kester, B. H. Ketchum, I. W. Duedall, and P. K. Parks (eds.), John Wiley and Sons, New York, pp 71-88.
- Peddicord, R. K., Lee, C. R., Palermo, M. R., and Francingues, N. R. 1986. "General Decisionmaking Framework for Management of Dredged Material: Example Application to Commencement Bay, Washington," Miscellaneous Paper, US Army Engineer Waterways Experiment Station, Vicksburg, MS.
- Sangalang, G. B., Freeman, H. C., and Crowell, R. 1981. "Testicular Abnormalities in Cod (*Gadus morhua*) Fed Aroclor 1254," *Archives Environmental Contamination and Toxicology*, Vol 10, pp 617-626.
- Schimmel, S. C., Patrick, J. M., Jr., Faas, L. F., Oglesby, J. L., and Wilson, A. J., Jr. 1979. "Kepone: Toxicity and Bioaccumulation in Blue Crabs," *Estuaries*, Vol 2, No. 1, pp 9-15.
- Schimmel, S. C., Patrick, J. M., Jr., and Forester, J. 1977. "Uptake and Toxicity of Toxaphene in Several Estuarine Organisms," *Archives Environmental Contamination and Toxicology*, Vol 5, pp 353-367.
- Tagatz, M. E. 1976. "Effect of Mirex on Predator-prey Interaction in an Experimental Estuarine Ecosystem," *Transactions, American Fisheries Society*, Vol 105, pp 546-549.
- Tagatz, M. E., Borthwick, P. W., Ivey, J. M., and Knight, J. 1976. "Effects of Leached Mirex on Experimental Communities of Estuarine Animals," *Archives Environmental Contamination and Toxicology*, Vol 4, pp 435-442.
- Thomas, P., Carr, R. S., and Neff, J. M. 1981. "Biochemical Stress Responses of Mullet *Mugil cephalus* and Polychaete Worms *Neanthes virens* to Pentachlorophenol," In: *Biological Monitoring of Marine Pollutants*, A. Calabrese, F. P. Thurberg, F. J. Vernberg, and W. B. Vernberg (eds.), Academic Press, New York, pp 73-104.
- US Environmental Protection Agency. 1980. "Water Quality Criteria Documents," Office of Water Regulations and Standards, Criteria and Standards Division, (WH-585), Washington, DC.
- Von Westernhagen, H., Rosenthal, H., Dethlefsen, V., Ernst, W., Harms, U., and Hansen, P.-D. 1981. "Bioaccumulating Substances and Reproductive Success in Baltic Flounder *Platichthys flesus*," *Aquatic Toxicology*, Vol 1, pp 85-99.

Table 1
List of Scientific Journals and Data Base Search Services Used To
Identify Published Papers

Journals

Aquatic Toxicology
Archiv fur Hydrobiologie
Archives of Environmental Contamination and Toxicology
Australian Journal of Biological Science
Australian Journal of Marine and Freshwater Research
Australian Journal of Zoology
Biological Bulletin
Bulletin of Environmental Contamination and Toxicology
California Fish and Game
Canadian Journal of Fisheries and Aquatic Sciences
Canadian Journal of Zoology
Chemosphere
Comparative Biochemistry and Physiology
Critical Reviews in Environmental Control
Crustaceana
Developmental Biology
Ecotoxicology and Environmental Safety
Environmental Biology of Fish
Environmental Pollution Series A, B, and C
Environmental Research
Environmental Science and Technology
Environmental Toxicology and Chemistry
Estuaries
Fisheries
Fisheries Bulletin, U.S.
Hydrobiologia
International Review of Invertebrate Reproduction and Development
International Revue der Gesamten Hydrobiologia
Journal Applied Ecology
Journal of Crustacean Biology
Journal of Experimental Biology
Journal of Experimental Zoology
Journal of Fish Biology
Journal of Invertebrate Pathology
Journal of Pesticide Science
Journal of Plankton Research
Journal of Toxicological and Environmental Research
Journal of Water Pollution Control Federation
Journal of Zoology
Limnology and Oceanography
Marine Environmental Research
New York Fish and Game
New Zealand Journal of Marine and Freshwater Research
Oecologia
Oikos

(Continued)

Table 1 (Concluded)

Journals (Concluded)

Pesticide Biochemistry and Physiology
Pesticides Science
Physiological Zoology
The Progressive Fish Culturist
Quarterly Review of Biology
Science
Science of the Total Environment
Transactions, American Fisheries Society
US Environmental Protection Agency's Ecological Research Series
Water, Air and Soil Pollution
Water Pollution, Research and Control
Water Quality International
Water Research
Water Resources Research

Computerized Data Base Searches Services

Biosis
Water Resources Abstract
Aquatic Sciences and Fisheries Abstract
Chemical Abstracts
Life Sciences Collection
Zoological Record
NTIS (National Technical Information Service)
Dissertation Abstracts
Conference Papers Index
Pollution Abstracts

Table 2
 Summary of Published Papers on the Biological Effects of Chlorinated Environmental
 Contaminants on Marine Organisms and Associated Tissue Residues

| Parameter | Contaminant | Organism | Exposure | | Tissue Concentration** | Biological Effect | Reference |
|-----------|--------------|----------------------|----------|--------------------|------------------------|---|--------------------------------------|
| | | | Time | Concentration* | | | |
| Growth | Kepone | Sheepshead minnow | 36 days | 0.08-6.60 | 1-22 | Growth inversely related to concentration | Hansen, Goodman, and Wilson 1977 |
| Growth | Kepone | Sheepshead minnow | 160 days | 0-0.12 | 0-0.86 | No effect on growth | Goodman et al. 1982 |
| Growth | Kepone | Blue crab | 28 days | 0.39-0.78 | 1.1-5.0 | Reduction in growth | Schimmel et al. 1979 |
| Growth | Kepone | Blue crab | 65 days | 0.15 µg/g food | 0.069 | Reduction in growth | Schimmel et al. 1979 |
| Growth | Kepone | Blue crab | 65 days | 0.36-2.5 µg/g food | 0.38-4.16 | No effect on growth, ratio of carapace thickness to width inversely proportional to concentration | Fisher, Bender, and Roberts 1983 |
| Growth | Methoxychlor | Crab | 10 days | 0.7 | 0.51 | Reduction in growth | Bookout, Costlow, and Monroe 1976 |
| Growth | Endrin | Sheepshead minnows | 22 weeks | 0-1.31 | 0.94 | No effect on growth | Hansen, Schimmel, and Forester 1977 |
| Behavior | Toxaphene | Killifish (embryo) | 28 days | ND†-0.6 | No data | Reduction in survival at all concentrations | Schimmel, Patrick, and Forester 1977 |
| | | | | 1.3-6.5 | No data | Erratic swimming, loss of equilibrium | |
| | | Killifish (fry) | 28 days | ND-0.6 | ND-8.0 | Reduction in survival at all concentrations | |
| | | | | 1.3-6.5 | 34-no data | Reduction in survival at all behavior, loss of equilibrium | |
| | | Killifish (juvenile) | 28 days | ND-0.8 | ND-24.7 | Reduction in survival at all concentrations | |
| | | | | 1.7-3.4 | 102-no data | Erratic swimming, loss of equilibrium | |

(Continued)

* Exposure concentrations are expressed in units of micrograms per litre (µg/l) unless noted otherwise.

** Tissue concentrations are expressed in units of micrograms per gram (µg/g) wet weight whole animals unless noted otherwise.

† ND - Nondetectable, <0.2 µg/l in water, <0.2 µg/g in tissue.

(Sheet 1 of 5)

Table 2 (Continued)

| Parameter | Contaminant | Organism | Exposure | | Tissue Concentration | Biological Effect | Reference |
|--------------|--------------|--------------------|-----------------|---------------------------------------|--|--|--|
| | | | Time | Concentration | | | |
| Behavior | Kepone | Killifish (adults) | 28 days | ND-0.9 1.7-3.8 | ND-6.1 No data | Reduction in survival Erratic swimming, loss of equilibrium | Hansen, Goodman, and Wilson 1977 |
| Behavior | Kepone | Sheepshead minnow | 28 days | 0-1.9 | 0.26-11 | Erratic swimming behavior and a reduction in feeding rate both increased as concentrations increased | Fisher, Bender, and Roberts 1983 |
| Behavior | Kepone | Blue crab | 65 days | 0.36-1.64 μ g/g food | 0.38-1.73 | No effect on behavior | Fisher, Bender, and Roberts 1983 |
| Behavior | Methoxychlor | Crab | 15 days | 1.8-32 | 0.11-1.59 | Excitable behavior during feeding, reduced ability to locate and consume food | Armstrong et al. 1976 |
| Behavior | Mirex | Grass shrimp | 14-16 days | 0.011-0.130 | 0.02-0.20 | Diminished ability to avoid predation | Tagatz 1976 |
| Behavior | Mirex | Oyster | 10 weeks | 0.038 | 1.3-28 | Diminished ability to withstand predation | Tagatz et al. 1976 |
| Behavior | PCB DDT | Fish | Field collected | No data | 110 7 | Diminished ability to withstand predation Decreased ability to maintain position while swimming in a current | Oloffson and Lindahl 1979 |
| Reproduction | Kepone | Sheepshead minnow | 90-133 days | 0.041-0.074 | 0.15-0.56 | Increased number of eggs/female/day; fertility unaffected | Goodman et al. 1982 |
| Reproduction | Kepone | Sheepshead minnow | 28 days | 0.12-0.39 0.78 0.05-0.80 1.9 | 0.86-3.0 5.0-6.8 0.26-4.7 11 (Continued) | Number of eggs/female/day unaffected; fertility unaffected Decreased number of eggs/female/day, reduced fertility Production of normal embryos Production of abnormal embryos | Hansen, Goodman, and Wilson 1977 (Sheet 2 of 5) |

Table 2 (Continued)

| Parameter | Contaminant | Organism | Exposure | | Tissue Concentration | Biological Effect | Reference |
|--------------|--------------------|-------------------|------------------------|--------------------|---|---|--|
| | | | Time | Concentration | | | |
| Reproduction | PCB (Aroclor 1254) | Cod | 5-1/2 months | 1-50 µg/g wet food | 0.06-5.3 (testes) | Disruption in production of sex steroids from testes | Freeman, Sangalang, and Flemming 1982 |
| Reproduction | PCB (Aroclor 1254) | Sheepshead minnow | 4 weeks | 0-10.0 | 0.52-170 | No effect on number of eggs fertilized | Hansen, Schimmel, and Forester 1973 |
| Reproduction | PCB (Aroclor 1016) | Sheepshead minnow | 29 days | 1-10 | 5.4-110 (adults) 4.2-66 (eggs) | No effect on egg fertility, hatching, or subsequent survival of progeny | Hansen, Schimmel, and Forester 1975 |
| Reproduction | Endrin | Sheepshead minnow | 23 weeks, 1 generation | 0.027-0.12 | 200-1,100 (adults) 0.20-1.0 (adults) 0.09-0.87 (eggs) | 100 percent mortality in adults No effect on reproduction | Hansen, Schimmel, and Forester 1977 |
| Reproduction | PCB | Flounder | Field collected | No data | 0.94 (adults) 1.80 (eggs) | Reduced fertilization and early hatching, high mortalities | |
| | | | | 0.72 | No data | | |
| Reproduction | PCB | Fish | Field collected | No data | 5.0-317 ng/g (ovaries) | Reduced viable hatch at PCB tissue concentrations above 120 ng/g | Von Westernhagen et al. 1981 |
| | DDD | | | | 3.0-30.3 ng/g (ovaries) | Hatch viability not correlated with tissue concentration of any other contaminant | |
| | DDE | | | | 0.1-62.0 (ovaries) | | |
| | Hexachlorobenzene | | | | 0.06-2.0 (ovaries) | | |
| | Dieldrin | | | | 0.1-49.0 (ovaries) | | |
| | Heptachlorepoxyde | | | | 0.08-3.0 (ovaries) | | |
| Reproduction | PCB | Fish | Field collected | No data | 19-241 ng/g (ovaries) | Reduced viable hatch at PCB tissue concentrations above 18 ng/g | Hansen, Von Westernhagen, and Rosenthal 1985 |
| | DDD | | | | <1-16.0 (ovaries) | Hatch viability not correlated with tissue concentration of any other contaminant | |
| | DDE | | | | <1-34.0 (ovaries) | | |
| | Dieldrin | | | | <1-8.1 (ovaries) | | |
| | Hexachlorobenzene | | | | <1-8.6 (ovaries) | | |
| | Heptachlorepoxyde | | | | <1-8.9 (ovaries) | | |

(Continued)

(Sheet 3 of 5)

Table 2 (Continued)

| Parameter | Contaminant | Organism | Time | Exposure Concentration | Tissue Concentration | Biological Effect | Reference |
|------------|--|-----------|--------------|--------------------------|---|--|---------------------------------------|
| | α Hexachlorocyclohexane γ Hexachlorocyclohexane | | | | <1-9.2 (ovaries) <1-12.1 (ovaries) | | |
| Morphology | PCB (Aroclor 1254) | Cod | 5-1/2 months | 1-50 μ g/g wet food | 0.04-2.1 (kidney) 0.02-0.98 (muscle) 10.1-374 (liver) | Disruption in production of adrenal hormones from kidney No effect on histopathology of kidney Degeneration of liver's fatty tissue | Freeman, Sangalang, and Flemming 1982 |
| Morphology | PCB (Aroclor 1254) | Cod | 5-1/2 months | 1-50 μ g/g food | 0.06-5.3 (testes) | Response intensified as tissue concentration increased and progressed from testicular fibrosis to inhibition of spermatogenesis and finally to complete disintegration of the testes | Sangalang, Freeman, and Crowell 1981 |
| Morphology | PCB (Aroclor 1254) | Cod | 5-1/2 months | 1-50 μ g/g wet food | 0.02-0.98 (muscle) | Hyperplasia of gills with disrupted blood spaces | Freeman, Sangalang, and Flemming 1982 |
| Morphology | PCB (Aroclor 1254) | Shrimp | 35 days | 0.6-0.7 | 2 (muscle) 21 (hepatopancreas) | Increased occurrence of viral pathogen | Couch and Courtney 1977 |
| Morphology | Kepone | Crab | 65 days | 0.36-2.50 μ g/g food | 0.38-4.61 | Carapace thickness-to-width ratio inversely related to concentration | Fisher, Bender, and Roberts 1983 |
| Morphology | Kepone | Killifish | 28 days | 0-1.9 | 0.26-11 | Response intensified as tissue concentration increased. Response progressed from deformed vertebral column, hemorrhaging near brain, and darkened posterior to increased hemorrhaging and fin rot | Hansen, Goodman, and Wilson 1977 |
| Morphology | Dieldrin | Oyster | 43 days | 1-100 | 25.6-2,685* | No effect on fibrous or cellular components of gills, gut, or mantle, no inflammation or infiltration of leucocytes | Emanuelson, Lincer, and Rifkin 1978 |

(Continued)

* Data originally reported on a dry-weight basis were converted to wet weight assuming 80 percent body water.

Table 2 (Concluded)

| Parameter | Contaminant | Organism | Exposure | | Tissue Concentration | Biological Effect | Reference |
|----------------|--|----------------|----------|-------------------------------|-------------------------------------|--|----------------------------------|
| | | | Time | Concentration | | | |
| Osmoregulation | Pentachlorophenol | Fish | 5 days | 100 | 37.1 | Reduction in total osmotic pressure | Thomas, Carr, and Neff 1981 |
| Osmoregulation | Methoxychlor | Crab | 7 days | 10 | 0.31 2.0 (gill) | No effect on total osmotic pressure | Caldwell 1974 |
| Osmoregulation | DDT | Crab | 14 days | 10 | 1.0 2.5 (gill) | No effect on total osmotic pressure, sodium or potassium regulation but magnesium regulation was disrupted | |
| Osmoregulation | DDT | Crab | 50 hr | Single injection of 100 µg/kg | 0.06 (gill) 1.5 (hepatopancreas) | Sodium and potassium regulation in the gill disrupted | Neufeld and Pritchard 1979 |
| Metabolism | PCB (Aroclor 1016) | Horseshoe crab | 96 days | 0.35-71.5 | 0.08-92.8 | No ecologically significant change in oxygen consumption | Neff and Giam 1977 |
| Metabolism | Halowax 1099 (chlorinated naphthalene) | Horseshoe crab | 96 days | 22-70 | 0.51-5.7 | Highly variable oxygen consumption | Neff and Giam 1977 |
| Metabolism | Kepone | Blue crab | 65 days | 0.36-1.64 µg/g food | 0.38-1.73 | No effect on oxygen consumption | Fisher, Bender, and Roberts 1983 |
| | | | | 2.26-2.50 µg/g food | 2.54 | Elevated rates of oxygen consumption | |