



# **Head, Face, and Neck Injuries During Operation Iraqi Freedom II: Results From the US Navy and Marine Corps Combat Trauma Registry**

*Amber L. Wade, MPH*

*Judy L. Dye, MS*

*Charlene R. Mohrle, MS*

*Michael R. Galarneau, MS*



***Naval Health Research Center***

---

***Technical Report 06-01***

*Approved for public release: distribution unlimited*

*Naval Health Research Center  
P.O. BOX 85122  
San Diego, California 92186-5122*

**Head, Face, and Neck Injuries During Operation Iraqi Freedom II:  
Results From the US Navy and Marine Corps Combat Trauma Registry**

Amber L. Wade, MPH<sup>1</sup>  
Judy L. Dye, MS<sup>1</sup>  
Charlene R. Mohrle, MS<sup>1</sup>  
Michael R. Galarneau, MS<sup>2</sup>

<sup>1</sup>Science Applications International Corporation  
10260 Campus Point Drive  
San Diego, CA 92121

<sup>2</sup>Naval Health Research Center  
P.O. Box 85122  
San Diego, CA 92186-5122

Report No. 06-01 supported by the Office of Naval Research, Arlington, VA, under Work Unit No. 63706N-M0095.005-60332. The views expressed in this article are those of the authors and do not necessarily reflect the official policy or position of the Department of the Navy, Department of Defense, nor the U.S. Government. Approved for public release; distribution is unlimited. This research has been conducted in compliance with all applicable federal regulations governing the protection of human subjects in research.

**Table of Contents**

Abstract..... 4

Introduction..... 5

Methods..... 6

Results..... 8

Discussion..... 13

References..... 16

### **Abstract**

**Background:** This study assessed characteristics of head, face, and neck injury among US military casualties from the stability and support phase of Operation Iraqi Freedom.

**Methods:** A retrospective analysis of trauma registry data was performed. Data were collected from the US Navy and Marine Corps Combat Trauma Registry for the time period of 1 March to 30 September 2004.

**Results:** Four hundred and forty-five casualties with head, face, or neck injury were identified for analysis. Of these, 140 (31%) sustained multiple wounds to the head, face, and neck; 177 (40%) had head wounds; 336 (76%) had facial wounds; and 84 (19%) had neck wounds. Head, face, and neck wounds accounted for 61% of the total 1,666 injuries incurred. Improvised explosive devices were the most frequent cause of injury among battle casualties, whereas motor vehicle crashes were the most common cause of injury among nonbattle casualties. The overall mean abbreviated injury severity scores for head, face, and neck injuries were 2.1, 1.1, and 1.2, respectively.

**Conclusions:** Although the majority of casualties incurred wounds to the face, head and neck injuries were generally more severe, especially among those who died of wounds. The high frequency of facial wounding suggests, however, that the face may be more vulnerable during armed conflict than the head or neck.

**Key Words:** combat, casualty, head injury, Operation Iraqi Freedom

## Introduction

Traumatic injuries to the head, face, and neck have been a major contributor to the mortality and morbidity of US armed forces injured during hostile action.<sup>1-7</sup> Although the head, face, and neck comprise only 12% of the total body surface area exposed during combat,<sup>1</sup> these body areas sustained proportionately more injuries and proportionately more injuries leading to death during the major conflicts of the 20th century. During World War II, the Korean War, and Vietnam War, head, face, and neck wounds accounted for more than 40% of deaths (more than any other body region) among US armed forces.<sup>2-4</sup> The mean relative incidence of head, face, and neck injury was approximately 16% based on a meta-analysis of major armed conflicts from 1914 to 1986.<sup>8</sup> This pattern remains evident in current warfare. During the major combat phase of Operation Iraqi Freedom (OIF-I), US Navy and Marine Corps forces wounded in action sustained approximately 19% of all wounds to the head, face, or neck.<sup>9</sup>

Despite the prevalence of head, face, and neck injuries during war, few studies have addressed the severity and types of these injuries. An analysis of wounds to the head, face, and neck would facilitate the determination of resource needs of future operations as well as potential improvements to modern individual body armor with respect to the head, face, and neck. The present study provides such an assessment of head, face, and neck injuries sustained by US military forces during the stability and support phase of Operation Iraqi Freedom (OIF-II) from March to September 2004.

This paper also identifies the primary causes of battle and nonbattle injury to the head, face, and neck during OIF-II. Research on previous conflicts<sup>6,10</sup> has found that most injuries sustained during hostile action are due to small arms and explosive munitions. During OIF-I, explosive munitions such as rocket-propelled grenades accounted for the majority of head, face,

and neck wounds sustained in action.<sup>9</sup> Conversely, researchers<sup>2,10</sup> reported motor vehicle crashes and falls as the primary causes of nonbattle injuries during armed conflict. This study will examine whether the same causes of injury were observed in OIF-II.

### **Methods**

Data were obtained from the US Navy and Marine Corps Combat Trauma Registry (Navy–Marine Corps CTR) in a retrospective review of head, face, and neck injuries incurred during the time period from 1 March to 30 September 2004 of OIF-II. The Navy–Marine Corps CTR is a data warehouse maintained by the Naval Health Research Center (NHRC), San Diego, CA, to collect and store personal health information on US service members who receive initial care for battle or nonbattle injuries at Navy–Marine Corps military medical treatment facilities (MTFs). Examples of Navy–Marine Corps MTFs include battalion aids stations (ie, Level I), which provide initial resuscitative intervention; and Shock Trauma Platoons, Forward Resuscitative Surgical Systems, and Surgical Companies, which provide advanced resuscitative intervention and damage control surgical procedures (ie, Level II). Information regarding patient demographics, incident details, injuries, treatment, and outcomes is collected by the patient's medical provider(s) in the form of a clinical record. Clinical records are then sent to NHRC for data management and analysis. The Navy–Marine Corps CTR clinical research personnel compile the medical records of each patient and evaluate the severity of these patients' injuries.

In the present study, the clinicians identified patients with head, face, and neck injury for analysis. Head injuries included trauma to the skull and brain; facial injuries included trauma to the facial bones, eyes, ears, nose, and mouth; and neck injuries included trauma to cervical area and cervical spine.

Descriptive measures included casualty types, causes of injury, injuries, and injury severity scores. Casualties were divided into two main categories: battle injury, defined as those who suffered an injury to the head, face, or neck as a direct result of hostile action; and nonbattle injury, defined as those who sustained an injury to the head, face, or neck that was not due to hostile action. Battle injury casualties were further defined as died of wounds (ie, those who died from wounds received in action following treatment at an MTF); or wounded in action (ie, those who survived wounds sustained in action). Wounded in action and nonbattle injury casualties were analyzed as a function of their disposition: returned to duty from a Level I or Level II MTF; or medically evacuated to a Level III MTF (ie, Combat Support Hospital) or higher level of care. Each of the above-mentioned categories were mutually exclusive.

Injury severity was assessed with the Abbreviated Injury Scale (AIS) 2005<sup>12</sup> and the Injury Severity Score (ISS).<sup>13</sup> The AIS provided anatomically specific severity scores for each injury (ie, head, face, or neck injury), whereas the ISS was used to assess the overall injury severity of each patient, which included injuries from all anatomical locations. Analyses were also performed for protective equipment worn at the time of injury, disposition, anatomical location of injury, *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM)<sup>14</sup> codes, and the Glasgow Coma Scale (GCS).

Differences in AIS, ISS, and GCS scores were compared between medically evacuated and returned to duty populations among battle and nonbattle injury casualties. The 2 sample *t* test for independent samples and Mann-Whitney *U* test were used, as appropriate. All statistical analyses were performed using SPSS software version 12.0.2 (SPSS Inc., Chicago, IL); all tests were 2-tailed and  $P \leq .05$  was used to determine statistical significance. Mean values are presented as means  $\pm$  SD unless otherwise indicated.

## Results

Four hundred and forty-five US military casualties with head, face, or neck injuries were identified for analysis, which accounted for 39% of all injury casualties in the Navy-Marine Corps CTR from 1 March to 30 September 2004. Of the 445 casualties, four Marine Corps incurred two separate injury events during the study period; each injury event was counted as 1 casualty. Average age (with 69 unknown) was  $24 \pm 5.9$  years, with a range of 18 to 48 years. All but 6 casualties were male. Of the 445 casualties, 381 were Marine Corps, 45 were Army, 14 were Navy, and 1 was Air Force. Military service category was unknown for 4 casualties.

Among the 445 head, face, and neck injury casualties, 19 (4%) were classified as died of wounds, 315 (71%) were wounded in action, and 111 (25%) sustained nonbattle injuries. Of the wounded in action, 144 (46%) were medically evacuated to a Level III MTF or higher level of care, and 171 (54%) of the casualties were returned to duty from a Level I or Level II MTF. Of the nonbattle injury casualties, 26 (23%) were medically evacuated and 85 (77%) were returned to duty.

As shown in Table 1, the most common causes of head, face, and neck injury among battle casualties were improvised explosive devices (64%), mortars (11%), gunshot wounds or bullets (7%), and other explosives (6%).

Table 1. Causes of Battle Injury ( $n = 334$ )

| Cause of Injury             | Casualties |
|-----------------------------|------------|
|                             | No. (%)    |
| Improvised explosive device | 214 (64)   |
| Mortar                      | 36 (11)    |
| Gunshot wound/bullet        | 22 (7)     |
| Explosive (NOS)             | 21 (6)     |
| Rocket-propelled grenade    | 14 (4)     |
| Fragmentation (NOS)         | 13 (4)     |
| Landmine                    | 8 (2)      |
| Aircraft crash              | 4 (1)      |
| Grenade                     | 1 (<1)     |
| Unknown                     | 1 (<1)     |

NOS, not otherwise specified.

As shown in Table 2, nonbattle injury casualties sustained head, face, and neck injury from primarily vehicle accidents (31%), unspecified blunt trauma (19%), sports recreation and training (14%), flying debris (11%), and falls (10%).

Table 2. Causes of Nonbattle Injury (*n* = 111)

| Cause of Injury            | Casualties |      |
|----------------------------|------------|------|
|                            | No.        | (%)  |
| Motor vehicle crash        | 34         | (31) |
| Blunt trauma (NOS)         | 21         | (19) |
| Sports recreation/training | 15         | (14) |
| Flying debris              | 12         | (11) |
| Fall                       | 11         | (10) |
| Assault/altercation        | 8          | (7)  |
| Crush                      | 3          | (3)  |
| Explosive (NOS)            | 3          | (3)  |
| Hot object/liquid          | 3          | (3)  |
| Knife/edge                 | 1          | (1)  |

NOS, not otherwise specified.

More than 80% of battle casualties (279/334; with 21 unknown) were reported wearing helmets at the point of injury. Only 68% (228/334; with 52 unknown), however, were reported wearing protective eyewear at the point of injury. As shown in Table 3, casualties reported not wearing helmets were injured primarily by mortars (indirect fire), whereas casualties reported not wearing eyewear were injured primarily by improvised explosive devices (direct fire).

Table 3. Protective Equipment Reported as Not Worn, by Cause of Injury\*

| Cause of Injury             | Equipment Not Worn         |                             |
|-----------------------------|----------------------------|-----------------------------|
|                             | Helmet<br>( <i>n</i> = 21) | Eyewear<br>( <i>n</i> = 52) |
| Mortar                      | 13 (62)                    | 16 (31)                     |
| Improvised explosive device | 4 (19)                     | 22 (42)                     |
| Explosive (NOS)             | 2 (10)                     | 4 (8)                       |
| Fragmentation (NOS)         | 1 (5)                      | 4 (8)                       |
| Rocket-propelled grenade    | 1 (5)                      | 2 (4)                       |
| Gunshot wound/bullet        | 0                          | 2 (4)                       |
| Landmine                    | 0                          | 2 (4)                       |

NOS, not otherwise specified.

\*Values are number (percentage). Because of rounding, percentages may not total 100.

Overall, casualties incurred a total of 1666 injuries. Of these, 61% (1011) were injuries to the head, face, or neck. Of the 445 casualties, 140 (31%) sustained multiple wounds to the head, face, and neck; 177 casualties (40%) had head wounds; 336 (76%) casualties had facial wounds; and 84 (19%) had neck wounds. As shown in Table 4, the most frequent location of injury among those who died of wounds was the head. Among the wounded in action and nonbattle injury casualties, the most frequent location of injury was the face.

Table 4. Distribution of Anatomical Location of Injuries by Disposition

| Injury Location | No. (%) of Injuries* |                               |                      |                  |                     |                  |
|-----------------|----------------------|-------------------------------|----------------------|------------------|---------------------|------------------|
|                 | All<br>(N = 1011)    | Died of<br>Wounds<br>(n = 60) | Wounded in Action    |                  | Nonbattle           |                  |
|                 |                      |                               | MEDEVAC<br>(n = 478) | RTD<br>(n = 296) | MEDEVAC<br>(n = 47) | RTD<br>(n = 128) |
| Head            | 259 (26)             | 29 (48)                       | 115 (24)             | 68 (23)          | 13 (28)             | 34 (26)          |
| Face            | 653 (65)             | 20 (33)                       | 319 (67)             | 201 (68)         | 32 (68)             | 81 (63)          |
| Neck            | 99 (10)              | 11 (18)                       | 44 (9)               | 27 (9)           | 2 (4)               | 15 (12)          |

MEDEVAC, medically evacuated; RTD, returned to duty.

\*Because of rounding, percentages may not all total 100.

The distribution of specific injury descriptions and injury codes, as provided by ICD-9-CM, are presented in Table 5. Approximately 45% (450/1011) of all head, face, and neck injuries were coded as open wounds (ICD-9-CM codes 870 to 897). “Intracranial injury” was the most common diagnosis among injuries specific to the head. The most common facial injury diagnosis was “other open wounds of the head,” which included only those ICD-9-CM codes anatomically specific to the face (873 to 873.9). Notably, there were 156 injuries of the eye and ocular adnexa. “Open wound of the neck” was the most common neck injury.

Table 5. Distribution of Injuries and ICD-9-CM Codes by Anatomical Region by Disposition

| Injury Description                            | ICD-9-CM Code(s) | No. (%) of Injuries* |                               |                      |                  |                     |                  |
|---|------------------|----------------------|-------------------------------|----------------------|------------------|---------------------|------------------|
|   |                  | All<br>(N = 1011)    | Died of<br>Wounds<br>(n = 60) | Wounded in Action    |                  | Nonbattle           |                  |
|   |                  |                      |                               | MEDEVAC<br>(n = 478) | RTD<br>(n = 296) | MEDEVAC<br>(n = 47) | RTD<br>(n = 128) |
| <b>Head</b>                                   |                  |                      |                               |                      |                  |                     |                  |
| Intracranial injury, excluding skull fracture | 850-854          | 128 (13)             | 18 (30)                       | 56 (12)              | 32 (11)          | 7 (15)              | 15 (12)          |
| Other open wound of head                      | 873              | 52 (5)               | 4                             | 21                   | 14               | 1                   | 12               |
| Fracture of base and vault of skull           | 800,801          | 33 (3)               | 6 (10)                        | 24 (5)               | 2 (<1)           | 1 (2)               | 0                |
| Injury to other cranial nerve(s)              | 951              | 26 (3)               | 1 (2)                         | 7 (2)                | 18 (6)           | 0                   | 0                |
| Other conditions of the brain                 | 348              | 1 (<1)               | 1 (2)                         | 0                    | 0                | 0                   | 0                |
| <b>Face</b>                                   |                  |                      |                               |                      |                  |                     |                  |
| Other open wound of head                      | 873              | 184 (18)             | 7 (12)                        | 90 (19)              | 60 (20)          | 6 (13)              | 21 (16)          |
| Open wound of ear                             | 872              | 94 (9)               | 2 (3)                         | 40 (8)               | 49 (17)          | 1 (2)               | 2 (2)            |
| Fracture of face bones                        | 802              | 65 (6)               | 1 (2)                         | 57 (12)              | 0                | 5 (11)              | 0                |
| Open wound of ocular adnexa and eyeball       | 870,871          | 64 (6)               | 4 (7)                         | 47 (10)              | 3 (1)            | 6 (13)              | 4 (3)            |
| Superficial injury of the eye and adnexa      | 918              | 39 (4)               | 0                             | 24 (5)               | 5 (2)            | 3 (6)               | 7 (6)            |
| Contusion of eye and adnexa                   | 921              | 36 (4)               | 3 (5)                         | 22 (5)               | 2 (1)            | 6 (13)              | 3 (2)            |
| Foreign body on external eye                  | 930              | 5 (<1)               | 0                             | 0                    | 0                | 0                   | 5 (4)            |
| Burn confined to eye and adnexa               | 940              | 5 (<1)               | 1 (2)                         | 1 (<1)               | 0                | 0                   | 3 (2)            |
| Injury to optic nerve and pathways            | 950              | 3 (<1)               | 0                             | 2 (<1)               | 1 (<1)           | 0                   | 0                |
| Other disorders of the eye                    | 379              | 3 (<1)               | 0                             | 3 (1)                | 0                | 0                   | 0                |
| <b>Neck</b>                                   |                  |                      |                               |                      |                  |                     |                  |
| Open wound of neck                            | 874              | 56 (6)               | 7 (12)                        | 30 (6)               | 19 (6)           | 0                   | 0                |
| Sprains and strains of neck                   | 847.0            | 22 (2)               | 0                             | 2 (<1)               | 5 (2)            | 1 (2)               | 14 (11)          |
| Fracture of vertebral column                  | 805,806          | 6 (1)                | 0                             | 6 (1)                | 0                | 0                   | 0                |
| Other, multiple, and ill-defined dislocations | 839.0            | 1 (<1)               | 0                             | 1 (<1)               | 0                | 0                   | 0                |
| <b>Other, multiple, and ill-defined</b>       |                  |                      |                               |                      |                  |                     |                  |
| Superficial injury of face, neck, and scalp   | 910              | 61 (6)               | 0                             | 12 (3)               | 26 (9)           | 4 (9)               | 19 (15)          |
| Injury, other and unspecified                 | 959.0            | 54 (5)               | 0                             | 3 (1)                | 48 (16)          | 1 (2)               | 2 (2)            |
| Burn of face, head, and neck                  | 941              | 35 (3)               | 2 (3)                         | 16 (3)               | 5 (2)            | 2 (4)               | 10 (8)           |
| Contusion of face, scalp, and neck            | 920              | 27 (3)               | 0                             | 9 (2)                | 6 (2)            | 3 (6)               | 9 (7)            |
| Injury to blood vessels of head and neck      | 900              | 7 (1)                | 3 (5)                         | 4 (1)                | 0                | 0                   | 0                |
| Burns according to extent of body surface     | 948              | 4 (<1)               | 0                             | 1 (<1)               | 1                | 0                   | 2 (2)            |

ICD-9-CM, *International Classification of Diseases, 9th Revision, Clinical Modification*<sup>14</sup>; MEDEVAC, medically evacuated; RTD, returned to duty.

\*Because of rounding, percentages may not all total 100.

A higher AIS score, on a scale of 1 to 6, indicates more severe injury. The overall mean AIS scores (with 13 unknown) were  $2.1 \pm 1.3$  for the head,  $1.1 \pm 0.3$  for the face, and  $1.2 \pm 0.7$  for the neck. Comparisons of injury severity by AIS body systems across dispositions are shown in Table 6. Among casualties wounded in action, the mean AIS scores for head, face, and neck body systems were significantly higher among medically evacuated patients than for those who were returned to duty. There were no statistical differences for mean AIS scores between medically evacuated and returned to duty nonbattle injuries.

Table 6. Comparison of Abbreviated Injury Scale (AIS) 2005<sup>12</sup> for Injuries by Body System Across Dispositions\*

| AIS Body System | Died of Wounds | Wounded in Action |           |          | Nonbattle |           |            |
|-----------------|----------------|-------------------|-----------|----------|-----------|-----------|------------|
|                 |                | MEDEVAC           | RTD       | <i>P</i> | MEDEVAC   | RTD       | <i>P</i> † |
| Head            | 4.1 (1.7)      | 2.6 (1.3)         | 1.4 (0.6) | <.001    | 1.3 (0.7) | 1.3 (0.4) | .72        |
| Face            | 1.1 (0.2)      | 1.2 (0.4)         | 1.0 (0.1) | <.001    | 1.1 (0.3) | 1.0 (0.2) | .19        |
| Neck‡           | 2.0 (1.3)      | 1.3 (0.8)         | 1.0 (0.0) | .02      | 1.0 (NA)  | 1.0 (0.0) | ...        |

MEDEVAC, medically evacuated; RTD, returned to duty.

\*Values are mean (SD) unless otherwise indicated. A score of 1 is considered minor injury; the maximum score is 5 (critical injury).

†Ellipses indicate *P* value not computed.

‡Includes spine (cervical) AIS.

The ISS takes values from 1 to 75. A higher ISS score indicates more severe injuries. Of the 445 casualties, 373 (84%) had an ISS of 0 to 8; 30 (7%) had an ISS of 9 to 15; 14 (3%) had an ISS of 16 to 24; 24 (5%) had an ISS of 25 or greater; and 4 (1%) were unable to be coded. The median ISS for casualties who died of wounds was 25 (range, 1–57). Among wounded in action, the median ISS was higher for the medically evacuated compared with the returned to duty casualties (5 vs. 2,  $P < .001$ ). Similarly, for nonbattle injury casualties, the median ISS was higher among the medically evacuated than the returned to duty (2 vs. 1,  $P = .01$ ).

Of the 409 casualties who had a documented GCS score on admission to a Navy–Marine Corps MTF, the majority (91%) had minor or no evidence of head injury (GCS 13 to 15). The mean GCS score on admission among those who later died of wounds was  $5.2 \pm 3.6$ , indicating

severe head injury. Among the wounded in action, the mean GCS score was lower for medically evacuated than returned to duty casualties ( $14.2 \pm 2.3$  vs.  $15.0 \pm 0.2$ ;  $P < .001$ ). Among nonbattle injury casualties, the mean GCS score was also lower for medically evacuated than returned to duty, but not statistically different ( $14.2 \pm 2.0$  vs.  $15.0 \pm 0.4$ ,  $P = .11$ ). The effect size for the combination of these groups, however, was small ( $h^2 = .00$ ,  $P = .80$ , 95% confidence interval =  $-0.89, 0.69$ ). Thus, there is no evidence that the effect of disposition (ie, medically evacuated or returned to duty) on GCS score differs between wounded in action and nonbattle injury casualties.

### **Discussion**

Consistent with research on OIF-I,<sup>9</sup> the primary causes of head, face, and neck battle injuries in the present study were explosive munitions such as improvised explosive devices and mortars. Although more than 80% of battle casualties wore helmets, only 68% were reported wearing protective eyewear at the point of injury. Mortars were the primary cause of injury among those not wearing helmets and improvised explosive devices were the primary cause of injury among those not wearing protective eyewear. Speculatively, casualties who were not wearing their helmets at the point of injury may have been injured more frequently by indirect fire, such as mortars, rather than direct fire because they perceived themselves to be in an environment safe from hostile action.

Motor vehicle crashes were the most common cause of injury among nonbattle casualties. These findings are consistent with previous research from OIF-I indicating that motor vehicle crashes cause the majority of nonbattle injuries during hostile action,<sup>11</sup> and result in the highest number of nonbattle injury admissions.<sup>7</sup> These prior assessments also indicate that falls have been a notable cause of nonbattle injuries and nonbattle injury admissions, which included

injuries from all anatomical locations. In our analysis falls were not a major contributor to nonbattle head, face, and neck injury suggesting that these regions may have been less susceptible to injury from a fall than other anatomical locations during OIF-II.

Wounding patterns among battle injury deaths are known to differ from that of battle injury survivors.<sup>15</sup> In the present study, the face was the predominant site of injury among the wounded survivors. The head, however, was the predominant site of injury among casualties who died of wounds. More than one third of all casualties presented with multiple injuries to the head, face, or neck. Consistent with research based on prior conflicts,<sup>2,3,9,11</sup> open (or penetrating) wounds were altogether the most common injury diagnosis.

The severity of specific injuries (ie, AIS) and overall severity scores (ie, ISS) were expectedly higher for battle casualties who were medically evacuated compared with those returned to duty given that medical evacuation is protocol for the more severely injured patients. Similarly, among nonbattle injury casualties, ISS was higher for medically evacuated than returned to duty. Although the AIS scores were generally higher among the medically evacuated than returned to duty nonbattle casualties, they were not statistically different.

There have been relatively few descriptions of head, face, and neck injuries incurred during hostile action. The present results extend the existing literature in several ways. This study assessed the types of head, face, and neck injuries as well as the corresponding severity levels of each injury. The Navy–Marine Corps CTR is the only current US trauma registry to comprehensively collect injury-related personal health information at Level I and Level II MTFs. Furthermore, this sample from the CTR allowed us to analyze injuries among casualties who were returned to duty from the Level I and Level II MTFs as well as describe the frequency of protective equipment worn at the time of injury.

While this broad-based analysis provides important information in the assessment of the types of injuries and may aid in future military medical planning, it also presents a few limitations including the retrospective nature of the data, missing data, and the lack of denominator data, which precludes risk assessments. Although each of the US Armed Forces was represented in this examination, the large majority of casualties were Marines. Due to inherent differences in force operations and body armor, these data may not generalize to other military forces.

Despite these limitations, our results do indicate that casualty medical care resource allocation and planning should take into consideration the impact of facial injuries, especially with regard to improvements to modern individual body armor. Further research is required to identify the risks associated with head, face, and neck injury during hostile action. Outcome assessments are also needed in order to define the implications of survived head, face, and neck injury.

## References

1. Carey ME. Learning from traditional combat mortality and morbidity data used in the evaluation of combat medical care. *Mil Med.* 1987;152:6-13.
2. Reister FA. *Medical Statistics in World War II.* Washington, DC: Department of the Army, Office of the Surgeon General;1975:21.
3. Reister FA. *Battle Casualties and Medical Statistics: US Army Experience in the Korean War.* Washington, DC: Department of the Army, Office of the Surgeon General; 1973. Chapter 3.
4. Maughon JS. An inquiry into the nature of wounds resulting in killed in action in Vietnam. *Mil Med.* 1970;135:8-13.
5. Carey ME. Analysis of wounds incurred by U.S. Army Seventh Corps personnel treated in Corps hospitals during Operation Desert Storm, February 20 to March 10, 1991. *J Trauma.* 1996;40(suppl):S165-S169.
6. Mabry RL, Holcomb JB, Baker AM, et al. United States Army Rangers in Somalia: an analysis of combat casualties on an urban battlefield. *J Trauma.* 2000;49:515-529.
7. Montgomery SP, Swiecki CW, Shriver CD. The evaluation of casualties from Operation Iraqi Freedom on return to the continental United States from March to June 2003. *J Am Coll Surg.* 2005;201:7-13.
8. Dobson JE, Newell MJ, Shepherd JP. Trends in maxillofacial injuries in war-time (1914-1986). *Br J Oral Maxillofac Surg.* 1989;27:441-450.
9. Zouris JM, Walker GJ, Dye J, et al. Wounding patterns of United States Marines and Sailors during Operation Iraqi Freedom: major combat phase. *Mil Med.* In press.

10. Leedham CS, Blood CG, Newland C. A descriptive analysis of wounds among U.S. Marines treated at second-echelon facilities in the Kuwaiti theater of operations. *Mil Med.* 1993;158:508-512.
11. Galarneau MR, Hancock WC, Konoske P, et al. *U.S. Navy-Marine Corps Combat Trauma Registry Operation Iraqi Freedom-1: Preliminary Findings.* San Diego, CA: Naval Health Research Center; 2004. NHRC Report No. 04-20.
12. Association for the Advancement of Automotive Medicine: The Abbreviated Injury Scale (AIS) 2005. Barrington, IL.
13. Baker SP, O'Neill B, Haddon W, et al. The Injury Severity Score: a method for describing patients with multiple injuries and evaluating emergency care. *J Trauma.* 1974;14:187-196.
14. Commission on Professional Hospital Activities. *International Classification of Diseases, 9th Revision, Clinical Modification.* Ann Arbor, MI: Edwards Brothers; 1977.
15. Champion HF, Bellamy RF, Roberts CP, et al. A profile of combat injury. *J Trauma.* 2003;54(suppl):S13-S19.

# REPORT DOCUMENTATION PAGE

The public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing the burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB Control number. **PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.**

|  |   |  |
|--|---|--|
| <b>1. Report Date (DD MM YY)</b><br>12-01-2006 | <b>2. Report Type</b><br>Technical Report | <b>3. DATES COVERED (from - to)</b><br>Mar 2004 – Jan 2006 |
|--|---|--|

|   |   |
|---|---|
| <b>4. TITLE AND SUBTITLE</b><br>Head, Face, and Neck Injuries during Operation Iraqi Freedom II: Results from the US Navy and Marine Corps Combat Trauma Registry | <b>5a. Contract Number:</b><br><b>5b. Grant Number:</b><br><b>5c. Program Element:</b> 0603729<br><b>5d. Project Number:</b> 02914<br><b>5e. Task Number:</b><br><b>5f. Work Unit Number:</b> 60332 |
|---|---|

|   |  |
|---|--|
| <b>6. AUTHORS</b><br>Amber L. Wade, Judy L. Dye, Charlene R. Mohrle, Michael R. Galarneau |  |
|---|--|

|   |   |
|---|---|
| <b>7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES)</b><br>Naval Health Research Center<br>P.O. Box 85122<br>San Diego, CA 92186-5122 | <b>9. PERFORMING ORGANIZATION REPORT NUMBER</b><br>Report No. 06-01 |
|---|---|

|  |  |
|--|--|
| <b>8. SPONSORING/MONITORING AGENCY NAMES(S) AND ADDRESS(ES)</b><br><br>Office of Naval Research                      Marine Corps Systems Command<br>800 N Quincy Street                            2006 Hawkins Avenue<br>Arlington, VA 22217-5660                      Quantico, CA 22134-5010<br><br>Marine Corps Warfighting Lab<br>3255 Meyers Avenue<br>Quantico, CA 22134 | <b>10. Sponsor/Monitor's Acronyms(s)</b><br>ONR, MCSC, MCWL<br><b>11. Sponsor/Monitor's Report Number(s)</b> |
|--|--|

|  |
|--|
| <b>12 DISTRIBUTION/AVAILABILITY STATEMENT</b><br>Approved for public release; distribution is unlimited. |
|--|

|                                |
|--------------------------------|
| <b>13. SUPPLEMENTARY NOTES</b> |
|--------------------------------|

|   |
|---|
| <b>14. ABSTRACT (maximum 200 words)</b><br><br>A retrospective analysis was performed to assess characteristics of head, face, and neck injury among US military casualties from Operation Iraqi Freedom Phase Two. Data were collected from the US Navy and Marine Corps Combat Trauma Registry for the time period of 1 March to 30 September 2004. Four hundred and forty-five casualties with head, face, or neck injury were identified for analysis. Of these, 140 (31%) sustained multiple wounds to the head, face, and neck; 177 (40%) had head wounds; 336 (76%) had facial wounds; and 84 (19%) had neck wounds. Head, face, and neck wounds accounted for 61% of the total 1,666 injuries incurred. Improvised explosive devices were the most frequent cause of injury among battle casualties, whereas motor vehicle accidents were the most common cause of injury among nonbattle casualties. The overall mean abbreviated injury severity scores for head, face, and neck injuries were 2.1, 1.1, and 1.2, respectively. Although the majority of casualties incurred wounds to the face, head and neck injuries were generally more severe, especially among those who died of wounds. The high frequency of facial wounding suggests, however, that the face may be more vulnerable during armed conflict than the head or neck. |
|---|

|   |
|---|
| <b>15. SUBJECT TERMS</b><br>Navy–Marine Corps Combat Trauma Registry, head, face, neck, injury, Operation Iraqi Freedom |
|---|

|  |                            |                             |   |                                  |  |
|--|----------------------------|-----------------------------|---|----------------------------------|--|
| <b>16. SECURITY CLASSIFICATION OF:</b> |                            |                             | <b>17. LIMITATION OF ABSTRACT</b><br>UNCL | <b>18. NUMBER OF PAGES</b><br>16 | <b>19a. NAME OF RESPONSIBLE PERSON</b><br>Commanding Officer                   |
| <b>a. REPORT</b><br>UNCL               | <b>b. ABSTRACT</b><br>UNCL | <b>c. THIS PAGE</b><br>UNCL |   |                                  | <b>19b. TELEPHONE NUMBER (INCLUDING AREA CODE)</b><br>COMM/DSN: (619) 553-8429 |